## LARES/WEBER-SAT and the equivalence principle

L. Iorio<sup>1</sup>

<sup>1</sup> *Viale Unit* a` *di Italia 68, 70125, Bari (BA), Italy tel. 0039 328 6128815 e-mail: lorenzo.iorio@libero.it*

PACS  $04.80.-y$  – Experimental studies of gravity PACS 04.80.Cc – Experimental tests of gravitational theories PACS 91.10.Sp – Satellite orbits

Abstract. - It has often been claimed that the proposed Earth artificial satellite LARES/WEBER-SAT−whose primary goal is, in fact, the measurement of the general relativistic Lense-Thirring effect at a some percent level–would allow greatly improving, among (many) other things, the present–day (10<sup>-13</sup>) level of accuracy in testing the equivalence principle as well. Recent claims point towards even two orders of magnitude better, i.e.  $10^{-15}$ . In this note we show that such a goal is, in fact, unattainable by many orders of magnitude being, instead, the achievable level  $\approx 10^{-9}$ .

The LAGEOS III/LARES/WEBER-SAT mission. – In 1976 Van Patten and Everitt [1, 2] suggested measuring the general relativistic Lense-Thirring node precession [3] with a pair of counter-orbiting spacecraft to be placed in terrestrial polar orbits and endowed with active, drag-free apparatus to counter-act the non-gravitational perturbations. In 1977-1978 Cugusi and Proverbio [4,5] proposed using the existing passive geodetic satellite LAGEOS and, more generally, the Satellite Laser Ranging (SLR) technique to measure the Lense-Thirring effect in the terrestrial gravitational field with the existing artificial satellites. In 1986 Ciufolini [6] put forth a strategy somewhat equivalent to the ones by Van Patten and Everitt and Cugusi and Proverbio involving the launch of a passive, LAGEOSlike satellite−named LAGEOS III after the launch of LAGEOS II−in an orbit identical to that of LAGEOS (semimajor axis  $a = 12,270$  km, inclination of the orbital plane to the Earth's equator  $i = 110$  deg, eccentricity  $e = 0.0045$ , apart from the inclination which should have been equal to  $i = 70$  deg. The observable originally proposed was the sum of the nodes of LAGEOS and LAGEOS III: the particular orbital configuration of the new spacecraft was chosen to cancel out the aliasing impact of the even zonal harmonic coefficients  $J_{\ell}$ ,  $\ell = 2, 4, ...$  of the multipolar expansion of the Newtonian part of the Earth's gravitational potential which induce node precessions qualitatively identical to the relativistic ones but much larger. Although extensively studied by various groups [7, 8], such an idea has not yet been implemented. In 1998, after its originally proposed eccentricity was augmented by almost one order of magnitude in order to use the perigee as well [9], renamed LAser RElativity Satellite (LARES), it was rejected by the Agenzia Spaziale Italiana (ASI,

Italian Space Agency). With the name of WEBER-SAT<sup>[1](#page-1-0)</sup>, it is currently under examination by the Istituto Nazionale di Fisica Nucleare<sup>[2](#page-1-1)</sup> (INFN, National Institute of Nuclear Physics).

In addition to the original, by far principal, purpose of measuring the Lense-Thirring effect at some percent level, many other goals of fundamental physics have been recently added to the LARES mission in order to enhance its chances of being finally approved. In doing so it has gone too far by often making unrealistic claims. A typical example of such a policy is represented by the alleged possibility [10] of measuring the perigee precession [11] induced by the multi-dimensional braneworld model by Dvali, Gabadadze and Porrati (DGP) [12]. Amounting to  $4 \times 10^{-3}$  milliarcseconds per year (mas yr<sup>-1</sup>), it has been proven to be undetectable in [13]. In such a work motivations mainly concerning the gravitational systematic errors were considered: here we will yield further, simple arguments related to the non-gravitational perturbations which the perigees of LAGEOS-like satellites are particularly sensitive to. In [14] it has been reported that the INFN team should be able to reduce the impact of the non-gravitational perturbations of thermal origin down to  $\approx 10^{-3}$ of the Lense-Thirring effect on LARES. Although not explicitly stated in [14], let us assume that this will be true for the perigee as well, which is certainly not an easy task to be implemented: the Lense-Thirring perigee precession for the originally proposed LARES orbital configuration amounts to about 30 mas yr−<sup>1</sup> , so that a mismodelled precession of  $\approx 10^{-2}$  mas yr<sup>-1</sup> of non-gravitational origin would be left, i.e. a bias just one order of magnitude larger than the DGP effect itself.

The equivalence principle. – Another example is the equivalence principle. Indeed, in several works [15–19] it has been recently claimed that LARES/WEBER-SAT would be able to greatly improve the accuracy level in testing such a cornerstone of general relativity and of all other competing metric theories of gravity. E.g., in [17] we find: "An additional physics goal of LARES is the improvement of the limits on the violation of the Einstein Equivalence Principle." More precisely, in [15] it is explicitly written: "LARES would improve, by about two orders of magnitude, the accuracy in testing the equivalence principle [...]". Now, since the present−day level of accuracy in testing it is 10−<sup>13</sup> [20], it must be argued that Ciufolini is claiming in [15] that a level of  $10^{-15}$  is to be expected from the successful implementation of the LARES mission. Unfortunately, the authors of such claims nowhere explicitly explain how they would achieve such notable goals.

By contrast, the situation is quite different, and by many orders of magnitude. The possibility of testing the equivalence principle with artificial Earth satellites of different compositions was tackled in  $[21-23]$ . In  $[22]$  just the originally proposed configuration of LARES  $(a = 12,270 \text{ km}, i = 70 \text{ deg})$  was examined: the existing LAGEOS satellite would be used in conjunction with LARES/WEBER-SAT which would substantially differ from it by composition, weight and manufacturing. Both the orbital periods  $T$  and the secular precessions of the nodes were considered, finding that the choice of the orbital periods would be better by one order of magnitude. In the former case, one should measure, after many revolutions, the difference of time spans which are multiple  $N$  of the orbital periods  $\Delta T_N = T_N^{(2)} - T_N^{(1)}$  of the pair of satellites, here denoted with the superscripts (1) and (2), orbiting the Earth along paths with almost the same semimajor axes  $a$ , to be measured from SLR data as well

$$
a^{(1)} \equiv a, \ a^{(2)} = a + d. \tag{1}
$$

Indeed, the equivalence principle-violating parameter  $\Delta \psi$  can be written in terms of  $\Delta T_N$ as [22]

<span id="page-1-2"></span>
$$
\Delta\psi_N = \frac{\frac{\sqrt{GM}}{N\pi} \Delta T_N - 3d\sqrt{a}}{3d\sqrt{a} + 2\sqrt{a^3}}.
$$
\n(2)

<span id="page-1-0"></span><sup>&</sup>lt;sup>1</sup>In memory of Dr. J. Weber, US Naval Academy (USNA) Class of 1940

<span id="page-1-1"></span><sup>2</sup>See on the Internet http://www.lnf.infn.it/acceleratori/lares/ and http://cadigweb.ew.usna.edu/ webersat/

The major limiting factor is mainly represented by the fact that, due to the unavoidable orbital injection errors, it would not be possible to insert the new satellite in an orbit with exactly the same semimajor axis of LAGEOS and that such difference d could be known only with a finite precision. For  $d \approx 5$  km and an uncertainty<sup>[3](#page-2-0)</sup>  $\delta d \approx 1$  cm the obtainable accuracy would be  $10^{-9}$  only, i.e. four orders of magnitude less than the present-day accuracy and six orders of magnitude less than that claimed in [15]. Similar conclusions were recently reached in [23] in which LAGEOS and its perigee, along with another similar SLR target were examined. Nobili et al. write in [23] that "it would need about 120,000 yr of LAGEOS data to get this limit down to the 10−<sup>12</sup> level of EP test already reached by torsion balances!!".

Alternative orbital configurations for LARES have been recently examined [24, 25] involving much smaller semimajor axes than that originally proposed. Let us consider the case of a low-altitude LARES with  $a = 7,878$  km [25] (and  $i \approx 90$  deg), which was, incidentally, already proven to be unsuitable to measure the Lense-Thirring effect [26]. According to eq.  $(2)$ , where, in this case, a is the semimajor axis of LARES and d is the difference between it and the one of LAGEOS, i.e.  $d = 4,392$  km, the errors due to  $\delta a$  and  $\delta d$  are

<span id="page-2-1"></span>
$$
\begin{cases}\n[\delta(\Delta\psi_N)]_a \le \left| \frac{-\frac{3d}{2\sqrt{a}} (3d\sqrt{a} + 2\sqrt{a^3}) + 3d\sqrt{a} \left(\frac{3d}{2\sqrt{a}} + 3\sqrt{a}\right)}{(3d\sqrt{a} + 2\sqrt{a^3})^2} \right| \delta a = 1 \times 10^{-10} \text{ cm}^{-1} \delta a, \\
[\delta(\Delta\psi_N)]_d \le \left| \frac{-6a^2}{(3d\sqrt{a} + 2\sqrt{a^3})^2} \right| \delta d = 2 \times 10^{-11} \text{ cm}^{-1} \delta d.\n\end{cases} (3)
$$

By realistically assuming  $\delta a \approx \delta d \approx 1$  cm, eq. [\(3\)](#page-2-1) tells us that we are still three orders of magnitude above even the present−day level. It is important to note that the results of eq. [\(3\)](#page-2-1) are optimistic: indeed, for the sake of simplicity we neglected the impact of the first even zonal harmonic  $J_2$  which, instead, was considered in [22] along with the impact of the Earth's GM as well which can be kept under control for a sufficiently high number of orbital revolutions.

Conclusions. – In conclusion, after having yielded further arguments concerning the impossibility of using the perigee of LARES/WEBER-SAT for testing the DGP braneworld model, we have shown that, whatever orbital configuration may be devised for the new spacecraft, it will not be able to obtain any improvement in testing the equivalence principle, remaining far from even the present−day level of accuracy by three−four orders of magnitude.

## REFERENCES

- [1] VAN PATTEN R. A. and EVERITT C. W. F., *Phys. Rev. Lett.*, **36** (1976) 629.
- [2] VAN PATTEN R. A. and EVERITT C. W. F., Celest. Mech. Dyn. Astron., 13 (1976) 429.
- [3] LENSE J. and THIRRING H., *Phys. Z.*, **19** (1918) 156. Translated and discussed in MASHHOON B., HEHL F. W. and THEISS D. S., Gen. Relativ. Gravit., 16 (1984) 711. Reprinted in RUFFINI R. J. and SIGISMONDI C. (Editors), Nonlinear Gravitodynamics (World Scientific, Singapore)2003, pp. 349-388.
- [4] CUGUSI L. and PROVERBIO E., *J. of Geodesy*, **51** (1977) 249.
- [5] CUGUSI L. and PROVERBIO E., Astron. Astrophys., **69** (1978) 321.
- [6] CIUFOLINI I., *Phys. Rev. Lett.*, **56** (1986) 278.
- [7] IORIO L., LUCCHESI D. M. and CIUFOLINI I., Class. Quantum Grav., 19 (2002) 4311.
- [8] RIES J. C., EANES R. J., WATKINS M .M. and TAPLEY B., Joint NASA/ASI Study on Measuring the Lense-Thirring Precession Using a Second LAGEOS Satellite, Vol. CSR-89-3 (Center for Space Research, The University of Texas at Austin)1989.

<span id="page-2-0"></span><sup>3</sup>Recall that both the existing LAGEOS and the proposed LARES are not endowed with any satelliteto-satellite tracking system or any other apparatus apart from retroreflectors.

- [9] CIUFOLINI I. ET AL., LARES Phase A (University La Sapienza, Rome) 1998.
- [10] Ciufolini I., LARES/WEBER-SAT, frame-dragging and fundamental physics, http://arxiv.org/abs/gr-qc/0412001, 2004.
- [11] LUE A. and STARKMANN, G., *Phys. Rev. D*, **67** (2003) 064002.
- [12] Dvali G., Gabadadze G. and Porrati M., Phys. Lett. B, 485 (2000) 208.
- [13] IORIO L., *J. Cosmol. Astropart. Phys.*, **7** (2005) 8.
- [14] Cantone C. for the LARES and ETRUSCO Collaborations, The INFN-LNF Space Climatic Facility for the LARES mission and the ETRUSCO project, talk delivered at International Workshop on "ADVANCES IN PRECISION TESTS AND EXPERIMENTAL GRAV-ITATION IN SPACE" Florence, ITALY, 28-30, Sep. 2006, http://www.fi.infn.it/GGI-gravspace/egs\_w.html, 2006.
- [15] CIUFOLINI I., Gen. Relativ. Gravit., **36** (2004) 2257.
- [16] CIUFOLINI I., CURRIE D. G. and PAOLOZZI A., The lares mission for testing the dynamics of general relativity, Aerospace Conference, 2003. Proceedings. 2003 IEEE, 2, 2 693- 2 703, 2005.
- [17] Dell'Agnello S., Delle Monache G., Franceschi M. A. and Paolozzi A., LARES, http://www.lnf.infn.it/acceleratori/lares/general%20description files/LARESrapatt05.ps, 2005.
- [18] Smith Jr. B. R. (Lt. Col. USAF) and Currie D.G., Laser Relativity Satellite: A Search for the Dragging of Inertial Frames (the Lense-Thirring Effect), http://cadigweb.ew.usna.edu/∼webersat/amos2.pdf
- [19] SMITH JR. B. R. (LT. COL. USAF), USNA-5: Laser Relativity Satellite (WeberSat-LARES), 2003. http://cadigweb.ew.usna.edu/∼webersat/AMOS.ppt
- [20] Will C. M., Living Rev. Relativity, 9 (2006) 3. http://www.livingreviews.org/lrr-2006-3
- [21] Blaser J. P., Class. Quantum Grav., 18 (2001) 2509.
- [22] Iorio L., Gen. Relativ. Gravit., 36 (2004) 361.
- [23] Nobili A., Comandi G. L., Doravari S., Lucchesi D. M. and Maccarone F., Limitations to testing the Equivalence Principle with Satellite Laser Ranging, talk given at Rencontres de Moriond-La Thuile, Italia March 11-18 2007, 2007. http://eotvos.dm.unipi.it/documents/generalpapers/lathuile2007.pdf
- [24] IORIO L., New. Astron., **10** (2005b) 616.
- [25] Ciufolini I., On the orbit of the LARES satellite, http://arxiv.org/abs/gr-qc/0609081, 2006.
- [26] IORIO L., *Planet. Space Sci.*, **55** (2007) 1198.