

# UNDECIDABILITY IN FUNCTION FIELDS OF POSITIVE CHARACTERISTIC

KIRSTEN EISENTRÄGER AND ALEXANDRA SHLAPENTOKH

ABSTRACT. We prove that the first-order theory of any function field  $K$  of characteristic  $p > 2$  is undecidable in the language of rings without parameters. When  $K$  is a function field in one variable whose constant field is algebraic over a finite field, we can also prove undecidability in characteristic 2. The proof uses a result by Moret-Bailly about ranks of elliptic curves over function fields.

## 1. INTRODUCTION

The current investigation started as an attempt by the authors to resolve Hilbert's Tenth Problem for all function fields of positive characteristic. Hilbert's Tenth Problem in its original form was to find an algorithm to decide, given a polynomial equation  $f(x_1, \dots, x_n) = 0$  with coefficients in the ring  $\mathbb{Z}$  of integers, whether it has a solution with  $x_1, \dots, x_n \in \mathbb{Z}$ . Matiyasevich ([12]), building on earlier work by Davis, Putnam, and Robinson ([2]), proved that no such algorithm exists, i.e. Hilbert's Tenth Problem is undecidable.

Since then, analogues of this problem have been studied by asking the same question for polynomial equations with coefficients and solutions in other recursive commutative rings. Perhaps the most important unsolved question in this area is Hilbert's Tenth Problem over the field of rational numbers.

The function field analogue turned out to be much more tractable. We know that Hilbert's Tenth Problem for the function field  $k$  of a curve over a finite field is undecidable. This was proved by Pheidas for  $k = \mathbb{F}_q(t)$  with  $q$  odd ([14]), and then extended to all global function fields in [25, 19, 4]. We also have undecidability of Hilbert's Tenth Problem for certain function fields over possibly infinite constant fields of positive characteristic ([20, 18, 4, 9]). The results of [4] and [20] also generalize to higher transcendence degree (see [21]) and give undecidability of Hilbert's Tenth Problem for finite extensions of  $\mathbb{F}_q(t_1, \dots, t_n)$  with  $n \geq 2$ . In [6] the problem was shown to be undecidable for finite extensions of  $k(t_1, \dots, t_n)$  with  $n \geq 2$  and  $k$  algebraically closed of odd characteristic.

So all known undecidability results for Hilbert's Tenth Problem in positive characteristic either require that the constant field not be algebraically closed or that we are dealing with a function field in at least 2 variables. The big open question that remains is whether Hilbert's Tenth Problem for a one-variable function field over an algebraically closed field of constants is undecidable.

The current methods for proving undecidability of Hilbert's Tenth Problem for function fields  $K$  of positive characteristic  $p$  usually require showing that the following sets are existentially definable in the language of rings :  $\{(x, x^{p^s}) : x \in K, s \in \mathbb{Z}_{\geq 0}\}$  and

---

*Key words and phrases.* Undecidability, elliptic curves, Hilbert's Tenth Problem.

K. Eisenträger was partially supported by a National Science Foundation postdoctoral fellowship. A. Shlapentokh was partially supported by NSF grants DMS-0354907 and DMS-0650927.

$\{x \in K : \text{ord}_{\mathfrak{p}}x \geq 0\}$  for some nontrivial prime  $\mathfrak{p}$  of  $K$ . In this paper we show that we can existentially define one of these sets for a large class of fields: we will prove that the set of  $p$ -th powers is existentially definable in *any* function field  $K$  of characteristic  $p > 2$  whose constant field has transcendence degree at least one over  $\mathbb{F}_p$ .

By a *function field (in  $n$  variables)* over a field  $F$  we mean a field  $K$  containing  $F$  and  $n$  elements  $x_1, \dots, x_n$ , algebraically independent over  $F$ , such that  $K/F(x_1, \dots, x_n)$  is a finite algebraic extension. The algebraic closure of  $F$  in  $K$  is called the constant field of  $K$ , and it is a finite extension of  $F$ .

Given the present difficulties of showing that Hilbert's Tenth Problem, or, equivalently, the existential theory of an arbitrary function field of positive characteristic is undecidable, one can also consider a weaker result, namely proving the undecidability of the first-order theory of these fields. Duret showed that the first-order theory of function fields (in  $n$  variables) over algebraically closed fields of positive characteristic is undecidable ([3]). In [1] Cherlin showed that the first-order theory of  $F(t)$  is undecidable for infinite perfect fields  $F$  of positive characteristic. In [15] Pheidas extended this result to rational function fields  $F(t)$  for any field  $F$  of characteristic  $p \geq 5$ , but he had to add a transcendental parameter  $t$  to the ring language to prove undecidability.

In this paper we generalize Duret's and Pheidas' results and prove that the first-order theory in the language of rings without parameters of *any* function field over a field of characteristic greater than 2 is undecidable. In the case the field of constants is algebraic over a finite field, we can also treat the case of characteristic 2.

The paper is organized as follows. We first show that the first-order theory for function fields  $K$  of positive characteristic is undecidable in the language of rings with finitely many parameters. This is done by defining a model of the nonnegative integers with addition and multiplication in  $K$ . Then, using a result of R. Robinson, we show that this also gives us the undecidability of the theory of  $K$  in the language of rings without parameters. The details of this argument are discussed in Section 5.

We endeavored to make the presentation as uniform as possible across all the different types of fields. Thus, in the second section of the paper we first show that in order to establish the first-order undecidability of a function field of characteristic  $p > 0$ , it is enough to show that  $p$ -th powers of a specific field element are first-order definable. To define  $p$ -th powers of a specific element, the techniques we use depend on the constant field. When the constant field is algebraic over a finite field we generalize equations that have previously been used in [20, 4] to reduce the problem to the rational function field case. This is done in Section 3. When the constant field has transcendence degree  $\geq 1$  over  $\mathbb{F}_p$ , things are more complicated. In Section 4 we show that the  $p$ -th powers we want to define occur as  $x$ -coordinates of the  $K$ -rational points on a certain elliptic curve. We then use a theorem by Moret-Bailly about the rank of elliptic curves in extensions function fields to reduce to the rational function field case. The theorem by Moret-Bailly was also used in [5, 7, 13] to obtain undecidability results. Moret-Bailly's theorem only holds in odd characteristic, and so for higher transcendence degree we obtain undecidability for function fields of odd positive characteristic.

2. USING  $p$ -TH POWERS TO CONSTRUCT A MODEL OF THE POSITIVE INTEGERS

**2.1. Statement of results.** The main result that we will prove in the first four sections is the following:

**Theorem 2.1.** *Let  $K$  be a function field of characteristic  $p > 2$  or a function field in one variable of characteristic 2 whose constant field is algebraic over  $\mathbb{F}_2$ . There exists a finite set of parameters  $\{z_1, \dots, z_k\} \subseteq K$  (depending on  $K$ ) such that first-order theory of  $K$  is undecidable in the language of rings augmented by  $\{z_1, \dots, z_k\}$ .*

From the result in Section 5 we obtain a strengthening of Theorem 2.1:

**Theorem 2.2.** *Let  $K$  be as in Theorem 2.1. Then the first-order theory of  $K$  is undecidable in the language of rings without parameters.*

**2.2. Idea of proof.** In this section we show that to prove Theorem 2.1, it is enough to define  $p$ -th powers of an element in the field with at least one simple pole or zero. To prove that this is enough we use a result of J. Robinson that shows how to define multiplication of positive integers in terms of addition and divisibility.

*Remark 2.3.* Since we are only interested in the function field  $K$  and not the underlying field  $F$ , we can always replace  $F$  with  $F(x_2, \dots, x_n)$  and view a function field  $K/F$  in  $n$  variables as a function field  $K/F(x_2, \dots, x_n)$  in one variable. So in the following all the function fields we consider will be function fields in one variable.

*Notation 2.4.* Let  $K$  be a function field (in one variable) of positive characteristic  $p$  over a field of constants  $F$ . Let  $F_0$  be the algebraic closure of a finite field in  $F$ . Let  $t \in K \setminus F$ , and let  $n = [K : F(t)]$ .

The following result is due to J. Robinson (see [17]).

**Lemma 2.5.** *There exists a first-order formula  $\mathcal{F}$  in the language  $\langle \mathbb{Z}_{>0}, +, | \rangle$  such that for integers  $k, m, n$ , we have  $k = mn \iff \mathcal{F}(k, m, n)$ . Here  $a | b$  means “ $a$  divides  $b$ ” for positive integers  $a, b$ .*

An immediate corollary of this lemma is the fact that the first-order theory of  $\langle \mathbb{Z}_{>0}, +, | \rangle$  is undecidable. So to prove the undecidability of the first-order theory of  $K$  it is enough to construct a model of the positive integers with addition and divisibility in  $K$ .

We say that we have a *model* of  $\langle \mathbb{Z}_{>0}, +, | \rangle$  in  $K$  if there is a bijection  $\phi : \mathbb{Z}_{>0} \rightarrow D$  between  $\mathbb{Z}_{>0}$  and a definable subset  $D$  of  $K^d$  (for some  $d \geq 1$ ), such that the graphs of  $+$  and  $|$  on  $D$  induced by  $\phi$  correspond to definable subsets of  $D^3$  and  $D^2$ , respectively.

As we will see in Theorem 2.9 below, we can construct a model of  $\langle \mathbb{Z}_{>0}, +, | \rangle$  if we can define  $p$ -th powers of a specific element.

**2.3. From  $p$ -th powers of a special element to arbitrary  $p$ -th powers.** The known definitions of  $p$ -th powers in general are produced in the following manner: first define  $p$ -th powers of a specific element, then use  $p$ -th powers of this element to produce  $p$ -th powers of arbitrary elements.

We have to distinguish two cases: the case where the constant field is perfect and when it is not perfect. The function fields  $K$  considered in Section 3 have constant fields which are algebraic over finite fields. These constant fields are always perfect. However, this will not be necessarily true of the constant fields in Section 4. The constant field will be relevant in

two places: in the definition of the  $p$ -th powers of the special element in Lemma 4.6 and also in the proof of Proposition 2.6 below.

**Proposition 2.6.** *Let  $K$  be a function field of positive characteristic. Suppose the set  $p(K, t) := \{x \in K : \exists s \in \mathbb{Z}_{>0}, x = t^{p^s}\}$  is definable in  $K$  for some  $t \in K \setminus F$  which is not a  $p$ -th power in  $K$ . Then the following subset of  $K^2$  is also definable in  $K$ :*

$$\mathcal{X}(K) := \{(x, x^{p^s}) : s \in \mathbb{Z}_{>0}, x \in K\}.$$

*Proof.* When the constant field is perfect we can follow the arguments of Section 8.4 of [23] which covers all positive characteristics. While the function fields considered in [23] are global function fields, the only condition that is really required is that the constant field is perfect. The main ingredient in the proof is that we can define a global derivation and use it to determine whether certain elements of the field have simple zeroes and poles.

When the field of constants is not perfect we obtain the result from Lemma 2.10 and Corollary 2.11 of [20]. In both cases we might need to enlarge the field of constants to avoid ramifying valuations as zeros and poles of elements whose  $p$ -th powers we define. These constant field extensions are covered by Proposition 7.1 and Lemma 7.2 from the appendix. So the proposition holds for any field of constants.  $\square$

Both [23] and [20] also prove the following corollary which will be needed below.

**Corollary 2.7.** *Let  $t \in K \setminus F$  and assume that  $t$  is not a  $p$ -th power in  $K$ . If  $p(K, t)$  is definable in  $K$ , then the set*

$$\mathcal{B}(K, t) := \{(t^{p^s}, x^{p^s}, x) : s \in \mathbb{Z}_{>0}, x \in K\}$$

*is definable in  $K$ .*

**2.4. Constructing a model with addition and divisibility.** The final result we will need to construct a model of  $\langle \mathbb{Z}_{>0}, +, | \rangle$  is the the following proposition.

**Proposition 2.8.** *Let  $t$  be as above. If  $\mathcal{X}(K) = \{(x, x^{p^s}), x \in K, s \in \mathbb{Z}_{>0}\}$  is definable in  $K$ , then the set*

$$\mathcal{C}(K, t) := \{(t^{p^a}, t^{p^b}, t^{p^{a+b}}) : a, b > 0\}$$

*is definable in  $K$ .*

*Proof.* Consider the following system of equations:

$$(1) \quad \begin{cases} x - 1 = t^{p^a} \\ \exists l \in \mathbb{Z}_{>0} : z = ((t + 1)t^{p^b})^{p^l} \\ \exists j \in \mathbb{Z}_{>0} : z/x = t^{p^j} \end{cases}$$

We claim that for any  $a, b > 0$ , if this system has solutions  $x, z \in K$  then  $x/z = t^{p^{a+b}}$ . Indeed, from the first equation we conclude that  $x = (t + 1)^{p^a}$ . From the second equation we get that  $z = (t + 1)^{p^l} t^{p^{b+l}}$ . Finally, from the third equation we have that  $(t + 1)^{p^l - p^a} t^{p^{b+l}} = t^{p^j}$ . The only way this equality can hold is for  $l = a$  and  $j = a + b$ . Conversely, we can always satisfy the system if  $x = (t + 1)^{p^a}$  and  $z = (t + 1)^{p^a} t^{p^{b+a}}$ .  $\square$

We are now ready for the following theorem.

**Theorem 2.9.** *Assume that  $t$  has a simple pole or a simple zero. Suppose that the set  $p(K, t)$  is definable in  $K$ . Then  $\langle \mathbb{Z}_{>0}, +, | \rangle$  has a model over  $K$ .*

*Proof.* We map  $s > 0$  to  $t^{p^s}$ . Then  $s = s_1 + s_2 \Leftrightarrow (t^{p^{s_1}}, t^{p^{s_2}}, t^{p^s}) \in \mathcal{C}(K, t)$ . Further  $s_1 \mid s_2$  if and only if  $(p^{s_1} - 1) \mid (p^{s_2} - 1)$  if and only if there exists  $x \in K$  such that

$$(2) \quad x^{p^{s_1}-1} = t^{p^{s_2}-1},$$

since at least one pole or zero of  $t$  is simple. Indeed suppose that the equality holds and let  $\mathfrak{q}$  be a simple pole or zero of  $t$ . Then

$$p^{s_2} - 1 = \text{ord}_{\mathfrak{q}} t^{p^{s_2}-1} = \text{ord}_{\mathfrak{q}} x^{p^{s_1}-1} \equiv 0 \pmod{p^{s_1} - 1}.$$

Conversely, if  $p^{s_2} - 1 = l(p^{s_1} - 1)$  for some  $l \in \mathbb{Z}_{>0}$ , then we can set  $x = t^l$  and (2) will hold.

Hence  $s_1 \mid s_2$  if and only if

$$\exists x, y \in K \left( (t^{p^{s_1}}, y, x) \in \mathcal{B}(K, t) \wedge y/x = t^{p^{s_2}}/t \right).$$

The result now follows from the fact that the sets  $\mathcal{p}(K, t)$ ,  $\mathcal{B}(K, t)$  and  $\mathcal{C}(K, t)$  are all definable in  $K$ .  $\square$

**2.5. Defining  $p$ -th powers of one special element.** We now address the issue of defining  $p$ -th powers of one specific element when the constant field is perfect. In the next proposition we observe that if we avoid ramified zeros and poles and consider rational functions only, we have the desired result.

**Proposition 2.10.** *Assume  $t$  has no zeros or poles ramifying in the extension  $K/F(t)$ . Let  $r = 1$  if  $p > 2$  and let  $r = 2$  if  $p = 2$ . Assume that  $F$  is perfect and for some element  $w \in F(t)$ , having no poles or zeros at the primes ramifying in the extension  $K/F(t)$ , there exists  $u, v \in K$  such that the following system is satisfied.*

$$(3) \quad \begin{cases} \frac{1}{t} - \frac{1}{w} = u^{p^r} - u \\ t - w = v^{p^r} - v \end{cases}$$

*Then for some  $s \in \mathbb{Z}_{\geq 0}$  we have that  $w = t^{p^{rs}}$ . Conversely, if  $w = t^{p^{rs}}$ ,  $s \geq 0$ , then there exist  $u, v \in F(t)$  satisfying (3). (For the last assertion we do not need the requirement that  $F$  is perfect.)*

*Proof.* Given our assumptions on  $F$  and  $w$  the proof of this proposition is identical to the proofs of Lemma 8.3.3, Corollary 8.3.4, and Proposition 8.3.8 of [23].  $\square$

Unfortunately, we cannot always assume that an arbitrary rational field element  $w$  avoids all ramified poles and zeros. However, this problem can be solved rather easily if we modify the equations. The next remark and the proposition below deal with an arbitrary element  $w \in F(t)$ .

*Remark 2.11.* We recall from Notation 2.4 that  $K/F$  was a function field of positive characteristic  $p$  and  $F_0$  denoted the algebraic closure of a finite field in  $F$ . By Proposition 7.1 and Lemma 7.2 from the appendix we can enlarge the constant field and assume that  $F_0$  contains elements  $c_0 = 0, c_1, \dots, c_{2n(\alpha)+2}$  such that when  $i \neq j$  we have for all  $k \in \mathbb{Z}_{\geq 0}$  that  $c_i^{p^k} \neq c_j$ . Here  $n(\alpha)$  is the constant that is defined in Lemma 7.2 from the appendix.

We can now prove the following proposition.

**Proposition 2.12.** *Assume  $F$  is perfect and  $t$  is not a  $p$ -th power in  $K$ . Let  $c_0, \dots, c_{2n(\alpha)+2}$  be as in Remark 2.11, and let  $V_i = \{c_i^{p^k}, k \in \mathbb{Z}_{\geq 0}\}$ .*

*Let  $r = 1$  if  $p > 2$  and let  $r = 2$  if  $p = 2$ . Let  $w \in F(t)$  and suppose that for all  $i \neq j \in \{1, \dots, 2n(\alpha) + 2\}$  for some  $a \in V_i, b \in V_j$  there exist  $u_{i,j,a,b}, v_{i,j,a,b} \in K$  such that*

$$(4) \quad \begin{cases} \frac{t-c_i}{t-c_j} - \frac{w-a}{w-b} = u_{i,j,a,b}^{p^r} - u_{i,j,a,b} \\ \frac{t-c_j}{t-c_i} - \frac{w-b}{w-a} = v_{i,j,a,b}^{p^r} - v_{i,j,a,b} \end{cases}$$

*Then for some  $s \in \mathbb{Z}_{\geq 0}$  we have that  $w = t^{p^{rs}}$ . Conversely, if  $w = t^{p^{rs}}$  for some  $s \in \mathbb{Z}_{\geq 0}$  then the equations can be satisfied as specified above (even if  $F$  is not perfect).*

*Proof.* First of all, using an argument similar to the one used in Lemma 8.3.10 of [23], we conclude that for some  $c_i, c_j$  and  $a \in V_i, b \in V_j$ , we have that  $t - c_i, t - c_j, w - a, w - b$  do not have zeros at any prime ramifying in the separable extension  $K/F(t)$  and therefore  $\frac{t - c_i}{t - c_j}, \frac{w - a}{w - b} \in F(t)$  do not have zeros or poles at any primes ramifying in the extension

$K/F(t)$ . Now applying Proposition 2.10 we conclude that either  $\frac{w - a}{w - b} = \left(\frac{t - c_i}{t - c_j}\right)^{p^{rs}}$ ,  $s > 0$

or  $\frac{w - a}{w - b} = \frac{t - c_i}{t - c_j}$ . In the first case we can take the  $p^r$ -th ‘‘root’’ of all our equations as in

Lemma 8.3.1 and Lemma 8.3.2 of [23]. In the second case we obtain  $w = a_1 t + a_2$  for some  $a_1, a_2 \in F_0$ . However, if we plug in this expression for  $w$  into our equations with  $c_0 = 0$  we obtain a contradiction unless  $a_1 = 1$  and  $a_2 = 0$ . Thus, we conclude that  $w = t^{p^{rs}}$ . Finally, the satisfiability assertion follows as before from Proposition 8.3.8 of [23].  $\square$

To define  $p$ -th powers of a special element  $t$  over fields of transcendence degree one and higher transcendence degree we will use some of the equations that were used in [20] and [4]. What we need to make the same arguments go through in our more general setup is a set of equations over  $K$  that forces its solutions to be in the rational function field  $F(t)$  and which are satisfied by all elements  $t^{p^s}$ ,  $s \in \mathbb{Z}_{>0}$ . I.e., we want a set  $\mathcal{S}$  which is definable in  $K$  such that  $p(K, t) \subseteq \mathcal{S} \subseteq F(t)$  and thus we can apply Proposition 2.12. This will be accomplished in the next two sections. For the transcendence degree one case, the equations defining  $\mathcal{S}$  are given in Corollary 3.5. For higher transcendence degree, they are given in Proposition 4.7 below.

### 3. DEFINING $p$ -TH POWERS FOR FUNCTION FIELDS WHOSE CONSTANT FIELD IS ALGEBRAIC

Let  $K/F$  be a function field in one variable of positive characteristic  $p$  with  $F$  algebraic over a finite field. When  $F$  has an extension of degree  $p$ , the results in [20] and [4] show that the existential theory of  $K$  and hence also the first-order theory of  $K$  are undecidable. Hence we can make additional assumptions about the field of constants for the algebraic case and assume that we are in a situation that is not covered by [20] or [4].

*Notation 3.1.*

Let  $K/F$  be a function field in one variable of positive characteristic  $p$ . Assume that  $F$  is algebraic over a finite field and has no extension of degree  $p$ .

Let  $t$  be a fixed element of  $K \setminus F$  which is not a  $p$ -th power in  $K$ .

We write  $g_K$  for the genus of  $K$ , and when  $f \in F[X, Y]$  defines a plane curve  $\mathcal{C}$  over  $F$ , we denote by  $g_f$  the genus of the function field of  $\mathcal{C}$  and also refer to this as the genus of  $f$ .

In this section we will show how to define  $p$ -th powers of the element  $t$  under the above assumptions.

**Lemma 3.2.** *For a pair of positive integers  $k = p^l, u$ , let*

$$f_{k,u}(X, Y) = Y^{p^k} - Y + \frac{1}{\prod_{i=1}^u (X - c_i)}.$$

*Then for any  $a \in F$  there exists  $b \in F$  such that  $f_{k,u}(a, b) = 0$ .*

*Proof.* Fix an  $a \in F$  and let  $\alpha_1, \alpha_2$  be roots of  $f_{k,u}(a, Y)$  in the algebraic closure of  $F$ . Then  $(\alpha_1^{p^k} - \alpha_2^{p^k}) - (\alpha_1 - \alpha_2) = 0$ . Thus,  $\alpha_1 - \alpha_2 = c$  is of degree  $k = p^l$  over a field of  $p$  elements and therefore  $c \in F$ . Since  $F$  is algebraic over a finite field, the extension  $F(\alpha_1)/F$  is cyclic. Assume that  $[F(\alpha_1) : F] = m > 1$  and let  $\sigma \in \text{Gal}(F(\alpha_1)/F)$  be a generator. Then for some  $c \in F$  we have that  $\sigma(\alpha_1) = \alpha_1 + c$  and  $\text{id}(\alpha_1) = \sigma^m(\alpha_1) = \alpha_1 + mc = \alpha_1$ . Thus  $m \equiv 0 \pmod{p}$ , and  $F(\alpha)/F$  has a subextension of degree  $p$  over  $F$ , contradicting our assumption on  $F$ .  $\square$

**Lemma 3.3.** *There exists a set  $A \subset K^2$ , diophantine over  $K$  such that  $A \subset F^2$  and for all  $a \in F$  there exists  $c \in F$  such that  $(a, c) \in A$ .*

*Proof.* Before we proceed with the proof we should note that using the effective version Chebotarev Density Theorem (see [8], Proposition 6.4.8) one could show that any infinite field algebraic over a finite field is anti-Mordellic and therefore one could use a result of Poonen and Pop to see that  $F$  is first-order definable in  $K$  (see [16]). However in our case we can give a very simple existential definition of  $F$  along the lines of [3], [10] and [22].

The idea is to construct an equation  $f$  whose genus is greater than the genus of  $K$  and then use the Riemann-Hurwitz formula to show that all the  $K$ -rational solutions must be  $F$ -rational. We also have to ensure that  $f$  has enough solutions over  $F$ . Consider an equation

$$f_{k,u}(X, Y) = Y^{p^k} - Y + \prod_{i=1}^u \frac{1}{(X - c_i)},$$

where  $k$  and  $u$  are as above and  $c_1, \dots, c_u$  are all distinct and in  $F$ . For sufficiently high  $k$  and  $u$  the genus of this equation is higher than the genus of  $K$ . To see that this is so, assume  $(u, p) = 1$  and consider the field extension  $F_{k,u}(X, Y)$  of  $F(X)$  where  $f_{k,u}(X, Y) = 0$ . It is clear that in this extension the primes corresponding to  $(X - c_1), \dots, (X - c_u)$  are completely ramified. It is also clear from considering the difference between any two roots of this equation as in the lemma above, that no other prime of  $F(X)$  is ramified in the extension  $F_{k,u}(X, Y)/F(X)$ . Furthermore, the  $F_{k,u}(X, Y)$ -factor of  $(X - c_i)$  is of relative degree 1 and also of degree 1 in  $F(X, Y)$ . Let  $g_X = 0$  be the genus of  $F(X)$ , and let  $g_{f_{k,u}}$  be the genus of  $f_{k,u}(X, Y)$ . Then by the Riemann-Hurwitz formula and Remark 3.5.7 of [8], we have that

$$2g_{f_{k,u}} - 2 \geq p^k(g_X - 2) + \deg \sum_{i=1}^u (p^k - 1)\mathfrak{P}_i,$$

where for  $i = 1, \dots, u$ , we let  $\mathfrak{P}_i$  denote the prime above  $X - c_i$ . Thus,

$$g_{f_{k,u}} \geq \frac{1}{2}(u(p^k - 1) - 2p^k + 2) = \frac{1}{2}(p^k u - u - 2p^k + 2) = \frac{(u - 2)(p^k - 1)}{2}.$$

Now choose  $k_0, u_0$  large enough so that  $g_{f_{k_0, u_0}}$  is greater than  $g_K$ , the genus of  $K$ . Let  $f := f_{k_0, u_0}$ , let  $F(X, Y)$  the corresponding field extension of  $F(X)$ , and  $g_f$  its genus.

Now assume that there exists a solution  $x, y \in K \setminus F$  to  $f(X, Y) = 0$ . Then  $F(X, Y) \simeq F(x, y)$ , so  $F(X, Y)$  can be viewed as a subfield of  $K$ . If  $x$  is not a  $p$ -th power in  $K$ , then the extension  $K/F(x)$  is separable (see Chapter VI of [11]) and as a consequence of the Riemann-Hurwitz formula,  $g_K \geq g_f$  contradicting the hypothesis.

If  $x$  is a  $p$ -th power in  $K$ , then  $\prod_{i=1}^u (x - c_i)$  is also a  $p$ -th power in  $K$  since  $F$  is algebraic over a finite field, and therefore all the coefficients  $c_i$  of  $f$  are also  $p$ -th powers. Consequently  $y$  is also a  $p$ -th power in  $K$ . Thus, by replacing all the terms of  $f$  by their  $p$ -th roots we can obtain a new equation  $f^{(1)}(X, Y) = 0$  which is a “ $p$ -th root” of  $f$ . The equation  $f^{(1)}$  has the same genus as  $f$  because its genus only depends on the values  $k_0, u_0$  that were chosen above. Since  $x$  and  $y$  were both  $p$ -th powers in  $K$ , the equation  $f^{(1)}(X, Y) = 0$  also has a non-constant solution in  $K$ . Thus, at some point we will have an equality  $f^{(\ell)}(\tilde{x}, \tilde{y}) = 0$ , with the genus of  $f^{(\ell)}$  higher than the genus of  $K$  and  $\tilde{x}$  not a  $p$ -th power in  $K$ . Consequently,  $f(X, Y) = 0$  can have constant solutions in  $K$  only. At the same time, by Lemma 3.2, for all  $x \in F$  we have  $y \in F$  so that  $f(x, y) = 0$ .  $\square$

**Proposition 3.4.** *Suppose for some  $w \in K$ , for infinitely many primes  $\mathfrak{P}$  of  $F(t)$  we have that*

$$(5) \quad w \equiv a(\mathfrak{P}) \pmod{\mathfrak{P}},$$

where  $a(\mathfrak{P}) \in F$ . Then  $w \in F(t)$ .

*Proof.* This proof follows from an argument similar to the argument in the proof of Theorem 10.1.1 of [23]. The main difference is that we do not assume that the prime  $\mathfrak{P}$  is inert in the extension  $K/F(t)$ . However, as long as the equivalence (5) holds for all the factors of the given prime below, the argument is unchanged.  $\square$

**Corollary 3.5.** *Suppose that for some  $w \in K$  and infinitely many  $(a, b) \in F^2$  we have that the following system has a solution  $u_a$  in  $K$ :*

$$(6) \quad \frac{1}{t-a} - \frac{1}{w-b} = u_a^p - u_a$$

Then  $w \in F(t)$ . Conversely, if for some positive integer  $s$  we have that  $w = t^{p^s}$ , then for any  $a \in F$ , there exist  $b \in F, u_a \in F(t)$  such that equation (6) is satisfied.

*Proof.* First of all observe that the extension  $K/F(t)$  is separable since  $t$  is not a  $p$ -th power in  $F(t)$ . (See Lemma B1.32 of [23].) Thus only finitely many primes ramify in the extension  $K/F(t)$ . Therefore for all but finitely many  $a \in F$ , for any factor  $\mathfrak{p}_a$  of the rational prime  $\mathfrak{P}_a$  which is the zero of  $t - a$  in  $F(t)$ , it is the case that  $\text{ord}_{\mathfrak{p}_a}(t - a) = 1$  in  $K$ . On the other hand, for any pole  $\mathfrak{q}$  of  $u_a$  in  $K$  we have that  $\text{ord}_{\mathfrak{q}}(u_a^p - u_a) \equiv 0 \pmod{p}$ . Thus, for all but finitely many  $a \in F$ , for all factors  $\mathfrak{p}_a$  of the rational prime  $\mathfrak{P}_a$  in  $K$  we have that  $\text{ord}_{\mathfrak{p}_a}(w - b) > 0$ . In other words, for infinitely many  $(a, b) \in F^2$  we have that  $w \equiv b \pmod{\mathfrak{P}_a}$ , where  $\mathfrak{P}_a$  is, as above, the zero divisor in  $K$  and  $F(t)$  of  $t - a$ . Now the first assertion of the corollary follows by Proposition 3.4. The second assertion of the corollary follows from Proposition 8.3.8 of [23].  $\square$

Finally note that we have assembled all the parts (i.e. Proposition 2.12, Lemma 3.3 and Corollary 3.5) for the main result of this section.



**Theorem 3.6.** *Let  $K$  be a function field (in one variable) whose constant field  $F$  is algebraic over a finite field of characteristic  $p > 0$ . Let  $t$  be an element of  $K \setminus F$  which is not a  $p$ -th power in  $K$ . Then the set  $p(K, t) = \{x \in K : \exists s \in \mathbb{Z}_{>0} x = t^{p^s}\}$  is first-order definable in  $K$ .*

#### 4. DEFINING $p$ -TH POWERS OVER FIELDS OF HIGHER TRANSCENDENCE DEGREE

Let  $K$  be a function field of characteristic  $p > 2$  with constant field  $F$ , and assume that  $F$  has transcendence degree at least one over a finite field. To define  $p$ -th powers of a suitable element  $t$  we will use a theorem by Moret-Bailly ([13]). Here, we quickly review his notation and state the theorem in the form we need.

**Definition 4.1.** Let  $u : A \rightarrow B$  be a morphism of abelian groups. We say that  $u$  is *almost bijective* if  $u$  is injective and  $\text{Coker } u$  is a finite  $p$ -group.

By [13, Theorem 1.8], the following theorem holds:

**Theorem 4.2.** *Let  $F$  be a field of characteristic  $p > 2$ , and assume that  $F$  contains an element which is transcendental over  $\mathbb{F}_p$ . Let  $K$  be a function field in one variable with constant field  $F$ , and let*

$$E : y^2 = P(x)$$

*be an elliptic curve which is defined over a finite field contained in  $F$ . There exists a non-constant element  $t \in K$  such that  $t$  is not a  $p$ -th power in  $K$  and the elliptic curve  $\mathcal{E}$  given by*

$$\mathcal{E} : P(t)y^2 = P(x)$$

*has the property that the natural homomorphism  $\mathcal{E}(F(t)) \hookrightarrow \mathcal{E}(K)$  induced by the inclusion  $F(t) \hookrightarrow K$  is almost bijective.*

*Notation 4.3.* From now on, let  $P(x)$ ,  $E$  and  $t$  be as in Theorem 4.2. Let  $s$  be an element in a quadratic extension of  $K$  satisfying  $s^2 = P(t)$ . Let  $q = p^r$  be the size of a finite field containing all the coefficients of the equation defining  $E$ .

Let  $F'$  be an algebraic closure of  $F$ , and let  $K' = F'K$ .

By [13, Theorem 1.8], it follows that the natural homomorphism  $\mathcal{E}(F(t)) \hookrightarrow \mathcal{E}(K')$  is still almost bijective.

**Proposition 4.4.** *The set  $\mathcal{E}(F(T))$  is diophantine over  $K$  and over  $K'$ .*

*Proof.* Let  $A := \mathcal{E}(F(T))$  and  $B := \mathcal{E}(K)$ . The set  $B$  is clearly diophantine over  $K$ . By Theorem 4.2,  $A$  is a subgroup of finite index in  $B$  and  $B/A$  is a finite  $p$ -group.

Hence for some integer  $k$  we have that  $p^k B \subseteq A$  and  $p^k B$  has finite index in  $B$ . Since  $B$  is diophantine over  $K$ , and since multiplication by  $p^k$  is given by explicit equations, the set  $p^k B$  is diophantine over  $K$ . It is easy to see that this implies that  $A$  is diophantine over  $K$ :

Let  $Q_1, \dots, Q_\ell$  be coset representatives for  $p^k B$  in  $A$ . Then for  $P \in \mathcal{E}(K)$

$$P \in A \Leftrightarrow (\exists S \in p^k B)(P = S + Q_1) \vee \dots \vee (P = S + Q_\ell).$$

The same argument with  $K$  replaced by  $K'$  shows that  $A$  is also diophantine over  $K'$ .  $\square$

From the proposition above we also obtain the following easy corollary.

**Corollary 4.5.** *There exists a polynomial equation  $R(u, v, x_1, \dots, x_l) \in K[u, v, x_1, \dots, x_l]$  such that  $R(u, v, x_1, \dots, x_l) = 0$  for some  $u, v, x_1, \dots, x_l \in K'$  implies  $(u, v)$  are affine coordinates of a point in  $\mathcal{E}(F(t))$ . Conversely, if  $(u, v)$  are affine coordinates of a point in  $\mathcal{E}(F(t))$  the equation  $R(u, v, x_1, \dots, x_l) = 0$  can be satisfied with  $x_1, \dots, x_l \in K$ .*

Next we observe that  $p$ -th powers occur as affine coordinates of points of  $\mathcal{E}$ .

**Lemma 4.6.** *Let  $\mathcal{E}, s, t, q$  be as in Notation 4.3. The point  $(t^{q^m}, s^{q^m-1}) \in \mathcal{E}(F(t))$ .*

*Proof.* Observe that  $(P(t))^{q^m} = P(t^{q^m})$ . Thus,  $P(t)(s^{q^m-1})^2 = (P(t))^{q^m} = P(t^{q^m})$ . Also  $q^m - 1$  is even, so the point  $(t^{q^m}, s^{q^m-1})$  has coordinates in the ground field.  $\square$

We conclude with the proposition defining  $p$ -th powers of  $t$  for the case of  $K$  of transcendence degree greater than one.

**Proposition 4.7.** *Assume that for some  $z, w, u, v \in K$  the following system is satisfied over  $K'$ .*

$$(7) \quad \begin{cases} R(w, z, x_1, \dots, x_l) = 0 \\ \forall i, j \in \{1, \dots, 2n(\alpha) + 2\} \exists a \in V_i, b \in V_j : \\ \frac{t-c_i}{t-c_j} - \frac{w-a}{w-b} = u_{i,j,a,b}^p - u_{i,j,a,b} \\ \frac{t-c_j}{t-c_i} - \frac{w-b}{w-a} = v_{i,j,a,b}^p - v_{i,j,a,b} \end{cases}$$

*Then for some  $s \in \mathbb{Z}_{\geq 0}$  we have that  $w = t^{p^s}$ . Conversely, if  $w = t^{p^s}$ , then the system has solutions in  $K$ .*

Finally we note that  $t$  selected so that Theorem 4.2 holds might have zeros and poles which are not simple. Let  $t' = \frac{t-a}{t-b}$  be such that that all of its poles and zeros in  $K$  are simple. Observe that  $F(t) = F\left(\frac{t-a}{t-b}\right)$  and we can generate  $p$ -th powers of  $t'$ . Thus in Theorem 2.9 and Proposition 2.6 we can replace  $t$  by  $t'$  if necessary.

*Remark 4.8.* The proof of Proposition 2.6 that  $p$ -th powers of the special element  $t$  allow us to define  $p$ -th powers of arbitrary elements  $x$  in  $K$  only used equations involving existential quantifiers. The proof of Proposition 4.7 which defined  $p$ -th powers of  $t$  also only used existential quantifiers. So when the constant field of  $K$  contains transcendental elements over  $\mathbb{F}_p$ , the set  $\{(x, x^{p^s}) : x \in K, s \in \mathbb{Z}_{\geq 0}\}$  is actually *existentially* definable in  $K$ .

## 5. RING LANGUAGE

In this section we address the issue of the language needed to produce an undecidable set of sentences. We have already shown that we can construct a model of the positive integers  $\mathbb{Z}_{>0}$  (and hence also of  $\mathbb{Z}_{\geq 0}$ ) in the function field  $K$ . We used this model to prove Theorem 2.1.

In this section we will show that this easily implies Theorem 2.2 by using a result of R. Robinson, so we obtain undecidability of the first-order theory of  $K$  in the language of rings *without* parameters.

**Corollary 5.1** (Theorem 2.2). *Let  $K$  be a function field of characteristic  $p > 2$ . Then the first-order theory is undecidable in the language of rings without parameters. When  $K$  is a function field in one variable whose constant field is algebraic over a finite field, then we also obtain undecidability in characteristic 2.*

*Proof.* From the previous sections it follows that the equations we used to construct a model of  $\mathbb{Z}_{\geq 0}$  are in the language  $L_{\bar{d}} = \langle +, \cdot; 0, 1, \{d_1, \dots, d_r\} \rangle$ , for fixed elements  $d_1, \dots, d_r$  of  $K$ . In other words, we are working in the language of rings with finitely many parameters. To show that we can achieve undecidability in the ring language *without parameters*, we use a result of R. Robinson who gave an example of a finitely axiomatizable and essentially undecidable theory  $Q$  ([24, p. 32]). A theory is *essentially undecidable* if any consistent extension of it is also undecidable. Since  $Q$  is a subtheory of  $\mathbb{Z}_{\geq 0}$  ([24, p. 51]), the axioms of  $Q$  hold in  $\mathbb{Z}_{\geq 0}$ . Let  $Ax(\mathbb{Z}_{\geq 0})$  be the conjunction of all the axioms of  $Q$ . For a sentence  $\psi$  in the language  $L = \langle +, \cdot, 0, 1 \rangle$  let  $\phi_K(\psi, \bar{d})$  be its translation in our model, and consider the set of all  $L$ -sentences  $\psi$  for which

$$(8) \quad \forall \bar{w} (\phi_K(Ax(\mathbb{Z}_{\geq 0}), \bar{w}) \rightarrow \phi_K(\psi, \bar{w}))$$

is true in  $K$ . This set contains the axioms of  $Q$  and therefore the theory generated by these sentences is an extension of  $Q$ . The extension is consistent. Suppose not. Then for some  $\psi$  as above we have that  $Ax(\mathbb{Z}_{\geq 0}) \rightarrow \neg\psi$  holds in  $\mathbb{Z}_{\geq 0}$ , and hence  $\phi_K(Ax(\mathbb{Z}_{\geq 0}), \bar{d}) \rightarrow \neg\phi_K(\psi, \bar{d})$  holds in  $K$ . But this contradicts (8). Since the collection of all  $L$ -formulas  $\psi$  satisfying (8) in  $K$  is undecidable, the set of all formulas of the form (8) true in  $K$  is also undecidable. Finally we note that the formulas in this set are in  $L$ .  $\square$

## 6. OPEN QUESTIONS

Even though we proved that the first-order theory of function fields of characteristic  $p > 2$  is undecidable, we needed transcendental elements in the construction of the model of the nonnegative integers. So the following questions arise naturally:

**Question 6.1.** Let  $K$  be a function field as in Theorem 2.1. Does  $K$  admit a model of  $\langle \mathbb{Z}_{\geq 0}, +, \cdot \rangle$  in which the equations defining the model have integer coefficients?

**Question 6.2.** Is the degree of unsolvability of the first-order theory of  $K$  at least that of the first-order theory of  $\mathbb{Z}$ ?

## 7. APPENDIX

The following proposition and lemma are used to handle constant field extensions of function fields.

**Proposition 7.1.** *Let  $Q_i$  be either “ $\forall$ ” or “ $\exists$ ”. Let  $M/K$  be a finite extension of fields with  $M$  not algebraically closed. Let  $P(t_1, \dots, t_r, x_1, \dots, x_k) \in M_0[t_1, \dots, t_r, x_1, \dots, x_k]$ . Let*

$$(9) \quad A_M = \{(t_1, \dots, t_r) \in M^r : Q_1 x_1 \in M \dots Q_k x_k \in M : P(t_1, \dots, t_r, x_1, \dots, x_k) = 0\}$$

*be a first-order definable set. Then there exists a polynomial  $T(u_1, \dots, u_m, y_1, \dots, y_l) \in K_0[u_1, \dots, u_m, y_1, \dots, y_l]$ , and a first-order definable set*

$$(10) \quad A_K = \{(u_1, \dots, u_m) \in K^m : Q_{k+1} y_1 \in K \dots Q_{k+l} y_l \in K : T(u_1, \dots, u_m, y_1, \dots, y_l) = 0\}$$

*such that for any  $(t_1, \dots, t_r) \in M^r$  we have that  $(t_1, \dots, t_r) \in A_M$  if and only if for some  $m$ -tuple  $(u_1, \dots, u_m) \in K^m$  we have that  $(u_1, \dots, u_m) \in A_K$ . Thus, if  $M$  has a first-order model of  $\mathbb{Z}$  in the language of rings augmented by finitely many parameters from  $M$ , then  $K$  has a first-order model of  $\mathbb{Z}$  in the language of rings with finitely many parameters from  $K$ .*

The proof of the proposition requires standard “rewriting” techniques utilizing a basis of  $M$  over  $K$  and the fact that over a field which is not algebraically closed we can replace a finite set of equations by a single equivalent equation.

Proposition 7.1 will play a role in case we need to extend the field of constants to ensure that we have “enough” conjugacy classes of constants algebraic over a finite field relative to the number of primes ramifying in  $K/F(t)$  or  $K'/F'(t)$ , where  $F'$  is the algebraic closure of  $F$  and  $K' = F'K$ . In this connection we have the following lemma.

**Lemma 7.2.** *Let  $\alpha$  be any generator of  $K$  over  $F(t)$  with  $K/F(t)$  separable. Let  $h(T) = a_0 + a_1T + \dots + T^n$  be the monic irreducible polynomial of  $\alpha$  over  $F(t)$ . Let  $D(\alpha) = \mathbf{N}_{K/F(t)}(h'(\alpha))$  where  $h'(T)$  is the derivative of  $h(T)$  with respect to  $T$ . Let  $P(\alpha)$  be the pole divisor of  $\prod_{i=0}^{n-1} a_i$ . Since  $F(t)$  is a rational function field,  $D(\alpha)$  and  $P(\alpha)$  are both polynomials in  $t$ . Let  $n(\alpha)$  be the degree of the polynomial  $D(\alpha)P(\alpha)$ . Let  $\hat{F}$  be any algebraic extension of  $F$ . Then the number of  $\hat{F}(t)$  primes ramifying in the extension  $\hat{F}K/\hat{F}(t)$  is less or equal to  $n(\alpha)$ .*

*Proof.* Since there is no constant field extension in the extension  $K/F(t)$  (see Notation 2.4), we have that  $K$  and  $\hat{F}(t)$  are linearly disjoint over  $F(t)$ . Thus,  $\hat{F}K = \hat{F}(t)(\alpha)$  and  $[\hat{F}K : \hat{F}(t)] = [K : F(t)] = n$ . Therefore if  $\mathfrak{q}$  is a prime of  $\hat{F}(t)$  ramified in the extension  $\hat{F}K/\hat{F}(t)$  we have two options: either  $\mathfrak{q}$  divides the discriminant  $D(\alpha)$  of the power basis of  $\alpha$  or  $\alpha$  is not integral at  $\mathfrak{q}$  and  $\mathfrak{q}$  divides  $P(\alpha)$ . In either case  $\mathfrak{q}$  divides  $D(\alpha)P(\alpha)$ . Since  $P(\alpha)D(\alpha)$  is a polynomial, its degree is invariant under any constant field extension and therefore the number of primes dividing  $P(\alpha)D(\alpha)$  in  $\hat{F}$  is bounded by the degree of this polynomial.  $\square$

#### ACKNOWLEDGMENTS

The authors would like to thank Arno Fehm and Stephen Simpson for helpful discussions leading to Corollary 5.1.

#### REFERENCES

- [1] Gregory L. Cherlin. Undecidability of rational function fields in nonzero characteristic. In *Logic colloquium '82 (Florence, 1982)*, volume 112 of *Stud. Logic Found. Math.*, pages 85–95. North-Holland, Amsterdam, 1984.
- [2] Martin Davis, Hilary Putnam, and Julia Robinson. The decision problem for exponential diophantine equations. *Ann. of Math. (2)*, 74:425–436, 1961.
- [3] Jean-Louis Duret. Sur la théorie élémentaire des corps de fonctions. *J. Symbolic Logic*, 51(4):948–956, 1986.
- [4] Kirsten Eisenträger. Hilbert’s tenth problem for algebraic function fields of characteristic 2. *Pacific J. Math.*, 210(2):261–281, 2003.
- [5] Kirsten Eisenträger. Hilbert’s tenth problem for function fields of varieties over  $\mathbb{C}$ . *Int. Math. Res. Not.*, (59):3191–3205, 2004.
- [6] Kirsten Eisenträger. Hilbert’s Tenth Problem for function fields of varieties over algebraically closed fields of positive characteristic, 2006. Preprint.
- [7] Kirsten Eisenträger. Hilbert’s Tenth Problem for function fields of varieties over number fields and  $p$ -adic fields. *Journal of Algebra*, 310:775–792, 2007.
- [8] Michael D. Fried and Moshe Jarden. *Field arithmetic*, volume 11 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer-Verlag, Berlin, second edition, 2005.
- [9] H. K. Kim and F. W. Roush. Diophantine unsolvability for function fields over certain infinite fields of characteristic  $p$ . *Journal of Algebra*, 152(1):230–239, 1992.

- [10] Jochen Koenigsmann. Defining transcendentals in function fields. *J. Symbolic Logic*, 67(3):947–956, 2002.
- [11] R. C. Mason. *Diophantine Equations over Function Fields*, volume 96 of *London Mathematical Society Lecture Notes*. Cambridge University Press, Cambridge, UK, 1996.
- [12] Yu. V. Matijasevič. The Diophantineness of enumerable sets. *Dokl. Akad. Nauk SSSR*, 191:279–282, 1970.
- [13] Laurent Moret-Bailly. Elliptic curves and Hilbert’s Tenth Problem for algebraic function fields over real and  $p$ -adic fields. *Journal für Reine und Angewandte Mathematic*, 587:77–143, 2006.
- [14] Thanases Pheidas. Hilbert’s tenth problem for fields of rational functions over finite fields. *Inventiones Mathematicae*, 103:1–8, 1991.
- [15] Thanases Pheidas. Endomorphisms of elliptic curves and undecidability in function fields of positive characteristic. *J. Algebra*, 273(1):395–411, 2004.
- [16] Bjorn Poonen and Florian Pop. First order definitions in function fields over anti-Mordellic fields. Preprint.
- [17] Julia Robinson. Definability and decision problems in arithmetic. *Journal of Symbolic Logic*, 14:98–114, 1949.
- [18] Alexandra Shlapentokh. Diophantine undecidability for some function fields of infinite transcendence degree and positive characteristic. *Zapiski Seminarov POMI*, 304:141–167.
- [19] Alexandra Shlapentokh. Diophantine undecidability of algebraic function fields over finite fields of constants. *Journal of Number Theory*, 58(2):317–342, June 1996.
- [20] Alexandra Shlapentokh. Hilbert’s tenth problem for algebraic function fields over infinite fields of constants of positive characteristic. *Pacific Journal of Mathematics*, 193(2):463–500, 2000.
- [21] Alexandra Shlapentokh. Diophantine undecidability of function fields of characteristic greater than 2 finitely generated over a field algebraic over a finite field. *Compositio Mathematica*, 132(1):99–120, May 2002.
- [22] Alexandra Shlapentokh. First-order definitions of rational functions and  $\mathcal{S}$ -integers over holomorphy rings of algebraic functions of characteristic 0. *Ann. Pure Appl. Logic*, 136(3):267–283, 2005.
- [23] Alexandra Shlapentokh. *Hilbert’s Tenth Problem: Diophantine Classes and Extensions to Global Fields*. Cambridge University Press, 2006.
- [24] Alfred Tarski. Undecidable theories. *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Company, Amsterdam, 1953. In collaboration with Andrzej Mostowski and Raphael M. Robinson.
- [25] Carlos Videla. Hilbert’s tenth problem for rational function fields in characteristic 2. *Proceedings of the American Mathematical Society*, 120(1):249–253, January 1994.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA.

DEPARTMENT OF MATHEMATICS, EAST CAROLINA UNIVERSITY, GREENVILLE, NC 27858, USA.