# ADDENDUM TO THE PAPER "TWO-DIMENSIONAL INFINITE PRANDTL NUMBER CONVECTION: STRUCTURE OF BIFURCATED SOLUTIONS, J. NONLINEAR SCI., 17(3), 199-220, 2007"

## TIAN MA, JUNGHO PARK, AND SHOUHONG WANG

The main objective of this addendum to the mentioned article [49] by Park is to provide some remarks on bifurcation theories for nonlinear partial differential equations (PDE) and their applications to fluid dynamics problems. We only wish to comment and list some related literatures, without any intention to provide a complete survey.

For steady state PDE bifurcation problems, the often used classical bifurcation methods include 1) the Lyapunov-Schmidt procedure, which reduces the PDE problem to a finite dimensional algebraic system, 2) the Krasnoselskii theorem for bifurcations crossing an eigenvalue of odd algebraic multiciplicity [28], 3) the Krasnoselskii theorem for potential operators, 4) the Rabinowitz global bifurcation theorem [51], 5) Crandall and Rabinowitz theorem for bifurcations crossing a simple eigenvalue [11], and 6) bifurcation from higher-order terms, regardless of the multiplicity of the eigenvalues [31, 32]. We also refer the interested readers to, among many others, [47, 8, 20, 21, 35, 42] for more comprehensive discussions. Nirenberg have a beautiful survey paper [46] on topological and variational methods for nonlinear problems, which has influenced a whole generation of nonlinear analysts.

The Hopf bifurcation, also called Poincaré-Andronov-Hopf bifurcation, was independently studied and discovered by Andronov in 1929 and Hopf in 1942 and Poincaré in 1892 for ordinary differential equations. In particular, in his paper [24], Hopf also indicated the possible application of the Hopf bifurcation theorem to bifurcation of time periodic solutions for the Navier-Stokes equations. The Hopf bifurcation was generalized to infinite dimensional setting for PDEs by Crandall and Rabinowitz [12], Marsden and McCracken [44], and Henry [23]. We mention in particular the last two references using the center manifold reduction procedure to reduce the problem to a finite dimensional problem.

<sup>1991</sup> Mathematics Subject Classification. 35Q, 35B, 37L, 76E.

Key words and phrases. Rayleigh-Bénard convection, attractor bifurcation, structure of solutions, structural stability, infinite Prandtl number.

The work was supported in part by the Office of Naval Research and by the National Science Foundation.

Over the last 30 years or so, there have been extensive studies using the bifurcation theory with symmetry methods and applications; see, among many others, D. Sattinger [53, 54, 55, 56], M. Golubitsky, I. Stewart, and D. Schaeffer [20, 21], and M. Field [14].

Recently, Ma and Wang have developed a bifurcation theory for nonlinear PDEs [36, 33]. This bifurcation theory is centered at a new notion of bifurcation, called attractor bifurcation for nonlinear evolution equations, and is synthesized in two recent books by Ma and Wang [35, 42]. Furthermore, this new bifurcation theory has been further developed by Ma and Wang into a complete new dynamic transition theory for nonlinear problems; see two recent books by Ma and Wang [42, 40] and the references therein for a more detailed account of the theory. These new theories has been used in many problems in sciences and engineering, including the Bénard convection problem and the Taylor problem in the classical fluid dynamics, doubly-diffusive convections and rotating Boussinesq equations in geophysical fluid dynamics, many phase transition problems in statistical physics, biology and chemistry; see [30, 34, 38, 35, 42, 40] and the references therein.




FIGURE 1. A bifurcated attractor containing 4 nodes (the points a, b, c, and d), 4 saddles (the points e, f, g, h), and orbits connecting these 8 points.

The main purpose of this new bifurcation theory is to study transitions of one state to another in nonlinear problems. We illustrate the concept by a simple example. For  $x = (x_1, x_2) \in \mathbb{R}^2$ , the system  $\dot{x} = \lambda x - (x_1^3, x_2^3)$ bifurcates from  $(x, \lambda) = (0, 0)$  to an attractor  $\Sigma_{\lambda} = S^1$ . This bifurcated attractor is as shown in Figure 1, and contains exactly 4 nodes (the points a, b, c, and d), 4 saddles (the points e, f, g, h), and orbits connecting these 8 points. From the physical transition point of view, as  $\lambda$  crosses 0, the new state after the system undergoes a transition is represented by the

 $\mathbf{2}$ 

whole bifurcated attractor  $\Sigma_{\lambda}$ , rather than any of the steady states or any of the connecting orbits. Note that the global attractor is the 2D region enclosed by  $\Sigma_{\lambda}$ . We point out here that the bifurcated attractor is different from the study on global attractors of a dissipative dynamical system-both finite and infinite dimensional. Global attractor studies the global long time dynamics ( see among others [15, 1, 9, 10]), while the bifurcated attractor provides a natural object for studying dynamical transitions [35, 42, 40].

Now we return to the Rayleigh-Bénard convection problem. This is classical problem in fluid dynamics. The study for this problem on the one hand plays an important role in understanding the turbulent behavior of fluid flows, and on the other hand often leads to new insights and methods toward solutions of other problems in sciences and engineering.

Linear theory of the Rayleigh-Bénard problem were essentially derived by physicists; see, among others, Chandrasekhar [3] and Drazin and Reid [13]. Bifurcating solutions of the nonlinear problem were first constructed formally by Malkus and Veronis [43]. The first rigorous proofs of the existence of bifurcating solutions were given by Yudovich [59, 60] and Rabinowitz [50]. Yudovich proved the existence of bifurcating solutions by a topological degree argument. Earlier, however, Velte [58] had proved the existence of branching solutions of the Taylor problem by a topological degree argument as well. The application of group-theoretic bifurcation theory to the Bénard convection are explored in last thirties years or so by many authors. To the best knowledge of authors, Sattinger's papers [52, 53, 55] are the first ones to use the group-theoretic point of view to study this problem in combination with the Lyapunov-Schmidt reduction procedure. Then group-theoretic methods are used, in conjunction with center manifold reduction and leading to the amplitude equations, to fluid problems by, among others, M. Golubitsky, I. Stewart, and D. Schaeffer [21], Chossat and his collaborators [6, 7], Iooss and his collaborators [25, 57], and the references therein. We would like also to mention the Hopf bifurcation result obtained by Chen and Price [4], where they use the continued fractions method first introduced by Meshalkin and Sinai [45]. Recently, Ma and Wang have used their newly developed attractor bifurcation theory mentioned earlier to study the Bénard convection problem. In [33], they proved that the Rayleigh-Bénard problem bifurcates from the basic state to an attractor  $A_R$  when the Rayleigh number R crosses the first critical Rayleigh number  $R_c$  for all physically sound boundary conditions, regardless of the geometry of the domain and the multiplicity of the eigenvalue  $R_c$  for the linear problem. Furthermore detailed characterization of solutions in the bifurcated attractor  $A_R$  in both the physical and the phase spaces for special geometries of the domain is given in [41]. Also, the bifurcated attractor  $A_R$  attracts any bounded set in  $H \setminus \Gamma$ , where H is the whole phase space and  $\Gamma$  is the stable manifold of the basic solution. Finally, we would like to point out that the bifurcation and stability analysis discussed in this and the above mentioned articles

are for viscous flows, and we refer to Friedlander and Yudovich [17] and Friedlander, Strauss, and Vishik [16] and the references therein for details.

The work presented in Park [48, 49] is an application of the aforementioned new attractor bifurcation theory to the infinite Prandtl number Bénard convection. Here are some specific comments on the results in these articles and some further developments.

FIRST, the results obtained are motivated by the attractor bifurcation theory [35] and the geometric theory for incompressible flows [37], both developed recently by Ma and Wang [35]. Without the new insights from these theories, one does not come up with the theorems proved in [48, 49] and in other related articles, as evidenced by the fact that no such theorems have been stated in the vast existing literature on the Rayleigh-Bénard convection.

SECOND, the types of solutions in this  $S^1$  attractor depend on the boundary conditions. With the periodic boundary condition in the  $x_1$  direction in this article, the bifurcated attractor consists of only steady states.

In fact, when the boundary conditions for the velocity field are free slip boundary conditions and the spatial domain is  $\Omega = (0, L)^2 \times (0, 1)$  with  $0 < L^2 < (2 - \sqrt[3]{2})/(\sqrt[3]{2} - 1)$ , Ma and Wang [41, 40] prove that the bifurcated attractor is still an  $S^1$ , consisting of exactly eight singular steady states (with four saddles and four minimal attractors) and eight heteroclinic orbits connecting these steady states. The bifurcated attractor and its detailed classification provide a global dynamic transition in both the physical and phase spaces.

Furthermore, again in a more general three-dimensional (3D) domain  $\Omega = (0, L_1) \times (0, L_2) \times (0, 1)$ , with doubly-periodic boundary conditions in the horizontal directions and the free-boundary conditions on the top and bottom, it is proved [40] that the bifurcated attractor is either  $\Sigma_R = S^5$ if  $L_2 = \sqrt{k^2 - 1}L_1$  for  $k = 2, 3, \cdots$ , or else  $\Sigma_R = S^3$ . It is clear that these bifurcated attractors contains many more solutions than the solutions derived by any group-theoretic methods.

Hence, we iterate here that the method and ideas developed by Ma and Wang are crucial to obtain these results, which can not be obtained using only the classical bifurcation theories. For the case studied in this article, the classical bifurcation theory with symmetry arguments implies that the bifurcated attractor *contains* a circle of steady states. We need, however, the new bifurcated theory to prove in particular that the bifurcated attractors are *exactly* an  $S^1$ .

Furthermore, it is also much obvious, as partially indicated, that the 3D results can not be derived by group-theoretic methods. In addition, for general boundary conditions such as the free-slip boundary conditions mentioned above, no symmetry can be used, and the classical amplitude equation methods fails to derive the dynamics.

4

In short, although group-theoretic bifurcation methods are useful in different problems, the need for other more general methods is obvious and inevitable as suggested by, among others, D. Sattinger [53, 55], one of the pioneers of the group-theoretic bifurcation methods, and B. Grünbaun [22].

THIRD, it is proved in [33, 41, 48, 49] that the bifurcated attractor in the Bénard convection problem attracts **any bounded set** in  $H \setminus \Gamma$ , where H is the whole phase space and  $\Gamma$  is the stable manifold of the basic solution. To the best of the knowledge of authors, this "global" stability result can not be derived from any existing methods. It is obtained by using a new stability result proved by Ma and Wang [33], which is derived using a combination of energy estimates and topological arguments.

As Kirchgässner indicated in [27], "an ideal stability theorem would include all physically meaningful perturbations and establish the local stability of a selected class of stationary solutions, and today we are still far from this goal." On the other hand, fluid flows are normally time dependent. Therefore bifurcation analysis for steady state problems provides in general only partial answers to the problem, and is not enough for solving the stability problem. Hence it appears that the right notion of asymptotic stability after the first bifurcation should be best described by the attractor near, but excluding, the trivial state. It is one of our main motivations for introducing attractor bifurcation theory and the dynamic transition theory [35, 42, 40].

FOURTH, the geometric theory for incompressible flows recently developed by Ma and Wang [37] is crucial for the structure and its stability of the solutions in the physical spaces obtained in the main theorems in [49]. Also, we note that a special structure with rolls separated by a cross channel flow derived in [41] has not been rigorously examined in the Bénard convection setting although it has been observed in other physical contexts such as the Branstator-Kushnir waves in the atmospheric dynamics [2, 29].

For completeness, we mention that this geometric theory for incompressible flows consists of research in two steps: 1) the study of the structure and its transitions/evolutions of divergence-free vector fields, and 2) the study of the structure and its transitions of velocity fields for 2-D incompressible fluid flows governed by the Navier-Stokes equations or the Euler equations. The original motivation of this research program was to understand the dynamics of the ocean currents in the physical space, and it turns out that there is a much richer new mathematical theory with more applications than the original motivation from the Oceanography. Among other results, for example, the theory leads to a rigorous characterization of boundary layer separation on when, where, and how the separation occurs and to make connections between the time and location of the separation; see Ghil, Ma and Wang [18, 19], Ma and Wang [37], and the references therein. This is a long standing problem in fluid mechanics going back to the pioneering work of Prandtl (1904); see also Chorin and Marsden [5], and Jäger, Lax and Morawetz [26].

With this characterization of the boundary layer-separation in our disposal, Ma and Wang [39] are able to derive a rigorous characterization of the boundary-layer and interior separations in the Taylor-Couette-Poiseuille flow. The results obtained provide a rigorous characterization on how, when and where the propagating Taylor vortices (PTV) are generated. In particular, contrary to what is commonly believed, it is shown that the PTV do not appear after the first dynamical bifurcation, and they appear only when the Taylor number is further increased to cross another critical value so that a structural bifurcation occurs. This structural bifurcation corresponds to the boundary-layer and interior separations of the flow structure in the physical space.

#### References

- A. V. BABIN AND M. I. VISHIK, Regular attractors of semigroups and evolution equations, J. Math. Pures Appl. (9), 62 (1983), pp. 441–491 (1984).
- G. W. BRANSTATOR, A striking example of the atmosphere's leading traveling pattern, J. Atmos. Sci., 44 (1987), pp. 2310–2323.
- [3] S. CHANDRASEKHAR, Hydrodynamic and Hydromagnetic Stability, Dover Publications, Inc., 1981.
- [4] Z.-M. CHEN AND W. G. PRICE, Remarks on: "Time dependent periodic Navier-Stokes flows on a two-dimensional torus" [Comm. Math. Phys. 179 (1996), no. 3, 577–597; MR1400753 (97g:58152)], Comm. Math. Phys., 207 (1999), pp. 81–106.
- [5] A. CHORIN AND J. MARSDEN, A Mathematical Introduction to Fluid Mechanics, Springer-Verlag, 1997.
- [6] P. CHOSSAT AND G. IOOSS, *The Couette-Taylor problem*, vol. 102 of Applied Mathematical Sciences, Springer-Verlag, New York, 1994.
- [7] P. CHOSSAT AND R. LAUTERBACH, Methods in equivariant bifurcations and dynamical systems, vol. 15 of Advanced Series in Nonlinear Dynamics, World Scientific Publishing Co. Inc., River Edge, NJ, 2000.
- [8] S. N. CHOW AND J. K. HALE, Methods of bifurcation theory, vol. 251 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science], Springer-Verlag, New York, 1982.
- [9] P. CONSTANTIN AND C. FOIAS, Global Lyapunov exponents, Kaplan-Yorke formulas and the dimension of the attractors for 2D Navier-Stokes equations, Comm. Pure Appl. Math., 38 (1985), pp. 1–27.
- [10] P. CONSTANTIN, C. FOIAS, AND R. TEMAM, Attractors representing turbulent flows, Mem. Amer. Math. Soc., 53 (1985), pp. vii+67.
- [11] M. G. CRANDALL AND P. H. RABINOWITZ, Bifurcation from simple eigenvalues, J. Functional Analysis, 8 (1971), pp. 321–340.
- [12] —, The Hopf bifurcation theorem in infinite dimensions, Arch. Rational Mech. Anal., 67 (1977), pp. 53–72.
- [13] P. DRAZIN AND W. REID, Hydrodynamic Stability, Cambridge University Press, 1981.
- [14] M. FIELD, Lectures on bifurcations, dynamics and symmetry, vol. 356 of Pitman Research Notes in Mathematics Series, Longman, Harlow, 1996.
- [15] C. FOIAŞ AND R. TEMAM, Some analytic and geometric properties of the solutions of the evolution Navier-Stokes equations, J. Math. Pures Appl. (9), 58 (1979), pp. 339– 368.
- [16] S. FRIEDLANDER, W. STRAUSS, AND M. VISHIK, Nonlinear instability in an ideal fluid, Ann. Inst. H. Poincaré Anal. Non Linéaire, 14 (1997), pp. 187–209.

- [17] S. FRIEDLANDER AND V. YUDOVICH, Instabilities in fluid motion, Notices Amer. Math. Soc., 46 (1999), pp. 1358–1367.
- [18] M. GHIL, T. MA, AND S. WANG, Structural bifurcation of 2-D incompressible flows, Indiana Univ. Math. J., 50 (2001), pp. 159–180. Dedicated to Professors Ciprian Foias and Roger Temam (Bloomington, IN, 2000).
- [19] \_\_\_\_\_, Structural bifurcation of 2-D nondivergent flows with Dirichlet boundary conditions: applications to boundary-layer separation, SIAM J. Appl. Math., 65 (2005), pp. 1576–1596 (electronic).
- [20] M. GOLUBITSKY AND D. G. SCHAEFFER, Singularities and groups in bifurcation theory. Vol. I, vol. 51 of Applied Mathematical Sciences, Springer-Verlag, New York, 1985.
- [21] M. GOLUBITSKY, I. STEWART, AND D. G. SCHAEFFER, Singularities and groups in bifurcation theory. Vol. II, vol. 69 of Applied Mathematical Sciences, Springer-Verlag, New York, 1988.
- [22] B. GRÜNBAUN, Periodic ornamentation of the fabric plane: Lessons from peruvian fabrics, Symmetry, 1 (1990), pp. 45 68.
- [23] D. HENRY, Geometric theory of semilinear parabolic equations, vol. 840 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1981, early version appeared as University of Kentucky Lecture notes in 1974.
- [24] E. HOPF, Abzweigung einer periodischen Ibsung von einer stationaren lösung eines differentialsystems, Ber. Math.-Phys. K. Sachs. Akad. Wiss. Leipzig, 94 (1942), pp. 1– 22.
- [25] G. IOOSS AND M. ADELMEYER, Topics in bifurcation theory and applications, vol. 3 of Advanced Series in Nonlinear Dynamics, World Scientific Publishing Co. Inc., River Edge, NJ, second ed., 1998.
- [26] W. JÄGER, P. LAX, AND C. S. MORAWETZ, Olga Oleinik (1925–2001), Notices Amer. Math. Soc., 50 (2003), pp. 220–223.
- [27] K. KIRCHGÄSSNER, Bifurcation in nonlinear hydrodynamic stability, SIAM Rev., 17 (1975), pp. 652–683.
- [28] M. A. KRASNOSEL'SKII, Topologicheskie metody v teorii nelineinykh integralnykh uravnenii, Gosudarstv. Izdat. Tehn.-Teor. Lit., Moscow, 1956.
- [29] K. KUSHNIR, Retrograding wintertime low-frequency disturbances over the north pacific ocean, J. Atmos. Sci., 44 (1987), pp. 2727–2742.
- [30] T. MA, J. PARK, AND S. WANG, Dynamic bifurcation of the Ginzburg-Landau equation, SIAM J. Appl. Dyn. Syst., 3 (2004), pp. 620–635 (electronic).
- [31] T. MA AND S. WANG, Bifurcation of nonlinear equations. I. Steady state bifurcation, Methods Appl. Anal., 11 (2004), pp. 155–178.
- [32] \_\_\_\_\_, Bifurcation of nonlinear equations. II. Dynamic bifurcation, Methods Appl. Anal., 11 (2004), pp. 179–209.
- [33] —, Dynamic bifurcation and stability in the Rayleigh-Bénard convection, Commun. Math. Sci., 2 (2004), pp. 159–183.
- [34] —, Bifurcation and stability of superconductivity, J. Math. Phys., 46 (2005), pp. 095112, 31.
- [35] —, Bifurcation Theory and Applications, World Scientific, 2005.
- [36] —, Dynamic bifurcation of nonlinear evolution equations, Chinese Ann. Math. Ser. B, 26 (2005), pp. 185–206.
- [37] —, Geometric theory of incompressible flows with applications to fluid dynamics, vol. 119 of Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 2005.
- [38] \_\_\_\_\_, Stability and bifurcation of the Taylor problem, Arch. Ration. Mech. Anal., 181 (2006), pp. 149–176.
- [39] —, Boundary layer and interior separations in the Taylor–Couette–Poiseuille flow, submitted; see also ArXiv, math-ph/0701073, (2007).

- [40] —, Phase Transition Dynamics in Nonlinear Sciences, in preparation, 2007.
- [41] —, Rayleigh-benard convection: Dynamics and structure in the physical space, Communications in Math. Sci., 5:3 (2007), pp. 553–574.
- [42] \_\_\_\_\_, Stability and Bifurcation of Nonlinear Evolution Equations, Chinese Academic Press, 2007.
- [43] W. V. R. MALKUS AND G. VERONIS, Finite amplitude cellular convection, J. Fluid Mech., 4 (1958), pp. 225–260.
- [44] J. E. MARSDEN AND M. MCCRACKEN, The Hopf bifurcation and its applications, Springer-Verlag, New York, 1976. With contributions by P. Chernoff, G. Childs, S. Chow, J. R. Dorroh, J. Guckenheimer, L. Howard, N. Kopell, O. Lanford, J. Mallet-Paret, G. Oster, O. Ruiz, S. Schecter, D. Schmidt and S. Smale, Applied Mathematical Sciences, Vol. 19.
- [45] L. D. MEŠALKIN AND J. G. SINAĬ, Investigation of the stability of a stationary solution of a system of equations for the plane movement of an incompressible viscous liquid, J. Appl. Math. Mech., 25 (1961), pp. 1700–1705.
- [46] L. NIRENBERG, Variational and topological methods in nonlinear problems, Bull. Amer. Math. Soc. (N.S.), 4 (1981), pp. 267–302.
- [47] —, Topics in nonlinear functional analysis, vol. 6 of Courant Lecture Notes in Mathematics, New York University Courant Institute of Mathematical Sciences, New York, 2001. Chapter 6 by E. Zehnder, Notes by R. A. Artino, Revised reprint of the 1974 original.
- [48] J. PARK, Dynamic bifurcation theory of Rayleigh-Bénard convection with infinite Prandtl number, Discrete Contin. Dyn. Syst. Ser. B, 6 (2006), pp. 591–604 (electronic).
- [49] \_\_\_\_\_, Two-dimensional infinite Prandtl number convection: structure of bifurcated solutions, J. Nonlinear Sci., 17 (2007), pp. 199–220.
- [50] P. H. RABINOWITZ, Existence and nonuniqueness of rectangular solutions of the Bénard problem, Arch. Rational Mech. Anal., 29 (1968), pp. 32–57.
- [51] P. H. RABINOWITZ, Some global results for nonlinear eigenvalue problems, J. Functional Analysis, 7 (1971), pp. 487–513.
- [52] D. H. SATTINGER, Selection mechanisms for pattern formation, Arch. Rational Mech. Anal., 66 (1977), pp. 31–42.
- [53] —, Group representation theory, bifurcation theory and pattern formation, J. Funct. Anal., 28 (1978), pp. 58–101.
- [54] —, *Group-theoretic methods in bifurcation theory*, vol. 762 of Lecture Notes in Mathematics, Springer, Berlin, 1979. With an appendix entitled "How to find the symmetry group of a differential equation" by Peter Olver.
- [55] —, Bifurcation and symmetry breaking in applied mathematics, Bull. Amer. Math. Soc. (N.S.), 3 (1980), pp. 779–819.
- [56] —, Branching in the presence of symmetry, vol. 40 of CBMS-NSF Regional Conference Series in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1983.
- [57] A. VANDERBAUWHEDE AND G. IOOSS, Center manifold theory in infinite dimensions, in Dynamics reported: expositions in dynamical systems, vol. 1 of Dynam. Report. Expositions Dynam. Systems (N.S.), Springer, Berlin, 1992, pp. 125–163.
- [58] W. VELTE, Stabilität and verzweigung stationärer lösungen der davier-stokeschen gleichungen, Arch. Rat. Mech. Anal., 22 (1966), pp. 1–14.
- [59] V. I. YUDOVICH, Free convection and bifurcation, J. Appl. Math. Mech., 31 (1967), pp. 103–114.
- [60] —, Stability of convection flows, J. Appl. Math. Mech., 31 (1967), pp. 272–281.

8

(TM) DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU, P. R. CHINA AND DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405

(SW) Department of Mathematics, Indiana University, Bloomington, IN 47405

 $E\text{-}mail\ address:\ \texttt{junjupar@indiana.edu}$ 

(SW) Department of Mathematics, Indiana University, Bloomington, IN 47405

*E-mail address*: showang@indiana.edu