

COMMENT ON $\mathbf{GL}(2, \mathbb{R})$ GEOMETRY OF 4TH ORDER ODE'S

PAWEŁ NUROWSKI

ABSTRACT. We describe 4th order ODEs satisfying two contact invariant conditions of Bryant in terms of the Ricci tensor of a certain $\mathfrak{gl}(2, \mathbb{R})$ valued connection. We also provide nonhomogeneous examples of such ODEs.

MSC classification: 53A40, 53B05, 34C30, 34C31

CONTENTS

1. Introduction	1
2. The closed system	4
3. Proof of the main theorem	5
4. $\mathbf{GL}(2, \mathbb{R})$ geometry on the solution space	9
5. Examples	12
5.1. Equations with symmetric Ricci tensor	12
5.2. Inhomogeneous examples	16
References	16

1. INTRODUCTION

Recently there is a growing interest in the geometrization program of ODEs [5, 6, 7, 8, 11, 10]. Although the program may be traced back to S. Lie [9] and M. A. Tresse [12], and although it was formulated by E. Cartan and S. S. Chern in the 1940s [2, 3, 4], it was not very popular until the works of R. Bryant (see e.g. [1]) on the invariants of the fourth order ODEs. In the present note we restate some of the results of [1] in terms of the invariants of the recently discussed $\mathbf{GL}(2, \mathbb{R})$ geometry of ODEs [8]. In particular we interpret Bryant's results in terms of the Ricci tensor of a certain $\mathfrak{gl}(2, \mathbb{R})$ connection, which characterises the ODEs satisfying contact invariant conditions of Bryant [1].

Our starting point is the following well known

Proposition 1.1. *Ordinary differential equation*

$$y^{(4)} = 0$$

has $\mathbf{GL}(2, \mathbb{R}) \times_{\rho} \mathbb{R}^4$ as its group of contact symmetries. Here $\rho : \mathbf{GL}(2, \mathbb{R}) \rightarrow \mathbf{GL}(4, \mathbb{R})$ is the 4-dimensional irreducible representation of $\mathbf{GL}(2, \mathbb{R})$.

Date: November 2, 2018.

This research was supported by the KBN grant 1 P03B 07529.

The representation ρ , at the level of the Lie algebra $\mathfrak{gl}(2, \mathbb{R})$, is given in terms of the Lie algebra generators

$$(1.1) \quad \begin{aligned} E_+ &= \begin{pmatrix} 0 & 3 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & E_- &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 \end{pmatrix}, & E_0 &= \begin{pmatrix} -3 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix}, \\ E &= - \begin{pmatrix} 3 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix}. \end{aligned}$$

These matrices satisfy the $\mathfrak{gl}(2, \mathbb{R})$ commutation relations

$$[E_0, E_+] = -2E_+ \quad , \quad [E_0, E_-] = 2E_- \quad , \quad [E_+, E_-] = -E_0 \quad ,$$

where the commutator in the $\mathfrak{gl}(2, \mathbb{R}) = \text{Span}_{\mathbb{R}}(E_-, E_+, E_0, E) \subset \text{End}(\mathbb{R}^4)$ is the usual commutator of matrices.

Now, we consider a general 4-th order ODE

$$(1.2) \quad y^{(4)} = F(x, y, y', y'', y^{(3)}).$$

To simplify the notation, we introduce the coordinates $x, y, y_1 = y', y_2 = y'', y_3 = y^{(3)}$ on the 5-dimensional *jet space* J . Introducing the four *contact forms*

$$(1.3) \quad \begin{aligned} \omega^0 &= dy - y_1 dx \\ \omega^1 &= dy_1 - y_2 dx \\ \omega^2 &= dy_2 - y_3 dx \\ \omega^3 &= dy_3 - F(x, y, y_1, y_2, y_3) dx, \end{aligned}$$

and an additional 1-form

$$w_+ = dx,$$

we define a *contact transformation* to be a diffeomorphism $\phi : J \rightarrow J$ which transforms the above five one-forms via:

$$(1.4) \quad \begin{aligned} \phi^* \omega^0 &= \alpha_0^0 \omega^0 \\ \phi^* \omega^1 &= \alpha_0^1 \omega^0 + \alpha_1^1 \omega^1 \\ \phi^* \omega^2 &= \alpha_0^2 \omega^0 + \alpha_1^2 \omega^1 + \alpha_2^2 \omega^2 \\ \phi^* \omega^3 &= \alpha_0^3 \omega^0 + \alpha_1^3 \omega^1 + \alpha_2^3 \omega^2 + \alpha_3^3 \omega^3 \\ \phi^* w_+ &= \alpha_0^4 \omega^0 + \alpha_1^4 \omega^1 + \alpha_4^4 w_+. \end{aligned}$$

Here α_j^i , $i, j = 0, 1, 2, 3, 4, 5$, are real functions on J such that

$$\alpha_0^0 \alpha_1^1 \alpha_2^2 \alpha_3^3 \alpha_4^4 \neq 0.$$

The contact equivalence problem for the 4th order ODEs (1.2) can be studied in terms of the invariant forms $(\theta^0, \theta^1, \theta^2 \theta^3, \Omega_+)$ defined by

$$(1.5) \quad \begin{pmatrix} \theta^0 \\ \theta^1 \\ \theta^2 \\ \theta^3 \\ \Omega_+ \end{pmatrix} = \begin{pmatrix} \alpha_0^0 & & & & \\ \alpha_0^1 & \alpha_1^1 & & & \\ \alpha_0^2 & \alpha_1^2 & \alpha_2^2 & & \\ \alpha_0^3 & \alpha_1^3 & \alpha_2^3 & \alpha_3^3 & \\ \alpha_0^4 & \alpha_1^4 & & & \alpha_4^4 \end{pmatrix} \begin{pmatrix} \omega^0 \\ \omega^1 \\ \omega^2 \\ \omega^3 \\ w_+ \end{pmatrix}.$$

Among all ODEs (1.2) considered modulo contact transformations (1.4) there is a remarkable class for which the invariant forms satisfy

$$\begin{aligned}
 d\theta^0 &= 3(\Omega + \Omega_0) \wedge \theta^0 - 3\Omega_+ \wedge \theta^1 \\
 d\theta^1 &= -\Omega_- \wedge \theta^0 + (3\Omega + \Omega_0) \wedge \theta^1 - 2\Omega_+ \wedge \theta^2 \\
 d\theta^2 &= -2\Omega_- \wedge \theta^1 + (3\Omega - \Omega_0) \wedge \theta^2 - \Omega_+ \wedge \theta^3 \\
 d\theta^3 &= -3\Omega_- \wedge \theta^2 + 3(\Omega - \Omega_0) \wedge \theta^3.
 \end{aligned}
 \tag{1.6}$$

This system is defined on an 8-dimensional $\mathbf{GL}(2, \mathbb{R})$ principal fibre bundle P over the solution space M^4 for the corresponding ODE (1.2). The invariant forms $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+)$ together with the additional three 1-forms $(\Omega_-, \Omega_0, \Omega)$ constitute a well defined coframe on P .

As noted by Bryant [1], the class of ODEs having forms $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+, \Omega_-, \Omega_0, \Omega)$ of system (1.6), is distinguished by the demand that their defining functions $F = F(x, y, y_1, y_2, y_3)$ satisfy the following two conditions:

$$\begin{aligned}
 4D^2F_3 - 8DF_2 + 8F_1 - 6DF_3F_3 + 4F_2F_3 + F_3^3 &= 0, \\
 160D^2F_2 - 640DF_1 + 144(DF_3)^2 - 352DF_3F_2 + 144F_2^2 - \\
 80DF_2F_3 + 160F_1F_3 - 72DF_3F_3^2 + 88F_2F_3^2 + 9F_3^4 + 16000F_y &= 0.
 \end{aligned}
 \tag{1.7}$$

Here $F_i = \frac{\partial F}{\partial y_i}$ and $D = \partial_x + y_1\partial_y + y_2\partial_{y_1} + y_3\partial_{y_2} + F\partial_{y_3}$. Bryant's conditions (1.7), considered simultaneously, are *contact invariant*; if the ODE undergoes contact transformation of its variables, the conditions (1.7) are preserved. Examples are known of ODEs satisfying these conditions [1], the simplest being

$$y^{(4)} = (y^{(3)})^{(4/3)}.
 \tag{1.8}$$

The purpose of this note is to establish a theorem on speciality of a $\mathfrak{gl}(2, \mathbb{R})$ -valued connection defined by such ODEs on their solution spaces.

2. THE CLOSED SYSTEM

Let us make the following choice

$$\begin{aligned}
\alpha_0^0 &= -3\alpha_1^1\alpha_4^4 \\
\alpha_0^2 &= -\frac{(\alpha_0^1)^2}{3\alpha_1^1\alpha_4^4} + \frac{\alpha_1^1}{240\alpha_4^4}(-24DF_3 + 36F_2 + 11F_3^2) \\
\alpha_1^2 &= -\frac{2\alpha_0^1}{3\alpha_4^4} + \frac{\alpha_1^1}{12\alpha_4^4}F_3 \\
\alpha_2^2 &= -\frac{\alpha_1^1}{2\alpha_4^4} \\
\alpha_0^3 &= \frac{(\alpha_0^1)^3}{9(\alpha_1^1\alpha_4^4)^2} + \frac{\alpha_0^1}{240(\alpha_4^4)^2}(24DF_3 - 36F_2 - 11F_3^2) + \\
(2.1) \quad &\frac{\alpha_1^1}{720(\alpha_4^4)^2}(36(DF_2 - 4F_1) + 18(DF_3 - 2F_2)F_3 - 7F_3^3) \\
\alpha_1^3 &= \frac{(\alpha_0^1)^2}{3\alpha_1^1(\alpha_4^4)^2} - \frac{\alpha_0^1}{12(\alpha_4^4)^2}F_3 + \frac{\alpha_1^1}{240(\alpha_4^4)^2}(36DF_3 - 84F_2 - 19F_3^2) \\
\alpha_2^3 &= \frac{\alpha_0^1}{2(\alpha_4^4)^2} - \frac{\alpha_1^1}{4(\alpha_4^4)^2}F_3 \\
\alpha_3^3 &= \frac{\alpha_1^1}{2(\alpha_4^4)^2} \\
\alpha_0^4 &= -\frac{\alpha_4^4}{60}(12DF_{33} - 6F_{23} + F_3F_{33}) \\
\alpha_1^4 &= \frac{\alpha_4^4}{6}F_{33}
\end{aligned}$$

for the group parameters defining forms $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+)$ of (1.5). Then we have the following

Theorem 2.1. *If a fourth order ODE*

$$(2.2) \quad y^{(4)} = F(x, y, y', y'', y^{(3)})$$

satisfies contact invariant conditions

$$\begin{aligned}
(2.3) \quad &4D^2F_3 - 8DF_2 + 8F_1 - 6DF_3F_3 + 4F_2F_3 + F_3^3 = 0, \\
&160D^2F_2 - 640DF_1 + 144(DF_3)^2 - 352DF_3F_2 + 144F_2^2 - \\
&80DF_2F_3 + 160F_1F_3 - 72DF_3F_3^2 + 88F_2F_3^2 + 9F_3^4 + 16000F_y = 0
\end{aligned}$$

then the manifold P parametrised by $(x, y, y_1, y_2, y_3, \alpha_0^1, \alpha_1^1, \alpha_4^4)$ is a principal $\mathbf{GL}(2, \mathbb{R})$ bundle $P \rightarrow M^4$ over the solution space M^4 of (2.2) and forms $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+)$, together with additional three 1-forms $(\Omega_-, \Omega_0, \Omega)$, constitute an invariant coframe

on P satisfying

$$(2.4) \quad \begin{aligned} d\theta^0 &= 3(\Omega + \Omega_0) \wedge \theta^0 - 3\Omega_+ \wedge \theta^1 \\ d\theta^1 &= -\Omega_- \wedge \theta^0 + (3\Omega + \Omega_0) \wedge \theta^1 - 2\Omega_+ \wedge \theta^2 \\ d\theta^2 &= -2\Omega_- \wedge \theta^1 + (3\Omega - \Omega_0) \wedge \theta^2 - \Omega_+ \wedge \theta^3 \\ d\theta^3 &= -3\Omega_- \wedge \theta^2 + 3(\Omega - \Omega_0) \wedge \theta^3 \end{aligned}$$

$$(2.5) \quad \begin{aligned} d\Omega_+ &= 2\Omega_0 \wedge \Omega_+ + \frac{1}{12}(-3a_0 + 4b_1)\theta^0 \wedge \theta^1 + \\ &\quad \frac{1}{4}(a_1 + 2b_2)\theta^0 \wedge \theta^2 + \frac{1}{24}(3a_2 + 4b_3)\theta^0 \wedge \theta^3 + \\ &\quad \frac{1}{8}(-5a_2 + 4b_3)\theta^1 \wedge \theta^2 + \frac{1}{6}b_4\theta^1 \wedge \theta^3 \\ d\Omega_- &= -2\Omega_0 \wedge \Omega_- + \frac{1}{6}b_0\theta^0 \wedge \theta^2 + \\ &\quad \frac{1}{24}(-3a_0 + 4b_1)\theta^0 \wedge \theta^3 + \frac{1}{8}(5a_0 + 4b_1)\theta^1 \wedge \theta^2 + \\ &\quad \frac{1}{4}(-a_1 + 2b_2)\theta^1 \wedge \theta^3 + \frac{1}{12}(3a_2 + 4b_3)\theta^2 \wedge \theta^3 \\ d\Omega_0 &= \Omega_+ \wedge \Omega_- - \frac{1}{6}b_0\theta^0 \wedge \theta^1 + \\ &\quad \frac{1}{24}(-3a_0 - 4b_1)\theta^0 \wedge \theta^2 + \frac{1}{4}a_1\theta^0 \wedge \theta^3 - \frac{1}{4}a_1\theta^1 \wedge \theta^2 + \\ &\quad \frac{1}{24}(-3a_2 + 4b_3)\theta^1 \wedge \theta^3 + \frac{1}{6}b_4\theta^2 \wedge \theta^3 \\ d\Omega &= -\frac{1}{6}b_0\theta^0 \wedge \theta^1 - \frac{1}{3}b_1\theta^0 \wedge \theta^2 - \frac{1}{6}b_2\theta^0 \wedge \theta^3 - \\ &\quad \frac{1}{2}b_2\theta^1 \wedge \theta^2 - \frac{1}{3}b_3\theta^1 \wedge \theta^3 - \frac{1}{6}b_4\theta^2 \wedge \theta^3. \end{aligned}$$

The coefficients $a_0, a_1, a_2, b_0, b_1, b_2, b_3, b_4$ are totally determined by (2.2) and are expressible in terms of the derivatives of function F and the coordinates. The simplest of these coefficients are:

$$\begin{aligned} b_4 &= -2 \frac{(\alpha_4^4)^3}{(\alpha_1^1)^2} F_{333} \\ b_3 &= -\frac{(\alpha_4^4)^2}{12(\alpha_1^1)^2} (6DF_{333} + 5F_3F_{333}) - \frac{2\alpha_0^1(\alpha_4^4)^2}{3(\alpha_1^1)^3} F_{333} \\ b_2 &= -\frac{2\alpha_4^4(\alpha_0^1)^2}{9(\alpha_1^1)^4} F_{333} - \frac{\alpha_4^4\alpha_0^1}{18(\alpha_1^1)^3} (6DF_{333} + 5F_3F_{333}) + \\ &\quad \frac{\alpha_4^4}{360(\alpha_1^1)^2} [60(2DF_{233} + 4F_{133} - 2F_{223} + DF_{333}F_3) + (-36DF_3 + 204F_2 + 79F_3^2)F_{333}] \\ a_2 &= -\frac{(\alpha_4^4)^2}{45(\alpha_1^1)^2} (18DF_{333} + 24F_{233} + 4F_{33}^2 + 27F_3F_{333}). \end{aligned}$$

Other coefficients are given in the next two sections.

The proof of this theorem is a lengthy calculation based on a variant of Cartan's equivalence method. In the next section we outline the main points of the proof.

3. PROOF OF THE MAIN THEOREM

The basic idea in the proof of Theorem 2.1 is to force 1-forms (1.5) to satisfy system (1.6). This requirement makes restrictions on the free parameters α^i_j and, more importantly, on the possible functions $F = F(x, y, y', y'', y^{(3)})$ defining the ODE.

The main steps when imposing (1.6) on (1.5) are:

- 1) equation $d\theta^0 \wedge \theta^0 \wedge \theta^2 = 3\Omega_+ \wedge \theta^0 \wedge \theta^1 \wedge \theta^2$ requires $\alpha^0_0 = -3\alpha^1_1\alpha^4_4$,
- 2) the first equation (1.6) gives a relation between Ω , Ω_0 , $d\alpha^4_4$ and $d\alpha^1_1$,
- 3) similarly, equation $d\theta^1 \wedge \theta^1 \wedge \theta^2 = -\Omega_- \wedge \theta^0 \wedge \theta^1 \wedge \theta^2$ gives a relation between Ω_- , $d\alpha^1_0$, $d\alpha^1_1$,
- 4) equation $d\theta^1 \wedge \theta^0 \wedge \theta^1 = -2\Omega_+ \wedge \theta^0 \wedge \theta^1 \wedge \theta^2$ gives $\alpha^2_2 = -\frac{\alpha^1_1}{2\alpha^4_4}$,
- 5) equation $d\theta^1 \wedge \theta^0 \wedge \theta^2 = -(3\Omega + \Omega_0) \wedge \theta^0 \wedge \theta^1 \wedge \theta^2$ gives a relation between Ω , Ω_0 and $d\alpha^1_1$,
- 6) now, the expressions for $d\theta^2 \wedge \theta^0 \wedge \theta^1 \wedge \theta^3$, $d\theta^2 \wedge \theta^0 \wedge \theta^1 \wedge \theta^3$, $d\theta^3 \wedge \theta^0 \wedge \theta^1 \wedge \theta^2$ enable us to fix α^3_2 , α^3_3 , and α^2_1 respectively,
- 7) considering successively $d\theta^3 \wedge \theta^0 \wedge \theta^1$, $d\theta^2 \wedge \theta^0$, $d\theta^2 \wedge \theta^1 \wedge \theta^2 \wedge \theta^3$ we fix α^2_0 , α^3_1 , α^3_0 ,
- 8) now the requirement $d\theta^3 \wedge \theta^0 \wedge \theta^2 \wedge \theta^3 = 0$ gives the first Bryant condition $4D^2F_3 - 8DF_2 + 8F_1 - 6DF_3F_3 + 4F_2F_3 - F_3^3 = 0$,
- 9) the second of Bryant's conditions (1.7) is equivalent to the requirement that $d\theta^3 \wedge \theta^1 \wedge \theta^2 \wedge \theta^3 = 0$,
- 10) now, having Bryant's conditions determined, it is straightforward to obtain the required system (1.6) and express all the α^i_j s in terms of α^1_0 , α^1_1 and α^4_4 only,
- 11) the expressions for α^i_j s are given by (2.1); inserting them to (1.5) we get the invariant forms $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+)$
- 12) forms $\Omega_0, \Omega_-, \Omega$ are determined by the linear relations from points 2), 3) and 5).

In this way one finds the explicit expressions for the invariant coframe satisfying system (1.6). Instead of giving these formulae we present formulae for $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+)$ evaluated at $(\alpha^1_0, \alpha^1_1, \alpha^4_4) = (0, 1, 1)$. Denoting these forms by $(\theta^0_0, \theta^1_0, \theta^2_0, \theta^3_0, \Omega^0_+)$, we have

$$\begin{aligned}
\theta^0_0 &= -3\omega^0 \\
\theta^1_0 &= \omega^1 \\
\theta^2_0 &= \frac{1}{240}(-24DF_3 + 36F_2 + 11F_3^2)\omega^0 + \frac{1}{12}F_3\omega^1 - \frac{1}{2}\omega^2 \\
\theta^3_0 &= \frac{1}{720}\left(36(DF_2 - 4F_1) + 18(DF_3 - 2F_2)F_3 - 7F_3^2\right)\omega^0 + \\
&\quad \frac{1}{240}(36DF_3 - 84F_2 - 19F_3^2)\omega^1 - \frac{1}{4}F_3\omega^2 + \frac{1}{2}\omega^3 \\
\Omega^0_+ &= -\frac{1}{60}(12DF_{33} - 6F_{23} + F_3F_{33})\omega^0 + \frac{1}{6}F_{33}\omega^1 + w_+.
\end{aligned}$$

The remaining three 1-forms $(\Omega_0, \Omega_-, \Omega)$, when written in the gauge $(\alpha^1_0, \alpha^1_1, \alpha^4_4) = (0, 1, 1)$, read:

$$\begin{aligned}
\Omega_0^0 &= \frac{1}{4320}(72DF_{23} + 432F_{13} - 288F_{22} + 60DF_{33}F_3 - 216F_{23}F_3 - \\
&\quad 108DF_3F_{33} + 324F_2F_{33} + 47F_3^2F_{33})\theta_0^0 + \frac{1}{180}(3DF_{33} - 9F_{23} - F_3F_{33})\theta_0^1 + \\
&\quad \frac{1}{6}F_{33}\theta_0^2 - \frac{1}{12}F_3\theta_0^4 \\
\Omega_-^0 &= \frac{1}{64800}(720DF_{22} + 288DF_3DF_{33} - 2160F_{12} - 432DF_{33}F_2 + 216DF_3F_{23} + \\
&\quad 216F_2F_{23} + 720DF_{23}F_3 - 1080F_{13}F_3 - 360F_{22}F_3 + 48DF_{33}F_3^2 - \\
&\quad 174F_{23}F_3^2 - 360DF_2F_{33} + 1440F_1F_{33} + 24DF_3F_3F_{33} + 324F_2F_3F_{33} + \\
&\quad 29F_3^3F_{33} + 3600F_{3y})\theta_0^0 + \\
&\quad \frac{1}{1080}(-108DF_{23} - 288F_{13} + 252F_{22} - 54DF_{33}F_3 + 186F_{23}F_3 + \\
&\quad 66DF_3F_{33} - 252F_2F_{33} - 31F_3^2F_{33})\theta_0^1 + \frac{1}{90}(12DF_{33} - 6F_{23} + F_3F_{33})\theta_0^2 + \\
&\quad \frac{1}{360}(-24DF_3 + 36F_2 + 11F_3^2)\theta_0^4 \\
\Omega^0 &= \frac{1}{4320}(120DF_{23} + 240F_{13} - 240F_{22} + 36DF_{33}F_3 - 168F_{23}F_3 - \\
&\quad 36DF_3F_{33} + 204F_2F_{33} + 17F_3^2F_{33})\theta_0^0 + \\
&\quad \frac{1}{12}(-DF_{33} + F_{23})\theta_0^1 - \frac{1}{6}F_{33}\theta_0^2 + \frac{1}{12}F_3\theta_0^4.
\end{aligned}$$

All the eight forms $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+^0, \Omega_0^0, \Omega_-^0, \Omega^0)$ satisfy system (2.4)-(2.5), with corresponding coefficients $(a_0^0, a_1^0, a_2^0, b_0^0, b_1^0, b_2^0, b_3^0, b_4^0)$ given by:

$$\begin{aligned}
b_4^0 &= -2F_{333} \\
b_3^0 &= \frac{1}{12}(-6DF_{333} - 5F_3F_{333}) \\
b_2^0 &= \frac{1}{360}(120DF_{233} + 240F_{133} - 120F_{223} + 60DF_{333}F_3 - 36DF_3F_{333} + 204F_2F_{333} + 79F_3^2F_{333}) \\
b_1^0 &= \frac{1}{1080}(-180DF_{223} - 540F_{123} + 360F_{222} - 90DF_{33}F_{23} + 270F_{23}^2 + 90DF_3F_{233} - 540F_2F_{233} \\
&\quad - 180DF_{233}F_3 - 270F_{133}F_3 + 360F_{223}F_3 - 45DF_{333}F_3^2 - 45F_{233}F_3^2 + 90DF_{23}F_{33} + \\
&\quad 90F_{13}F_{33} - 360F_{22}F_{33} - 90F_{23}F_3F_{33} + 90F_2F_{33}^2 + 18DF_2F_{333} - 72F_1F_{333} + \\
&\quad 54DF_3F_3F_{333} - 288F_2F_3F_{333} - 71F_3^3F_{333} - 180F_{33y})
\end{aligned}$$

$$\begin{aligned}
b_0^0 &= \frac{1}{129600}(-8640DF_{233}DF_3 - 12960DF_{23}DF_{33} - 4320DF_2DF_{333} + 43200DF_{33y} + \\
&17280DF_{333}F_1 + 129600F_{113} - 64800F_{122} - 34560DF_{33}F_{13} - 86400DF_3F_{133} + \\
&12960DF_{233}F_2 + 194400F_{133}F_2 + 30240DF_{33}F_{22} + 32400DF_3F_{223} - 64800F_2F_{223} + \\
&6480DF_{23}F_{23} - 25920F_{13}F_{23} - 15120F_{22}F_{23} + 2160DF_2F_{233} - 8640F_1F_{233} - \\
&64800F_{23y} - 6480DF_{33}^2F_3 - 6480DF_3DF_{333}F_3 - 21600F_{123}F_3 + 10800DF_{333}F_2F_3 - \\
&10800F_{222}F_3 + 25560DF_{33}F_{23}F_3 - 18360F_{23}^2F_3 + 13320DF_3F_{233}F_3 - 25920F_2F_{233}F_3 + \\
&3960DF_{233}F_3^2 + 50400F_{133}F_3^2 - 28800F_{223}F_3^2 + 2820DF_{333}F_3^3 - 10980F_{233}F_3^3 - \\
&18000DF_{22}F_{33} + 6480DF_3DF_{33}F_{33} + 86400F_{12}F_{33} - 28080DF_{33}F_2F_{33} - \\
&11880DF_3F_{23}F_{33} + 10800F_2F_{23}F_{33} - 19080DF_{23}F_3F_{33} + 18720F_{13}F_3F_{33} + \\
&16920F_{22}F_3F_{33} - 8100DF_{33}F_3^2F_{33} + 7200F_{23}F_3^2F_{33} + 7560DF_2F_{33}^2 - 30240F_1F_{33}^2 - \\
&11520F_2F_3F_{33}^2 - 1620F_3^3F_{33}^2 + 11664DF_3^2F_{333} - 63072DF_3F_2F_{333} + 76464F_2^2F_{333} - \\
&2520DF_2F_3F_{333} + 10080F_1F_3F_{333} - 17712DF_3F_3^2F_{333} + 42768F_2F_3^2F_{333} + 5299F_3^4F_{333} - \\
&18000F_3F_{33y} - 75600F_{33}F_{3y} + 43200F_{333}F_y) \\
a_2^0 &= \frac{1}{45}(-18DF_{333} - 24F_{233} - 4F_{33}^2 - 27F_3F_{333}) \\
a_1^0 &= \frac{1}{540}(-72DF_{233} - 432F_{133} + 216F_{223} - 36DF_{333}F_3 + 96F_{233}F_3 + 48DF_{33}F_{33} + 16F_3F_{33}^2 + \\
&108DF_3F_{333} - 324F_2F_{333} - 81F_3^2F_{333}) \\
a_0^0 &= \frac{1}{4050}(-180DF_{223} + 288DF_{33}^2 - 4860F_{123} + 2520F_{222} - 378DF_{33}F_{23} + 1782F_{23}^2 + \\
&810DF_3F_{233} - 2700F_2F_{233} - 180DF_{233}F_3 - 2430F_{133}F_3 + 2880F_{223}F_3 - 45DF_{333}F_3^2 + \\
&435F_{233}F_3^2 + 810DF_{23}F_{33} + 810F_{13}F_{33} - 2520F_{22}F_{33} + 408DF_{33}F_3F_{33} - 594F_{23}F_3F_{33} + \\
&810F_2F_{33}^2 + 122F_3^2F_{33}^2 - 270DF_2F_{333} + 1080F_1F_{333} + 270DF_3F_3F_{333} - \\
&1080F_2F_3F_{333} - 135F_3^3F_{333} + 2700F_{33y}).
\end{aligned}$$

One can use these, relatively simple, formulae to generate expressions for the invariant forms on P . This may be achieved by means of a matrix

$$(3.1) \quad m = \begin{pmatrix} \alpha_1^1 \alpha_4^4 & 0 & 0 & 0 \\ -\frac{\alpha_0^1}{3} & \alpha_1^1 & 0 & 0 \\ \frac{(\alpha_0^1)^2}{9\alpha_1^1 \alpha_4^4} & -\frac{2\alpha_0^1}{3\alpha_4^4} & \frac{\alpha_1^1}{\alpha_4^4} & 0 \\ \frac{-(\alpha_0^1)^3}{27(\alpha_1^1 \alpha_4^4)^2} & \frac{(\alpha_0^1)^2}{3\alpha_1^1 (\alpha_4^4)^2} & -\frac{\alpha_0^1}{(\alpha_4^4)^2} & \frac{\alpha_1^1}{(\alpha_4^4)^2} \end{pmatrix}.$$

Then the expression for the invariant 1-forms $(\theta^i) = (\theta^0, \theta^1, \theta^2, \theta^3)$ can be written as

$$(3.2) \quad \theta^i = m_j^i \theta_0^j, \quad i, j = 0, 1, 2, 3.$$

The residual group $G = \{m \mid \alpha_1^1, \alpha_4^4 \neq 0, \alpha_0^1 \in \mathbb{R}\}$ has the Lie algebra $\mathfrak{g} = \mathfrak{h}_2 \oplus \mathfrak{h}_1$ isomorphic to the direct sum of the 2-dimensional noncommuting Lie algebra \mathfrak{h}_2 and a 1-dimensional Lie algebra \mathfrak{h}_1 . Algebra \mathfrak{h}_2 is related to the parameters (α_0^1, α_4^4) and algebra \mathfrak{h}_1 is associated with α_1^1 .

The action of G on θ_0^i , induces its action on $(\Omega_+^0, \Omega_0^0, \Omega_-^0, \Omega^0)$. Indeed, defining

$$\overset{\circ}{\Gamma} = \Omega_-^0 E_- + \Omega_+^0 E_+ + \Omega_0^0 E_0 + \Omega^0 E,$$

and

$$\Gamma = \Omega_- E_- + \Omega_+ E_+ + \Omega_0 E_0 + \Omega E,$$

where 4×4 matrices (E_-, E_+, E_0, E) are the generators of the Lie algebra $\mathfrak{gl}(2, \mathbb{R})$ given in (1.1), we find that

$$\Gamma = m \overset{\circ}{\Gamma} m^{-1} + m d m^{-1}.$$

This enables us to find the explicit expressions for the invariant forms $(\Omega_+, \Omega_0, \Omega_-, \Omega)$.

The transformation rule for Γ resembles transformation rule for a connection. Since Γ is $\mathfrak{gl}(2, \mathbb{R})$ -valued, it is reasonable to look for a $\mathbf{GL}(2, \mathbb{R})$ principal fibre bundle associated with corresponding ODE (2.2).

Due to properties of system (2.4)-(2.5) the desired bundle is just P of Theorem 2.1. To see this, note that equations (2.4) ensure that $(\theta^1, \theta^2, \theta^3, \theta^4)$ form a closed differential ideal. Thus a 4-dimensional distribution \mathcal{V} on P such that $\mathcal{V} \lrcorner \theta^i = 0$, $\forall i = 0, 1, 2, 3$, is integrable. As a consequence, the manifold P is foliated by 4-dimensional integral leaves of this distribution. Looking at equations (2.5) we see that *on each leaf* of \mathcal{V} the forms $(\Omega_+, \Omega_0, \Omega_-, \Omega)$ satisfy the Maurer-Carathéodory equations for the $\mathbf{GL}(2, \mathbb{R})$ group. This means that P is a principal $\mathbf{GL}(2, \mathbb{R})$ bundle over the leaf space $M^4 = P/\mathcal{V}$. This 4-dimensional space may be identified with a solutions space of ODE (2.2).

Remark 3.1. For local calculations, it may be convenient to pass from coordinates $(x, y, y_1, y_2, y_3, \alpha_0^1, \alpha_1^1, \alpha_4^4)$ on P to coordinates $(c_0, c_1, c_2, c_3, s, \alpha_0^1, \alpha_1^1, \alpha_4^4)$ on P , where (c_0, c_1, c_2, c_3) are the integration constants of ODE (2.2), and s is a real parameter such that the total differential vector field $D = \partial_s$. In such parametrisation $(s, \alpha_0^1, \alpha_1^1, \alpha_4^4)$ constitute coordinates on the leaves of \mathcal{V} and (c_0, c_1, c_2, c_3) parametrise the solution space M^4 .

4. $\mathbf{GL}(2, \mathbb{R})$ GEOMETRY ON THE SOLUTION SPACE

Using matrices $\Gamma = (\Gamma_j^i)$, $i, j = 0, 1, 2, 3$, and part $\theta^i = (\theta^0, \theta^1, \theta^2, \theta^3)$ of the invariant coframe we rewrite equations (2.4) in a compact form as:

$$(4.1) \quad d\theta^i + \Gamma_j^i \wedge \theta^j = 0,$$

and equation (2.5) in a compact form as:

$$(4.2) \quad d\Gamma_k^i + \Gamma_j^i \wedge \Gamma_k^j = \frac{1}{2} R_{jkl}^i \theta^j \wedge \theta^l.$$

The coefficients R_{jkl}^i appearing in this last equation can be easily read off from (2.5). They are linear combinations of the coefficients $a_0, a_1, a_2, a_3, b_0, b_1, b_2, b_3, b_4$ of (2.5). The meaning of equations (4.1)-(4.2) is obvious: they constitute, respectively, the first and the second Cartan's structure equations, for a $\mathfrak{gl}(2, \mathbb{R})$ -valued connection Γ on the principal fibre bundle $\mathbf{GL}(2, \mathbb{R}) \rightarrow P \rightarrow M^4$. Due to the first equation, (4.1), this connection has no torsion. The second equation, (4.2), determines the curvature of Γ ; the coefficients R_{jkl}^i are the curvature tensor coefficients for Γ .

Given the curvature tensor R_{jkl}^i of Γ we define its 'Ricci' tensor R_{jl} by

$$R_{jl} = R_{jil}^i.$$

Recalling that the curvature of Γ is totally expressible in terms of $a_0, a_1, a_2, a_3, b_0, b_1, b_2, b_3, b_4$ and performing a purely algebraic manipulation on the curvature tensor coefficients R^i_{jkl} we get a remarkable

Theorem 4.1. *Every 4th order ODE satisfying conditions (2.3) uniquely defines a principal fibre bundle $\mathbf{GL}(2, \mathbb{R}) \rightarrow P \rightarrow M^4$ over the space of its solutions M^4 and a torsionless $\mathfrak{gl}(2, \mathbb{R})$ -connection Γ on P with curvature R^i_{jkl} having the Ricci tensor R_{jl} in the form*

$$R_{jl} = \begin{pmatrix} 0 & b_0 & a_0 + 2b_1 & -a_1 + b_2 \\ -b_0 & -2a_0 & a_1 + 3b_2 & a_2 + 2b_3 \\ a_0 - 2b_1 & a_1 - 3b_2 & -2a_2 & b_4 \\ -a_1 - b_2 & a_2 - 2b_3 & -b_4 & 0 \end{pmatrix}.$$

Its respective symmetric and antisymmetric parts read:

$$R_{(jl)} = \begin{pmatrix} 0 & 0 & a_0 & -a_1 \\ 0 & -2a_0 & a_1 & a_2 \\ a_0 & a_1 & -2a_2 & 0 \\ -a_1 & a_2 & 0 & 0 \end{pmatrix},$$

and

$$R_{[jl]} = \begin{pmatrix} 0 & b_0 & 2b_1 & b_2 \\ -b_0 & 0 & 3b_2 & 2b_3 \\ -2b_1 & -3b_2 & 0 & b_4 \\ -b_2 & -2b_3 & -b_4 & 0 \end{pmatrix}.$$

Thus the entire curvature tensor R^i_{jkl} is encoded in the Ricci tensor.

Remark 4.2. Note that we also have $R^i_{ikl} = 2R_{[kl]}$.

Now we can use matrix m of the previous section to find explicit formulae for the coefficients $a_0, a_1, a_2, a_3, b_0, b_1, b_2, b_3, b_4$. It follows that if we evaluate R_{ij} for $(\alpha^1_0, \alpha^1_1, \alpha^4_4) = (0, 1, 1)$, denoting the calculated R_{ij} by R^0_{ij} , then the full Ricci tensor R_{ij} is related to R^0_{ij} via

$$(4.3) \quad R_{ij} = R^0_{kl} m^{-1k}{}_i m^{-1l}{}_j.$$

Here $m^{-1} = (m^{-1j}{}_j)$ is the inverse matrix to m . From this expression we can calculate explicit form of $a_0, a_1, a_2, a_3, b_0, b_1, b_2, b_3, b_4$. The resulting formulae involve coefficients $a^0_0, a^0_1, a^0_2, a^0_3, b^0_0, b^0_1, b^0_2, b^0_3, b^0_4$ of the previous section and parameters $\alpha^1_0, \alpha^1_1, \alpha^4_4$ and read:

$$\begin{aligned} b_4 &= \frac{(\alpha^4_4)^3}{(\alpha^1_1)^2} b^0_4 \\ b_3 &= \frac{(\alpha^4_4)^2}{(\alpha^1_1)^2} b^0_3 + \frac{\alpha^1_0 (\alpha^4_4)^2}{3(\alpha^1_1)^3} b^0_4 \\ b_2 &= \frac{\alpha^4_4}{(\alpha^1_1)^2} b^0_2 + \frac{2\alpha^1_0 \alpha^4_4}{3(\alpha^1_1)^3} b^0_3 + \frac{(\alpha^1_0)^2 \alpha^4_4}{9(\alpha^1_1)^4} b^0_4 \\ b_1 &= \frac{1}{(\alpha^1_1)^2} b^0_1 + \frac{\alpha^1_0}{(\alpha^1_1)^3} b^0_2 + \frac{(\alpha^1_0)^2}{(3\alpha^1_1)^4} b^0_3 + \frac{(\alpha^1_0)^3}{27(\alpha^1_1)^5} b^0_4 \\ b_0 &= \frac{1}{(\alpha^1_1)^2 \alpha^4_4} b^0_0 + \frac{4\alpha^1_0}{3(\alpha^1_1)^3 \alpha^4_4} b^0_1 + \frac{2(\alpha^1_0)^2}{3(\alpha^1_1)^4 \alpha^4_4} b^0_2 + \frac{4(\alpha^1_0)^3}{27(\alpha^1_1)^5 \alpha^4_4} b^0_3 + \frac{(\alpha^1_0)^4}{81(\alpha^1_1)^6 \alpha^4_4} b^0_4, \end{aligned}$$

$$(4.4) \quad \begin{aligned} a_2 &= \frac{(\alpha_4^4)^2}{(\alpha_1^1)^2} a_2^0 \\ a_1 &= \frac{\alpha_4^4}{(\alpha_1^1)^2} a_1^0 - \frac{\alpha_0^1 \alpha_4^4}{3(\alpha_1^1)^3} a_2^0 \\ a_0 &= \frac{1}{(\alpha_1^1)^2} a_0^0 - \frac{2\alpha_0^1}{3(\alpha_1^1)^3} a_1^0 + \frac{(\alpha_0^1)^2}{9(\alpha_1^1)^4} a_2^0. \end{aligned}$$

This, in particular, means that the respective spaces consisting of $(b_0^0, b_1^0, b_2^0, b_3^0, b_4^0)$ and of (a_0^0, a_1^0, a_2^0) constitute a 5-dimensional and 3-dimensional representation of G and, as a consequence of $\mathbf{GL}(2, \mathbb{R})$.

Due to (4.3) the vanishing of any of the two determinants:

$$\det(R_{(ij)}) \quad \text{and} \quad \det(R_{[ij]})$$

is a contact invariant property of the corresponding 4th order ODE (1.2). These two determinants, when expressed in terms of the eight curvature coefficients $a_0, a_1, a_2, b_0, b_1, b_2, b_3, b_4$, are

$$\det(R_{(ij)}) = (a_1^2 - a_0 a_2)^2$$

and

$$\det(R_{[ij]}) = (3b_2^2 - 4b_1 b_3 + b_0 b_4)^2.$$

Thus they are expressible in terms of the two well known $\mathbf{GL}(2, \mathbb{R})$ -invariant polynomials

$$I_2 = a_1^2 - a_0 a_2 \quad \text{and} \quad I_3 = 3b_2^2 - 4b_1 b_3 + b_0 b_4.$$

Remark 4.3. In this context it is interesting to note that function $F = (y_3)^{(4/3)}$ of the well known example (1.8), provides a contact equivalent class of ODEs that has both invariants I_2 and I_3 vanishing.

Interestingly, the next $\mathbf{GL}(2, \mathbb{R})$ -invariant polynomial

$$I_4 = -3(\theta^1)^2(\theta^2)^2 + 4\theta^0(\theta^2)^3 + 4(\theta^1)^3\theta^3 - 6\theta^0\theta^1\theta^2\theta^3 + (\theta^0)^2(\theta^3)^2,$$

when thought as defined on P in terms of forms $(\theta^0, \theta^1, \theta^2, \theta^3)$ of the invariant coframe $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+, \Omega_-, \Omega_0, \Omega)$, has the following property:

$$\mathcal{L}_X I_4 = 12(X \lrcorner \Omega) I_4,$$

where $X \in \mathcal{V}$ is any vertical vector field on $\mathbf{GL}(2, \mathbb{R}) \rightarrow P \rightarrow M^4$. Thus I_4 descends to a well defined conformal symmetric tensor of fourth degree on the solution space M^4 of the ODE [1]. Let us denote the descended to M^4 tensor I_4 by Υ . It is also worthwhile to mention that, for the vertical vectors $X \in \mathcal{V}$, we have

$$\mathcal{L}_X \Omega = d(X \lrcorner \Omega).$$

This means that on the solution space M^4 the form Ω is defined up to a *gradient*. It is convenient to rescale Ω and to define a 1-form A on P equal to

$$A = -12\Omega.$$

This form is also defined up to a gradient on the solutions space M^4 . Thus, a solution space M^4 of any 4th order ODE satisfying (2.3) is equipped with a sort of Weyl geometry $[\Upsilon, A]$. This consists of class of pairs (Υ, A) , in which Υ is a 4th order symmetric tensor field, A is a 1-form on M^4 , and two pairs (Υ, A) and (Υ', A') represent the same class iff

$$\Upsilon' = e^{4\phi} \Upsilon, \quad A' = A - 4d\phi.$$

In the context of this gauge freedom, it is worthwhile to note that the vanishing of $R_{[ij]}$ corresponds to the $[\Upsilon, A]$ geometries on M^4 with form A that can be gauged to $A = 0$. Such situation occurs if and only if $b_i = 0$ for all $i = 0, 1, 2, 3, 4$.

Remark 4.4. In terms of the Weyl-like geometry $[\Upsilon, A]$ on the solution space M^4 , the $\mathfrak{gl}(2, \mathbb{R})$ -valued connection may be defined as the unique torsionless connection satisfying

$$\nabla_X \Upsilon = -A(X)\Upsilon.$$

Thus we have the following

Theorem 4.5. *Every 4th order ODE $y^{(4)} = F(x, y, y', y'', y^{(3)})$ satisfying Braynt's conditions (2.3) uniquely defines a conformal Weyl-like geometry $[\Upsilon, A]$ on its solution space M^4 . The Weyl-like geometry $[\Upsilon, A]$ consists of a symmetric 4th rank tensor Υ and a 1-form A given up to transformations*

$$\Upsilon' = e^{4\phi}\Upsilon, \quad A' = A - 4d\phi.$$

Its corresponding $\mathfrak{gl}(2, \mathbb{R})$ -valued connection has no torsion and very special curvature tensor described by Theorem 4.1.

5. EXAMPLES

5.1. Equations with symmetric Ricci tensor. There is only one contact equivalence class of ODEs (2.2) having an 8-dimensional group of contact symmetries. This is equivalent to $y^{(4)} = 0$ and the symmetry group is $\mathbf{GL}(2, \mathbb{R}) \times_\rho \mathbb{R}^4$. For this class of equation the $\mathfrak{gl}(2, \mathbb{R})$ -valued connection of Theorem 4.1 is flat.

In this section we focus on the equivalence classes of ODEs (2.2) for which the Maxwell form $dA = -12d\Omega$ of this connection is flat $dF = 0$. In such case we have $b_0 = b_1 = b_2 = b_3 = b_4 = 0$.

Let us assume that we are in this situation.

Looking at the transformation properties (4.4) of the curvature coefficient a_2 we see that there are essentially two distinct cases distinguished by the *vanishing or not* of the expression $a_2^0 = \frac{1}{45}(-18DF_{333} - 24F_{233} - 4F_{33}^2 - 27F_3F_{333})$.

We analyse the more easy case $a_2^0 = 0$ first.

If $a_2^0 = 0$ then also $a_2 = 0$. Thus we have $a_2 = 0$ everywhere on P with the full system (2.4)-(2.5) of eight independent 1-forms $\theta^1, \theta^2, \theta^3, \Omega_+, \Omega_0, \Omega_-, \Omega$ there. Imposing $(d^2\Omega_+) \wedge \theta^1 \wedge \theta^2 = 0$ on (2.4)-(2.5) quickly leads to $a_1 = 0$ and, consequently, by imposition of $(d^2\Omega_+) \wedge \theta^1 = 0$, to $a_0 = 0$. This shows that if $a_2^0 = 0$ then the corresponding ODEs (2.2) are contact equivalent to $y^{(4)} = 0$.

Now we assume that $a_2^0 \neq 0$. Then the choice

$$\alpha_1^1 = \frac{\sqrt{2}}{4}\alpha_4^4\sqrt{|a_2^0|}$$

brings a_2 to the form

$$a_2 = 8\epsilon_1,$$

where $\epsilon_1 = \text{sgn}(a_2^0)$. Then the choice

$$\alpha_1^0 = \frac{3\sqrt{2}}{4}\epsilon_1\alpha_4^4\frac{a_1^0}{\sqrt{|a_2^0|}}$$

makes

$$a_1 = 0.$$

After these two normalisations we get

$$a_0 = \frac{8\epsilon_1}{(\alpha_4^4 a_2^0)^2} (a_0^0 a_2^0 - (a_1^0)^2).$$

Thus again we have two cases, depending on the vanishing or not of the invariant $I_2^0 = (a_1^0)^2 - a_0^0 a_2^0$.

It follows that the $I_2^0 = 0$ case, which under our assumptions is the same as $a_0 = 0$, corresponds to only *one* nonequivalent class of equations. They are defined by $\epsilon_1 = 1$ (the $\epsilon_1 = -1$ case is not compatible with system (2.4)-(2.5)), and are described by the following

Theorem 5.1. *All ODEs $y^{(4)} = F(x, y, y', y'', y^{(3)})$ satisfying Bryant's conditions (2.3), having symmetric Ricci tensor, and invariants $I_2 = 0$ and $a_2 \neq 0$, are in local one-to-one correspondence with coframes $(\theta^0, \theta^1, \theta^2, \theta^3, \Omega_+, \Omega)$ on a 6-manifold satisfying:*

$$\begin{aligned} d\theta^0 &= 12\Omega \wedge \theta^0 - 3\Omega_+ \wedge \theta^1 + \frac{3\sqrt{2}}{2}\theta^0 \wedge \theta^2 \\ d\theta^1 &= 6\Omega \wedge \theta^1 - 2\Omega_+ \wedge \theta^2 + \frac{\sqrt{2}}{2}(\theta^0 \wedge \theta^3 + \theta^1 \wedge \theta^2) \\ d\theta^2 &= -\Omega_+ \wedge \theta^3 + \sqrt{2}\theta^1 \wedge \theta^3 \\ d\theta^3 &= -6\Omega \wedge \theta^3 + 3\sqrt{2}\theta^2 \wedge \theta^3 \\ d\Omega_+ &= 6\Omega \wedge \Omega_+ + \sqrt{2}\Omega_+ \wedge \theta^2 + \theta^0 \wedge \theta^3 - 5\theta^1 \wedge \theta^2 \\ d\Omega &= 0. \end{aligned}$$

The forms Ω and Ω_0 are given by

$$\Omega_- = \frac{\sqrt{2}}{2}\theta^3, \quad \Omega_0 = 3\Omega - \frac{\sqrt{2}}{2}\theta^2.$$

All the equations having such invariant forms are equivalent to an ODE defined by

$$F = \frac{4}{3} \frac{y_3^2}{y_2}.$$

This class has strictly 6-dimensional group of contact symmetries.

Now we pass to the $I_2^0 \neq 0$ case. We introduce $\epsilon_2 = \pm 1$, which encodes the sign of I_2^0 . This is defined by $\epsilon_1 \epsilon_2 (a_0^0 a_2^0 - (a_1^0)^2) > 0$. Now we chose

$$\alpha_4^4 = \sqrt{\frac{\epsilon_1 \epsilon_2 (a_0^0 a_2^0 - (a_1^0)^2)}{(a_2^0)^2}}.$$

This normalises a_0 to

$$a_0 = 8\epsilon_2.$$

Under such normalisations system (2.4)-(2.5) descends from P to the 5-dimensional jet space J . There, it reads:

$$\begin{aligned}
d\theta^0 &= 3(\Omega + \Omega_0) \wedge \theta^0 - 3\Omega_+ \wedge \theta^1 \\
d\theta^1 &= -\Omega_- \wedge \theta^0 + (3\Omega + \Omega_0) \wedge \theta^1 - 2\Omega_+ \wedge \theta^2 \\
d\theta^2 &= -2\Omega_- \wedge \theta^1 + (3\Omega - \Omega_0) \wedge \theta^2 - \Omega_+ \wedge \theta^3 \\
d\theta^3 &= -3\Omega_- \wedge \theta^2 + 3(\Omega - \Omega_0) \wedge \theta^3 \\
\\
d\Omega_+ &= 2\Omega_0 \wedge \Omega_+ - 2\epsilon_2\theta^0 \wedge \theta^1 + \epsilon_1(\theta^0 \wedge \theta^3 - 5\theta^1 \wedge \theta^2) \\
d\Omega_- &= -2\Omega_0 \wedge \Omega_- + \epsilon_2(-\theta^0 \wedge \theta^3 + 5\theta^1 \wedge \theta^2) + 2\epsilon_1\theta^2 \wedge \theta^3 \\
d\Omega_0 &= \Omega_+ \wedge \Omega_- - \epsilon_2\theta^0 \wedge \theta^2 - \epsilon_1\theta^1 \wedge \theta^3 \\
d\Omega &= 0
\end{aligned}$$

To close this system it is convenient to eliminate form Ω . This can be achieved by an introduction of new forms $(\sigma^0, \sigma^1, \sigma^2, \sigma^3)$ related to $(\theta^0, \theta^1, \theta^2, \theta^3)$ via:

$$\sigma^0 = e^w\theta^0, \quad \sigma^1 = e^w\theta^1, \quad \sigma^2 = e^w\theta^2, \quad \sigma^3 = e^w\theta^3,$$

where w is a function on J such that $\Omega = -\frac{1}{3}dw$. The local existence of such function is guaranteed by $d\Omega = 0$. In terms of the new variables $(\sigma^0, \sigma^1, \sigma^2, \sigma^3)$, w , the reduced system takes a form in which the 1-form Ω is not present:

$$\begin{aligned}
d\sigma^0 &= 3\Omega_0 \wedge \sigma^0 - 3\Omega_+ \wedge \sigma^1 \\
d\sigma^1 &= -\Omega_- \wedge \sigma^0 + \Omega_0 \wedge \sigma^1 - 2\Omega_+ \wedge \sigma^2 \\
d\sigma^2 &= -2\Omega_- \wedge \sigma^1 - \Omega_0 \wedge \sigma^2 - \Omega_+ \wedge \sigma^3 \\
d\sigma^3 &= -3\Omega_- \wedge \sigma^2 - 3\Omega_0 \wedge \sigma^3 \\
(5.1) \quad d\Omega_+ &= 2\Omega_0 \wedge \Omega_+ + e^{-2w} \left(-2\epsilon_2\sigma^0 \wedge \sigma^1 + \epsilon_1(\sigma^0 \wedge \sigma^3 - 5\sigma^1 \wedge \sigma^2) \right) \\
d\Omega_- &= -2\Omega_0 \wedge \Omega_- + e^{-2w} \left(\epsilon_2(-\sigma^0 \wedge \sigma^3 + 5\sigma^1 \wedge \sigma^2) + 2\epsilon_1\sigma^2 \wedge \sigma^3 \right) \\
d\Omega_0 &= \Omega_+ \wedge \Omega_- - e^{-2w} \left(\epsilon_2\sigma^0 \wedge \sigma^2 + \epsilon_1\sigma^1 \wedge \sigma^3 \right).
\end{aligned}$$

As we can see the price paid for elimination of Ω is an introduction of nonconstant function w appearing explicitly in these equations.

Now the remarkable fact is that system (5.1) closes on J and is described by the following Theorem.

Theorem 5.2. *All ODEs $y^{(4)} = F(x, y, y', y'', y^{(3)})$ satisfying Bryant's conditions (2.3), having symmetric Ricci tensor, and invariants $I_2 \neq 0$ and $a_2 \neq 0$, are in local one-to-one correspondence with coframes $(\sigma^0, \sigma^1, \sigma^2, \sigma^3, \Omega_+)$ on a 5-manifold satisfying system (5.1) with:*

$$\begin{aligned}
\Omega_0 &= w_0\sigma^0 - (w_1 + 4\epsilon_1\epsilon_2w_3)\sigma^1 + (4\epsilon_1\epsilon_2w_0 + w_2)\sigma^2 - w_3\sigma^3 \\
(5.2) \quad \Omega_- &= -\epsilon_1\epsilon_2\Omega_+ - 2(\epsilon_1\epsilon_2w_1 + 2w_3)\sigma^0 + 2w_0\sigma_1 + \\
&\quad 2\epsilon_1\epsilon_2w_3\sigma^2 - 2(2\epsilon_1\epsilon_2w_0 + w_2)\sigma^3.
\end{aligned}$$

Functions w, w_0, w_1, w_2, w_3 appearing here are defined by:

$$(5.3) \quad dw = w_0\sigma^0 + w_1\sigma^1 + w_2\sigma^2 + w_3\sigma^3.$$

They satisfy

$$\begin{aligned}
dw_0 &= -\epsilon_1 \epsilon_2 w_1 \Omega_+ + \frac{1}{4}(-\epsilon_2 e^{-2w} + 4w_0^2 + 16w_1 w_3 + 32\epsilon_1 \epsilon_2 w_3^2) \sigma^0 + \\
&\quad 3w_0 w_1 \sigma^1 - (-\epsilon_1 \epsilon_2 w_{13} - 11w_0 w_2 - 4\epsilon_1 \epsilon_2 w_2^2 + 5\epsilon_1 \epsilon_2 w_1 w_3 + 12w_3^2) \sigma^2 + \\
&\quad (11w_0 + 4\epsilon_1 \epsilon_2 w_2) w_3 \sigma^3 \\
dw_1 &= (3w_0 - 2\epsilon_1 \epsilon_2 w_2) \Omega_+ + (-3w_0 w_1 - 4\epsilon_1 \epsilon_2 w_1 w_2 - 12\epsilon_1 \epsilon_2 w_0 w_3 - 8w_2 w_3) \sigma^0 - \\
&\quad \frac{1}{4}(3\epsilon_1 e^{-2w} + 24\epsilon_1 \epsilon_2 w_0^2 - 20w_1^2 + 8\epsilon_1 \epsilon_2 w_{13} + 64w_0 w_2 + 32\epsilon_1 \epsilon_2 w_2^2 - \\
&\quad 120\epsilon_1 \epsilon_2 w_1 w_3 - 192w_3^2) \sigma^1 - \\
(5.4) \quad &\quad (12w_0 w_1 \epsilon_1 \epsilon_2 + w_1 w_2 + 30w_0 w_3 + 4\epsilon_1 \epsilon_2 w_2 w_3) \sigma^2 + w_{13} \sigma^3 \\
dw_2 &= (2w_1 - 3\epsilon_1 \epsilon_2 w_3) \Omega_+ + \\
&\quad \frac{1}{2}(24\epsilon_1 \epsilon_2 w_0^2 + 2\epsilon_1 \epsilon_2 w_{13} + 30w_0 w_2 + 8\epsilon_1 \epsilon_2 w_2^2 - 26\epsilon_1 \epsilon_2 w_1 w_3 - 48w_3^2) \sigma^0 - \\
&\quad (8\epsilon_1 \epsilon_2 w_0 w_1 + w_1 w_2 + 24w_0 w_3 + 12\epsilon_1 \epsilon_2 w_2 w_3) \sigma^1 + \\
&\quad \frac{1}{4}(-3\epsilon_2 e^{-2w} + 96w_0^2 - 8w_{13} - 12w_2^2 + 40w_1 w_3 + 96\epsilon_1 \epsilon_2 w_3^2) \sigma^2 - \\
&\quad 3(8\epsilon_1 \epsilon_2 w_0 + 3w_2) w_3 \sigma^3 \\
dw_3 &= w_2 \Omega_+ + (4\epsilon_1 \epsilon_2 w_0 w_1 + 2w_1 w_2 + 11w_0 w_3 + 4\epsilon_1 \epsilon_2 w_2 w_3) \sigma^0 + \\
&\quad (w_{13} + 8\epsilon_1 \epsilon_2 w_0 w_2 + 4w_2^2 - 4w_1 w_3 - 12\epsilon_1 \epsilon_2 w_3^2) \sigma^1 + w_2 w_3 \sigma^2 + \\
&\quad \frac{1}{4}(-\epsilon_1 e^{-2w} + 32\epsilon_1 \epsilon_2 w_0^2 + 32w_0 w_2 + 8\epsilon_1 \epsilon_2 w_2^2 + 4w_3^2) \sigma^3,
\end{aligned}$$

with a function w_{13} satisfying

$$\begin{aligned}
dw_{13} &= (-12\epsilon_1 \epsilon_2 w_0 w_1 - w_1 w_2 + 45w_0 w_3 + 30\epsilon_1 \epsilon_2 w_2 w_3) \Omega_+ + \\
&\quad \frac{1}{2}(-6\epsilon_2 w_0 e^{-2w} - 240w_0^3 + 40\epsilon_1 \epsilon_2 w_0 w_1^2 - 16w_0 w_{13} + 5\epsilon_1 w_2 e^{-2w} - \\
&\quad 472\epsilon_1 \epsilon_2 w_0^2 w_2 + 20w_1^2 w_2 - 16\epsilon_1 \epsilon_2 w_{13} w_2 - 304w_0 w_2^2 - 64\epsilon_1 \epsilon_2 w_3^3 + \\
&\quad 384w_0 w_1 w_3 + 192\epsilon_1 \epsilon_2 w_1 w_2 w_3 + 552\epsilon_1 \epsilon_2 w_0 w_3^2 + 272w_2 w_3^2) \sigma^0 - \\
&\quad \frac{1}{4}(20\epsilon_2 w_1 e^{-2w} - 256w_0^2 w_1 - 28w_1 w_{13} - 416\epsilon_1 \epsilon_2 w_0 w_1 w_2 - 144w_1 w_2^2 + \\
(5.5) \quad &\quad 15\epsilon_1 w_3 e^{-2w} - 840\epsilon_1 \epsilon_2 w_0^2 w_3 + 20w_1^2 w_3 - 24\epsilon_1 \epsilon_2 w_{13} w_3 - 1440w_0 w_2 w_3 - \\
&\quad 480\epsilon_1 \epsilon_2 w_2^2 w_3 - 40\epsilon_1 \epsilon_2 w_1 w_3^2 - 192w_3^3) \sigma^1 - \\
&\quad \frac{1}{2}(-15\epsilon_1 w_0 e^{-2w} + 480\epsilon_1 \epsilon_2 w_0^3 - 24\epsilon_1 \epsilon_2 w_0 w_{13} - 2\epsilon_2 w_2 e^{-2w} + 544w_0^2 w_2 - \\
&\quad 16w_{13} w_2 + 184\epsilon_1 \epsilon_2 w_0 w_2^2 + 16w_3^3 + 240\epsilon_1 \epsilon_2 w_0 w_1 w_3 + 80w_1 w_2 w_3 + \\
&\quad 588w_0 w_3^2 + 184\epsilon_1 \epsilon_2 w_2 w_3^2) \sigma^2 - \\
&\quad \frac{1}{4}(5\epsilon_1 w_1 e^{-2w} - 160\epsilon_1 \epsilon_2 w_0^2 w_1 - 160w_0 w_1 w_2 - 40\epsilon_1 \epsilon_2 w_1 w_2^2 + \\
&\quad 36\epsilon_2 w_3 e^{-2w} - 1152w_0^2 w_3 - 28w_{13} w_3 - 1152\epsilon_1 \epsilon_2 w_0 w_2 w_3 - \\
&\quad 288w_2^2 w_3 + 20w_1 w_3^2) \sigma^3.
\end{aligned}$$

System (5.1)-(5.5) is closed, meaning that $d^2 = 0$ does not implies any further relations between forms $\sigma^0, \sigma^1, \sigma^2, \sigma^3, \Omega_+$ and functions $w, w_0, w_1, w_2, w_3, w_{13}$.

We easily see that the assumption that *all* $w, w_0, w_1, w_2, w_3, w_{13}$ are *constant* is incompatible with system (5.1)-(5.5). Finding *any* solution to system (5.1)-(5.5) is a difficult task.

5.2. Inhomogeneous examples. Here we present examples of contact equivalent classes of 4th order ODEs satisfying Bryant's conditions (2.3) which are *not homogeneous*. By this we mean they do *not* admit a transitive contact symmetry group of dimension greater than *four*. We consider an ansatz in which function F depends in a special way on only two coordinates y_2 and y_3 . Explicitly:

$$(5.6) \quad F = (y_2)^2 q\left(\frac{y_3^2}{y_2}\right),$$

where $q = q(z)$ is a sufficiently differentiable real function of its argument

$$z = \frac{y_3^2}{y_2}.$$

Imposing Bryant's conditions (2.3) on (5.6) we find the following

Proposition 5.3. *Function F of (5.6) satisfies Bryant's conditions (2.3) if and only if*

a) *either:*

$$6z(3z - 2q)q'' + 3zq'^2 - 6qq' + 4q = 0,$$

b) *or:*

$$6z(3z - 2q)q'' + 3zq'^2 - 6qq' + 14q - 15z = 0.$$

The special solutions of a) are: $q(z) = 0$ and $q(z) = \frac{4}{3}z$. In case b) we have $q(z) = 3z$ and $q(z) = \frac{5}{3}z$ as special solutions. Writing these four solutions as $q(z) = cz$ we remark that in cases $c = 0$ and $c = 3$ function F defines a 4th order ODE which is contact equivalent to $y^{(4)} = 0$. Cases $c = \frac{4}{3}$ and $c = \frac{5}{3}$ define two different F s, but the corresponding 4th order ODEs are contact equivalent. They both are equivalent to the ODE described by Theorem 5.1.

We emphasise that apart from the singular solutions $q = cz$, each equation a) or b) admits a 2-parameter family of solutions. Every solution $q = q(z)$ from these two families leads to a 4th order ODE which satisfies Bryant's conditions (2.3) and which is *inhomogeneous*. Remarkably all Bryant's F s which are defined by the ansatz (5.6) have $I_3 = I_4 = 0$, but $a_2 \neq 0$ and $b_4 \neq 0$. Thus, in particular, $dA \neq 0$ for them.

We were unable to find any example of Bryant's ODEs for which at least one of I_2 or I_3 is not vanishing.

REFERENCES

- [1] Bryant R L (1991) "Two exotic holonomies in dimension four, path geometries, and twistor theory" *Proc. Symp. Pure Math.* Vol. **53**, 33-88
- [2] Cartan E (1924) "Varietes a connexion projective" *Bull. Soc. Math.* **LII** 205-241
- [3] Cartan E (1941) "La geometria de las ecuaciones diferenciales de tercer orden" *Rev. Mat. Hispano-Aamer.* **4** 1-31
- [4] Chern S S (1940) "The geometry of the differential equations $y''' = F(x, y, y', y'')$ " *Sci. Rep. Nat. Tsing Hua Univ.* **4** 97-111
- [5] Doubrov B (2007) "Generalized Wilczynski invariants for non-linear ordinary differential equations", [arXiv:math/0702251](https://arxiv.org/abs/math/0702251)

- [6] Dunajski M, Tod P (2006) "Paraconformal geometry of n th order ODEs, and exotic holonomy in dimension four", *J. Geom. Phys.* **56**, 1790-1809
- [7] Godlinski M, Nurowski P (2006) Third-order ODEs and four-dimensional split signature Einstein metrics *Journ. Geom. Phys.* **56** 344-357
- [8] Godlinski M, Nurowski P (2007) $\mathbf{GL}(2, \mathbb{R})$ geometry of ODEs [arXiv:math/07100297](https://arxiv.org/abs/math/07100297)
- [9] Lie, S. (1924) "Klassifikation und Integration von gewöhnlichen Differentialgleichungen zwischen x, y , die eine Gruppe von Transformationen gestatten III", in *Gesammelte Abhandlungen*, Vol. 5, Teubner, Leipzig
- [10] Nurowski P (2005) Differential equations and conformal structures *Journ. Geom. Phys.* **55** 19-49
- [11] Nurowski P, Sparling G (2003) 3-dimensional CR structures and 2nd order ordinary differential equations *Class. Q. Grav.* **20** 4995-5016
- [12] Tresse M A, (1896) *Determinations des invariants ponctuels de l'equation differentielle ordinaire du second ordre $y'' = \omega(x, y, y')$* , Hirzel, Leipzig.

INSTYTUT FIZYKI TEORETYCZNEJ, UNIwersYTET WARSZAWSKI, UL. HOZA 69, WARSZAWA,
POLAND

E-mail address: nurowski@fuw.edu.pl