

A la Fock-Goncharov coordinates for $\mathrm{PU}(2,1)$

Julien Marché, Pierre Will
 Institut de Mathématiques
 Université Pierre et Marie Curie
 4, place Jussieu
 F-75252 Paris Cedex 05
 email : `marche,will@math.jussieu.fr`

November 7, 2018

Abstract

In this work, we describe a set of coordinates on the $\mathrm{PU}(2,1)$ -representation variety of the fundamental group of an oriented punctured surface Σ with negative Euler characteristic. The main technical tool we use is a set of geometric invariants of a triple of flags in the complex hyperbolic plane $\mathbf{H}_{\mathbb{C}}^2$. We establish a bijection between a set of decorations of an ideal triangulation of Σ and a subset of the $\mathrm{PU}(2,1)$ -representation variety of $\pi_1(\Sigma)$.

1 Introduction

In their work [4], Fock and Goncharov have described a coordinate system on the representation variety of the fundamental group of a punctured surface Σ in a split semi-simple real Lie group. The typical example is $\mathrm{PSL}(n, \mathbb{R})$. When n equals 2 (resp. 3), they identify the Teichmüller space of Σ (resp. the moduli space of convex projective structures on Σ) within the representation variety. The main goal of our work is to describe an analogous coordinate system for representations in $\mathrm{PU}(2,1)$, which is the group of holomorphic isometries of the complex hyperbolic plane $\mathbf{H}_{\mathbb{C}}^2$. Note that $\mathrm{PU}(2,1)$ is not split and thus does not belong to the family studied by Fock and Goncharov. The preprint [3] in which the cases of $\mathrm{PSL}(2, \mathbb{R})$ and $\mathrm{PSL}(3, \mathbb{R})$ are dealt with separately has been our main source of inspiration (see also [5]).

Throughout this article, we will use the following notation. Let $\Sigma_{g,p}$ be a genus g surface with p punctures x_1, \dots, x_p , assuming $p > 0$ and $2 - 2g - p < 0$. We denote by $\pi_{g,p}$ its fundamental group and use the following standard presentation where the c_i 's are homotopy classes of curves enclosing the x_i 's:

$$\pi_{g,p} = \langle a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_p \mid \prod_{i=1}^g [a_i, b_i] \prod_{j=1}^p c_j \rangle.$$

We will call *flag* a pair (C, p) where C is a complex line of $\mathbf{H}_{\mathbb{C}}^2$ (see definition 2) and p is a boundary point of C . Our goal is to parametrize the variety

$$\mathfrak{R}_{g,p} = \{\rho, \mathcal{F}\} / \text{PU}(2,1)$$

where

- ρ is a representation of $\pi_{g,p}$ in $\text{PU}(2,1)$
- $F = (F_1, \dots, F_p)$ is a p -tuple of flags such that $\rho(c_i)$ stabilizes F_i .
- The group $\text{PU}(2,1)$ acts on ρ by conjugation and on F by isometries.

Remark 1. In [3, 4], the authors use an alternative description of $\mathfrak{R}_{g,p}$ which is equivalent to the above one but appears to be more efficient for certain aspects. We recall it briefly for later use. Let $\widehat{\Sigma}_{g,p}$ be the universal covering of $\Sigma_{g,p}$. The surface $\widehat{\Sigma}_{g,p}$ may be seen as a topological disk with an action of $\pi_{g,p}$ and an invariant family X of boundary points projecting onto the x_i 's. The space $\mathfrak{R}_{g,p}$ is in bijection with the space of couples (ρ, F) up to conjugation, where ρ is a representation of $\pi_{g,p}$ in $\text{PU}(2,1)$ and F is an equivariant map from X to the space of flags, that is, for all g in $\pi_{g,p}$ and x in X one has $F(g.x) = \rho(g).F(x)$.

Let us give some rough indications about the equivalence of the two definitions. The curve c_i determines a preferred lift of x_i in X which we denote by \widehat{x}_i . Given an equivariant map F , we just set $F_i = F(\widehat{x}_i)$. Reciprocally, if we have a p -tuple (F_1, \dots, F_p) of flags, we construct a map F by setting $F(\widehat{x}_i) = F_i$ and extend it by the equivariance property.

To build the coordinate system, we start from an ideal triangulation T of $\Sigma_{g,p}$ (see definition 15). Each triangle Δ of T lifts to $\widehat{\Sigma}_{g,p}$ as a triangle whose vertices are denoted by $x, y, z \in X$. Given a pair (ρ, F) , the triple of flags $(F(x), F(y), F(z))$ is well defined up to isometry. In definition 10, we introduce a notion of genericity for a triple of flags. We will say that a couple (ρ, F) is *generic with respect to T* if the triple of flags associated to any triangle of T is generic. We will denote by $\mathfrak{R}_{g,p}^T$ the subset of $\mathfrak{R}_{g,p}$ containing those class of pairs (ρ, F) that are generic with respect to T .

Next, we associate to Δ a family of invariants which parametrizes generic triples of flags up to isometry. The definition and study of these invariants is a crucial point of the article. To represent a geometric configuration, the invariants associated to adjacent triangles must satisfy compatibility relations. We will call *decoration* of a triangulation T the following data: a family of invariants for each triangle of T such that the compatibility conditions are satisfied (see definition 14).

The main result of the article is the following

Theorem 1. *Let T be an ideal triangulation of $\Sigma_{g,p}$. There is a bijection between $\mathfrak{R}_{g,p}^T$ and $\mathcal{X}(T)$, the set of decorations of T .*

Note that some isometries of $\mathbf{H}_{\mathbb{C}}^2$ do not preserve any flag. These isometries are unipotent parabolic and are conjugate in $\text{PU}(2,1)$ to non-vertical Heisenberg translations (see chap. 4 in [7]). As a consequence, $\mathfrak{R}_{g,p}$ do not contain all representations of $\pi_{g,p}$ in $\text{PU}(2,1)$.

To any ideal triangulation, Fock and Goncharov associate a coordinate system on the representation variety. The transition from a triangulation to another may be done by a succession of elementary moves, the so-called flips, which allow them to forget about the initial choice of a triangulation. In the cases treated by Fock and Goncharov, the introduction of coordinate systems gives rise to a special class of representations called positive. By computing the coordinate changes associated to the flips, they show first that the positivity of a representation is independent of the choice of triangulation and

second that the positive representations are discrete and faithful. These coordinates appear *a posteriori* to be a quick and elegant way to study Teichmüller spaces and their generalizations. Such a treatment of discreteness in the case of $\mathrm{PU}(2,1)$ seems to be still out of reach.

The study of representations of surface groups in $\mathrm{PU}(2,1)$ began in the eighties with Goldman and Toledo among others (see [6, 13]). However, many natural questions still do not have received a complete answer. Apart from a few general results about rigidity and flexibility (see [6, 10, 13]) most of the results are dealing with examples or families of examples (see [1, 8, 16]).

Up to this day, no example of a $\mathrm{PU}(2,1)$ -moduli space of discrete and faithful representations of a given surface group has been described. The only infinite group of finite type for which all the discrete and faithful representations in $\mathrm{PU}(2,1)$ are known is the modular group $\mathrm{PSL}(2, \mathbb{Z})$ (see [2]). In the case of closed surfaces, Parker and Platis have described in [9] coordinates analogous to Fenchel-Nielsen coordinates in the setting of $\mathrm{PU}(2,1)$.

In this article, we have chosen to introduce all the notions of complex hyperbolic geometry we are using. Some of the invariants we are dealing with are very classical. As an example, the invariant φ of a pair of complex lines (see 2.3.1) is treated in [7] and the classification of triples of complex lines (see 2.3.2) appears in [11]. We decided to include the definitions and proofs about these invariants for the convenience of the reader. Nevertheless, the invariants m and δ (see definitions 8 and 9) are specially adapted to pairs of flags and do not appear elsewhere to our knowledge.

The article is organized as follows:

- The section 2 is devoted to the exposition of notions of complex hyperbolic geometry. We describe totally geodesic subspaces of the complex hyperbolic plane and introduce invariants of pairs and triples of complex lines.
- In section 3, we describe the main technical tools which are the invariants m and δ . The main result of this section is the theorem 2 which classifies triples of flags up to isometry.
- In section 4, we define the *standard configuration* of a flag and a complex line. Using the invariants described in the previous section, we provide two explicit matrices that are the elementary pieces necessary to construct the representations from the invariants. These matrices may be useful for numerical applications.
- The section 5 is devoted to the definition of the decoration space and to the proof of theorem 1.
- We prove in section 6 that the compatibility equations involved in the decoration space always have solutions. The main tool is the lemma 4: it shows that once the Φ and m invariants of a triple of flag are known, there exist generically 2 possible triples of δ invariants, which correspond to the fixed points of an antiholomorphic isometry in the boundary of a disk. As a consequence of this lemma, we obtain in proposition 11 that $\mathfrak{R}_{g,p}^T$ is a 2^N cover of a simpler space denoted by $\mathcal{M}_{g,p}^T$, which is an auxiliary decoration space of the triangulation T , given by the Φ and m invariants.
- We give in section 7 some indications about how to control the isometry type of the images of the boundary curves by the representation constructed from a decorated triangulation of $\Sigma_{g,p}$. We first deal with the case of an arbitrary punctured surface, and move then to the case of the 1-punctured torus.

2 Complex hyperbolic geometry

2.1 Generalities

Consider the hermitian form of signature $(2, 1)$ in \mathbb{C}^3 given by the formula $\langle v, w \rangle = v^T J \bar{w}$ where J is the matrix given by

$$J = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

We define the following subsets of \mathbb{C}^3 :

$$\begin{aligned} V_0 &= \{v \in \mathbb{C}^3 \setminus \{0\}, \langle v, v \rangle = 0\} \\ V_- &= \{v \in \mathbb{C}^3 \setminus \{0\}, \langle v, v \rangle < 0\} \\ V_+ &= \{v \in \mathbb{C}^3 \setminus \{0\}, \langle v, v \rangle > 0\} \end{aligned}$$

Let $\mathbf{P} : \mathbb{C}^3 \setminus \{0\} \rightarrow \mathbb{C}P^2$ be the canonical projection onto the complex projective space.

Definition 1. The complex hyperbolic plane $\mathbf{H}_{\mathbb{C}}^2$ is the set $\mathbf{P}(V_-)$ equipped with the Bergman metric.

The boundary of $\mathbf{H}_{\mathbb{C}}^2$ is $\mathbf{P}(V_0)$. The distance function associated to the Bergman metric is given in terms of Hermitian product by

$$\cosh^2 \left(\frac{d(m, n)}{2} \right) = \frac{\langle \mathbf{m}, \mathbf{n} \rangle \langle \mathbf{n}, \mathbf{m} \rangle}{\langle \mathbf{m}, \mathbf{m} \rangle \langle \mathbf{n}, \mathbf{n} \rangle}, \quad (1)$$

where \mathbf{m} and \mathbf{n} are lifts of m and n to \mathbb{C}^3 . It follows from (1) that $U(2,1)$, the unitary group associated to J , acts on $\mathbf{H}_{\mathbb{C}}^2$ by holomorphic isometries. The full isometry group of $\mathbf{H}_{\mathbb{C}}^2$ is generated by $PU(2,1)$ and the complex conjugation. The usual trichotomy of isometries for $PSL(2, \mathbb{R})$ holds here also: an isometry is elliptic if it has a fixed point inside $\mathbf{H}_{\mathbb{C}}^2$, parabolic if it has a unique fixed point on $\partial \mathbf{H}_{\mathbb{C}}^2$, and loxodromic if it has exactly two fixed points on $\partial \mathbf{H}_{\mathbb{C}}^2$, and this exhausts all possibilities.

2.2 Subspaces of $\mathbf{H}_{\mathbb{C}}^2$.

There are two types of maximal totally geodesic subspaces of $\mathbf{H}_{\mathbb{C}}^2$, which are both of (real) dimension 2: complex lines and \mathbb{R} -planes. We give now a few indications about these. More details may be found in [7].

2.2.1 Complex lines

Definition 2. We call *complex line* in $\mathbf{H}_{\mathbb{C}}^2$ the intersection with $\mathbf{H}_{\mathbb{C}}^2$ of the projectivization of a 2-dimensional subspace of \mathbb{C}^3 which intersects V_- . Such a subspace is orthogonal to a one-dimensional subspace contained in V_+ : we call *polar vector* of the complex line any generator of this subspace.

Note that a complex line is an isometric embedding of the complex hyperbolic line $\mathbf{H}_{\mathbb{C}}^1$. To any complex line C is associated a unique holomorphic involution fixing pointwise C , which we shall refer to as the *complex symmetry* with respect to C . The group $PU(2,1)$ acts transitively on the set of complex lines of $\mathbf{H}_{\mathbb{C}}^2$.

Definition 3. We call *flag* a pair (C, p) where C is a complex line and p is a point in $C \cap \partial\mathbf{H}_{\mathbb{C}}^2$.

Lemma 1. $PU(2,1)$ acts transitively on the set of flags of $\mathbf{H}_{\mathbb{C}}^2$.

2.2.2 \mathbb{R} -planes

Definition 4. An \mathbb{R} -plane is the intersection with $\mathbf{H}_{\mathbb{C}}^2$ of the projection of a vectorial Lagrangian subspace of $\mathbb{C}^{2,1}$.

Every \mathbb{R} -plane P is fixed pointwise by a unique antiholomorphic isometric involution I_P , which is the projectivization of the Lagrangian symmetry with respect to any lift of P as a vectorial Lagrangian. We will refer to I_P as the *Lagrangian reflection about P* . The standard example is the set of points of $\mathbf{H}_{\mathbb{C}}^2$ with real coordinates, which is fixed by the complex conjugation. We will refer to this \mathbb{R} -plane as $\mathbf{H}_{\mathbb{R}}^2 \subset \mathbf{H}_{\mathbb{C}}^2$. It is an embedding of the real hyperbolic plane into $\mathbf{H}_{\mathbb{C}}^2$.

As a consequence, we obtain

Proposition 1. Let Q be an \mathbb{R} -plane. There exists a matrix $M_Q \in SU(2,1)$ such that

$$M_Q \overline{M_Q} = 1, \text{ and } I_Q(m) = \mathbf{P}(M_Q \cdot \overline{\mathbf{m}}) \text{ for any } m \text{ in } \mathbf{H}_{\mathbb{C}}^2 \text{ with lift } \mathbf{m}. \quad (2)$$

Proof. Let \mathbf{Q} be a vectorial lift of Q , and choose $\mathbb{R}^3 \subset \mathbb{C}^{2,1}$ as a vectorial lift of $\mathbf{H}_{\mathbb{R}}^2$. Since the group $U(2,1)$ acts transitively on the Lagrangian Grassmanian of $\mathbb{C}^{2,1}$, there exists a matrix $A \in U(2,1)$ such that $A\mathbb{R}^3 = \mathbf{Q}$. The matrix $M_Q = A\overline{A}^{-1}$ belongs to $SU(2,1)$ and satisfies the condition (2). \square

Remark 2. If I_1 and I_2 are two Lagrangian reflections with associated matrices \mathbf{M}_1 and \mathbf{M}_2 , then their composition, which is a holomorphic isometry, admits the matrix $\mathbf{M}_1 \overline{\mathbf{M}_2}$ as a lift to $SU(2,1)$.

2.3 Classical invariants

2.3.1 Invariant of two complex lines

Definition 5. Let C_1 and C_2 be two complex lines of $\mathbf{H}_{\mathbb{C}}^2$, with polar vectors \mathbf{c}_1 and \mathbf{c}_2 . We set

$$\varphi(C_1, C_2) = \frac{|\langle \mathbf{c}_1, \mathbf{c}_2 \rangle|^2}{\langle \mathbf{c}_1, \mathbf{c}_1 \rangle \langle \mathbf{c}_2, \mathbf{c}_2 \rangle}.$$

Clearly, $\varphi(C_1, C_2)$ does not depend on the choice of the lift in the pair of polar vectors, and is $PU(2,1)$ -invariant. We recall the geometric interpretation of φ , and we refer to [7] for details:

- $\varphi(C_1, C_2) > 1$ if C_1 and C_2 are disjoint in $\mathbf{H}_{\mathbb{C}}^2$. In this case, the distance d between C_1 and C_2 is given by the formula $\varphi(C_1, C_2) = \cosh^2(d/2)$.
- $\varphi(C_1, C_2) = 1$ if C_1 and C_2 are either identical or asymptotic, by which we mean that they meet in $\partial\mathbf{H}_{\mathbb{C}}^2$.
- $\varphi(C_1, C_2) < 1$ if C_1 and C_2 intersect. The angle θ of their intersection is given by the relation $\varphi(C_1, C_2) = \cos^2(\theta)$.

Note that two complex lines are orthogonal if and only if $\varphi(C_1, C_2) = 0$. The φ -invariant classifies pairs of distinct complex lines up to isometries.

Proposition 2. *Let C_1, C_2, D_1, D_2 be 4 complex lines such that $C_1 \neq C_2$ and $D_1 \neq D_2$. There exists an isometry $g \in PU(2, 1)$ such that $D_1 = gC_1$ and $D_2 = gC_2$ if and only if $\varphi(C_1, C_2) = \varphi(D_1, D_2)$.*

Proof. It is clear that if the two pairs are isometric, their invariant is the same. Reciprocally, choose polar vectors $\mathbf{c}_1, \mathbf{c}_2, \mathbf{d}_1, \mathbf{d}_2$ with norm 1 and such that $\langle \mathbf{c}_1, \mathbf{c}_2 \rangle$ and $\langle \mathbf{d}_1, \mathbf{d}_2 \rangle$ are in $\mathbb{R}_{\geq 0}$.

As $C_1 \neq C_2$, the vectors \mathbf{c}_1 and \mathbf{c}_2 are independent and the same is true for \mathbf{d}_1 and \mathbf{d}_2 . With the assumption on the φ -invariant, the Gram matrices of $(\mathbf{c}_1, \mathbf{c}_2)$ and $(\mathbf{d}_1, \mathbf{d}_2)$ are identical. It means that one can find an isometry which maps \mathbf{c}_1 on \mathbf{d}_1 and \mathbf{c}_2 on \mathbf{d}_2 . This ends the proof. \square

We will need the following lemma.

Lemma 2. *Let C_1 and C_2 be two non orthogonal distinct complex lines, and p_1 a point in $\partial\mathbf{H}_{\mathbb{C}}^2 \cap C_1$ which is not in C_2 . Except for the identity, no isometry preserves C_1 and C_2 and fixes p_1 .*

Proof. Pick \mathbf{c}_1 and \mathbf{c}_2 two vectors polar to C_1 and C_2 of norm 1 and \mathbf{p}_1 a lift of p_1 . Writing $\langle \mathbf{p}_1, \mathbf{c}_2 \rangle = a$ and $\langle \mathbf{c}_1, \mathbf{c}_2 \rangle = b$, the hermitian form has the following matrix in the basis $(\mathbf{p}_1, \mathbf{c}_1, \mathbf{c}_2)$

$$H = \begin{bmatrix} 0 & 0 & a \\ 0 & 1 & b \\ \bar{a} & \bar{b} & 1 \end{bmatrix}$$

An isometry having the requested property has a diagonal lift to $SU(2,1)$ in this basis. The result is obtained by writing the isometry condition ${}^t\overline{M}HM = H$, and by using the fact that b is non zero since C_1 and C_2 are not orthogonal. \square

2.3.2 Invariants of three complex lines

Let C_1, C_2 and C_3 be three complex lines in $\mathbf{H}_{\mathbb{C}}^2$. We will say that they are in *generic position* if their polar vectors form a basis of \mathbb{C}^3 . There are three invariants of the triple (C_1, C_2, C_3) given by the φ -invariant of all pairs of complex lines. We will need a fourth one (given in the following definition) to classify all triples up to isometry.

Definition 6. Let C_1, C_2, C_3 be three complex lines in $\mathbf{H}_{\mathbb{C}}^2$ with respective polar vectors $\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3$. Then we set

$$\Phi(C_1, C_2, C_3) = \frac{\langle \mathbf{c}_1, \mathbf{c}_2 \rangle \langle \mathbf{c}_2, \mathbf{c}_3 \rangle \langle \mathbf{c}_3, \mathbf{c}_1 \rangle}{\langle \mathbf{c}_1, \mathbf{c}_1 \rangle \langle \mathbf{c}_2, \mathbf{c}_2 \rangle \langle \mathbf{c}_3, \mathbf{c}_3 \rangle}.$$

The importance of this invariant should be clear from the following two propositions (see [11]):

Proposition 3. *Let C_1, C_2, C_3 be three complex lines of $\mathbf{H}_{\mathbb{C}}^2$ in generic position. For simplicity, we denote by φ_{ij} the φ -invariant of C_i and C_j and by Φ_{ijk} the Φ -invariant of C_i, C_j, C_k . These invariants enjoy the following properties.*

1. For all distinct $i, j, k \in \{1, 2, 3\}$, the following relations are satisfied.

$$\varphi_{ij} = \varphi_{ji}, \Phi_{ijk} = \Phi_{jki} = \overline{\Phi_{ikj}} \text{ and } \Phi_{ijk}\Phi_{ikj} = \varphi_{ij}\varphi_{jk}\varphi_{ki}. \quad (3)$$

2. The four invariants satisfy to the inequality

$$1 - \varphi_{12} - \varphi_{23} - \varphi_{31} + \Phi_{123} + \Phi_{132} < 0. \quad (4)$$

Proof. 1. These relations are straightforward from the definitions of the invariants φ and Φ .

2. Let C_1, C_2, C_3 be three complex lines in generic position. Let $\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3$ be three polar vectors associated to these lines. Let G be the Gram matrix of the basis $(\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3)$. A direct computation shows that the left-hand side of relation (4) is equal to

$$\Delta(C_1, C_2, C_3) = \frac{\det G}{\langle \mathbf{c}_1, \mathbf{c}_1 \rangle \langle \mathbf{c}_2, \mathbf{c}_2 \rangle \langle \mathbf{c}_3, \mathbf{c}_3 \rangle}.$$

This number is an invariant of the triple (C_1, C_2, C_3) . As the Gram matrix represents the hermitian form, it has signature $(2, 1)$ and its determinant is negative. \square

Proposition 4. Consider non negative real numbers φ_{ij} and complex numbers Φ_{ijk} satisfying the relations of proposition 3. There exists a triple C_1, C_2, C_3 in generic position, unique up to isometry, such that for all distinct i, j, k in $\{1, 2, 3\}$ the relations $\varphi(C_i, C_j) = \varphi_{ij}$ and $\Phi_{ijk} = \varphi(C_i, C_j, C_k)$ hold.

Proof. Consider real numbers φ_{ij} and complex numbers Φ_{ijk} satisfying the relations (3) and (4), and \mathbb{C}^3 with its canonical basis (e_1, e_2, e_3) . We define a hermitian form h on it by setting

$$\begin{aligned} h(e_i, e_i) &= 1 \text{ for all } i, & h(e_1, e_2) &= \sqrt{\varphi_{12}}, \\ h(e_2, e_3) &= \sqrt{\varphi_{23}}, & h(e_3, e_1) &= \frac{\Phi_{123}}{\sqrt{\varphi_{12}\varphi_{23}}}. \end{aligned}$$

The matrix of h in the basis (e_1, e_2, e_3) has unit diagonal entries – thus positive trace – and according to the relation (4), it has negative determinant. As a consequence, h has signature $(2, 1)$. By the classification of hermitian forms, this model is conjugate to the standard one, and the vectors e_1, e_2, e_3 map to the polar vectors of the desired complex lines. Moreover, the few choices we made disappear projectively, hence the triple of complex lines is unique up to isometry. \square

3 Invariants of flags

3.1 Invariant of two flags

Definition 7. Let (C_1, p_1) and (C_2, p_2) be two flags in $\mathbf{H}_{\mathbb{C}}^2$. We will say that they are in *generic position* if p_1 does not belong to C_2 , p_2 does not belong to C_1 and C_1 is not orthogonal to C_2 .

Remark 3. The condition of non-orthogonality of C_1 and C_2 will be needed to define the elementary isometries associated to a triple of flags in a unique way (see propositions 7 and 8).

Definition 8. Let (C_1, p_1) and (C_2, p_2) be two flags in generic position. Let $\mathbf{c}_1, \mathbf{c}_2$ be polar vectors of C_1, C_2 and $\mathbf{p}_1, \mathbf{p}_2$ be representatives of p_1, p_2 . We set

$$m[(C_1, p_1), (C_2, p_2)] = \frac{\langle \mathbf{c}_1, \mathbf{c}_2 \rangle \langle \mathbf{p}_1, \mathbf{p}_2 \rangle}{\langle \mathbf{c}_1, \mathbf{p}_2 \rangle \langle \mathbf{p}_1, \mathbf{c}_2 \rangle}.$$

This invariant is a complex generalization of the φ -invariant of two complex lines. Its properties are summed up in the following proposition.

Proposition 5. Let (C_1, p_1) and (C_2, p_2) be two flags in generic position, and m_{12} their invariant $m[(C_1, p_1), (C_2, p_2)]$.

1. The two invariants $\varphi(C_1, C_2)$ and m_{12} are linked by the relation

$$\varphi(C_1, C_2) = \left| \frac{m_{12}}{m_{12} - 1} \right|^2. \quad (5)$$

2. For any complex number $m_{12} \in \mathbb{C} \setminus \{0, 1\}$ there exists a pair of flags $(C_1, p_1), (C_2, p_2)$ in generic position such that $m[(C_1, p_1), (C_2, p_2)] = m_{12}$. This pair is unique up to isometry.

Proof. 1. Let (C_1, p_1) and (C_2, p_2) be two flags in generic position, $\mathbf{c}_1, \mathbf{c}_2$ be polar vectors of C_1, C_2 and $\mathbf{p}_1, \mathbf{p}_2$ be representatives of p_1, p_2 . The family of vectors $(\mathbf{c}_1, \mathbf{c}_2, \mathbf{p}_1, \mathbf{p}_2)$ is linearly dependent, hence the determinant of its Gram matrix vanishes. Computing this determinant and dividing by non vanishing factors, we obtain the relation $|m_{12} - 1|^2 \varphi_{12} = |m_{12}|^2$. From that relation we see that m_{12} cannot be equal to 1 since the two flags are in generic position and therefore φ_{12} is non-zero. This proves relation (5).

2. In order to prove the last part of the proposition, we make the following observation: given two flags (C_1, p_1) and (C_2, p_2) in generic position, there is a unique complex line C_3 joining p_1 and p_2 . Following proposition 4, the triple (C_1, C_2, C_3) is determined by its φ -invariants, hence, we can classify couples of flags using φ -invariants.

More precisely, as C_1 and C_3 are asymptotic (they meet on $p_1 \in \partial \mathbf{H}_{\mathbb{C}}^2$), their φ invariant φ_{13} equals 1. For the same reason, $\varphi_{23} = 1$. As a consequence of relation (3), we obtain the equality $|\Phi_{123}|^2 = \varphi_{12}$. Plugging these values into the relation (4) yields

$$\Delta_{123} = -1 - \varphi_{12} + \Phi_{123} + \Phi_{132} = -|1 - \Phi_{123}|^2. \quad (6)$$

Suppose that we have $\Phi_{123} \neq 1$, then the complex lines C_1, C_2, C_3 are in generic position. Let $\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3$ be polar vectors of these lines. They form a basis of \mathbb{C}^3 and the linear forms $\langle \cdot, \mathbf{c}_1 \rangle, \langle \cdot, \mathbf{c}_2 \rangle, \langle \cdot, \mathbf{c}_3 \rangle$ form a linear basis of the dual of \mathbb{C}^3 . One can find a unique anti-dual basis $\mathbf{d}_1, \mathbf{d}_2, \mathbf{d}_3$ such that for all i, j in $\{1, 2, 3\}$ one has $\langle \mathbf{d}_i, \mathbf{c}_j \rangle = \delta_{ij}$. A direct computation shows that the Gram matrix of the hermitian form in the basis $(\mathbf{d}_1, \mathbf{d}_2, \mathbf{d}_3)$ is the inverse of the Gram matrix of $(\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3)$. Moreover \mathbf{d}_1 being

orthogonal to \mathbf{c}_2 and \mathbf{c}_3 , it is a representative of p_2 and \mathbf{d}_2 is a representative of p_1 . Using these representatives we get (see remark 4 above)

$$\begin{aligned} m_{12} &= \langle \mathbf{c}_1, \mathbf{c}_2 \rangle \langle \mathbf{d}_1, \mathbf{d}_2 \rangle = \frac{\langle \mathbf{c}_1, \mathbf{c}_2 \rangle (\langle \mathbf{c}_3, \mathbf{c}_1 \rangle \langle \mathbf{c}_2, \mathbf{c}_3 \rangle - \langle \mathbf{c}_2, \mathbf{c}_1 \rangle \langle \mathbf{c}_3, \mathbf{c}_3 \rangle)}{\langle \mathbf{c}_1, \mathbf{c}_1 \rangle \langle \mathbf{c}_2, \mathbf{c}_2 \rangle \langle \mathbf{c}_3, \mathbf{c}_3 \rangle \Delta_{123}} = \frac{\Phi_{123} - \varphi_{12}}{\Delta_{123}} \\ &= \frac{\Phi_{123} - |\Phi_{123}|^2}{-|1 - \Phi_{123}|^2} = \frac{\Phi_{123}}{\Phi_{123} - 1}. \end{aligned}$$

This proves that $m_{12} = 1$ if and only if $\Phi_{123} = 1$ and that m_{12} classifies couples of flags as Φ_{123} does. \square

Remark 4. Note that this anti-dual basis is usually used in the literature under a slightly different form, using the so-called hermitian cross-product. The vector \mathbf{d}_2 is proportionnal to the hermitian cross-product of \mathbf{c}_1 and \mathbf{c}_3 , denoted by $\mathbf{c}_1 \boxtimes \mathbf{c}_3$. It is a simple computation using hermitian cross-product to check that $\langle \mathbf{d}_1, \mathbf{d}_2 \rangle$ equals $(\langle \mathbf{c}_3, \mathbf{c}_1 \rangle \langle \mathbf{c}_2, \mathbf{c}_3 \rangle - \langle \mathbf{c}_2, \mathbf{c}_1 \rangle \langle \mathbf{c}_3, \mathbf{c}_3 \rangle) \cdot (\langle \mathbf{c}_1, \mathbf{c}_1 \rangle \langle \mathbf{c}_2, \mathbf{c}_2 \rangle \langle \mathbf{c}_3, \mathbf{c}_3 \rangle \Delta_{123})^{-1}$. See [7] for details.

3.2 Invariant of a flag and two complex lines

Definition 9. Let (C_1, p_1) be a flag and C_2, C_3 be two complex lines such that the three complex lines C_1, C_2 and C_3 are in generic position and such that p_1 does not belong to C_2 nor C_3 . Take $\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3$ three polar vectors of C_1, C_2, C_3 and \mathbf{p}_1 a representative of p_1 . Then we set:

$$\delta[(C_1, p_1), C_2, C_3] = \delta_{23}^1 = \frac{\langle \mathbf{c}_2, \mathbf{c}_3 \rangle \langle \mathbf{p}_1, \mathbf{c}_2 \rangle}{\langle \mathbf{c}_2, \mathbf{c}_2 \rangle \langle \mathbf{p}_1, \mathbf{c}_3 \rangle}. \quad (7)$$

This invariant may be viewed as a coordinate of p_1 knowing C_1, C_2 and C_3 . Its main properties are summed up in the following proposition.

Proposition 6. *Let (C_1, p_1) be a flag and C_2, C_3 be two complex lines such that the three complex lines C_1, C_2 and C_3 are in generic position and such that p_1 does not belong to C_2 nor C_3 . The invariants δ_{23}^1 and δ_{32}^1 satisfy the following equations:*

$$\varphi_{23} = \delta_{23}^1 \delta_{32}^1 \quad (8)$$

$$0 = (1 - \varphi_{13}) |\delta_{23}^1|^2 + 2\text{Re} [(\Phi_{132} - \varphi_{23}) \delta_{23}^1] + \varphi_{23} (1 - \varphi_{12}) \quad (9)$$

Reciprocally, take C_1, C_2, C_3 three complex lines in generic position. Any non zero value of δ_{23}^1 which satisfies the second equation corresponds to a unique point p_1 in C_1 which is not on C_2 nor on C_3 .

Proof. The first equation is a direct consequence of the definition. For the second one, let $(\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3)$ be a basis of \mathbb{C}^3 formed by polar vectors for C_1, C_2, C_3 . Let $(\mathbf{d}_1, \mathbf{d}_2, \mathbf{d}_3)$ be its anti-dual basis. We will use the latter basis to prove relation (9). We recall that the matrix of the Hermitian form in the basis $(\mathbf{d}_1, \mathbf{d}_2, \mathbf{d}_3)$ is the inverse of the Gram matrix of $(\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3)$.

As p_1 belongs to C_1 , its representative is a linear combination of \mathbf{d}_2 and \mathbf{d}_3 , and we may thus write $\mathbf{p}_1 = a\mathbf{d}_2 + b\mathbf{d}_3$. The coordinates a and b can be recovered by computing the hermitian products $\langle \mathbf{p}_1, \mathbf{c}_2 \rangle = a$ and $\langle \mathbf{p}_1, \mathbf{c}_3 \rangle = b$. In particular, this implies

$$\delta_{23}^1 = \frac{\langle \mathbf{c}_2, \mathbf{c}_3 \rangle a}{\langle \mathbf{c}_2, \mathbf{c}_2 \rangle b}. \quad (10)$$

By expressing that \mathbf{p}_1 is in the isotropic cone of the Hermitian form, we obtain the relation (9).

On the other hand, if we know δ_{23}^1 , then according to relation (10), we know projective coordinates for \mathbf{p}_1 . If δ_{23}^1 satisfies (9), the vector \mathbf{p}_1 must be on the cone of the quadratic form. It proves that δ_{23}^1 determines the position of p_1 on C_1 as asserted. \square

3.3 Summary : invariants of three flags

In the remaining part of the article, we will be interested in the space of configurations of three flags. Let us sum up what are the relevant invariants for such configurations.

Definition 10. We will say that three flags $(C_i, p_i)_{i=1,2,3}$ are in *generic position* if they are pairwise in generic position, and if the triple of complex lines (C_1, C_2, C_3) is also in generic position, that is, if

- any two of the complex lines are distinct and non-orthogonal,
- any triple of vectors polar to the C_i 's is a basis of \mathbb{C}^3 .

We classify now the triples of flags up to $\text{PU}(2,1)$.

Theorem 2. *Let (C_1, p_1) , (C_2, p_2) and (C_3, p_3) be three flags in generic position. The configuration of these flags modulo holomorphic isometry is classified by the invariants φ_{ij} , Φ_{ijk} and δ_{jk}^i for all distinct i, j, k in $\{1, 2, 3\}$. These invariants satisfy the following equations for all i, j, k :*

$$(3) \quad \varphi_{ij} = \varphi_{ji} = \overline{\varphi_{ij}} > 0, \quad \Phi_{ijk} = \Phi_{jki} = \overline{\Phi_{ikj}} \quad \text{and} \quad \Phi_{ijk}\Phi_{ikj} = \varphi_{ij}\varphi_{jk}\varphi_{ki}.$$

$$(4) \quad \Delta_{ijk} = 1 - \varphi_{ij} - \varphi_{jk} - \varphi_{ki} + \Phi_{ijk} + \Phi_{ikj} < 0.$$

$$(8) \quad \delta_{jk}^i \delta_{kj}^i = \varphi_{ij}.$$

$$(9) \quad (1 - \varphi_{ik})|\delta_{jk}^i|^2 + 2\text{Re} \left[(\Phi_{ikj} - \varphi_{jk})\delta_{jk}^i \right] + \varphi_{jk}(1 - \varphi_{ij}) = 0.$$

The space of solutions is a manifold of dimension 7. Moreover, the invariants m_{ij} attached to pairs of flags are expressed in terms of the other invariants as follows :

$$\begin{aligned} m_{ij}\Delta_{ijk}\varphi_{ik}\varphi_{jk} &= \varphi_{ik}\varphi_{jk}(\Phi_{ijk} - \varphi_{ij}) + \varphi_{ik}(\varphi_{ij}\varphi_{jk} - \Phi_{ijk})\delta_{kj}^i \\ &\quad + \varphi_{jk}(\varphi_{ij}\varphi_{ik} - \Phi_{ijk})\overline{\delta_{ki}^j} + \Phi_{ijk}(1 - \varphi_{ij})\delta_{kj}^i\overline{\delta_{ki}^j} \end{aligned} \quad (11)$$

Proof. The first part of the proof is nothing but a summary of the preceding sections. Let us now compute m_{12} . The two other m -invariants are obtained in the same way. Choose $\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3$ polar vectors of C_1, C_2, C_3 and let $(\mathbf{d}_1, \mathbf{d}_2, \mathbf{d}_3)$ be the anti-dual basis as usual. Then, using the proof of proposition 6, one can find explicit coordinates for representatives of p_1 and p_2 in the basis $(\mathbf{d}_1, \mathbf{d}_2, \mathbf{d}_3)$. Precisely, we can choose

$$\begin{cases} \mathbf{p}_1 = \langle \mathbf{c}_3, \mathbf{c}_2 \rangle \mathbf{d}_2 + \langle \mathbf{c}_3, \mathbf{c}_3 \rangle \delta_{32}^1 \mathbf{d}_3 \\ \mathbf{p}_2 = \langle \mathbf{c}_3, \mathbf{c}_1 \rangle \mathbf{d}_1 + \langle \mathbf{c}_3, \mathbf{c}_3 \rangle \delta_{31}^2 \mathbf{d}_3. \end{cases}$$

To obtain a formula for m_{12} , we just need to replace \mathbf{p}_1 and \mathbf{p}_2 in the definition of m_{12} by the expressions above. We obtain the relation (11) after a computation. \square

4 Elementary isometries associated to a triple of flags

4.1 \mathbb{R} -planes associated to a triple of flags and elementary isometries

In this paragraph, we define the elementary isometries associated to a triple of flags. More precisely, we prove the

Proposition 7. *Let $F_i = (C_i, p_i)$ for $i = 1, 2, 3$ be a triple of flags in generic position such that any two complex lines are not asymptotic.*

1. *For any pair (i, j) with $i \neq j$, there exists a unique isometry E_{ij} exchanging C_i and C_j , and mapping p_j to p_i . It is called the exchange isometry associated to the pair of flags F_i and F_j .*
2. *There exists a unique isometry T_{jk}^i fixing p_i and preserving C_i which maps C_k to a complex line C'_k satisfying $R_{C'_k}(p_i) = R_{C_j}(p_i)$, where R_C is the complex symmetry with respect to the complex line C . It is called the transfer isometry associated to the ordered triple of flags (F_i, F_j, F_k) .*

We will give a geometric proof of this proposition, showing that the exchange and transfer isometry are obtained as products of Lagrangian reflections which are canonically associated to a triple of flags satisfying the assumption of proposition 7.

Proposition 8. *Let C_1 and C_2 be two complex lines which are neither orthogonal nor asymptotic.*

1. *Let p_1 be a point in ∂C_1 . There exists a unique \mathbb{R} -plane P such that I_P , the inversion in P , preserves both C_1 and C_2 , and fixes p_1 .*
2. *Let p_2 be a point in ∂C_2 . There exists a unique \mathbb{R} -plane Q such that I_Q , the inversion in Q , swaps C_1 and C_2 and maps p_1 to p_2 .*
3. *Let m and n be two points in the boundary of $\mathbf{H}_{\mathbb{C}}^2$, not belonging to ∂C_1 . There exists a unique Lagrangian reflection preserving C_1 and swapping m and n .*
4. *Let p_1, p_2 and p_3 be three points of $\partial \mathbf{H}_{\mathbb{C}}^2$, not contained in the boundary of a complex line. There exists a unique Lagrangian reflection fixing p_1 and swapping p_2 and p_3 .*

Proof. Let \mathbf{c}_k be a polar vector for C_k normalized so that $\langle \mathbf{c}_k, \mathbf{c}_k \rangle = 1$. Let \mathbf{p}_1 be a lift of p_1 . Rescaling if necessary, we may assume that both $a = \langle \mathbf{p}_1, \mathbf{c}_2 \rangle$ and $b = \langle \mathbf{c}_1, \mathbf{c}_2 \rangle$ are real (in fact $\langle \mathbf{c}_1, \mathbf{c}_2 \rangle$ is equal to $\sqrt{\varphi_{12}}$).

1. The hermitian form admits in the basis $(\mathbf{p}_1, \mathbf{c}_1, \mathbf{c}_2)$ the matrix

$$H = \begin{bmatrix} 0 & 0 & a \\ 0 & 1 & b \\ a & b & 1 \end{bmatrix}$$

The hermitian product b is non-zero since C_1 and C_2 are non-orthogonal. In this basis, any lift of a Lagrangian reflection fixing p_1 and preserving C_1 and C_2 must be diagonal. It follows after writing the isometry condition $M^*HM = H$ that there is only one such reflection, given in this basis by $\mathbf{m} \rightarrow \overline{\mathbf{m}}$.

2. This time, we use the basis $(\mathbf{c}_1, \mathbf{c}_2, \mathbf{d})$, where \mathbf{d} a vector orthogonal to \mathbf{c}_1 and \mathbf{c}_2 with norm $b^2 - 1$ (indeed, we are setting $\mathbf{d} = \mathbf{c}_1 \boxtimes \mathbf{c}_2$, see remark 4). The hermitian form is given by the matrix

$$H = \begin{bmatrix} 1 & b & 0 \\ b & 1 & 0 \\ 0 & 0 & b^2 - 1 \end{bmatrix} \quad (|b| = 1 \text{ iff } C_1 \text{ and } C_2 \text{ are asymptotic})$$

We may choose the lifts of p_1 and p_2 as follows :

$$\mathbf{p}_1 = \begin{bmatrix} -b \\ 1 \\ e^{i\theta_1} \end{bmatrix} \text{ and } \mathbf{p}_2 = \begin{bmatrix} 1 \\ -b \\ e^{i\theta_2} \end{bmatrix} \text{ with } \theta_i \in \mathbb{R}.$$

The fact that I_Q exchanges C_1 and C_2 implies that any matrix for I_Q has the form

$$\begin{bmatrix} 0 & \alpha & 0 \\ \beta & 0 & 0 \\ 0 & 0 & \gamma \end{bmatrix}.$$

Writing the isometry condition and the fact that $I_Q(p_1) = p_2$, provides relations determining α , β and γ . The result follows.

3. We may choose lifts \mathbf{m} and \mathbf{n} of m and n such that $\langle \mathbf{m}, \mathbf{n} \rangle = 1$ and a unit vector \mathbf{c} polar to the complex line containing m and n . In the basis $(\mathbf{m}, \mathbf{c}, \mathbf{n})$, where the hermitian form has matrix J , the complex line C_1 is polar to some vector $\mathbf{c}_1 = [\alpha \ \beta \ \gamma]^T$. It is a direct computation to check that a Lagrangian reflection swapping m and n and preserving C_1 lifts to the matrix below. Hence, it exists and is unique.

$$\begin{bmatrix} 0 & 0 & \alpha/\bar{\gamma} \\ 0 & \beta/\bar{\beta} & 0 \\ \gamma/\bar{\alpha} & 0 & 0 \end{bmatrix}$$

4. In the proof of the previous item, we have not used the fact that \mathbf{c}_1 was a positive vector. Thus the same result as \mathcal{B} remains true if we change C_1 to a boundary point, that is, \mathbf{c}_1 to a null vector. If the three points are in a complex line, then we lose the uniqueness. Note that this fourth part of the proposition is classical (see for instance lemma 7.17 of [7])

□

Proof of proposition 7. 1. Let h_1 and h_2 be two isometries having the requested properties. Then $h_2^{-1} \circ h_1$ preserves both C_i and C_j , and fixes p_i . According to the lemma 2, this implies that h_2 and h_1 are equal. This proves the uniqueness. To prove the existence part, we apply the first two items of proposition 8.

- There exists a unique Lagrangian reflection I_2 preserving C_i and C_j and fixing p_i (this follows from part 1 of proposition 8).

- There exists a unique Lagrangian reflection I_1 swapping C_i and C_j , and exchanging p_i and $I_2(p_j)$. This is part 2 of proposition 8, which may be applied since $I_2(p_j)$ belongs to C_j .

The isometry $E_{ij} = I_1 \circ I_2$ has the requested properties.

2. The uniqueness is proved in the same way as for 1. To prove the existence, we apply the third and fourth part of proposition 8.
 - The two points $R_{C_3}(p_1)$ and $R_{C_2}(p_1)$ do not belong to ∂C_1 since the three complex lines are non-asymptotic. Thus, there exists a unique Lagrangian reflection I_3 preserving C_1 and swapping $R_{C_3}(p_1)$ and $R_{C_2}(p_1)$ (this follows from part 3 of proposition 8). Note that I_3 does not fix p_1 .
 - The three points p_1 , $I_3(p_1)$ and $R_{C_2}(p_1)$ do not belong to a common complex line, for else C_1 and C_2 would be asymptotic. Thus we may apply the fourth part of proposition 8 to obtain a (unique) Lagrangian reflection I_4 fixing $R_{C_2}(p_1)$, and swapping p_1 and $I_3(p_1)$.

The isometry $I_4 \circ I_3$ has the requested properties (note that since I_4 swaps p_1 and $I_3(p_1)$ which both belong to C_1 , it preserves C_1).

□

4.2 Standard position of a triple of flags and elementary isometries

Definition 11. - Let (C_1, p_1) be a flag and C_2 be a complex line. We will say that they are in *generic position* if p_1 does not belong to C_2 , and if C_1 and C_2 are distinct and non-orthogonal.

- We say that (C_1, p_1) and C_2 are in *standard position* if p_1, C_1 and C_2 are respectively represented by the following vectors:

$$\mathbf{p}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{c}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \mathbf{c}_2 = \begin{bmatrix} a \\ \sqrt{2} \\ 1 \end{bmatrix} \text{ for } a \in (-1, +\infty).$$

The condition on C_2 is equivalent to saying that $R_{C_2}(p_1)$ is represented by the vector $[-1 \ \sqrt{2} \ 1]^T$. The motivation for this definition is the following proposition:

Proposition 9. *Let (C_1, p_1) be a flag and C_2 be a complex line in generic position.*

- *There exists a unique couple in standard position which is isometric to $((C_1, p_1), C_2)$.*
- *The parameter a is given by $\varphi(C_1, C_2) = (1 + a)^{-1}$*

Proof. Since $PU(2, 1)$ acts transitively on the set of flags of $\mathbf{H}_{\mathbb{C}}^2$, we can assume that \mathbf{p}_1 and \mathbf{c}_1 are in standard position. The isometries g in $PU(2, 1)$ stabilizing the standard flag admit lifts to $SU(2, 1)$ of the following form :

$$\mathbf{g} = \begin{bmatrix} \lambda & 0 & it\lambda \\ 0 & \bar{\lambda}/\lambda & 0 \\ 0 & 0 & 1/\bar{\lambda} \end{bmatrix} \text{ with } \lambda \in \mathbb{C} \setminus \{0\} \text{ and } t \in \mathbb{R}.$$

Note that λ is well-defined up to multiplication by a cubic root of 1. Now, a generic polar vector for C_2 and its image by g are given by

$$\mathbf{c}_2 = \begin{bmatrix} a \\ b \\ 1 \end{bmatrix} \text{ and } \mathbf{g}\mathbf{c}_2 \sim \begin{bmatrix} |\lambda|^2(a+it) \\ \bar{\lambda}^2 b/\lambda \\ 1 \end{bmatrix} \text{ with } |b|^2 + 2\text{Re}(a) > 0.$$

The assumption that \mathbf{c}_1 and \mathbf{c}_2 are not orthogonal, implies that $b \neq 0$. This means that there is only one isometry which stabilizes the standard flag and maps \mathbf{c}_2 in standard position. Namely, we have to set $t = -\text{Im}(a)$ and solve $\bar{\lambda}^2 b = \sqrt{2}\lambda$. This equation has three solutions in λ which represent the same element in $PU(2, 1)$. The value of $\varphi(C_1, C_2)$ is given by a straightforward computation. \square

Remark 5. Given three flags (C_1, p_1) , (C_2, p_2) , (C_3, p_3) , we can decide to put (C_1, p_1) and C_2 in standard position. However, we could have chosen (C_1, p_1) and C_3 or (C_2, p_2) and C_1 . All these configurations can be obtained one from the other by applying elementary isometries to the configuration.

As an example, assume that (C_1, p_1) and C_2 are in standard position, and apply the exchange E_{12} isometry swapping C_1 and C_2 and mapping p_2 to p_1 . Their images $(E_{12}(C_2), E_{12}(p_2))$ and $E_{12}(C_1)$ are in standard position.

In the same way, applying the transfer isometry T_{23}^1 to the triple (C_1, p_1) , (C_2, p_2) and (C_3, p_3) with (C_1, p_1) and C_2 in standard position makes (C_1, p_1) and C_3 in standard position.

Proposition 10. *Let (C_1, p_1) , (C_2, p_2) , (C_3, p_3) be a triple of flags in generic position and $\Theta : \mathbb{C} \rightarrow \mathbb{C}$ be the map defined by $\Theta(\rho e^{i\theta}) = \rho e^{i\theta/3}$ for $\rho \in [0, +\infty)$ and $\theta \in (-\pi, \pi]$. Assume that (C_1, p_1) and C_2 are in standard position.*

1. *The transfer isometry T_{23}^1 is given by its lift to $SU(2, 1)$:*

$$\mathbf{T}_{23}^1 = \begin{bmatrix} \mu & 0 & it\mu \\ 0 & \bar{\mu}/\mu & 0 \\ 0 & 0 & 1/\bar{\mu} \end{bmatrix} \text{ where } \mu = \Theta\left(\frac{\delta_{23}^1 \varphi_{13}}{\Phi_{123}}\right) \text{ and } t = \text{Im}\left(\frac{2\delta_{23}^1(\varphi_{23} - \Phi_{132})}{\varphi_{12}\varphi_{23}}\right)$$

2. *The exchange isometry E_{12} is given by its lift to $SU(2, 1)$:*

$$\mathbf{E}_{12} = \begin{bmatrix} \frac{\lambda(z - \bar{z} - |z|^2)}{4|z(z-1)|^2} & \frac{\sqrt{2}\bar{z}\lambda(z - \bar{z} - |z|^2)}{4|z(z-1)|^2} + \frac{\lambda}{\sqrt{2}(z-1)} & \frac{\lambda}{1-z} + \frac{\lambda(z - \bar{z} - |z|^2)^2}{4|z(z-1)|^2} \\ \frac{\bar{\lambda}}{\sqrt{2}\lambda(\bar{z}-1)} & \frac{\bar{\lambda}}{\lambda(\bar{z}-1)} & \frac{\bar{\lambda}(|z|^2 - z - \bar{z})}{\lambda(\bar{z}-1)\sqrt{2}} \\ 1 & \sqrt{2}\bar{z} & -|z|^2 + z - \bar{z} \\ \bar{\lambda} & \frac{\sqrt{2}\bar{z}}{\bar{\lambda}} & \frac{\bar{\lambda}}{\bar{\lambda}} \end{bmatrix}$$

where $z = 1/\bar{m}_{12}$ and $\lambda = 2\Theta(z(z-1))$.

Proof. 1. Suppose that the triple of flags (C_1, p_1) , (C_2, p_2) and (C_3, p_3) is in generic position as it is specified in the proposition, and suppose moreover that (C_1, p_1) and C_2 are in standard position. We can choose polar vectors $\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3$ and representatives $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3$ such that

$$\mathbf{p}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{c}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \mathbf{c}_2 = \begin{bmatrix} 1/\varphi_{12} - 1 \\ \sqrt{2} \\ 1 \end{bmatrix}.$$

The matrix we are interested in stabilizes C_1 and p_1 and sends C_3 to a standard complex line with polar vector

$$\mathbf{c}'_3 = \begin{bmatrix} 1/\varphi_{13} - 1 \\ \sqrt{2} \\ 1 \end{bmatrix}.$$

Call \mathbf{g} the inverse of the expected matrix, and compute the image of \mathbf{c}'_3 by \mathbf{g} :

$$\mathbf{g} = \begin{bmatrix} \lambda & 0 & it\lambda \\ 0 & \bar{\lambda}/\lambda & 0 \\ 0 & 0 & 1/\bar{\lambda} \end{bmatrix} \text{ and } \mathbf{c}_3 = \mathbf{g}\mathbf{c}'_3 = \begin{bmatrix} \lambda(1/\varphi_{13} - 1 + it) \\ \bar{\lambda}\sqrt{2}/\lambda \\ 1/\bar{\lambda} \end{bmatrix}.$$

Computing explicit expressions for $\delta_{23}^1, \varphi_{23}$ and Φ_{123} yields equations for λ and t . A direct computation gives the formulas of the proposition.

2. The second matrix is obtained in three steps: let (C_1, p_1) and (C_2, p_2) be two flags in generic position such that (C_1, p_1) and C_2 are in standard position. We look for a transformation which sends (C_2, p_2) and C_1 to a standard position. We find explicitly a first transformation which sends p_2 to p_1 . Then we compose it with a Heisenberg translation (see remark 6 below) which sends the image of C_2 by the first transformation to C_1 . It remains to find a matrix as in the first part which stabilize the standard flag (C_1, p_1) and sends the image of C_1 by the two first transformations to a standard complex line. The composition of these matrices gives the formula of the proposition. □

Remark 6. A Heisenberg translation is a unipotent parabolic isometry, given by the matrix

$$\begin{bmatrix} 1 & -\bar{w}\sqrt{2} & -|w|^2 + i\tau \\ 0 & 1 & w\sqrt{2} \\ 0 & 0 & 1 \end{bmatrix} \text{ with } w \in \mathbb{C} \text{ and } \tau \in \mathbb{R}.$$

It is an element of the maximal unipotent subgroup of $\text{PU}(2,1)$ fixing the vector $[1 \ 0 \ 0]^T$, which is a copy of the Heisenberg group of dimension 3.

Remark 7. If (C_1, p_1) and p_2 are in standard position, then the \mathbb{R} -plane provided by the first part of proposition 8 is $\mathbf{H}_{\mathbb{R}}^2$. The inversion in that plane is associated to the identity matrix. As a consequence of proposition 7, the associated exchange isometry admits a lift of the form $M_1 \circ \overline{Id} = M_1$ where M_1 is the matrix of a Lagrangian reflection. This shows that $\mathbf{E}_{12}\overline{\mathbf{E}_{12}} = 1$.

5 Decorated triangulations and representations of $\pi_{g,p}$

In this section, we will prove the theorem 1 that is stated in the introduction.

We denote by $\pi_{g,p}$ be the fundamental group of $\Sigma_{g,p}$, a surface of genus g with p punctures x_1, \dots, x_p , assuming $p > 0$. Recall that $\widehat{\Sigma}_{g,p}$ is the universal covering of $\Sigma_{g,p}$.

Provided that the inequality $2 - 2g - p < 0$ is satisfied, the surface $\widehat{\Sigma}_{g,p}$ is homeomorphic to a topological disk and the punctures lift to a $\pi_{g,p}$ -invariant subset X of the boundary of $\widehat{\Sigma}_{g,p}$.

Definition 12. We set

$$\mathfrak{R}_{g,p} = \{(\rho, F)\} / PU(2, 1),$$

where ρ is a morphism from $\pi_{g,p}$ to $PU(2,1)$ and F is a map from X to the set of flags in $\mathbf{H}_{\mathbb{C}}^2$ such that for any $x \in X$ and $g \in \pi_{g,p}$ one has $F(g.x) = \rho(g).F(x)$. The group $PU(2,1)$ acts on F by isometry on the target and acts on ρ by conjugation: this action corresponds to changing the base point in $\pi_{g,p}$.

For convenience, let us recall what will be called a triangulation of Σ , which is sometimes referred to as an ideal triangulation. A triangulation of Σ is an oriented finite 2-dimensional quasi-simplicial complex T with an homeomorphism h from the topological realization $|T|$ of T to Σ which maps vertices to punctures. By quasi-simplicial, we mean that two distinct triangles of T can share the same vertices. By a slight abuse of notation, we will nevertheless refer to a 2-simplex by its vertices.

Given a triangulation T of Σ , we can lift it to a triangulation of $\widehat{\Sigma}_{g,p}$. We thus obtain a triangulation of a disk with vertices on the boundary. Such a triangulation is isomorphic to the Farey triangulation which is a very nice and visual object (see [3]). We may think that any triangulated surface is a quotient of the Farey triangulation. Given a pair (ρ, F) and a triangle Δ of T , we can pick a lift of Δ which has three vertices x, y and z in X .

Definition 13. We will say that the pair (ρ, F) is *generic with respect to T* if for any lifts of triangles of T with vertices x, y and z , the triple of flags $(F(x), F(y), F(z))$ is generic in the sense of definition 10. We denote by $\mathfrak{R}_{g,p}^T$ the subset of $\mathfrak{R}_{g,p}$ made of pairs which are generic with respect to T .

Definition 14. Let T be a triangulation of Σ . We denote by $\mathcal{X}(T)$ the set of triples (φ, Φ, δ) where :

- φ is an $\mathbb{R}_{>0}$ -valued function defined on the set of unoriented edges of T ,
- Φ and δ are \mathbb{C} -valued functions defined on the set of ordered faces of T .

From these data, we define auxiliary invariants in the following way. For any ordered face (i, j, k) of T , we set:

$$\Delta_{ijk} = 1 - \varphi_{ij} - \varphi_{jk} - \varphi_{ik} + \Phi_{ijk} + \Phi_{ikj} \quad (12)$$

$$m_{ij}^k = \frac{1}{\Delta_{ijk}\varphi_{ik}\varphi_{jk}} [\varphi_{ik}\varphi_{jk}(\Phi_{ijk} - \varphi_{ij}) + \varphi_{ik}(\varphi_{ij}\varphi_{jk} - \Phi_{ijk})\delta_{kj}^i + \varphi_{jk}(\varphi_{ij}\varphi_{ik} - \Phi_{ijk})\overline{\delta_{ki}^j} + \Phi_{ijk}(1 - \varphi_{ij})\delta_{kj}^i\overline{\delta_{ki}^j}] \quad (13)$$

The maps φ, Φ and δ must satisfy the following relations for all ordered face (i, j, k) in T :

$$|\Phi_{ijk}|^2 = \varphi_{ij}\varphi_{jk}\varphi_{ki} \quad (14)$$

$$\Phi_{ijk} = \Phi_{jki} = \overline{\Phi_{ikj}} \quad (15)$$

$$\Delta_{ijk} < 0 \quad (16)$$

$$0 = |\delta_{jk}^i|^2(1 - \varphi_{ik}) + 2\text{Re}[\delta_{jk}^i(\Phi_{ikj} - \varphi_{jk})] + \varphi_{jk}(1 - \varphi_{ik}) \quad (17)$$

Moreover, for any edge (i, j) belonging to the faces (i, j, k) and (i, j, l) , we impose the relation

$$m_{ij}^k = m_{ij}^l. \quad (18)$$




Figure 1: A hexagon associated to the triangle (x, y, z) , and elementary matrices associated to its sides.

Before starting the proof of theorem 1, let us give some useful constructions:

Definition 15. Let T be a triangulation of $\Sigma_{g,p}$. By definition, T is a quasi-simplicial 2-complex and there is a homeomorphism h from $|T|$ to $\Sigma_{g,p}$. We consider the following sub-complex of $|T|$:

- vertices are combinations $V_{xy} = \frac{2}{3}x + \frac{1}{3}y$ where x and y belong to the same edge in T ,
- there are two types of simplicial edges: one from V_{xy} to V_{yx} for any edge (x, y) , and one from V_{xy} to V_{xz} for any two adjacent edges (x, y) and (x, z) .
- In each face of T , the edges constructed above draw an hexagon: we add to the sub-complex the corresponding 2-cell.

We denote by HT and call *hexagonation* of T the sub-complex we have obtained. It has the structure of a 2-dimensional CW-complex homeomorphic to $\Sigma_{g,p}$.

Let T be a decorated triangulation of $\Sigma_{g,p}$. We will define from these data a 1-cocycle A in $Z^1(HT, PU(2,1))$. Let s be an oriented edge of HT . Associate to s an elementary matrix A_s as follows:

- If $s = (V_{xy}, V_{yx})$ for some adjacent vertices of T , we set $A_s = E_{12}$ where we replaced m_{12} by m_{xy} .
- If $s = (V_{xy}, V_{xz})$, then we set $A_s = T_{23}^1$ where we replaced all invariants by the decorations corresponding to the bijection $1 \rightarrow x, 2 \rightarrow y, 3 \rightarrow z$.

Lemma 3. *Let T be a decorated triangulation of $\Sigma_{g,p}$. The mapping $s \rightarrow A_s$ is a 1-cocycle of HT with values in $PU(2,1)$.*

Proof. If (x, y, z) is a face of T and if $s_1 \cdots s_6$ are the sides of the associated hexagon, the product $\prod_{i=1}^6 A_{s_i}$ corresponds to an isometry of $\mathbf{H}_{\mathbb{C}}^2$ stabilizing a flag and a complex line. Hence, it is the identity map of $\mathbf{H}_{\mathbb{C}}^2$ (see lemma 2). \square

We now go to the proof of theorem 1.

Proof of theorem 1. We can finally prove the theorem by describing two mappings inverse one of each other. For this purpose, fix a triangulation T of $\Sigma_{g,p}$.

First, we associate to a decoration of T a representation of $\pi_{g,p}$ in $\text{PU}(2,1)$ and an equivariant map F . Assume that T is equipped with a decoration (φ, Φ, δ) and choose a vertex $v = V_{a,b}$ of HT as base point for the fundamental group of $\Sigma_{g,p}$.

Any loop l of $\pi_1(\Sigma_{g,p}, v)$ is homotopic to a sequence $s = s_1, \dots, s_k$ of oriented edges of HT . One can associate to l the element of $\text{PU}(2,1)$ corresponding to the product

$$A_{s_k} \cdots A_{s_1}.$$

Because of the cocycle condition given in lemme 3 above, this isometry does not depend on the choice of the simplicial path homotopic to l . This gives rise to a representation ρ of $\pi_1(\Sigma_{g,p}, v)$ into $\text{PU}(2,1)$. Let us now construct the map F . The choice of base point $v = V_{a,b}$ gives naturally a preferred lift of a and b in $\widehat{\Sigma}_{g,p}$ that we denote by \widehat{a} and \widehat{b} respectively. We choose $F(\widehat{a})$ and $F(\widehat{b})$ such that they are in standard position. Next, any element x of X is parametrized by a path from v to a vertex of HT . We can suppose that this path γ is simplicial. In that way, we set $F(x) = A_\gamma^{-1} \cdot F_0$ where F_0 is the standard flag given by the vectors $\mathbf{c}_1 = [0 \ 1 \ 0]^T$, $\mathbf{p}_1 = [1 \ 0 \ 0]^T$. One checks easily that this map F is equivariant and generic with respect to T and hence, the couple (ρ, F) gives an element of $\mathfrak{R}_{g,p}^T$.

Conversely, given a couple (ρ, F) generic with respect to T , we obtain an element of $\mathcal{X}(T)$ by the following construction. For all edges $[x, y]$ which lift to $[\widehat{x}, \widehat{y}]$ we set $\varphi_{x,y} = \varphi(F(\widehat{x}), F(\widehat{y}))$ and for all triangles $[x, y, z]$ which lift to $[\widehat{x}, \widehat{y}, \widehat{z}]$ we define the Φ and δ invariants of x, y, z as being equal to the corresponding invariants of the triple $(F(\widehat{x}), F(\widehat{y}), F(\widehat{z}))$. These data fit by construction as an element of $\mathcal{X}(T)$. The two maps we have constructed are inverse one of the other. This ends the proof. \square

6 Solving the equations

The aim of this part is to show how to construct solutions of the equations involved in $\mathcal{X}(T)$ in a systematic way. The key lemma is the following:

Lemma 4. *Let m_{12}, m_{23}, m_{31} be three complex number different from 0 and 1. From these numbers, define $\varphi_{i,j} = |m_{ij}/(m_{ij} - 1)|^2$ for all i, j . For any family $(\Phi_{ijk})_{i,j,k}$ of complex numbers satisfying the conditions*

$$\begin{aligned} \Phi_{ijk} &= \Phi_{jki} = \overline{\Phi_{ikj}} \\ |\Phi_{ijk}| &= \sqrt{\varphi_{ij}\varphi_{jk}\varphi_{ki}} \\ \Delta_{ijk} &= 1 - \varphi_{ij} - \varphi_{jk} - \varphi_{ki} + \Phi_{ijk} + \overline{\Phi_{ijk}} < 0 \end{aligned}$$

the following set of equations

$$\begin{aligned} \delta_{jk}^i \delta_{kj}^i &= \varphi_{jk} \\ m_{ij}^k &= \frac{1}{\Delta_{ijk} \varphi_{ik} \varphi_{jk}} (\varphi_{ik} \varphi_{jk} (\Phi_{ijk} - \varphi_{ij}) + \\ &\quad \varphi_{ik} (\varphi_{ij} \varphi_{jk} - \Phi_{ijk}) \delta_{kj}^i + \varphi_{jk} (\varphi_{ij} \varphi_{ik} - \overline{\Phi_{ijk}}) \delta_{ki}^j + \Phi_{ijk} (1 - \varphi_{ij}) \delta_{kj}^i \overline{\delta_{ki}^j}) \\ 0 &= |\delta_{jk}^i|^2 (1 - \varphi_{ik}) + 2\text{Re} [\delta_{jk}^i (\Phi_{ikj} - \varphi_{jk})] + \varphi_{jk} (1 - \varphi_{ik}) \end{aligned}$$

have two distinct solutions in the variables δ_{jk}^i provided that $\varphi_{ij} \neq 1$ for all i and j in $\{1, 2, 3\}$.

Proof. Geometrically, the lemma has the following interpretation: let C_1, C_2, C_3 be three complex lines in generic position. Their position is parametrized by the invariants φ_{ij} and Φ_{ijk} . The hypothesis on these invariants means that any two complex lines are neither orthogonal nor asymptotic.

The invariant m_{ij} specifies a Lagrangian reflexion I_{ij} swapping C_i and C_j in the following way: let p_1 and p_2 be two points in ∂C_1 and ∂C_2 respectively such that $m[(C_1, p_1), (C_2, p_2)] = m_{12}$. Then the second part of lemma 8 tells us that there is a unique lagrangian involution swapping C_1 and C_2 and sending p_1 on p_2 . This involution depends on p_1 and p_2 only through the data of m_{12} . In some sense, the involution I_{12} is the geometric realization of the invariant m_{12} .

A solution of the equations is equivalent to a triple of points p_1, p_2, p_3 lying respectively in $\partial C_1, \partial C_2$ and ∂C_3 such that for all i, j , $I_{ij}p_i = p_j$. Fixing a reference complex line, say C_1 , we see that a solution of the equations is given by a fixed point of the product $I_{31}I_{23}I_{12}$. This product is an anti-holomorphic isometry of C_1 preserving the boundary, hence it has two distinct fixed points on this circle. This proves the lemma.

If some of the φ_{ij} are equal to one, then the corresponding complex lines are asymptotic. The same argument as above applies but the points p_i may lie at the intersection of two complex lines which is not allowed in our settings. Hence, there are less than 2 admissible solutions but there are still some degenerate ones. □

One can apply the preceding lemma for each triangle of a triangulation at the same time. This is described in the following part.

Let T be an ideal triangulation of $\hat{\Sigma}_{g,p}$. We define a decoration space of T which is related to $\mathcal{X}(T)$ but which is somewhat simpler: let $\mathcal{M}(T)$ be the set of triple (φ, Φ, m) where:

- φ and m are functions defined on the set of oriented edges to \mathbb{C} satisfying the following relations:

$$\varphi_{ij} = \left| \frac{m_{ij}}{m_{ij} - 1} \right|^2 \quad \text{and} \quad m_{ji} = \overline{m_{ij}}.$$

Note that the φ invariant is redundant as it is a function of m but we keep it for the coherence of the notation.

- Φ is a \mathbb{C} valued function defined on ordered faces of T satisfying the following equations for all ordered faces (i, j, k) :

$$\Phi_{ijk} = \Phi_{jki} = \overline{\Phi_{ikj}} \quad \text{and} \quad \Delta_{ijk} < 0$$

We denote by $\mathcal{X}^{nd}(T)$ (resp. $\mathcal{M}^{nd}(T)$) the non-degenerate part of $\mathcal{X}(T)$ (resp. $\mathcal{M}(T)$) by which we mean the open set of triples (φ, Φ, δ) such that $\varphi_{ij} \neq 1$ for all i and j (resp. the triples (φ, Φ, m) such that $\varphi_{ij} \neq 1$ for all i, j).

The following proposition is a direct consequence of the preceding lemma.

Proposition 11. *The natural map $\mathcal{X}^{nd}(T) \rightarrow \mathcal{M}^{nd}(T)$ sending (φ, Φ, δ) to (φ, Φ, m) is a covering of order 2^N where N is equal to the number of triangles in T .*

This proposition explains that we can solve the equations in a simple way: we fix arbitrarily the φ and m invariants, and then we solve (with a computer) the remaining equations in δ . The important point given by the proposition is that we are sure to obtain 2^N solutions in the non-degenerate case. The simple structure of the map from $\mathcal{X}^{nd}(T)$ to $\mathcal{M}^{nd}(T)$ should allow us to describe precisely the representation space but it still does not seem to be an easy task and we do not have done it yet.

7 Controlling the holonomy of the cusps

7.1 The general case

Consider a pair $(\rho, F) \in \mathfrak{R}_{g,p}$ and denote as usual by c_i the curve in $\Sigma_{g,p}$ enclosing x_i . Since $\rho(c_i)$ stabilizes a flag $F_i = (C_i, p_i)$, it might be either

- loxodromic, in which case its second fixed point belongs to C_i ,
- parabolic, in which case p_i is its unique fixed point,
- a complex reflection, in which case its restriction to C_i is the identity.

We wish to determine the type of $\rho(c_i)$ in terms of the invariants φ , Φ and δ . The loop c_i encloses the vertex point x_i . It may be written $c_i = \gamma\nu\gamma^{-1}$, where γ is a path connecting the base point to one of the vertices of the hexagonation which is adjacent to the point x_i , and ν is a loop around x_i which is composed of a succession of edges of the hexagonation connecting two edges of the triangulation. As a consequence $\rho(c_i)$, may be written $M_\gamma N M_\gamma^{-1}$, where N is a product of elementary matrices which are all of transfer type (see proposition 10). Write

$$N = \mathbf{T}_k \dots \mathbf{T}_j \dots \mathbf{T}_1,$$

where \mathbf{T}_j is a matrix of transfer type:

$$\mathbf{T}_j = \begin{bmatrix} \mu_j & 0 & it_j \mu_j \\ 0 & \bar{\mu}_j / \mu_j & 0 \\ 0 & 0 & 1 / \bar{\mu}_j \end{bmatrix},$$

and the μ_j 's and t_j 's are written in terms of the invariants φ , Φ and δ as in proposition 10. Computing the product, we obtain

$$N = \begin{bmatrix} \mu & 0 & K \\ 0 & \bar{\mu} / \mu & 0 \\ 0 & 0 & 1 / \bar{\mu} \end{bmatrix},$$

where $\mu = \prod \mu_j$, and

$$K = i \sum_{j=1}^k t_j \frac{\prod_{l=j}^k \mu_l}{\prod_{l=1}^{j-1} \bar{\mu}_l}.$$

We obtain thus that

- N is loxodromic if and only if $|\mu| \neq 1$,
- N is parabolic if and only if $|\mu| = 1$ and $K \neq 0$,
- N is a complex reflection if and only if $|\mu| = 1$ and $K = 0$.

7.2 Type preserving representations of the 1-punctured torus

In this section, we focus on the special case of $\Sigma_{1,1}$, the 1-punctured torus. We first summarize the existing results about this case. We denote the fundamental group of $\Sigma_{1,1}$ by

$$\pi_{1,1} = \langle a, b, c \mid [a, b] \cdot c = 1 \rangle.$$

Recall that a representation of $\pi_{1,1}$ is said to be *type preserving* if and only if $\rho(c)$ is a parabolic isometry. Note that there are two main types of parabolic isometries (see [7, 14] for more details):

1. screw parabolic isometries. These parabolic elements preserve a complex, and thus a flag.
2. horizontal parabolic isometries, which preserves an \mathbb{R} -plane containing their fixed point. These isometries are also called non-vertical Heisenberg translations. No such parabolic isometry appears within the frame of the present work.

As a consequence, we can separate the type-preserving representations of $\pi_{1,1}$ into two types, according to whether $\rho(c)$ is screw parabolic or horizontal parabolic.

1. If $\rho(c)$ preserves a complex line, then it is in the frame of this work. In this case, all the examples known of a discrete, faithful and type preserving representation are obtained by passing to an index 6 subgroup in a discrete, faithful and type preserving representation of the modular group $\mathrm{PSL}(2, \mathbb{Z})$. The latter representations have all been described by Falbel and Parker in [2]. This family of examples consists up to $\mathrm{PU}(2,1)$ of 6 topological components, 4 of which are points, and the two other are segments.
2. If $\rho(c)$ does not preserve a complex line, then it is a consequence of [15] that there exists a unique triple of Lagrangian reflections (I_1, I_2, I_3) such that $\rho(a) = I_1 I_2$ and $\rho(b) = I_3 I_2$. In [14, 16], all these type preserving representations of $\pi_{1,1}$ are described, and, among them, a 3-dimensional family of discrete and faithful representations is identified.

We will now give necessary and sufficient conditions for $\rho(c)$ to be parabolic, in the case it preserves a flag.

A triangulation of a 1-punctured torus is made of two triangles, which we call α and β as on figure 2. We label the vertices as on figure 2. The decoration of this triangulation is given by :

- Triangle α (p_1, p_2, p_3): $\varphi_{12}, \varphi_{23}, \varphi_{31}, \Phi_{123}, \delta_{23}^1, \delta_{31}^2$ and δ_{12}^3 .
- Triangle β (p_1, p_2, p_4) : $\varphi_{12}, \varphi_{24}, \varphi_{41}, \Phi_{124}, \delta_{24}^1, \delta_{41}^2$ and δ_{12}^4 .

Note that because of the identification of the opposite sides of the square, the following relations hold:

$$\varphi_{23} = \varphi_{14} \text{ and } \varphi_{13} = \varphi_{24}.$$

We choose as a basepoint the vertex of the hexagonation which marked by B on figure 2. Let us call ν_{jk}^i the oriented edge of the hexagonation turning around the vertex p_i from the edge $p_i p_j$ edge to the edge $p_i p_k$. As an example, ν_{24}^1 is the oriented segment starting from the point B (see figure 2) and connecting the diagonal to the vertical side $p_1 p_4$. The




Figure 2: Ideal triangulation and hexagonation of a 1-punctured torus. The opposite sides of the square are identified.

homotopy class c is represented by the following sequence of edges, $\nu_{23}^1 \nu_{21}^4 \nu_{31}^2 \nu_{14}^2 \nu_{12}^3 \nu_{42}^1$, to which correspond the product of transfer type matrices $\mathbf{T} = \mathbf{T}_{42}^1 \mathbf{T}_{12}^3 \mathbf{T}_{14}^2 \mathbf{T}_{31}^2 \mathbf{T}_{21}^4 \mathbf{T}_{23}^1$. Denote by μ_{jk}^i and t_{jk}^i the two parameters in the matrix \mathbf{T}_{jk}^i given by proposition 10. The matrix \mathbf{T} is upper triangular, and according to proposition 10, its top left coefficient is

$$\mu = \Theta \left(\frac{\delta_{23}^1 \varphi_{13}}{\Phi_{123}} \frac{\delta_{21}^4 \varphi_{14}}{\Phi_{421}} \frac{\delta_{31}^2 \varphi_{12}}{\Phi_{231}} \frac{\delta_{14}^2 \varphi_{24}}{\Phi_{214}} \frac{\delta_{12}^3 \varphi_{32}}{\Phi_{312}} \frac{\delta_{42}^1 \varphi_{12}}{\Phi_{142}} \right). \quad (19)$$

We simplify this relation using the relations between the invariants ($\varphi_{ij} = \varphi_{ji}$, $|\Phi_{ijk}|^2 = \varphi_{ij} \varphi_{jk} \varphi_{ki}$, and $\delta_{jk}^i \delta_{ki}^j = \varphi_{ij}$). This yields:

$$|\mu|^2 = \frac{|\delta_{23}^1 \delta_{31}^2 \delta_{12}^3|^2}{|\delta_{42}^1 \delta_{14}^2 \delta_{12}^3|^2}.$$

We obtain as a direct consequence the following

Proposition 12. *Let (φ, Φ, δ) be a decorated triangulation of the punctured torus. The holonomy of a loop around the puncture is parabolic or a complex reflexion if and only if*

$$|\delta_{23}^1 \delta_{31}^2 \delta_{12}^3| = |\delta_{42}^1 \delta_{14}^2 \delta_{12}^3| \quad (20)$$

Moreover, the representation associated to the decoration is type preserving if and only if the relation (20) is satisfied and the following relation holds

$$\begin{aligned} 0 \neq & \mu_{42}^1 \mu_{12}^3 \mu_{14}^2 \mu_{31}^2 \mu_{21}^4 \mu_{23}^1 t_{23}^1 + \frac{\mu_{42}^1 \mu_{12}^3 \mu_{14}^2 \mu_{31}^2 \mu_{21}^4}{\mu_{23}^1} t_{21}^4 + \frac{\mu_{42}^1 \mu_{12}^3 \mu_{14}^2 \mu_{31}^2}{\mu_{21}^4 \mu_{23}^1} t_{31}^2 \\ & + \frac{\mu_{42}^1 \mu_{12}^3 \mu_{14}^2}{\mu_{31}^2 \mu_{21}^4 \mu_{23}^1} t_{14}^2 + \frac{\mu_{42}^1 \mu_{12}^3}{\mu_{14}^2 \mu_{31}^2 \mu_{21}^4 \mu_{23}^1} t_{12}^3 + \frac{\mu_{42}^1}{\mu_{12}^3 \mu_{14}^2 \mu_{31}^2 \mu_{21}^4 \mu_{23}^1} t_{42}^1 \end{aligned} \quad (21)$$

The relation (21) is just an explicit version of $K \neq 0$, with K as in the previous section.

References

- [1] E. Falbel and P.V. Koseleff. Rigidity and flexibility of triangle groups in complex hyperbolic geometry. *Topology*, 41, 2002.
- [2] E. Falbel and J. Parker. The moduli space of the modular group. *Inv. Math.*, 152, 2003.
- [3] V. Fock and A.B. Goncharov. Moduli spaces of convex projective structures on surfaces. *ArXiv:math.DG/0405348*, 2006.
- [4] V. Fock and A.B. Goncharov. Moduli spaces of local systems and higher Teichmüller theory. *Publ. Math. Inst. Hautes Etudes Sci.*, 103:1–211, 2006.
- [5] V. Fock and A.B. Goncharov. Dual Teichmüller and lamination spaces. Preprint arXiv:math/0510312v1
- [6] W. Goldman. Representations of fundamental groups of surfaces. In *Geometry and Topology (College Park, Md., 1983/84)*, pages 95–117. Springer, 1985.
- [7] W. Goldman. *Complex Hyperbolic Geometry*. Oxford University Press, Oxford, 1999.
- [8] W. Goldman, M. Kapovich, and B. Leeb. Complex hyperbolic manifolds homotopy equivalent to a Riemann surface. *Comm. Anal. Math.*, 9:61–95, 2001.
- [9] J. Parker and I. Platis. Complex hyperbolic Fenchel-Nielsen coordinates, *Preprint. www.maths.dur.ac.uk/~dma0jrp/*.
- [10] J. Parker and I. Platis. Open sets of maximal dimension in complex hyperbolic quasi-fuchsian space. *J. Diff. Geom.*, 73:319–350, 2006.
- [11] A. Pratoussevitch. Traces in complex hyperbolic triangle groups. *Geometriae Dedicata*, 111:159–185, 2005.
- [12] R. E. Schwartz. Ideal triangle groups, dented tori, and numerical analysis. *Ann. of Math. (2)*, 153:533–598, 2001.
- [13] D. Toledo. Representations of surface groups in complex hyperbolic space. *J. Differ. Geom.*, 29:125–133, 1989.
- [14] P. Will. *Groupes libres, groupes triangulaires et tore épointé dans $PU(2,1)$* . Thèse de l’université Paris VI.
- [15] P. Will. Traces, Cross-ratios and 2-generator Subgroups of $PU(2,1)$. *Preprint*
- [16] P. Will. The punctured torus and Lagrangian triangle groups in $PU(2,1)$. *J. reine angew. Math.*, 602:95–121, 2007.