# A GENERALISATION OF THE HOPF CONSTRUCTION AND HARMONIC MORPHISMS INTO  $\mathbb{S}^2$

S. MONTALDO AND A. RATTO

Abstract. In this paper we construct a new family of harmonic morphisms  $\varphi: V^5 \to \mathbb{S}^2$ , where  $V^5$  is a 5-dimensional open manifold contained in an ellipsoidal hypersurface of  $\mathbb{C}^4 = \mathbb{R}^8$ . These harmonic morphisms admit a continuous extension to the completion  $V^{*5}$ , which turns out to be an explicit real algebraic variety. We work in the context of a generalization of the Hopf construction and equivariant theory.

#### 1. INTRODUCTION

Harmonic maps are critical points of the energy functional

$$
E(\varphi) = \frac{1}{2} \int_M |d\varphi|^2 dv_g,
$$

where  $\varphi: (M, g) \to (N, h)$  is a smooth map between two Riemannian manifolds  $M$  and  $N$ . This is a very wide area of research, involving a rich interplay of geometry, analysis and topology. We refer to [\[3,](#page-9-0) [4\]](#page-9-1) for notation and background on harmonic maps and to [\[2\]](#page-9-2) fore a more recent bibliography.

A geometrically significant sub-family of harmonic maps is represented by the so-called harmonic morphisms. An exhaustive reference for this topic is the book of P. Baird and J.C. Wood [\[1\]](#page-9-3), where characterizing properties and existence of harmonic morphisms are presented in connection with central themes such as harmonic functions and potential theory, together with conformal mappings in the plane and holomorphic maps into a Riemann surface.

For an operational point of view the simplest way to characterize harmonic morphisms is to say that they are just harmonic maps with the additional property that the differential  $d\varphi$  is a horizontally weakly conformal map: this means that, at any point  $x \in M$ , either  $d\varphi_x$  vanishes or

$$
(1.1)\qquad \qquad d\varphi_x: (T_xM)_\mathcal{H} \to T_{\varphi(x)}N
$$

is surjective and conformal. More precisely, in  $(1.1)$ ,  $(T_xM)_{\mathcal{H}}$  denotes the horizontal space  $(\ker(d\varphi_x))^{\perp}$  and it is required that there exists a number  $\Lambda(x) > 0$  such that

<span id="page-0-0"></span>
$$
\Lambda(x)g(X,Y) = h(d\varphi_x(X), d\varphi_x(Y)), \quad \forall X, Y \in (T_xM)_{\mathcal{H}}
$$

<sup>2000</sup> Mathematics Subject Classification. 58E20.

Key words and phrases. The Hopf construction, harmonic maps, harmonic morphisms, equivariant theory.

The function  $\lambda(x) = \sqrt{\Lambda(x)}$  is called the *dilation* (of  $\varphi$  at x). In particular, if  $\varphi$  is a non constant harmonic morphism, then  $m \geq n$ , where  $m = \dim(M)$ and  $n = \dim(N)$ . Moreover, the set of singular points (i.e., those points where  $d\varphi_x = 0$ ) is a closed *polar set*, i.e., it has zero capacity (see [\[1\]](#page-9-3) for details).

In this paper we work in the context of equivariant theory: roughly speaking, that means that we restrict our attention to a class of mappings having enough symmetries to guarantee that harmonicity reduces to the study of a second order ordinary differential equation. We refer to [\[5\]](#page-9-4) for notation, background and examples. More specifically, we shall propose a generalisation of the Hopf construction which gives rise to a new family of harmonic morphisms from a 5-dimensional manifold with singularities onto the Euclidean 2-sphere  $\mathbb{S}^2$ .

## 2. Statement of the main results and related comments

In order to illustrate our framework, we first consider the 3-dimensional ellipsoid

(2.1) 
$$
Q^{*3}(a,b) = \left\{ [x,y] \in \mathbb{C} \times \mathbb{C} : \frac{|x|^2}{a^2} + \frac{|y|^2}{b^2} = 1 \right\}, (a,b > 0)
$$

In [\[5,](#page-9-4) chapter X] the authors study in detail a family of maps

<span id="page-1-2"></span>(2.2) 
$$
\varphi_{k,\ell}: Q^{*3}(a,b) \to \mathbb{S}^2 \subset \mathbb{R}^2 \times \mathbb{R}, \quad k,\ell \in \mathbb{Z}
$$

of the following form (Hopf's construction)

<span id="page-1-0"></span>(2.3) 
$$
[a\sin s \, e^{i\theta_1}, b\cos s \, e^{i\theta_2}] \mapsto [\sin \alpha(s) \, e^{i(k\theta_1 + \ell\theta_2)}, \cos \alpha(s)]
$$

where the function  $\alpha : (0, \pi/2) \to (0, \pi)$  satisfies the boundary conditions

<span id="page-1-1"></span>(2.4) (i) 
$$
\lim_{s \to 0^+} \alpha(s) = 0
$$
, (ii)  $\lim_{s \to \frac{\pi}{2}^-} \alpha(s) = \pi$ .

In particular, it turns out that the map  $\varphi_{k,\ell}$  is harmonic if and only if  $\alpha$  is a solution of the ODE

$$
\alpha'' + (\cot s - \tan s)\alpha' - \frac{h'(s)}{h(s)}\alpha' - h^2(s)\left(\frac{k^2}{a^2\sin^2 s} + \frac{\ell^2}{b^2\cos^2 s}\right)\sin\alpha\cos\alpha = 0
$$

where  $h^2(s) = a^2 \cos^2 s + b^2 \sin^2 s$ . Note, for future comparison, that the left member of [\(2.3\)](#page-1-0) represents a parametrisation of  $Q^{*3}(a, b)$   $(0 \le s \le \pi/2, 0 \le$  $\theta_1, \theta_2 < 2\pi$ ). The boundary conditions [\(2.4\)](#page-1-1) ensure that the solutions have a regular extension through the loci  $s = 0$  and  $s = \pi/2$ , where the coefficients of the ODE become singular. The map  $\varphi_{k,\ell}$  has topological significance because it has Hopf invariant  $k\ell$ , i.e., it represents  $k\ell \in \mathbb{Z} = \pi_3(\mathbb{S}^2)$ . In [\[5\]](#page-9-4) it is proved that, under the additional restriction

<span id="page-1-3"></span>
$$
\frac{a}{b} = \left| \frac{\ell}{k} \right|,
$$

there exist harmonic morphisms of the type [\(2.3\)](#page-1-0).

We are now in the right position to proceed to our generalisation of the previous examples.

We shall consider  $Q^{*5} = Q^{*5}(a_1, a_2, a_3, a_4) \subset \mathbb{C}^4$ ,  $a_1, a_2, a_3, a_4 > 0$ , where  $Q^{*5}$  is the subset of  $\mathbb{C}^4$  parametrised by

<span id="page-2-0"></span>(2.6) 
$$
[a_1 \sin s \, e^{i\theta_1}, a_2 \sin(s + \frac{\pi}{4}) \, e^{i\theta_2}, a_3 \cos s \, e^{i\theta_3}, a_4 \cos(s + \frac{\pi}{4}) \, e^{i\theta_4}]
$$

with  $0 \leq \theta_1, \theta_2, \theta_3, \theta_4 < 2\pi$ ,  $0 \leq s \leq \pi/4$ . Away from the loci  $s = 0$  and  $s = \pi/4$ , [\(2.6\)](#page-2-0) represents a Riemannian manifold

<span id="page-2-3"></span>
$$
(2.7) \t\t (Q5, g),
$$

where  $Q^5 = \mathbb{S}^1 \times \mathbb{S}^1 \times \mathbb{S}^1 \times (0, \pi/4)$ , and the Riemannian metric g induced by the Euclidean structure of  $\mathbb{C}^4(=\mathbb{R}^8)$  is given by

<span id="page-2-4"></span>(2.8) 
$$
g = [a_1^2 \sin^2 s] d\theta_1^2 + [a_2^2 \sin^2(s + \frac{\pi}{4})] d\theta_2^2 + [a_3^2 \cos^2 s] d\theta_3^2 + [a_4^2 \cos^2(s + \frac{\pi}{4})] d\theta_4^2 + h^2(s) ds^2,
$$

with

<span id="page-2-5"></span>(2.9) 
$$
h(s) = \sqrt{a_1^2 \cos^2 s + a_2^2 \cos^2 (s + \frac{\pi}{4}) + a_3^2 \sin^2 s + a_4^2 \sin^2 (s + \frac{\pi}{4})}
$$

**Remark 2.1.** The locus  $s = 0$  is not a topological singularity, since across it  $Q^{*5}$  is locally homeomorphic to  $U^2 \times \mathbb{S}^{\mathbb{I}} \times \mathbb{S}^{\mathbb{I}} \times \mathbb{S}^{\mathbb{I}}$ , where  $U^2$  is an open subset of  $\mathbb{R}^2$ . However, performing the change of local coordinates

 $x_1 = a_1 \sin s \cos \theta_1$ ,  $x_2 = a_1 \sin s \sin \theta_1$ ,  $\theta_2 = \theta_2$ ,  $\theta_3 = \theta_3$ ,  $\theta_4 = \theta_4$ 

it is not difficult to check that the coefficients of the metric tensor  $g$  are continuous, but not differentiable across  $s = 0$ . The same happens across  $s = \pi/4$ .

Bearing in mind the family of maps  $\varphi_{k,\ell}$  of [\(2.2\)](#page-1-2), we now define the class of equivarinat mappings we are interested in  $(k_i \in \mathbb{Z}, i = 1, \ldots, 4)$ :

<span id="page-2-1"></span>
$$
\varphi = \varphi_{k_1,k_2,k_3,k_4} : Q^5(a_1,a_2,a_3,a_4) \to \mathbb{S}^2 \subset \mathbb{R}^2 \times \mathbb{R}
$$

where, to simplify notation, we write  $w$  in place of the generic point  $(2.6)$  of  $Q^5$  and require

(2.10) 
$$
\varphi(w) = [\sin \alpha(s) \; e^{i(k_1 \theta_1 + k_2 \theta_2 + k_3 \theta_3 + k_4 \theta_4)}, \cos \alpha(s)].
$$

Here the function  $\alpha : (0, \pi/4) \to (0, \pi)$  must satisfy the boundary conditions (compare with [\(2.4\)](#page-1-1)):

<span id="page-2-2"></span>(2.11) 
$$
(i) \lim_{s \to 0^+} \alpha(s) = 0, \quad (ii) \lim_{s \to \pi/4^-} \alpha(s) = \pi.
$$

Such boundary conditions ensure that the map  $\varphi$  as in [\(2.10\)](#page-2-1) extends *continuously* across the loci  $s = 0$  and  $s = \pi/4$  in  $Q^{*5}$ . Harmonicity of such  $\varphi$  depends on an ODE for  $\alpha$ : we shall prove that this ordinary differential equation, under suitable restrictions of the type [\(2.5\)](#page-1-3), admits a prime integral which turns out to be essentially equivalent to the horizontal conformality of the map. A qualitative study of this prime integral will lead us to the existence of strictly increasing solutions satisfying [\(2.11\)](#page-2-2). More precisely, our result is

<span id="page-3-2"></span>**Theorem 2.2.** Assume that  $a_i = |k_i|, i = 1, \ldots, 4$ . Then there exist harmonic morphisms  $\varphi : Q^5 \to \mathbb{S}^2$  of the type [\(2.10\)](#page-2-1) which verify the boundary conditions  $(2.11)$ , so admitting a continuous extension to the whole  $Q^{*5}$ .

In view of the previous theorem, it is natural to study the manifold  $Q^{*5}$  a bit more accurately. First of all, we observe (routine verification using [\(2.6\)](#page-2-0)) that all the points of  $Q^{*5}$  satisfy the following set of polynomial equations

<span id="page-3-0"></span>
$$
\begin{cases} \frac{|z_1|^2}{a_1^2} + \frac{|z_3|^2}{a_3^2} = 1\\ \frac{|z_2|^2}{|z_3|^2} \end{cases} (i)
$$

(2.12) 
$$
\begin{cases} \frac{1}{a_1^2} + \frac{1}{a_3^2} = 1 & (i) \\ \frac{|z_2|^2}{a_2^2} + \frac{|z_4|^2}{a_4^2} = 1 & (ii) \\ \left[\frac{|z_3|^2}{a_3^2} - \frac{|z_1|^2}{a_1^2}\right]^2 + \left[\frac{|z_4|^2}{a_4^2} - \frac{|z_2|^2}{a_2^2}\right]^2 = 1 & (iii) \end{cases}
$$

where  $[z_1, z_2, z_3, z_4]$  denote complex coordinates on  $\mathbb{C}^4$ . Next, let us call  $V^{*5} = V^{*5}(a_1, a_2, a_3, a_4)$  the real algebraic variety defined by the set of equations [\(2.12\)](#page-3-0).

We have just remarked that  $Q^{*5} \subset V^{*5}$ . On the other hand, a combined inspection of [\(2.6\)](#page-2-0) and [\(2.12\)](#page-3-0) shows that a parametrisation of the whole  $V^{*5}$ requires, in [\(2.6\)](#page-2-0), to let s vary in the interval  $[0, \pi]$ . Now the singular loci are  $s = 0, s = \pi/4, s = \pi/2$  and  $s = 3\pi/4$   $(s = \pi \text{ coincide with } s = 0)$ . Away from these loci, we have a Riemannian manifold  $(V^5, g)$  as follows (compare with  $(2.7)$ :

(2.13) 
$$
V^5 = \mathbb{S}^1 \times \mathbb{S}^1 \times \mathbb{S}^1 \times \mathbb{S}^1 \times I
$$

where

$$
I = (0, \frac{\pi}{4}) \cup (\frac{\pi}{4}, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \frac{3\pi}{4}) \cup (\frac{3\pi}{4}, \pi)
$$

and the metric g is as in [\(2.8\)](#page-2-4), with  $s \in I$ .

<span id="page-3-1"></span>Next, it becomes natural to study maps

$$
(2.14) \qquad \varphi = \varphi_{k_1, k_2, k_3, k_4} : V^5(a_1, a_2, a_3, a_4) \to \mathbb{S}^2 \subset \mathbb{R}^2 \times \mathbb{R}
$$

defined again by [\(2.10\)](#page-2-1), with the only difference that this time the function  $\alpha : I \to (0, 4\pi)$  satisfies the following set of boundary conditions

(i) 
$$
\lim_{s \to 0^{+}} \alpha(s) = 0, \qquad \lim_{s \to \frac{\pi}{4}^{-}} \alpha(s) = \pi
$$
  
\n(ii) 
$$
\lim_{s \to \frac{\pi}{4}^{+}} \alpha(s) = \pi, \qquad \lim_{s \to \frac{\pi}{2}^{-}} \alpha(s) = 2\pi
$$
  
\n(iii) 
$$
\lim_{s \to \frac{\pi}{4}^{+}} \alpha(s) = 2\pi, \qquad \lim_{s \to \frac{\pi}{2}^{-}} \alpha(s) = 3\pi
$$

<span id="page-4-0"></span>
$$
\lim_{s \to \frac{\pi}{2}^+} \alpha(s) = 2\pi, \quad \lim_{s \to \frac{3\pi}{4}^-} \alpha(s) = 0
$$
\n
$$
\lim_{s \to \frac{3\pi}{4}^+} \alpha(s) = 3\pi, \quad \lim_{s \to \pi^-} \alpha(s) = 4\pi.
$$

(2.15)

**Remark 2.3.** The boundary conditions [\(2.15\)](#page-4-0) ensure that the maps  $\varphi$  in  $(2.14)$  extend continuously to the whole  $V^{*5}$ . In particular, such extension of the  $\varphi$ 's cover 4 times the 2-sphere  $\mathbb{S}^2$ : the singular loci  $s = 0$  and  $s = \pi/2$ are sent to the North pole, while  $s = \pi/4$  and  $s = 3\pi/4$  go into the South pole.

An adaptation of the proof of Theorem [2.2](#page-3-2) enables us to obtain the following

<span id="page-4-4"></span>**Corollary 2.4.** Assume that  $a_i = |k_i|$ ,  $i = 1, ..., 4$ . Then there exist harmonic morphisms  $\varphi: V^5 \to \mathbb{S}^2$  of the type [\(2.14\)](#page-3-1) which satisfy the boundary conditions [\(2.15\)](#page-4-0), so admitting a continuous extension to the whole  $V^{*5}$ .

#### 3. Proof of the results and final remarks

*Proof of Theorem [2.2.](#page-3-2)* We have to show that, under the assumption  $a_i =$  $|k_i|, i = 1, \ldots, 4$ , there exist a function  $\alpha : (0, \pi/4) \to (0, \pi)$  which satisfies the boundary conditions [\(2.11\)](#page-2-2) and such that the associated  $\varphi$  of [\(2.10\)](#page-2-1) is both harmonic and horizontally conformal. To this we need to express, explicitly, conditions which are equivalent to harmonicity and horizontal conformality.

By applying, for instance, the Reduction Theorem (4.13) of [\[5\]](#page-9-4), it is easy to verify that the harmonicity equation for maps  $\varphi$  as in [\(2.10\)](#page-2-1) is given by

<span id="page-4-3"></span>(3.1) 
$$
\alpha''(s) + D(s)\alpha'(s) - G(s, \alpha, \alpha') = 0,
$$

where the functions D and G can be calculated as in  $[5, p. 153]$ . One finds

<span id="page-4-1"></span>(3.2) 
$$
D(s) = \left[\frac{\cos s}{\sin s} + \frac{\cos(s + \frac{\pi}{4})}{\sin(s + \frac{\pi}{4})} - \frac{\sin s}{\cos s} - \frac{\sin(s + \frac{\pi}{4})}{\cos(s + \frac{\pi}{4})}\right] - \frac{h'(s)}{h(s)}
$$

where  $h(s)$  is the function defined in  $(2.9)$ . And also

<span id="page-4-2"></span>(3.3) 
$$
G(s, \alpha, \alpha') = h^2(s) \left[ \frac{k_1^2}{a_1^2 \sin^2 s} + \frac{k_2^2}{a_2^2 \sin^2(s + \frac{\pi}{4})} + \frac{k_3^2}{a_3^2 \cos^2 s} + \frac{k_4^2}{a_4^2 \cos^2(s + \frac{\pi}{4})} \right] \sin \alpha(s) \cos \alpha(s).
$$

Now we simplify [\(3.2\)](#page-4-1) and [\(3.3\)](#page-4-2). Indeed, using three times the trigonometrical identity

$$
(3.4) \t\cot x - \tan x = 2 \cot 2x
$$

we find that [\(3.2\)](#page-4-1) ca be rewritten as

<span id="page-5-0"></span>(3.5) 
$$
D(s) = 4 \cot(4s) - \frac{h'(s)}{h(s)}.
$$

Next, we use the hypothesis  $a_i = |k_i|, i = 1, \ldots, 4$ , in [\(3.3\)](#page-4-2). A routine computation then leads us to

<span id="page-5-1"></span>(3.6) 
$$
G(s, \alpha, \alpha') = h^2(s) \frac{16}{\sin^2(4s)} \sin \alpha(s) \cos \alpha(s).
$$

By way of summary, replacing  $(3.5)$  and  $(3.6)$  in  $(3.1)$ , we have proved that the harmonicity equation for the maps  $\varphi$  as in [\(2.10\)](#page-2-1) is given by

<span id="page-5-3"></span>(3.7) 
$$
\alpha''(s) + \left[4\cot(4s) - \frac{h'(s)}{h(s)}\right] \alpha'(s) - \frac{16h^2(s)}{\sin^2(4s)} \sin \alpha(s) \cos \alpha(s) = 0.
$$

The next step is to work out the condition for horizontal conformality. An orthonormal base of the tangent space  $T_w Q^5$  is

$$
\mathcal{B} = \{e_1, e_2, e_3, e_4, e_5\}
$$

where

<span id="page-5-2"></span>(3.8)  

$$
e_1 = \frac{1}{a_1 \sin s} \frac{\partial}{\partial \theta_1} \qquad e_2 = \frac{1}{a_2 \sin(s + \frac{\pi}{4})} \frac{\partial}{\partial \theta_2} \quad e_3 = \frac{1}{a_3 \cos s} \frac{\partial}{\partial \theta_3}
$$

$$
e_4 = \frac{1}{a_4 \cos(s + \frac{\pi}{4})} \frac{\partial}{\partial \theta_4} \quad e_5 = \frac{1}{h(s)} \frac{\partial}{\partial s}.
$$

We write the Riemannian metric on the range  $\mathbb{S}^2$  as

$$
(3.9) \qquad \qquad h = \sin^2 t \ d\gamma^2 + dt^2
$$

since in [\(2.10\)](#page-2-1) we are parametrising  $\mathbb{S}^2$ , as a subset of  $\mathbb{R}^2 \times \mathbb{R}$ , by means of

(3.10) 
$$
[\sin t \ e^{i\gamma}, \cos t] \quad t \in [0, \pi], \ \gamma \in [0, 2\pi).
$$

Now we can compute the differential for maps  $\varphi$  as in [\(2.10\)](#page-2-1). We find

<span id="page-6-0"></span>
$$
d\varphi(e_1) = \frac{k_1}{a_1 \sin s} \frac{\partial}{\partial \gamma} = \frac{1}{\sin s} \frac{\partial}{\partial \gamma}
$$
  

$$
d\varphi(e_2) = \frac{k_2}{a_2 \sin(s + \frac{\pi}{4})} \frac{\partial}{\partial \gamma} = \frac{1}{\sin(s + \frac{\pi}{4})} \frac{\partial}{\partial \gamma}
$$
  
(3.11)  

$$
d\varphi(e_3) = \frac{k_3}{a_3 \cos s} \frac{\partial}{\partial \gamma} = \frac{1}{\cos s} \frac{\partial}{\partial \gamma}
$$
  

$$
d\varphi(e_4) = \frac{k_4}{a_4 \cos(s + \frac{\pi}{4})} \frac{\partial}{\partial \gamma} = \frac{1}{\cos(s + \frac{\pi}{4})} \frac{\partial}{\partial \gamma}
$$
  

$$
d\varphi(e_5) = \frac{\alpha'(s)}{h(s)} \frac{\partial}{\partial t},
$$

where we have used  $a_i = k_i$ ,  $i = 1, ..., 4$  (the case  $a_i = -k_i$  can be handled similarly and it is left to the interested reader). Now, let  $v = \sum_{i=1}^{5} v_i e_i$  be a generic vector in  $T_w Q^5$ . By using [\(3.11\)](#page-6-0) and the linearity of  $d\varphi$  we find that

$$
(3.12) \quad v \in \ker(d\varphi) \Leftrightarrow \begin{cases} v_5 = 0\\ \frac{v_1}{\sin s} + \frac{v_2}{\sin(s + \frac{\pi}{4})} + \frac{v_3}{\cos s} + \frac{v_4}{\cos(s + \frac{\pi}{4})} = 0 \end{cases}
$$

Therefore, we deduce that a base for the horizontal part is

(3.13) 
$$
\{y, e_5\} \text{ with } y = \frac{e_1}{\sin s} + \frac{e_2}{\sin(s + \frac{\pi}{4})} + \frac{e_3}{\cos s} + \frac{e_4}{\cos(s + \frac{\pi}{4})}.
$$

Since

$$
||y||^2 = \frac{1}{\sin^2 s} + \frac{1}{\sin^2(s + \frac{\pi}{4})} + \frac{1}{\cos^2 s} + \frac{1}{\cos^2(s + \frac{\pi}{4})} = \frac{16}{\sin^2(4s)}
$$

we conclude that an orthonormal base for  $(T_w Q^5)$  is

$$
\mathcal{B}_{\mathcal{H}} = \{y^*, e_5\}
$$

where

<span id="page-6-1"></span>
$$
y^* = \frac{y}{\|y\|} = \frac{\sin(4s)}{4} y.
$$

Because  $d\varphi$  preserves orthogonality between  $y^*$  and  $e_5$ , horizontal conformality reduces to the condition

(3.14) 
$$
||d\varphi(y^*)||^2 = ||d\varphi(e_5)||^2.
$$

Now, using  $(3.8)$  and  $(3.11)$ , we easily find

$$
d\varphi(y^*) = \left[\frac{1}{\sin^2 s} + \frac{1}{\sin^2(s + \frac{\pi}{4})} + \frac{1}{\cos^2 s} + \frac{1}{\cos^2(s + \frac{\pi}{4})}\right] \frac{\sin(4s)}{4} \frac{\partial}{\partial \gamma}
$$
  
(3.15) =  $\frac{4}{\sin(4s)} \frac{\partial}{\partial \gamma}$ .

Consequently, we have

<span id="page-7-0"></span>(3.16) 
$$
\|d\varphi(y^*)\|^2 = \frac{16}{\sin^2(4s)} \left\| \frac{\partial}{\partial \gamma} \right\|^2 = \frac{16}{\sin^2(4s)} \sin^2 \alpha(s).
$$

Moreover, using [\(3.11\)](#page-6-0)

<span id="page-7-1"></span>(3.17) 
$$
\|d\varphi(e_5)\|^2 = \frac{\alpha'^2(s)}{h^2(s)} \left\| \frac{\partial}{\partial t} \right\|^2 = \frac{\alpha'^2(s)}{h^2(s)}.
$$

By way of summary, putting together  $(3.14)$ ,  $(3.16)$  and  $(3.17)$  we obtain that horizontal conformality is equivalent to

<span id="page-7-2"></span>(3.18) 
$$
\frac{16\sin^2\alpha(s)}{\sin^2(s)} = \frac{{\alpha'}^2(s)}{h^2(s)}.
$$

We are now in the right position to end the proof of Theorem [2.2.](#page-3-2) First, taking square roots and re-arranging the terms, we rewrite [\(3.18\)](#page-7-2) in a more convenient form

<span id="page-7-3"></span>(3.19) 
$$
\frac{\alpha'(s)}{\sin \alpha(s)} = \frac{4h(s)}{\sin(4s)}.
$$

Now, applying  $\frac{d}{ds}$  to both members of [\(3.19\)](#page-7-3) and computing, one finds that [\(3.19\)](#page-7-3) implies [\(3.7\)](#page-5-3), or, to say it in words, the condition [\(3.19\)](#page-7-3) of horizontal conformality is a prime integral for the harmonicity equation [\(3.7\)](#page-5-3). Thus it is enough to show that the first order ODE [\(3.19\)](#page-7-3) admits a solution  $\alpha$  :  $(0, \frac{\pi}{4})$  $\frac{\pi}{4}$   $\rightarrow$   $(0, \pi)$  which satisfies the prescribed boundary conditions [\(2.11\)](#page-2-2). Indeed, [\(3.19\)](#page-7-3) can be integrated by separation of variables and that yields a family of solutions

<span id="page-7-4"></span>(3.20) 
$$
\alpha(s) = 2 \tan^{-1} \left[ c \exp \left( \int_{\frac{\pi}{8}}^{s} 4 \frac{h(u)}{\sin(4u)} du \right) \right], \quad s \in (0, \frac{\pi}{4}), \ c > 0.
$$

Since the positive function  $h(u)$  is bounded and bounded away from zero, mere inspection of the solutions [\(3.20\)](#page-7-4) enables us to conclude that the boundary conditions [\(2.11\)](#page-2-2) are fulfilled, so ending the proof of the theorem.

**Remark 3.1.** The explicit meaning of the integration constant c in  $(3.20)$ is

$$
c = \left| \tan \left( \frac{\alpha(\frac{\pi}{8})}{2} \right) \right|.
$$

In particular, the most symmetric solution, i.e. the one with  $\alpha(\frac{\pi}{8})$  $\frac{\pi}{8}$ ) =  $\frac{\pi}{2}$ , corresponds to  $c = 1$ .

Proof of Corollary [2.4.](#page-4-4) Proceeding precisely as in the proof of Theorem [2.2,](#page-3-2) we have again that the horizontal conformality [\(3.19\)](#page-7-3) is a prime integral for the harmonicity equation [\(3.7\)](#page-5-3). Now one needs to show the existence of a solution  $\alpha: I \to (0, 4\pi)$  which satisfies the boundary conditions [\(2.15\)](#page-4-0). The construction of such  $\alpha$  is divided into 4 Steps: in Step 1, one obtains  $\alpha$ :  $(0, \pi/4) \rightarrow (0, \pi)$  which satisfies  $(2.15)$   $(i)$ . Of course, this was done during the proof of Theorem [2.2.](#page-3-2) Step 2 is the construction of a solution  $\alpha : (\pi/4, \pi/2) \to (\pi, 2\pi)$  satisfying [\(2.15\)](#page-4-0) (*ii*). That is achieved again by explicit integration of  $(3.19)$  which gives  $(c > 0)$ 

<span id="page-8-0"></span>(3.21) 
$$
\alpha(s) = 2\pi + 2 \tan^{-1} \left[ -c \exp \left( \int_{\frac{3\pi}{8}}^s 4 \frac{h(u)}{\sin(4u)} du \right) \right], \quad s \in (\frac{\pi}{4}, \frac{\pi}{2}).
$$

Once more, direct inspection of [\(3.21\)](#page-8-0) confirms that the boundary conditions  $(2.15)$  (ii) are verified. In a similar fashion, one constructs solutions  $\alpha$ :  $(\pi/2, 3\pi/4) \rightarrow (2\pi, 3\pi)$  satisfying  $(2.15)$  (iii), and finally  $\alpha : (3\pi/4, \pi) \rightarrow$  $(3\pi, 4\pi)$  which verify  $(2.15)$   $(iv)$ , so completing the proof of the corollary.

**Remark 3.2.** It follows easily from  $(2.12)$   $(i)$  and  $(ii)$  that  $V^{*5}$  is contained into the ellipsoidal hypersurface of equation

(3.22) 
$$
\frac{|z_1|^2}{a_1^2} + \frac{|z_2|^2}{a_2^2} + \frac{|z_3|^2}{a_3^2} + \frac{|z_4|^2}{a_4^2} = 2.
$$

Under the additional restriction

<span id="page-8-1"></span>(3.23)  $a_1 = a_3$  and  $a_2 = a_4$ 

the function  $h(s)$  of  $(2.9)$  is constant, i.e.,

(3.24) 
$$
h(s) \equiv \sqrt{a_1^2 + a_2^2} = A.
$$

In this case the solutions [\(3.20\)](#page-7-4) of Theorem [2.2](#page-3-2) (and, similarly, the solutions of Corollary [2.4\)](#page-4-4) can be made more explicit. For instance, performing an integration leads us to express [\(3.20\)](#page-7-4), under the hypothesis [\(3.23\)](#page-8-1), as

(3.25) 
$$
\alpha(s) = 2 \tan^{-1} \left[ c (\tan 2s)^A \right], \quad s \in (0, \frac{\pi}{4}).
$$

**Remark 3.3.** The integration constant  $c > 0$  in [\(3.20\)](#page-7-4) reflects the fact that our construction provides an infinite family of solutions. A similar situation had been observed in  $[5, p. 186]$ , and  $[1, Chapter 13]$ , where prime integrals occur and produce variations through equivariant harmonic morphisms. It should also be observed that all the solutions in Theorem [2.2](#page-3-2) and Corollary [2.4](#page-4-4) have  $\alpha'(s) > 0$ : the simplest way to verify this claim is direct inspection of [\(3.19\)](#page-7-3). In particular, only the singular loci are sent into the poles of  $\mathbb{S}^2$ .

Remark 3.4. For the purpose of comparison, we also point out that some other examples of harmonic morphisms defined on open subsets of compact manifolds can be found in [\[1,](#page-9-3) p. 410].

Remark 3.5. Since the range of our harmonic morphisms is 2-dimensional, their fibres provide a foliation of  $V^5$  made of 3-dimensional minimal submanifolds diffeomorphic to  $\mathbb{S}^1 \times \mathbb{S}^1 \times \mathbb{S}^1$ .

## **REFERENCES**

- <span id="page-9-3"></span>[1] P. Baird, J.C. Wood. Harmonic Morphisms between Riemannian Manifolds. Oxford Science Publications, 2003.
- <span id="page-9-2"></span>[2] The Harmonic maps Bibliography. <http://people.bath.ac.uk/masfeb/harmonic.html>
- <span id="page-9-1"></span><span id="page-9-0"></span>[3] J. Eells, L. Lemaire. A report on harmonic maps. Bull. London Math. Soc., 10 (1978), 1–68.
- [4] J. Eells, L. Lemaire. Another report on harmonic maps. Bull. London Math. Soc., 10 (1978), 1–68.
- <span id="page-9-4"></span>[5] J. Eells, A. Ratto. Harmonic Maps and Minimal Immersions with Symmetries: Methods of Ordinary Differential Equations Applied to Elliptic Variational Problems. Annals of Mathematics Studies (133), Princeton University Press, 1993.

UNIVERSITÀ DEGLI STUDI DI CAGLIARI, DIPARTIMENTO DI MATEMATICA E INFORMATica, Via Ospedale 72, 09124 Cagliari, Italia

E-mail address: montaldo@unica.it

UNIVERSITÀ DEGLI STUDI DI CAGLIARI, DIPARTIMENTO DI MATEMATICA E INFORMATica, Viale Merello 93, 09123 Cagliari, Italia

E-mail address: rattoa@unica.it