

Experimental magic state distillation for fault-tolerant quantum computing

Alexandre M. Souza¹, Jingfu Zhang¹, Colm A. Ryan¹ & Raymond Laflamme^{1,2}

¹*Institute for Quantum Computing and Department of Physics and Astronomy, University of Waterloo, Waterloo, Canada, ON N2L 3G1.*

²*Perimeter Institute for Theoretical Physics, Waterloo, Canada, ON N2J 2W9.*

Any physical quantum device for quantum information processing is subject to errors in implementation. In order to be reliable and efficient, quantum computers will need error correcting or error avoiding methods. Fault-tolerance achieved through quantum error correction will be an integral part of quantum computers. Of the many methods that have been discovered to implement it, a highly successful approach has been to use transversal gates and specific initial states. A critical element for its implementation is the availability of high-fidelity initial states such as $|0\rangle$ and the *Magic State*. Here we report an experiment, performed in a nuclear magnetic resonance (NMR) quantum processor, showing sufficient quantum control to improve the fidelity of imperfect initial magic states by distilling five of them into one with higher fidelity.

Quantum information processing (QIP)¹⁻⁴ promises a dramatic computational speed-up over classical computers for certain problems. In implementation, the physical quantum devices for QIP are subject to errors due to the effects of unwanted interactions with the environment or quantum control imperfections. In order to be reliable and efficient, quantum computers will need error

correcting or error avoiding methods. One method to achieve fault-tolerant quantum computation is to encode the state of a single quantum bit (qubit) into blocks of several qubits that are more robust to errors. Based on this idea, quantum error correction codes, the theory of fault-tolerant quantum computation and the accuracy threshold theorem have been developed⁵⁻⁷. A key element for fault-tolerant quantum computation is to avoid bad error propagation. One straightforward protocol is to use transversal gates where an error occurring on the k th qubit in one block can only propagate to the k th qubit in the other blocks. A highly successful approach to achieve fault-tolerant universal quantum computation is based on quantum error correcting codes with gates from the Clifford group that can be applied transversally^{8,9}. Unfortunately they are not universal^{10,11} and they must be supplemented with the preparation of not only the $|0\rangle$ state but also another type of state such as a *Magic State*¹²⁻¹⁶. Thus, a critical element for fault-tolerance is the availability of high-fidelity magic states. Consequently, in the pursuit of experimental fault-tolerant quantum computation, it is important to determine whether we have sufficient experimental control to prepare these magic states. In general these will be prepared with some imprecision. The states can be improved by distilling many magic states to produce a fewer number of them which have higher fidelity. Here we report an experiment, performed in a seven-qubit nuclear magnetic resonance (NMR) quantum processor, showing sufficient quantum control to implement a distillation protocol based on the five-bit quantum error correcting code^{12,17} which uses only Clifford gates. The fidelity of imperfect initial magic states is improved by distilling five of them into one with higher fidelity.

Results

Theoretical protocol. The Clifford group is defined as the group of operators that maps the Pauli group onto itself under conjugation. The Pauli group is defined as $^1 \{ \pm \mathbb{1}, \pm i \mathbb{1}, \pm \sigma_x, \pm i \sigma_x, \pm \sigma_y, \pm i \sigma_y, \pm \sigma_z, \pm i \sigma_z \}$ where $\sigma_x, \sigma_y, \sigma_z$, and $\mathbb{1}$ denote the Pauli matrices and identity operator, respectively. The Clifford group on n qubits is a finite subgroup of the unitary group $U(2^n)$ and can be generated by the Hadamard gate H , the phase-shift gate S_{ph} , and the controlled-not gate $CNOT$ represented as

$$H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, S_{ph} = \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}, CNOT = \begin{pmatrix} \mathbb{1} & 0 \\ 0 & \sigma_x \end{pmatrix} \quad (1)$$

in the computational basis $\{|0\rangle, |1\rangle\}$.

An arbitrary one-qubit state can be represented in the Bloch sphere as

$$\rho = (\mathbb{1} + p_x \sigma_x + p_y \sigma_y + p_z \sigma_z) / 2 \quad (2)$$

where p_x, p_y and p_z are the three polarization components of the state. The magic states ¹² are defined as the 8 states with $p_x = \pm 1/\sqrt{3}, p_y = \pm 1/\sqrt{3}, p_z = \pm 1/\sqrt{3}$ (T type) and the 12 states with $p_x = 0, p_y = \pm 1/\sqrt{2}, p_z = \pm 1/\sqrt{2}; p_y = 0, p_z = \pm 1/\sqrt{2}, p_x = \pm 1/\sqrt{2}; p_z = 0, p_x = \pm 1/\sqrt{2}, p_y = \pm 1/\sqrt{2}$ (H type). These states are called "magic" because of their ability, with Clifford gates, to enable universal quantum computation and the ability to be purified, when it has been prepared imperfectly, using only Clifford group operations ¹². In our current work we distill an imperfect magic state into a T -type magic state represented as

$$\rho_M = [\mathbb{1} + (\sigma_x + \sigma_y + \sigma_z)/\sqrt{3}] / 2. \quad (3)$$

To quantify how near a state ρ is to the magic state, we define the m-polarization (polarization in the direction of the magic state)

$$p = 2 \text{Tr}[\rho_M \rho] - 1 = \frac{1}{\sqrt{3}} (p_x + p_y + p_z). \quad (4)$$

The distillation algorithm requires five copies of a faulty magic state $\rho_{in} = \rho^{\otimes 5}$ as the input state. In the original proposal¹², the measurement of four stabilizers S_i ($i = 1, \dots, 4$) is applied to ρ_{in} , where $S_1 = \sigma_x \otimes \sigma_z \otimes \sigma_z \otimes \sigma_x \otimes \mathbb{1}$, $S_2 = \mathbb{1} \otimes \sigma_x \otimes \sigma_z \otimes \sigma_z \otimes \sigma_x$, $S_3 = \sigma_x \otimes \mathbb{1} \otimes \sigma_x \otimes \sigma_z \otimes \sigma_z$ and $S_4 = \sigma_z \otimes \sigma_x \otimes \mathbb{1} \otimes \sigma_x \otimes \sigma_z$. If the outcome of any of these observables is -1 , the state is discarded and the distillation fails. If the results of all the measurement are $+1$, corresponding to the trivial syndrome, one applies the decoding transformation for the five-qubit error correcting code¹⁷ to the measured state and obtains the output state $\rho_{dis} \otimes |0000\rangle\langle 0000|$ where ρ_{dis} has the output m-polarization p_{out} . If the input m-polarization $p_{in} > p_0 = \sqrt{3/7} \approx 0.655$, distillation is possible and $p_{out} > p_{in}$ and produces a state nearer to the magic one. In an iterative manner, it is possible to obtain the output m-polarization approaching 1.

As NMR QIP is implemented in an ensemble of spin systems, only the output of expectation values of ensemble measurements¹⁸ are available. Consequently the above projective measurement of the stabilizers cannot be implemented in our experiment. However, as the decoding operation is just a basis transformation from one stabilizer subspace to another, it is possible to evaluate the result of the distillation after decoding. Therefore, we directly apply the decoding operation to the input state ρ_{in} , and the output state becomes a statistical mixture of 16 possible outcomes

represented as

$$\rho_{out} = \sum_{i=0}^{15} \theta_i \rho_i \otimes |i\rangle\langle i| \quad (5)$$

where θ_i is the probability of each outcome, and $|i\rangle = |0000\rangle, |0001\rangle, \dots, |1111\rangle$, for $i = 0, 1, 2, \dots, 15$, noting $\rho_0 = \rho_{dis}$. Now measuring $|0\rangle$ on all four qubits in $|i\rangle$ indicates a successful purification. We can obtain θ_i and ρ_i using partial quantum state tomography¹⁹.

Experimental results. The data were taken with a Bruker 700 MHz spectrometer. We choose ^{13}C -labelled trans-crotonic acid dissolved in d6-acetone as a seven-qubit register. The structure of the molecule and the parameters of qubits are shown in Table 1. We prepare a labelled pseudo-pure state $\rho_s = \mathbf{00}\sigma_z\mathbf{0000}$ using the method in Ref.²⁰, where $\mathbf{0} = |0\rangle\langle 0|$ and the order of qubits is M, H₁, H₂, C₁, C₂, C₃, C₄. One should note that we are using the deviation density matrix formalism.

We prepare an initial imperfect magic state with three equal polarization components by depolarizing the state $\mathbf{0} = (\mathbb{1} + \sigma_z)/2$. First we apply a $\pi/2$ pulse to rotate the state $\mathbf{0}$ to $(\mathbb{1} + \sigma_x)/2$ and then another $\pi/2$ pulse along direction $[\cos a, \sin a, 0]$ is applied. We use phase cycling to average the x - and y - components of the state to zero, and therefore the polarization of the spin initially in the state $\mathbf{0}$ is reduced. The depolarized state is represented as

$$(\mathbb{1} - \sigma_z \sin a)/2. \quad (6)$$

Finally we apply a rotation with angle $\arccos(1/\sqrt{3})$ about the direction $[-1/\sqrt{2}, 1/\sqrt{2}, 0]$ to

obtain an imperfect magic state

$$\rho = \left[\mathbb{1} + p(\sigma_x + \sigma_y + \sigma_z)/\sqrt{3} \right] / 2 \quad (7)$$

where $p = -\sin a$. The evolution of σ_z in the preparing ρ is shown in Fig. 1. By doing the above operation for qubits M, C₁, C₂, C₃, C₄, respectively, we obtain five copies of the imperfect magic states $\rho_{in} = \rho^{\otimes 5}$. Exploiting partial state tomography, we measure p for each qubit and use the average as the input m-polarization p_{in} for ρ_{in} .

The circuit for the distillation operation is shown in Fig. 2. C₁ carries the distilled state after the distillation. With partial state tomography, we can determine θ_i and ρ_i in equation (5), where $\rho_0 = \rho_{dis}$, from which the output m-polarization p_{out} is obtained. The experimental results for magic state distillation for various p_{in} are shown in Fig. 3, where figures (a) and (b) show the measured p_{out} and θ_0 , respectively. The straight line in figure (a) represents the function $p_{out} = p_{in}$. The data points above the line show the states that have been distilled experimentally.

The implementation time of the distillation procedure is about 0.1 s, a non-negligible amount of time (10%) compared to coherence time (T_2 in Table 1). Hence the decay of the signals due to the limitation of coherence time is an important source of errors. We extract p_{out} by measuring the ratio of $\theta_0 p_{out}$ and θ_0 , where these two factors are obtained by various single coherent terms in a series of experiments (see details in Methods). We have assumed that the terms have the same amount of decoherence. The results of simulations with dephasing rates T_2^* and T_2 are shown in Fig. 3 as the blue squares and red triangles. The simulation results show that the decoherence rates are long enough to allow the distillation and suggest that the deviation of θ_0 from the theoretical

expectation can be mainly attributed to relaxation effects. Additionally, imperfection in the shaped pulses and inhomogeneities of magnetic fields also contribute to errors.

Discussion

We modify the original distillation protocol by avoiding the projective measurement, which is not possible to implement in the NMR QIP's. We exploit partial-state tomography to obtain each output in the mixture of the outcomes after the distillation, and only in a post-processing step do we choose the one we need. Although we could access the $|0000\rangle\langle 0000|$ subspace using a procedure similar to the pseudo-pure state preparation, the method would take a substantially longer amount of time and would be more error-prone. In this work we aim for a quantitative result, i.e. increasing the magic state purity. We need to minimize the readout manipulations to avoid control-error induced distortions of the inferred final state and associated purity. Hence, we limited ourselves to simple high-fidelity readout procedures.

In summary, we have implemented a protocol for distilling magic states based on the five qubit quantum error correction code. We exploit five qubits by controlling a seven-qubit NMR quantum information processor. The experiment shows that we have obtained enough control to purify faulty magic states through distillation.

Methods

Overview of the experiment. To implement the experiment, we exploit standard Isech and

Hermite-shaped pulses as well as numerically optimized GRAPE pulses ²¹ to implement single-spin operations. The GRAPE pulses are optimized to be robust to radio frequency (r.f.) inhomogeneities and chemical shift variations. All pulses are combined in a custom-built software compiler ²². The compiler loads the information about the internal Hamiltonian and the desired unitary transformation from simple predefined building blocks. The blocks are then systematically put together to form a pulse sequence ensuring that the errors in the building blocks do not propagate as the sequence progresses.

R.f. selection. The effect of pulse imperfections due to r.f. inhomogeneities is reduced by selecting signal based on r.f. power ²⁰. The signal selection is achieved by spatially selecting molecules from a small region in the sample. The method is similar to imaging methods ²³ and has been used in previous works ²⁰. Here we substitute the original pulse sequence proposed in Ref.²⁰ by a single GRAPE pulse to optimize the performance. Besides reducing r.f inhomogeneities, the spatial selection of spins can also reduce the static field inhomogeneities and therefore reduces the loss of signal during the experiment. We have found that the effective relaxation time (T_2^*) of spins after the r.f. selection increases significantly, e.g., up to a factor 2 for some spins.

Partial state tomography. We use the spectra obtained from the labelled pseudo-pure state $\rho_s = 00\sigma_z0000$ shown in Fig. 4 as a phase reference and to normalize the signals in C_1 and C_2 spectra, for measuring the initial and output m-polarization. To obtain the reduced density matrix of C_1 through the partial state tomography, we expand equation (5) as a sum of product operators ¹⁹, and represent ρ_i as

$$\rho_i = \frac{1}{2} (I + p_{i,x}\sigma_x + p_{i,y}\sigma_y + p_{i,z}\sigma_z). \tag{8}$$

In the expansion there are 128 terms that are required to determine by the experiment.

The coefficients of such expansion can be directly related to the measurable spectral amplitudes ¹⁹. On the other hand, such coefficients can also be related to the relevant parameters of (5), i.e, $p_{i,x}, p_{i,y}, p_{i,z}$ and θ_i for $i = 0, 1, 2, \dots, 15$. The relation between these parameters and the NMR observables can be expressed by the set of linear equations

$$C = A \times R. \quad (9)$$

The n th element, $C(n)$, of the column vector C is the coefficient related to the operator $\sigma_z^{\bar{n}_4} I \sigma_z s \sigma_z^{\bar{n}_1} \sigma_z^{\bar{n}_2} \sigma_z^{\bar{n}_3}$ with the order of qubits $M, H_1, H_2, C_1, C_2, C_3, C_4$, where s can be one element of the Pauli group $\{\sigma_x, \sigma_y, \sigma_z, I\}$ and the vector $\bar{n} = (\bar{n}_1, \bar{n}_2, \bar{n}_3, \bar{n}_4)$ is the four digit binary representation of the integer $n - 1$. For $s = \sigma_x, \sigma_y, \sigma_z$, and I , $R(n) = \theta_n p_{n,x}, \theta_n p_{n,y}, \theta_n p_{n,z}$ and θ_n , respectively. The elements of the matrix A are given by

$$A(k, m) = \prod_{i=1}^4 (-1)^{\bar{k}_i \bar{m}_i}. \quad (10)$$

Providing that we have all necessary coefficients measured, we can reconstruct the distilled states using the following approach. First, we fit the NMR spectral lines to the yield complex amplitudes for measuring all necessary coefficients ¹⁹. Fig. 5 illustrates the spectra of C_1 after the completion of distillation for $p_{in} = 0.95$, where the experimentally measured, fitted, and ideal spectra are shown as the red, black, and blue curves, respectively. Then the state (5) is reconstructed by solving the set of equations (9). Our calculation shows that four readout operations are sufficient to determine all coefficients: first, read out on C_1 ; second, read out on C_1 after the application of a $\pi/2$ pulse; third, read out on C_2 after the application of a $\pi/2$ pulse; and forth, read out on C_2 after

a polarization transfer from H_1 to C_2 . The last two readout operations are sufficient to measure all θ_i , and the first two are used to measure $\theta_i \rho_i$. The errors for the coefficients, as well errors for $p_{i,x}$, $p_{i,y}$, $p_{i,z}$ and θ_i , are estimated from the uncertainty of the fitting parameters. The measured initial and output m-polarization, as well as θ_i and ρ_i , are listed as Supplementary Tables S1-S10 in the Supplementary Information²⁵. The comparison of the various measured ρ_0 with the theory is shown as equations (1-7) in the Supplementary Methods in the Supplementary Information.

1. Nielsen, M. A. & Chuang, I. L. *Quantum Computation and Quantum Information*. Cambridge University Press, Cambridge, England (2000).
2. Preskill, J. *Introduction to Quantum Computation and Information*, edited by Lo, H. K., Popescu, S. & Spiller, T. World Scientific Publishing Company, Singapore (2006).
3. Kaye, P., Laflamme, R. & Mosca, M. *An Introduction to Quantum Computing*. Oxford University Press, Oxford, England (2007).
4. Preskill, J. Reliable quantum computers. *Proc. R. Soc. Lond. A* **454**, 385-410 (1998).
5. Aharonov, D. & Ben-Or, M. Fault tolerant quantum computation with constant error. *Proceedings of the 29th Annual ACM Symposium on Theory of Computing*, p. 176, ACM Press (1997).
6. Kitaev, A. Yu. Quantum computations: algorithms and error correction. *Russ. Math. Surv.* **52**, 1191-1249 (1997).
7. Knill, E., Laflamme, R. & Zurek, W. H. Resilient quantum computation. *Science* **279**, 342-345 (1998).

8. Knill, E. Quantum computing with realistically noisy devices. *Nature* **434**, 39-44 (2005).
9. Gottesman, D., Quantum Error Correction and Fault Tolerance. *Encyclopedia of Mathematical Physics*, edited by Francoise, J.-P., Naber, G. L. & Tsou, S. T., pp. 196-201, Elsevier, Oxford (2006).
10. Zeng, B., Cross, A. W. & Chuang, I. L., Transversality versus Universality for Additive Quantum Codes. e-print arXiv: 0706.1382 (2007).
11. Chen, X., Chung, H., Cross, A. W., Zeng, B. & Chuang, I. L. Subsystem stabilizer codes cannot have a universal set of transversal gates for even one encoded qudit. *Phys. Rev. A* **78**, 012353 (2008).
12. Bravyi, S. & Kitaev, A. Universal quantum computation with ideal Clifford gates and noisy ancillas. *Phys. Rev. A* **71**, 022316 (2005).
13. Campbell, E. T. & Browne, D. E., Bound States for Magic State Distillation in Fault-Tolerant Quantum Computation. *Phys. Rev. Lett.* **104**, 030503 (2010).
14. Campbell, E. T. & Browne, D. E., On the Structure of Protocols for Magic State Distillation. *Lecture Notes in Computer Science* **5906**, 20-32 (2009).
15. Dam, W. V., Howard, M., Tight Noise Thresholds for Quantum Computation with Perfect Stabilizer Operations. *Phys. Rev. Lett.* **103**, 170504 (2009).
16. Reichardt, B. W., Quantum universality by state distillation. *Quantum Information and Computation* **9**, 1030-1052 (2009).

17. Laflamme, R., Miquel, C., Paz, J. P. & Zurek, W. H. Perfect Quantum Error Correcting Code. *Phys. Rev. Lett.* **77**, 198-201 (1996).
18. Jones, J. A., NMR Quantum Computation: A Critical Evaluation. *Fortschr. Phys.* **48**, 909-924 (2000).
19. Leskowitz, G. M. & Mueller, L. J. State interrogation in nuclear magnetic resonance quantum-information processing. *Phys. Rev. A* **69**, 052302 (2004).
20. Knill, E., Laflamme, R., Martinez, R. and Tseng, C.-H. An algorithmic benchmark for quantum information processing. *Nature* **404**, 368-370 (2000).
21. Khaneja, N., Reiss, T., Kehlet, C., Schulte-Herbruggen, T. & Glaser, S. J. Optimal control of coupled spin dynamics: design of NMR pulse sequences by gradient ascent algorithms. *J. Magn. Reson.* **172**, 296-305 (2005).
22. Ryan, C. A., Negrevergne, C., Laforest, M., Knill, E. & Laflamme, R. Liquid-state nuclear magnetic resonance as a testbed for developing quantum control methods. *Phys. Rev. A* **78**, 012328 (2008).
23. Maffei, P., Elbayed, K., Broudeau, J. & Canet, D. A Slice Selection in NMR Imaging by Use of the B_1 Gradient along the Axial Direction of a Saddle-Shaped Coil. *J. Magn. Reson.* **95**, 82-386 (1991).
24. Hubbard, A. On Magic State Distillation using Nuclear Magnetic Resonance. *Master's thesis*, University of Waterloo, Waterloo, Canada (2008).

25. Available at <http://www.nature.com/naturecommunications>.

Acknowledgments We thank the Premier Discovery Award from the Government of Ontario, Industry Canada and the Canadian Institute for Advanced Research for financial support. We thank Nathan Babcock, Josh Slater, Stephanie Simmons and Martin Laforest for discussions and their contributions in the first stage of this experiment, and also thank Camille Negrevergne and Adam Hubbard for laying the groundwork for the experimental work in realization of distillation. J.-F.Z thanks Bei Zeng for helpful discussion.

Competing Interests The authors declare that they have no competing financial interests.

Correspondence Correspondence and requests for materials should be addressed to J.-F.Z. (j87zhang@iqc.ca).

Author contributions A.M.S. and J.-F.Z. performed the experiment and the numerical simulations, C. A.R. helped design the experimental scheme and wrote the control software; R.L. conceived the ideas and supervised the experiment. All authors contributed to the writing of the paper, discussed the experimental procedures and results.

	M	H ₁	H ₂	C ₁	C ₂	C ₃	C ₄
M	-1309						
H ₁	6.9	-4864					
H ₂	-1.7	15.5	-4086				
C ₁	127.5	3.8	6.2	-2990			
C ₂	-7.1	156.0	-0.7	41.6	-25488		
C ₃	6.6	-1.8	162.9	1.6	69.7	-21586	
C ₄	-0.9	6.5	3.3	7.1	1.4	72.4	-29398
T ₂ (s)	0.84	0.85	0.84	1.27	1.17	1.19	1.13
T ₂ [*] (s)	0.61	0.57	0.66	1.04	0.66	1.16	0.84

Table 1: **Characteristics of the molecule of trans-crotonic acid.** Molecular structure is shown as Fig. 6. The chemical shifts and J-coupling constants (in Hz) are on and below the diagonal in the table, respectively. The transversal relaxation times T_2 measured by a Hahn echo and T_2^* calculated by measuring the width of the peaks through fitting the spectra are listed at the bottom. The chemical shifts are given with respect to reference frequencies of 700.13 MHz (protons) and 176.05 MHz (carbons). The molecule contains nine weakly coupled spin half nuclei but consists of a seven qubit system since the methyl group can be treated as a single qubit using a gradient-based subspace selection ²⁰.




Figure 1: **The evolution of σ_z in the preparation of a faulty magic state in Bloch sphere, where arrows represent the states of the qubit.** **a**, A $\pi/2$ rotation along y-axis transforms σ_z (blue) to σ_x (green). **b**, Another $\pi/2$ rotation along direction $[\cos a, \sin a, 0]$ transforms σ_x (green) to $\sigma_x \cos^2 a + \sigma_y \cos a \sin a - \sigma_z \sin a$ (black). In phase cycling we apply the second $\pi/2$ rotation by changing a to $\pi + a$ to transform σ_x to $\sigma_x \cos^2 a + \sigma_y \cos a \sin a + \sigma_z \sin a$. After averaging the x - and y - components to zero, the polarization is reduced to $-\sigma_z \sin a$, shown as the yellow arrow in figure **c**, noting that $a \in [\pi, 3\pi/2]$. **d**, A final rotation with angle $\arccos(1/\sqrt{3})$ along $[-1/\sqrt{2}, 1/\sqrt{2}, 0]$ transforms $-\sigma_z \sin a$ (yellow) to $-\sin a(\sigma_x + \sigma_y + \sigma_z)/\sqrt{3}$ (red).




Figure 2: **Gate sequence for Magic State distillation.** The sequence is constructed from the five qubit error correction code ¹⁷ where $\pm X = \exp(\mp i\pi\sigma_x/4)$, $\pm Y = \exp(\mp i\pi\sigma_y/4)$, and $Z_\alpha = \exp(-i\alpha\sigma_z/2)$. Qubits labelled as M, C₁, C₂, C₃ and C₄ are used to encode the five copies of the initial state. Due to the nature of the algorithm, the carbon C₁ contains the distilled magic state only when M, C₂, C₃ and C₄ are in the $|0000\rangle$ state. It is important to emphasize that all gates are Clifford gates. The refocussing pulses (which also decouple H_1 and H_2) are not shown.




Figure 3: **Experimental results after the completion of magic state distillation.** Output m-polarization of the faulty magic state (a) and the probability θ_0 (b) of finding this state in the mixture of outcomes [see equation (5)] as a function of the input m-polarization of the initial faulty magic state. The experimental data are represented by the filled circles and the error bars are estimated from the uncertainty of the fitting parameters. The line in figure (a) represents the function $p_{out} = p_{in}$. The experimental points above the line show the states that have been distilled, while the points below the line show the states that cannot be distilled in the protocol. The theoretical prediction is represented by the black solid curves. The blue squares and red triangles, connected by dashed lines for visual convenience, are the simulation results where the dephasing rates are chosen as T_2^* and T_2 (see Table 1), respectively. The effective T_2 during the experiment should be similar to the Hahn echo T_2 . The deviation can be attributed to other error sources (see text). The dephasing times of H_1 and H_2 actually do not influence the results because H_1 and H_2 can be effectively assumed in $\mathbf{0}$ and σ_z during the whole experiment, respectively.




Figure 4: **NMR spectra.** **a,b**, Spectra of C_1 and C_2 obtained by $\pi/2$ readout pulses when the system lies in the labelled pseudo-pure state $\rho_s = 00\sigma_z0000$. The vertical axes have arbitrary but the same units.




Figure 5: **Spectra of C_1 after the completion of distillation for $p_{in} = 0.95$.** The spectra are divided in four different parts shown as figures (a-d) for better visualization. The vertical axes have arbitrary but the same units. The experimentally measured, fitting, and ideal spectra are shown as the red, black, and blue curves, respectively.




Figure 6: **Molecular structure of trans-crotonic acid.**