Nucleon-Nucleon Scattering and Large Nc QCD

Boris A. Gelman**

New York City College of Technology, The City University of New York E-mail: bgelman@citytech.cuny.edu

Nucleon-nucleon scattering observables are discussed in the context of large N_c QCD. As is well known, the baryon spectrum in the large Nc limit exhibits contracted $SU(2N_f)$ spin-flavor symmetry. This symmetry can be used to derive model-independent relations between proton-proton and proton-neutron total cross sections. These relations are valid in the kinematic regime in which the relative momentum of two nucleons is of order of N_c . In this semiclassical regime the nucleon-nucleon scattering can be described in the time-dependent mean field approximation. These model-independent results are compared to experimental data for spin-independent and polarized total nucleon-nucleon cross sections.

Xth Quark Confinement and the Hadron Spectrum, October 8-12, 2012 TUM Campus Garching, Munich, Germany

*Speaker.

)(TEED)

[†]The work on which this talk is based was done in collaboration with T.D Cohen [1, 2].

1. Introduction

Description of the nucleon-nucleon interaction, the basic ingredient of nuclear physics, directly from quantum chromodynamics (QCD) is a daunting task. The breakdown of the pertubative expansion in terms of the QCD gauge coupling g necessitates the use of alternative methods. One such method proposed by 't Hooft in 1974 is to consider a QCD-like theory with the number of colors N_c and the gauge group $SU(N_c)$, large $N_c QCD$ [3]. The observables in large $N_c QCD$ are expanded in powers of $1/N_c$ around the large N_c limit, $g \rightarrow 0$, $N_c \rightarrow \infty$ and finite g^2N_c . In addition, it is assumed that large $N_c QCD$ is a confining theory and the asymptotic states are $SU(N_c)$ singlets. Despite our inability at present to evaluate even the leading order terms, a great deal of insight comes from knowledge of the scaling of hadronic observables in powers of $1/N_c$. The phenomenological implications of large $N_c QCD$ are essentially topological in nature.

The description of the meson and baryon observables in the large N_C limit requires different methods. Formally, it is due to the fact that the correlation functions in the meson sector have a smooth expansion in powers of $1/N_c$ while the correlation functions in the baryon sector diverge in the large N_c limit. Physically, it is due to the fact that as shown by 't Hooft, the QCD in the large N_c limit is a theory of an infinite number of stable non-interacting mesons.¹ The meson masses and sizes are independent of N_c . This picture is phenomenologically satisfactory since in the real world mesons interact weakly.

QCD is also a theory of strongly interacting baryons, the states carrying quantum numbers of the odd number of quarks. As was shown by Witten [4], a consistent large N_c description of strongly interacting baryons is possible. Remarkably, the same feature which on the one hand makes an analysis of the baryons in the large N_c limit far more challenging than that of mesons, on the other hand allows one to apply a well-known method of nuclear physics, the semiclassical mean-field theory. Indeed, as argued by Witten, baryons in the large N_c limit contain N_c quarks with n-quark force scaling as N_c^{1-n} . Thus, one can treat this weakly interacting many-body state in a mean-field approximation. Unfortunately, the explicit treatment is only available for heavy non-relativistic quarks, in which case the mean-field treatment corresponds to the Hartree approximation [5]. The picture that arises from such a treatment is that of a baryon with a mass of order of N_c and a size and shape which are independent of N_c . Despite the fact that explicit mean-field treatment in the case of the light quarks is unknown, the large N_c scaling for baryon observables is expected to be valid.

Witten realized [4] that the above scaling of mesons and baryons indicates that the baryons in large N_c limit arise as quantized soliton-like configurations of mesonic fields. A particular model which satisfies the large N_c scaling is a well-known Skyrme model [6]. In this model the baryons appear as quantized skyrmions, the topological solitons of a particular non-linear mesonic lagrangian[7]. The stability of baryons as quantum solitons is due to the existence of conserved topological current [8, 9].

In addition to a single-baryon sector, it is of great interest to consider baryon-baryon interaction in the large N_c limit. Since the baryon mass diverges in the large N_c limit the baryon-baryon scattering observables don't have a smooth limit for scattering at fixed center-of-mass energy and momentum transfer. For such momentum, $p \sim N_c$, one instead focuses on the potential between

¹Additionally, the spectrum of large N_c QCD contains glueballs with vanishing mixing to mesons

two baryons which scales as N_c . In this context, the large N_c scaling rules had been used to analyze the spin-flavor structure of the nucleon-nucleon potential [10]. In particular, it was shown that at leading order the nucleon-nucleon potential is symmetric under contracted SU(4) spin-flavor symmetry [11]. In addition, one can also address a question of consistency of the meson-exchange picture of the nucleon-nucleon potential [12, 13].

The focus of the present talk is on the kinematic regime corresponding to a fixed center-ofmass velocity. In this case, both the kinetic and potential energy of two baryons are of order N_c , and thus one expects a smooth limit to exist for the scattering cross-section. As argued by Witten [4], here, as in the case of the single-baryon sector, the appropriate framework is the mean-field description. However, in this case one has to use *time dependent* mean-field theory (TDMFT). As was shown in [2], one can discuss the spin-flavor structure of the total nucleon-nucleon cross section at leading order in large N_c expansion. As in the case of the nucleon-nucleon potential, the emergent contracted SU(4) spin-flavor symmetry leads to certain relations between total protonproton and proton-neutron cross sections. It will be shown in section 3 that these relations satisfy the behavior of the experimental nucleon-nucleon cross sections at the center-of-mass energies of order of a few GeV.

2. Time Dependent Mean Field Theory Framework

The goal here is to show that time-dependent mean-field theory (TDMFT) framework valid in the large N_c limit, and contracted $SU(2N_f)$ symmetry, where N_f is the number of light quark flavors, to make model-independent predictions about the spin-isospin structure of the total nucleonnucleon cross sections. As discussed above, TDMFT treatment is a valid framework for the nucleon-nucleon scattering when the center-of-mass transfer momentum is $p \sim N_c$. Since the nucleon size and hence the size of the interaction region are of order of N_c^0 , the scattering in this kinematic regime is semi-classical.

The description of interaction by TDMFT methods requires time-averaging over all field configurations consistent with the initial state of two nucleons. This precludes one from being able to calculate the S-matrix elements [14, 15]. However, as shown in [2], there are certain inclusive nucleon-nucleon observables which can be evaluated in TDMFT framework. One such observable can be formed from conserved baryon current whose expectation value in the initial two-nucleon state can be in principle evaluated. The expectation values of the baryon current can be related to the inelastic differential cross section.

In TDMFT framework each quark and gluon field of two nucleons move in a time-dependent field created by all other quarks and gluons. These equations are not known explicitly. As a result, one can not determine the nucleon-nucleon cross section even in the large N_c limit. However, it is possible to determine the spin-isospin dependence of the cross section using the contracted SU(4) symmetry valid in the large N_c limit. Since the focus here on spin-isospin dependence of the nucleon-nucleon scattering, one can use the Skyrme model which encapsulates the spin-flavor structure of large N_c baryons [16].

In the Skyrme model the nucleon dynamics is described in terms of classical soliton configurations built out of pion fields. A convenient form for such a soliton is given by SU(2)-valued matrices $U_h(\vec{r}) = \exp(i\vec{\tau}\hat{n}F(r))$, where τ^a are Pauli matrices, $\hat{n} = \vec{r}/r$ and F(r) is the magnitude of the pion field, $F(r) = |\vec{\pi}(r)|$ [7]. Such classical configurations impose correlations between spacial and isospin rotations and they are referred to as *hedgehogs*. The baryons appear after the quantization of classical hedgehogs. This is done by quantizing the slow rotation of the hedgehog in isospin space, $A^{\dagger}(t)U_h(\vec{r})A(t)$, given by the time-dependent SU(2) matrix A(t). These rotations describe the slow collective degrees of freedom of the hedgehog, the zero-modes [17, 18]. After quantization, the generators of these vibrations proportional to $Tr(\tau^a A(t)A^{\dagger}(t))$ correspond to the spin and isospin quantum numbers of the the ground-state band of states, $I = J = 1/2, 3/2, ...N_c/2$ in the large N_c limit. The first two states correspond to nucleon and Δ baryons. The masses of these states are degenerate up to the terms $I(I+1)/M_B \sim 1/N_c$. This is a representation of the contracted spin-flavor symmetry SU(4) in the context of the Skyrme model. The spin-isospin dependence of the wave-function of these states is given by Wigner matrices $D_{m,m'}^{I=J}(A)$ where m,m' are the third components of spin and isospin respectively. Note that the Wigner matrices are functions of parameters of collective rotations A and not of $\dot{A}(t)$.

A crucial consequence of the above semi-classical analysis of the single-baryon sector in the Skyrme model is the appearance of the scale separation in the dynamics of the collective degrees of freedom described by the Wigner matrices $D_{m,m'}^{I=J}(A)$ at leading order in $1/N_c$ and intrinsic degrees of freedom which describe all other non-collective excitations which include excited states of the ground states baryons and mission of virtual and real mesons. The frequency of the collective excitations are of order $1/N_c$ while that of the *intrinsic* excitations are of order N_c^0 . This scale separation enables an adiabatic or treatment of the collective degrees in the context of TDMFT treatment of the rotational and vibrational excitations of molecules. There the slow degrees of freedom correspond to vibrations of atomic nuclei in the averaged field produced by electrons whose motion represent the intrinsic excitations.

To obtain an observable describing the nucleon-nucleon scattering one can start with a function which describes the initial state of two well separated hedgehogs corresponding to the initial state of two nucleons. As discussed in [2], in the context of the Skyrme model it is convenient to choose a conserved baryon current $J^{\mu}(\vec{r},t;A_1,A_2,v,\vec{b},\hat{n})$. In the Skyrme model it is a topological invariant. The dependence of the current on the collective degrees of freedom are described by the variables, $A_1, A_2, v, \vec{b}, \hat{n}$ where $A_1, A_2, v, \vec{b}, \hat{n}$ define the spin and isospin configuration, the center-of-mass speed, impact parameter and the unit vector along the direction separating the centers of the two hedgehogs at the initial moment. The initial distance is not indicated. Such parametrization corresponds to the semi-classical description of the scattering which as discussed above is valid in the large N_c limit. The functional dependence of this current can only be determined once the explicit form of TDMFT equations is known. However, as shown below one does not need to know these equations to determine the spin-isospin structure of the current under the spin and isospin rotations which is determined by the contracted sin-flavor symmetry.

As shown in [1, 2], the classical current can be turned into a differential cross section by integrating the current $J^{\mu}(\vec{r},t;A_1,A_2,v,\vec{b},\hat{n})$ over time and the impact parameters,

$$\frac{d\sigma_{\rm inc}(v,A_1,A_2;\theta,\phi)}{d\Omega} = \lim_{R \to \infty} R^2 \int_0^\infty db \left(2\pi b\right) \int_0^\infty dt \hat{r}(\theta,\phi) \cdot \vec{J}\left(R\hat{r}(\Omega),t;A_1,A_2,v,\vec{b},\hat{n}\right), \quad (2.1)$$

The above equations gives the probability for one hedgehog to emerge in a cone with a solid angle $d\Omega$ around a direction given by polar angles θ and ϕ . The current in Eq. (2.1) is normalized as to give the total baryon number two. The time t = 0 in Eq. (2.1) corresponds to the time at which two hedgehogs have the smallest separation. The integral in Eq. (2.1) can be explicitly evaluated only when TDMFT equations are known. It is also important to note that the probability in Eq. (2.1) is integrated over all outgoing meson degrees of freedom. In the final analysis it will give the inelastic cross section.

To turn the probability in Eq. (2.1) into a nucleon-nucleon cross section one needs to evaluate an expectation value of $d\sigma_{inc}/d\Omega$ in an initial two-nucleon state described by spin and isospin projections m_1, m_1^I and m_2, m_2^I on the direction given by the unit vector \hat{n} . It can be done due to the scale separation between the collective and intrinsic degrees of freedom discussed above. Indeed, the semiclassical quantization of the baryon current in Eq. (2.1) leads the appearance of the terms proportional to J^2/M_N and I^2/M_N , where M_N is the nucleon mass. These terms represent coupling between the collective and intrinsic excitations. However, as discussed above these terms are of order $1/N_c$ and do not contribute at leading order in the $1/N_c$ expansion. This result allows one to find expectation values of the inclusive cross section using the superposition of the initial hedgehogs described by the collective variables A_1 and A_2 weighted by the corresponding Wigner D-matrix. In other words, the spin-isospin part of the nucleon wave function in the initial state with given quantum numbers m, m^I is

$$|m^{J}, m^{I}\rangle = \int dA D_{m,m^{I}}^{1/2}(A) |A\rangle,$$
 (2.2)

where $|A\rangle$ represents a hedgehog with particular orientation in spin-isopsin space, $D^{1/2}$ is the Wigner D-matrix describing spin-isospin coordinates of the nucleon, and the integral is taken over the space of the collective coordinates.

Using Eq. 2.2 one can find the inclusive (integrated over all mesons in the final state) nucleonnucleon differential cross section at leading order in $1/N_c$ expansion,

$$\frac{d\sigma^{(m_1,m_1',m_2,m_2')}(v,\theta,\phi)}{d\Omega} = \int dA_1 dA_2 |D_{m_1,m_1'}^{1/2}(A_1)|^2 |D_{m_2,m_2'}^{1/2}(A_2)|^2 \frac{d\sigma_{\rm inc}(v,A_1,A_2;\theta,\phi)}{d\Omega}, \quad (2.3)$$

where the $d\sigma_{inc}(v,A_1,A_2;\theta,\phi)/d\Omega$ is given in Eq. (2.1). In Eq. (2.3), m_1,m_1^I,m_2,m_2^I are the spin and isospin components of two nucleons along the direction \hat{n} which can be taken as the beam axis.

It is possible now to find a general form of the inclusive differential cross section at leading order in $1/N_c$ by integrating over the impact parameter space in Eq. (2.1) and SU(2) measure in Eq. (2.3). The resulting expression found in [1] is

$$d\sigma^{(m_1,m_1^I,m_2,m_2^I)}(v,\theta,\phi)/d\Omega = a_0(v,\theta,\phi) + b_I(v,\theta,\phi) (\vec{\sigma}_1 \cdot \vec{\sigma}_2) (\vec{\tau}_1 \cdot \vec{\tau}_2) + c_I(v,\theta,\phi) (\vec{\sigma}_1 \cdot \vec{n}) (\vec{\sigma}_2 \cdot \vec{n}) (\vec{\tau}_1 \cdot \vec{\tau}_2) , \qquad (2.4)$$

where σ^i and τ^a are the spin and isospin Pauli matrices corresponding to the two initial nucleons, and the functions a_0 , b_I and c_I encode the leading order behavior at large N_c . In obtaining the result in Eq. 2.4 the following identity of the Wigner *D*-matrices was used,

$$\left(D_{m,n}^{J}\right)^{*} = (-1)^{m-n} D_{-m,-n}^{J}.$$
(2.5)

Equation (2.4) represents an inclusive differential cross section at leading order in $1/N_c$ expansion. However, more readily available is the data for total inelastic nucleon-nucleon cross section. To obtain the total cross section one should integrate the differential cross section over the whole solid angle. In doing so one obtains the following expression,

$$\sigma^{(m_1,m_1',m_2,m_2')}(v) = A_0(v) + B_I(v) \left(\vec{\sigma}_1 \cdot \vec{\sigma}_2\right) \left(\vec{\tau}_1 \cdot \vec{\tau}_2\right) + C_I(v) \left(\vec{\sigma}_1 \cdot \vec{n}\right) \left(\vec{\sigma}_2 \cdot \vec{n}\right) \left(\vec{\tau}_1 \cdot \vec{\tau}_2\right) , \qquad (2.6)$$

where

$$A_0(v) = 1/2 \int d\Omega a_0(v;\theta,\phi), \ B_I(v) = 1/2 \int d\Omega b_I(v;\theta,\phi), \ C_I(v) = 1/2 \int d\Omega c_I(v;\theta,\phi).$$

The factor of 1/2 in the above equations are due to the normalization of the baryon current.

However, while formally integrating over the solid angle to obtain Eq. (2.6) we did not consider that as discussed above the TDMFT treatment from which the differential cross section in Eq. (2.3) was derived strictly holds only in the semiclassical limit. As is well-known, [19] the semiclassical approximation breaks down for small forward angles for which $\theta \leq 1/Rp_N$, where $R \sim N_c^0$ is a size of the interaction potential which is of the order of the nucleon size, and $p_N \sim N_c$ is the center-ofmass momentum. Thus, the forward angle at which the semiclassical approximation breaks down is of order of $1/N_c$ in the kinematic region of interest here. However, the total cross section in Eq. (2.6) includes both forward and backward angles for which the semiclassical approximation breaks down.

However, as shown in details in [1] the contribution to the total cross section from the forward angles $\theta \leq 1/N_c$ vanishes as $1/N_c$ in the large N_c limit provided the scattering cross section is not anomalously peaked in a vanishingly small forward direction. The latter would require large elastic contribution. Thus, the total cross section given in Eq. (2.6) is valid up to corrections of order N_c .

The key result of the above discussion is the form of total nucleon-nucleon cross section given in Eq. (2.6) which is valid up to corrections of order $1/N_c$. While the Skyrme model has been used in the derivation, the result is independent of the details of the model and are valid in the large N_c limit. All the details of the dynamics for which the explicit TDMFT equations have to be used are in the functions $A_0(v)$, $B_0(v)$ and $C_0(v)$ which are of order one in the large N_c limit. These functions can not be determined explicitly at present time. Nevertheless, the form of the total cross section in Eq. (2.6) does contain testable predictions. Note that it is not the most general form which can be obtained based on parity and time reversal invariance. For example, it does not contain such terms as $A_I(\vec{\tau}_1 \cdot \vec{\tau}_2)$ and $B_0(\vec{\sigma}_1 \cdot \vec{\sigma}_2)$. These terms do not appear at leading order in $1/N_c$. This result does not depend on the details of the Skyrme but follows from the contracted spin-flavor symmetry SU(4) applied to the two-nucleon sector.

This result can be tested against the existing experimental data for total nucleon-nucleon cross sections. This is done in the next section [1].

3. Comparison with experimental data

The experimental data exists for the total spin-independent and polarized proton-proton and proton-neutron cross sections [20]. The kinematic regime in which the result obtained above is expected to be valid corresponds to the center-of-mass momentum above 1 *GeV*.




Figure 1: Spin-averaged proton-proton and neutron-proton total cross section as a function of beam momentum (Bugg et al., 1996).

The total nucleon-nucleon cross section given in Eq. (2.6) contains the total nucleon-nucleon cross sections for isosinglet and isotriplet initial states,

$$\sigma^{(m_1,m_1^I,m_2,m_2^I)}(v) = \left(egin{array}{cc} \sigma^{(I=0)} & 0 \ 0 & \sigma^{(I=1)} \end{array}
ight)$$

Using projection operators $(1 - \vec{\tau}_1 \cdot \vec{\tau}_2)/4$ and $(3 - \vec{\tau}_1 \cdot \vec{\tau}_2)/4$, one can extract $\sigma^{(I=0)}$ and $\sigma^{(I=1)}$ cross sections. Then one obtains the total proton-proton, $\sigma^{pp} = \sigma^{(I=1)}$, and neutron-proton cross sections, $\sigma^{np} = \frac{1}{2} (\sigma^{(I=1)} + \sigma^{(I=0)})$. Thus, at leading order in $1/N_c$ one has the following expressions,

$$\sigma^{(pp)} = A_0 + B_I \left(\vec{\sigma}_1 \cdot \vec{\sigma}_2 \right) + C_I \left(\vec{\sigma}_1 \cdot \vec{n} \right) \left(\vec{\sigma}_2 \cdot \vec{n} \right)$$

$$\sigma^{(np)} = A_0 - B_I \left(\vec{\sigma}_1 \cdot \vec{\sigma}_2 \right) - C_I \left(\vec{\sigma}_1 \cdot \vec{n} \right) \left(\vec{\sigma}_2 \cdot \vec{n} \right) .$$
(3.1)

Recall that in the above equations \vec{n} represents the beam axis. Using Eq. (3.1) and averaging over the spin-polarization of the beam and target nucleons, one can obtain spin-averaged total cross sections. Thus, at leading order we have the following relation,

$$\sigma_0^{(pp)} = \sigma_0^{(np)} \left(1 + \mathcal{O}(1/N_c) \right), \qquad (3.2)$$

where σ_0 's are the spin-averaged total cross sections. This prediction follows from the large- N_c analysis and *cannot* be obtained simply from isospin invariance. This large- N_c result is well satisfied by the experimental data shown in Fig. 1.

The total cross sections for the case when beam and target nucleons are transversely polarized relative to the beam direction can also be obtained. Two configurations are possible, $\uparrow\uparrow$ and $\uparrow\downarrow$.




Figure 2: $\Delta \sigma_T$ for neutron-proton (Fonteine et al., 1991) and proton-proton (Ditzler et al., 1983; Lesikar, J. D. 1981) scattering, and $\Delta \sigma_L$ for proton-proton (Auer et al., 1978), and neutron-proton (Sharov et al., 2008) scattering as functions of beam momentum.

These can be combined into an observable, $\Delta \sigma_T = -(\sigma(\uparrow\uparrow) - \sigma(\uparrow\downarrow))$ referred to as *delta sigma transverse*.

Analogously, for the the longitudinally polarized beam and target nucleons one can extract an observable, $\Delta \sigma_L = -(\sigma(\Rightarrow) - \sigma(\Rightarrow))$, the *delta sigma longitudinal*.

The large- N_c analysis, Eq. (3.1), predicts the following results for these observables,

$$\Delta \sigma_T^{(pp)} = -\Delta \sigma_T^{(np)} \left(1 + \mathcal{O}(1/N_c) \right),$$

$$\Delta \sigma_L^{(pp)} = -\Delta \sigma_L^{(np)} \left(1 + \mathcal{O}(1/N_c) \right).$$
(3.3)

Experimental data for these observables is shown in Figs. 2(a) and 2(b). One may conclude from Figs. 2(a) and 2(b) that the large N_c results given in Eq. (3.3) are not satisfied by data. However, the results are valid within corrections of order N_c . Indeed, according to Eq. (2.6) both σ_0 and $\Delta \sigma_T$ and $\Delta \sigma_L$ are of the same order in $1/N_c$. Experimentally however $\Delta \sigma_T$ and $\Delta \sigma_L$ are much smaller then σ_0 . The latter are about 40mb while the former are consistent with zero. The suppression of $\Delta \sigma_T$ and $\Delta \sigma_L$ are for reasons not predicted by large- N_c analysis. Nevertheless, qualitatively the predictions are valid since $\Delta \sigma_T$ and $\Delta \sigma_L$ are small for both pp and np scattering.

Acknowledgments

BAG would like to gratefully acknowledge the support of Professional Development Advisory Council of the New York City College of Technology, the Professional Staff Congress-City University of New York Research Award Program through the grant PSCREG-41-540, and the Center for Theoretical Physics, New York City College of Technology, The City University of New York. The work on which this talk is based was done in collaboration with T.D Cohen.

References

[1] T. D. Cohen and B. A. Gelman, Phys. Rev. C 85 (2012) 024001.

- [2] T. D. Cohen and B. A. Gelman, Phys. Lett. B 540 (2002) 227.
- [3] G. 't Hooft, Nucl. Phys. B 72 (1974) 461.
- [4] E. Witten, Nucl. Phys. B 160 (1979) 57.
- [5] T. D. Cohen, N. Kumar and K. K. Ndousse, Phys. Rev. C 84 (2011), 015204.
- [6] T.H.R. Skyrme, Proc. Roy. Soc. A 260 (1961) 127.
- [7] G. S. Adkins, C. R. Nappi and E. Witten, Nucl. Phys. B 228 (1983) 552.
- [8] I. Zahed and G. E. Brown, Phys. Rept. 142 (1986) 1.
- [9] B. Lucini and M. Panero, arXiv:1210.4997 (2012).
- [10] D. B. Kaplan and M. J. Savage, Phys. Lett. B 365 (1996) 244; D. B. Kaplan and A. V. Manohar, Phys. Rev. C 56 (1997) 76.
- [11] J. L. Gervais and B. Sakita, Phys. Rev. Lett. 52 (1984) 87; R. Dashen, E. Jenkins and A.V. Manohar, Phys. Rev. D 49 (1994) 4713; C. Carone, H. Gergi and S. Osofsky, Phys. Rev. Lett. B 322 (1994) 227; M. Luty and J. March-Russell, Nucl. Phys. B 426 (1994) 71.
- [12] M. K. Banerjee, T. D. Cohen and B. A. Gelman, Phys. Rev. C 65 (2002) 034011; A. V. Belitsky and T.D. Cohen, Phys. Rev. C 65 (2002) 064008; T. D. Cohen, Phys. Rev. C 66 (2002) 064003; A. Calle Cordon and E. Ruiz Arriola, Phys. Rev. C 78 (2008) 054002; Phys. Rev. C 80 (2009) 014002.
- [13] T. D. Cohen and D. C. Dakin, Phys. Rev. C 68 (2003) 017001.
- [14] J. J. Griffin and M. Dworzecka, Phys. Rev. Lett B 93 (1980) 235.
- [15] T.S. Walhout and J. Wambach, Phys. Rev. Lett. 67 (1991) 314; T. Gisiger and M. B. Paranjape, Phys. Rev. D 50 (1994) 1010.
- [16] M. P. Mattis, Phys. Rev. D 39 (1989) 994; E. Jenkins and R. F. Lebed, Phys. Rev. D 52 (1995) 282;
 C.L. Schat, J. L. Goity and N. N. Scoccola, Phys. Rev. Lett. 88 (2002) 102002.
- [17] S. Coleman, Aspects of Symmetry, Cambridge Univ. Press (1985).
- [18] R. Rajaraman, Solitons and Instantons: An Introduction to Solitons and Instantons in Quantum Field Theory, North-Holland Physics Publishing, 1987.
- [19] L. D. Landau and E. M. Lifshitz, Quantum Mechanics: non-relativistic theory, 3rd ed., Pergamon Press, 1977.
- [20] C. Lechanoine-LeLuc and F. Lehar, Rev. Mod. Phys. 65 (1993) 47.