

Factorization model for distributions of quarks in hadrons

B.I. Ermolaev

Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia

M. Greco

Department of Mathematics and Physics and INFN, University Roma Tre, Rome, Italy

S.I. Troyan

St.Petersburg Institute of Nuclear Physics, 188300 Gatchina, Russia

We consider distributions of unpolarized (polarized) quarks in unpolarized (polarized) hadrons. Our approach is based on QCD factorization. We begin with study of Basic factorization for the parton-hadron scattering amplitudes in the forward kinematics and suggest a model for non-perturbative contributions to such amplitudes. This model is based on the simple observation: after emitting an active quark by the initial hadron, the remaining set of quarks and gluons becomes unstable, so description of this colored state can approximately be done in terms of resonances, which leads to expressions of the Breit-Wigner type. Then we reduce these formulae to obtain explicit expressions for the quark-hadron scattering amplitudes and quark distributions in K_T - and Collinear factorizations.

PACS numbers: 12.38.Cy

I. INTRODUCTION

QCD factorization, i.e. separation of perturbative and non-perturbative QCD contributions, proved to be an efficient instrument for describing hadron reaction at high energies. Being first applied to processes in the hard kinematics in the form of Collinear factorization[1], it was soon extended to cover the forward kinematic region, with DGLAP[2] used to account for perturbative contributions. Then, in order to be able to use BFKL[3], a new kind of factorization, K_T -factorization was suggested in Ref.[4]. These kinds of factorization are usually illustrated by identical pictures. For instance, factorization of the DIS hadronic tensor $W_{\mu\nu}$ is conventionally depicted by the construction in Fig. 1 both in Collinear and in K_T -factorizations, where the upper, perturbative blob and the lower, non-perturbative blob are connected by two-parton state. The upper blob in Fig. 1 is calculated with regular perturbative means. On the




FIG. 1. Conventional illustration of QCD factorization. The s -cut of the graph is implied.

contrary, the lower blob is conventionally introduced from purely phenomenological considerations. Collinear and K_T -factorizations operate with different parametrizations for momentum k of the connecting partons and as a result, they are described by different formulae. Collinear factorization assumes that

$$k = \beta p, \quad (1)$$

while K_T -factorization allows for the transverse momentum in addition:

$$k = \beta p + k_{\perp}, \quad (2)$$

accounting therefore for one longitudinal and two transverse components of k . However as a matter-of-fact, k has four components: two of them are longitudinal and the other two are transverse. Accounting for the missing longitudinal component α (for definition of α see Eq. (3)) drove us to suggesting a new, more general factorization which we named in Ref. [5] Basic factorization. In contrast to K_T - and Collinear factorizations, the analytic expressions in Basic factorization can be obtained from the graphs of the type of the one in Fig. 1 with applying the standard Feynman rules.

It is worth reminding briefly our derivation of Basic factorization, for detail see Ref. [5]. Let us consider the Compton scattering amplitude off a hadron in the forward kinematics. It is depicted in Fig. 2.




FIG. 2. Amplitude for forward Compton scattering off a hadron target.

The blob in Fig. 2 denotes ensemble of perturbative and non-perturbative contributions. This blob can be expanded into an infinite series of terms, each of them is represented by two blobs connected with n parton lines, $n = 2, 3, \dots$ Considering only the simplest, two-parton state, we arrive to the graph similar to the one in r.h.s of Fig. 1 but without

the s -cut and with the both blobs accommodating perturbative and non-perturbative contributions at the same time. The integration of the convolution in Fig. 1 over momentum k now runs over the whole phase space and it is expected to bring a finite result. However, the propagators of the connecting partons become singular at $k^2 = 0$ (we neglect quark masses). Besides, the upper blob may contain IR-sensitive perturbative contributions $\sim \ln^n(2pk/k^2)$ (with $n = 1, 2, \dots$). In addition, it yields the factor $2qk/k^2$, when unpolarized gluon ladders are included into consideration. The only way to kill such IR singularity is to assume that the lowest, non-perturbative blob should tend to zero fast enough when $k^2 \rightarrow 0$. Doing so and repeating a similar procedure to regulate the UV singularity, we bring the convolution in Fig. 1 to agreement with the factorization concept: perturbative and non-perturbative contributions are located indifferent blobs. This is a new form of QCD factorization which we name Basic factorization.

We demonstrated in Ref. [5] that Basic factorization can be reduced step-by-step first to K_T - and then to Collinear factorizations. In Ref. [5] we began with considering Basic factorization for Compton scattering amplitudes in the forward kinematics, where integration over momentum k of the connecting partons in Fig. 1 runs over the whole phase space. Confronting two obvious facts that, on one hand, the integration over k should yield a finite results and that, on the other hand, the perturbative part in Fig. 1 (the upper, perturbative blob and propagators of the connecting partons) is divergent in both the infra-red (IR) and ultra-violet (UV) regions, allowed us to impose integrability restrictions on the lowest blob, which are necessary for the convolution in Fig. 1 to be finite. The obtained restrictions led us to theoretical constraints on the fits for the parton distributions to the DIS structure functions in Collinear and K_T - factorizations. In particular, we predicted the general form of the fits in K_T -factorization and excluded the factors x^{-a} from the fits in both K_T - and Collinear factorizations.

Another interesting object, where factorization is used, is distributions of partons in hadrons. In the present paper we examine their properties in IR and UV regions and suggest a simple resonance model for the non-perturbative contributions to the parton distributions. Our argumentation in favor of this model is as follows: after emitting an active quark by a hadron, the remains of the hadron, i.e. a set of quarks and gluons, acquires a color and therefore it becomes unstable. So, this colored state can be described in terms of resonances. We begin with considering amplitudes of the quark-hadron (QHA) and gluon-hadron (GHA) scattering in the forward kinematics. The Optical theorem relates such amplitudes to the parton distributions. Throughout the paper we use the standard Sudakov parametrization[6] for momentum k of the connecting partons:

$$k = -\alpha q' + \beta p' + k_{\perp}, \quad (3)$$

where momenta q' and p' are massless, $p'^2 \approx q'^2 \approx 0$, and they are made of the hadron momentum p and the parton momentum q :

$$p' = p + x_2 q, \quad q' = q + x_1 p, \quad (4)$$

where $x_2 = -p^2/w \equiv -M^2/w$, $x_1 = -q^2/w$, with $w = 2pq \approx 2p'q'$. In these terms

$$2pk = w(-\alpha - x_2\beta), \quad 2qk = w(\beta - x_1\alpha), \quad k^2 = -w\alpha\beta - k_{\perp}^2. \quad (5)$$

In Sect. II we introduce the quark-hadron scattering amplitudes in the forward kinematics and examine their IR and UV behavior. In Sect. III we consider separately the unpolarized and spin-dependent quark-hadron amplitudes in Basic factorization and suggest a model for non-perturbative contributions to the amplitudes. This model involves a spinor structure accompanied by invariant amplitudes $T^{(U)}$ and $T^{(S)}$. In Sect. III we specify the spinor structure of the non-perturbative contributions to the amplitudes and parton distributions. In Sect. IV we show how Basic factorization for the quark-hadron amplitudes and quark distributions in hadrons can be reduced to K_T - and Collinear factorizations. In Sect. V we focus on a model for the invariant amplitudes $T^{(U)}$ and $T^{(S)}$. The model is based on description of $T^{(U)}$ and $T^{(S)}$ in a quasi-resonant way and through the Optical theorem it easily leads to non-perturbative contributions to the parton distributions, with expressions of the Breit-Wigner kind both in Basic and in K_T - factorizations. Finally, Sect. VI is for concluding remarks.

II. QUARK-HADRON AMPLITUDES

In the factorization approach, the quark-hadron amplitudes (QHA) A_q are expressed through convolutions of perturbative amplitudes $A^{(pert)}$ and non-perturbative amplitudes T as shown in Fig. 3.

In the Born approximation $A^{(pert)}$ is depicted in Fig. 4 as a one-rung ladder. Adding more ladder rungs to it together with inclusion of non-ladder graphs and resumming all such graphs converts the Born amplitude into $A^{(pert)}$.




FIG. 3. Factorization of the quark-hadron amplitude.

In the present paper we do not consider mixing of quark and gluon ladder rungs, i.e. we consider the graphs where the vertical quark lines go from the bottom to the top without breaking.




FIG. 4. Born approximation for the factorized quark-hadron amplitude.

We begin consideration of the quark-hadron amplitudes A_q in Basic factorization, studying the simplest case depicted in Fig. 4, where the perturbative contributions are accounted in the Born approximation and denote such distributions B_q . In Basic factorization one can use the standard Feynman rules to write down the analytic expression corresponding to the graphs in Figs. 3,4. Doing so, we obtain that

$$B_q = -i4\pi\alpha_s C_F \int \frac{d^4k}{(2\pi)^4} \frac{\bar{u}(q)\gamma_\mu \hat{k} \hat{T}_q(k,p) \hat{k} \gamma_\nu u(q)}{k^2 k^2 (q+k)^2} d_{\mu\nu}, \quad (6)$$

where we have used the standard notations: $C_F = (N^2 - 1)/(2N) = 4/3$ and α_s is the QCD coupling. In Eq. (6) \hat{T}_q corresponds to the lowest blob in Fig. 3. It is altogether non-perturbative object. Throughout the paper we will address it as the primary quark-hadron amplitude¹. Choosing the Feynman gauge, where $d_{\mu\nu} = g_{\mu\nu}$, for the virtual gluon and the Sudakov parametrization (3) for the quark momentum k , we rewrite Eq. (6) as follows:

$$B_q = -i \frac{\alpha_s C_F}{8\pi^3} w \int d\alpha d\beta d^2k_\perp \frac{\bar{u}(q)\gamma_\mu \hat{k} \hat{T}_q(k,p) \hat{k} \gamma_\mu u(q)}{k^2 k^2 (q+k)^2}. \quad (7)$$

Throughout the paper, for the sake of simplicity, we will treat the external quarks with momentum q as on-shell ones, though our reasoning remains valid also when they are off-shell. Introducing the density matrix

$$\hat{\rho}(p)(q) = \frac{1}{2}(\hat{q} + m_q)(1 - \gamma_5 \hat{S}_q), \quad (8)$$

¹ In Ref. [7] non-perturbative contributions to parton distributions in the context of Collinear factorization were called intrinsic contributions.

with q , m_q and S_q being the quark momentum, mass and spin respectively, we bring Eq. (7) to the following form:

$$B_q \approx -i \frac{\alpha_s C_F}{8\pi^3} w \int d\alpha d\beta d^2 k_\perp \frac{\text{Tr} \left[\hat{\rho}(p)(q) \gamma_\mu \hat{k} \hat{T}_q(k, p) \hat{k} \gamma_\mu \right]}{k^2 k^2 (q+k)^2}. \quad (9)$$

We stress that the replacement of Eq. (7) by Eq. (9) is not necessary for us but it allows us to carry out a more detailed consideration of A_q^B . In particular, we can consider separately the spin-dependent, $B_q^{(spin)}$ and independent, $B_q^{(unpol)}$ quark-hadron amplitudes in a simple way:

$$B_q^{(unpol)} = -i \frac{\alpha_s C_F}{8\pi^3} w \int d\alpha d\beta d^2 k_\perp \frac{2(qk) \text{Tr} \left[\hat{k} \hat{T}_q^{(unpol)} \right] - k^2 \text{Tr} \left[\hat{q} \hat{T}_q^{(unpol)} \right]}{k^2 k^2 (q+k)^2}, \quad (10)$$

$$B_q^{(spin)} = \frac{\alpha_s C_F}{8\pi^3} m_q w \int d\alpha d\beta d^2 k_\perp \frac{2(S_q k) \text{Tr} \left[\gamma_5 \hat{k} \hat{T}_q^{(spin)} \right] - k^2 \text{Tr} \left[\gamma_5 \hat{S}_q \hat{T}_q^{(spin)} \right]}{k^2 k^2 (q+k)^2}. \quad (11)$$

In Eqs. (10,11) we have replaced the general primary amplitude \hat{T}_q by more specific amplitudes $\hat{T}_q^{(unpol)}$, $\hat{T}_q^{(spin)}$. In Eq. (10) we have neglected a contribution $\sim m$ in $\hat{\rho}(q)$ compared to the contribution $\sim \hat{q}$. Integrations in Eqs. (10,11) run over the whole phase space and it is supposed to yield finite results. However, there can be singularities in the integrands and they should be regulated. Regulating them with introducing various cut-offs would be unphysical, so the only way out is to impose appropriate constraints on the primary quark-hadron amplitudes $\hat{T}_q^{(unpol)}$, $\hat{T}_q^{(spin)}$ so that to kill the singularities. When the perturbative amplitude $A^{(pert)}$ is calculated in the Born approximation, the only possible singularities in Eqs. (10,11) are IR singularities at $k^2 = 0$ and UV singularities which we relate to integrations over α . However, when $A^{(pert)}$ is beyond the Born approximation, there appears another kind of singularities called in Ref. [8] rapidity divergences. Below we consider handling these singularities in the framework of Basic factorization.

A. Rapidity divergences of QHA

Rapidity divergences were investigated first in Ref. [8] and then in Ref. [9] in the context of K_T -factorization. Detailed investigation of this problem can be found in Ref. [10]. In the lowest order of the Perturbative QCD, the rapidity divergences come from the graphs in Fig. 5 (and symmetrical graphs as well), where the radiative corrections calculated in the first-loop approximation are convoluted with the unintegrated parton distribution $\tilde{\Phi}$. Let us stress that $\tilde{\Phi}$ accumulates both perturbative and non-perturbative corrections. When such convolutions are considered in K_T




FIG. 5. Graphs contributing to rapidity divergences in unintegrated parton distributions. The dashed lines denote cuts.

-factorization, each of the graphs in Fig. 5 acquires logarithmic divergences arising from integration over momentum l_+ (with $l_+ = (l_0 + l_z)/\sqrt{2}$). They are called rapidity divergences and they can be got rid of as shown in Ref. [8]

(when the Feynman gauge is used for the gluon propagators) and then in Ref. [9] for the case of the light-cone gauge. In Refs. [8, 9] the rapidity divergences are cured with redefining $\tilde{\Phi}$.

Now let us study this situation in Basic factorization. To this end we consider a contribution of the graph in Fig. 6 to the quark-hadron amplitude in Basic factorization. We remind that there are no cuts in Fig. 6 and the blob T accumulates non-perturbative contributions only. One of remarkable features here is that analytic expressions in Basic




FIG. 6. Graph contributing to quark-hadron amplitude .

factorization can be obtained by applying standard Feynman rules to the involved graphs. Second important point is that one is free to use any gauge for perturbative QCD calculations² in Basic factorization whereas the blob T in Fig. 6 is altogether non-perturbative and therefore it is insensitive to the choice of the gauge. Applying the Feynman rules to the graph in Fig. 6 and integrating over the loop momentum l , we immediately conclude that this integration yields a logarithmic UV-divergent contribution which, being complemented by a similar contribution from the symmetrical graph and self-energy graphs, in a conventional way leads to renormalization of the gluon-quark couplings. After absorption of such divergent contributions by the couplings, we obtain a renormalized amplitude which is free of divergences. Then, applying the Optical theorem to the this construction, we arrive at the parton distributions and they are also free of divergences. Obviously, the same treatment can be applied to other UV divergences coming from perturbative component $A^{(pert)}$ in higher loops: all of them can be absorbed by renormalizations. Now we focus on the divergences resulting from integration of the convolutions in Eqs. (10,11), where the perturbative amplitudes $A^{(pert)}$ are in the Born approximation.

B. IR and UV stability of QHA

First of all, let us note that the denominators in Eqs. (10,11) can become singular in the infra-red (IR) region, where $k^2 \sim 0$. In the case of purely perturbative QCD, IR singularities are conventionally regulated by introducing IR cut-offs. In our case there is not any physical reason for that, so we are left with the only way to kill these singularities: The primary quark-hadron amplitudes \hat{T}_q should become small at small k^2 :

$$\hat{T}_q^{(unpol)}, \hat{T}_q^{(spin)} \sim (k^2)^{1+\eta}, \quad (12)$$

when $k^2 \rightarrow 0$. Now let us consider the ultra-violet (UV) stability of the convolutions in Eqs. (10,11). The integration over α in Eqs. (10,11) runs from $-\infty$ to ∞ , so, at large $|\alpha|$ the integrands should decrease fast enough to guarantee UV stability. First of all we focus on the integration over α in Eq. (10). Taking into consideration that each factor in the denominator of Eq. (10) is $\sim \alpha$ makes that the denominator to be $\sim \alpha^3$. The term $2qk$ in the numerator depends on α because $2qk = w(\beta - x_1\alpha)$ and the factors k^2 and \hat{k} are $\sim \alpha$, which makes

$$\frac{2qk \hat{k}}{k^2 k^2 (q+k)^2} \sim \frac{\alpha^2}{\alpha^3}. \quad (13)$$

² For gauge invariance of Basic factorization see Ref. [5].

This divergence must be regulated by an appropriate decrease of $\hat{T}_q^{(unpol)}$ at large $|\alpha|$. The IR stability condition in Eq. (12) states that $\hat{T}_q^{(unpol)} \sim (k^2)^{1+\eta}$ at small k^2 but it can either disappear or be kept at large $|\alpha|$. Therefore we have two options:

(A) The factor $(k^2)^{1+\eta}$ survives at large $|\alpha|$.

(B) The factor $(k^2)^{1+\eta}$ disappears at large $|\alpha|$.

In the case (A), where IR and UV behaviors of $\hat{T}_q^{(unpol)}$ are related, $\hat{T}_q^{(unpol)}$ should behave at large $|\alpha|$ as follows:

$$\hat{T}_q^{(unpol)} \sim \alpha^{\eta-\chi} = (\alpha^{1+\eta}) [\alpha^{-1-\chi}], \quad (14)$$

with $\chi > \eta > 0$.

IR and UV behaviors of $\hat{T}_q^{(unpol)}$ are disconnected in the case (B). It converts Eq. (14) into

$$\hat{T}_q^{(unpol)} \sim \alpha^{-\chi}. \quad (15)$$

The first factor in Eq. (14) corresponds to the term $(k^2)^{1+\eta}$, while a contribution generating the asymptotic factor in the squared brackets has to be specified. We will do it in Sect. V. Now let us consider the spin-dependent amplitudes. In order to guarantee their IR stability, the primary spin-dependent amplitude $\hat{T}_q^{(spin)}$ should also be $\sim (k^2)^{1+\eta}$ at small k^2 but the situation with its UV stability is more involved than in the unpolarized case. Indeed, the quark spin S_q can be either in the plane formed by p and q , i.e. $S_q = S_q^{\parallel}$, or in the transverse space, where $S_q = S_q^{\perp}$. Depending on it, there are the longitudinal spin-dependent amplitude, B_q^{\parallel} and the transverse one, B_q^{\perp} . Now let us consider the term $2m_q S_q k$ in Eq. (11) for different orientations of the quark spin: When the spin is longitudinal,

$$2m_q S_q k = 2m_q S_q^{\parallel} k = w(\beta - x_1 \alpha) \quad (16)$$

and \hat{k} in the trace $Tr[\hat{k}\hat{T}_q]$ is also $\sim \alpha$. In contrast when the spin is transverse,

$$2m_q S_q k = -2m_q (\vec{S}_q^{\perp} \vec{k}_{\perp}) \quad (17)$$

and therefore $S_q^{\perp} k$ does not depend on α . Then, this \vec{k}_{\perp} should be accompanied by another \vec{k}_{\perp} from the trace in order to get a non-zero result at integration over the azimuthal angle, i.e. The first term in the numerator of Eq. (11) does not depend on α , while the second term is $\sim \alpha$. It means that, with \hat{T}_q^{\parallel} dropped, the explicit α -dependence of A_q^{\parallel} at large $|\alpha|$ coincides with the one in Eq. (13):

$$\frac{S_q^{\parallel} k \hat{k}}{k^2 k^2 (q+k)^2} \sim \frac{\alpha^2}{\alpha^3} \quad (18)$$

and

$$\frac{S_q^{\perp} k \hat{k}}{k^2 k^2 (q+k)^2} \sim \frac{\alpha}{\alpha^3}. \quad (19)$$

It follows from Eq. (18) states that the α -dependence of the amplitude \hat{T}_q^{\parallel} at large $|\alpha|$ is identical to the one of $\hat{T}_q^{(unpol)}$:

$$\hat{T}_q^{\parallel} \sim \hat{T}_q^{(unpol)} \sim (\alpha^{1+\eta}) [\alpha^{-1-\chi}] \quad (20)$$

in the case (A) and

$$\hat{T}_q^{\parallel} \sim \hat{T}_q^{(unpol)} \sim (\alpha^{-\chi}) \quad (21)$$

in the case (B). \hat{T}_q^{\perp} can decrease slower:

$$\hat{T}_q^{\perp} \sim (\alpha^{1+\eta}) [\alpha^{-\chi}] \quad (22)$$

in the case **(A)** and

$$\hat{T}_q^\perp \sim \alpha^{1-\chi} \quad (23)$$

in the case **(B)**. Eqs. (12,20 - 23) guarantee integrability of the convolutions for the quark-hadron amplitudes in Basic factorization. These integrability requirements can be used as general theoretical constraints on non-perturbative contributions to the amplitudes in Basic factorization (see Ref. [5] for detail) and we will use them in the present paper. Each of Eqs. (20,22) consists of two factors. The first factor in these equations is universally generated by the term $(k^2)^{1+\eta}$ while contributions generating the factors in squared brackets will be specified in Sect. V.

III. MODELING THE SPINOR STRUCTURE OF \hat{T}_q

Our next step is to simplify the traces in Eqs. (10,11). In order to do it, we have to specify the spinor structure of the primary QHA \hat{T}_q . By definition, \hat{T}_q is altogether non-perturbative, so specifying its spinor structure can only be done on basis of phenomenological considerations. However, any model expression for \hat{T}_q should respect the integrability conditions in Eqs. (12, 14,20,22). There is the well-known expression for the density matrix of an elementary fermion:

$$\hat{\rho}(p) = \frac{1}{2}(\hat{p} + M)(1 - \gamma_5 \hat{S}) \approx \frac{\hat{p}}{2} - \frac{1}{2}(\hat{p} + M)\gamma_5 \hat{S}, \quad (24)$$

where M and S are the fermion mass and spin. This expression drives us to approximate \hat{T}_q as follows:

$$\hat{T}_q = \hat{p} T_q^{(U)}(k^2, 2pk) - (\hat{p} + M)\gamma_5 \hat{S} T_q^{(S)}(k^2, 2pk), \quad (25)$$

where p , S are the hadron momentum and spin respectively and $T_q^{(U)}$, $T_q^{(S)}$ are scalar functions. Throughout the paper we will address them as invariant quark-hadron amplitudes. Substituting T_q of Eq. (12) in Eqs. (10,11) and calculating the traces, we arrive at the following expressions:

$$\begin{aligned} B_q^{(unpol)} &= -i \frac{1}{8\pi^3} \int d\alpha d\beta dk_\perp^2 \left[-g^2 C_F \frac{w}{[(q+k)^2 + i\epsilon]} \right] \left(\frac{k_\perp^2}{k^2 k^2} \right) \left(T_q^{(U)}(k^2, 2pk) \right) \\ &= -i \frac{1}{8\pi^3} \int d\alpha d\beta dk_\perp^2 \tilde{B}_q^{(unpol)}(q, k) \left(\frac{k_\perp^2}{k^2 k^2} \right) \left(T_q^{(U)}(k^2, 2pk) \right), \end{aligned} \quad (26)$$

where we have denoted $\tilde{B}_q^{(unpol)}$ the perturbative amplitude in the Born approximation for the forward annihilation of unpolarized quark-quark pair. We have neglected contributions $\sim x_{1,2}$ in the numerator of Eq. (26) and will do it in expressions for the spin-dependent amplitudes. These terms, if necessary, can easily be accounted for with more accurate implementation of Eq. (3) to Eqs. (26). Let us consider the structure of the integrand in Eq. (26) in more detail. The amplitude in the last brackets is entirely non-perturbative. It is supposed to mimic a transition from hadrons to quarks. The fraction in the middle corresponds to the convoluting the perturbative and non-perturbative amplitudes. The fraction in the first brackets corresponds to the perturbative amplitude for the forward scattering of quarks in the Born approximation. We explicitly wrote the factor $i\epsilon$ there to remind that this amplitude has the s -channel imaginary part. Doing similarly, we obtain an expression for the spin-dependent amplitudes:

$$B_q^{(spin)} = i \frac{g^2 C_F}{16\pi^4} 2m_q M w \int d\alpha d\beta d^2 k_\perp \frac{2(kS_q)(kS) - k^2(S_q S)}{k^2 k^2 [(q+k)^2 + i\epsilon]} T_q^{(S)} \quad (27)$$

Let us consider Eq. (27) for different orientation of the hadron spin:

- (i) The hadron spin S is in the plane formed by momenta p and q , so for this case we use the notation $S = S^\parallel$.
- (ii) The hadron spin is transverse to this plane. We denote this case as $S = S^\perp$.

Amplitude A_q^\parallel for the first case is given by the expression very close to the unpolarized amplitude:

$$\begin{aligned} B_q^{(\parallel)} &= -i \frac{1}{16\pi^3} \int d\alpha d\beta dk_\perp^2 \left[-g^2 C_F \frac{2mM(S_q^\parallel S^\parallel)}{(q+k)^2 + i\epsilon} \right] \left(\frac{k_\perp^2}{k^2 k^2} \right) T_q^{(\parallel)}(k^2, 2pk) \\ &= -i \frac{1}{8\pi^4} \int d\alpha d\beta d^2 k_\perp \tilde{B}_q^{(\parallel)}(q, k) \left(\frac{k_\perp^2}{k^2 k^2} \right) T_q^{(\parallel)}(k^2, 2pk), \end{aligned} \quad (28)$$

whereas the transverse amplitude is given by a different expression:

$$\begin{aligned} B_q^{(\perp)} &= -i \frac{1}{16\pi^3} \int d\alpha d\beta dk_{\perp}^2 \left[-g^2 C_F \frac{2mM(S_q^{\perp} S^{\perp})}{(q+k)^2 + i\epsilon} \right] \left(\frac{w\alpha\beta}{k^2 k^2} \right) T_q^{(\perp)}(k^2, 2pk) \\ &= -i \frac{1}{8\pi^3} \int d\alpha d\beta dk_{\perp}^2 \tilde{B}_q^{(\perp)}(q, k) \left(\frac{w\alpha\beta}{k^2 k^2} \right) T_q^{(\perp)}(k^2, 2pk), \end{aligned} \quad (29)$$

with $\tilde{B}_q^{(\parallel)}$, $\tilde{B}_q^{(\perp)}$ being the perturbative spin-dependent Born amplitudes. Accounting for perturbative QCD radiative corrections converts the Born amplitudes $\tilde{B}_q^{(unpol)}$, $\tilde{B}_q^{(\parallel)}$, $\tilde{B}_q^{(\perp)}$ in Eqs. (26,28,29) into perturbative dimensionless amplitudes $\tilde{A}_q^{(unpol)}$, $\tilde{A}_q^{(\parallel)}$, $\tilde{A}_q^{(\perp)}$, remaining the other factors unchanged:

$$\begin{aligned} A_q^{(unpol)}(p, q) &= -i \frac{1}{8\pi^3} \int d\beta \frac{dk_{\perp}^2}{k^2} d\alpha \tilde{A}_q^{(unpol)}(q, k) \left(\frac{k_{\perp}^2}{k^2} \right) T_q^{(U)}(k^2, 2pk), \\ A_q^{(\parallel)}(p, q) &= -i \frac{1}{8\pi^3} \int d\beta \frac{dk_{\perp}^2}{k^2} d\alpha \tilde{A}_q^{(\parallel)}(q, k) \left(\frac{k_{\perp}^2}{k^2} \right) T_q^{(\parallel)}(k^2, 2pk), \\ A_q^{(\perp)}(p, q) &= -i \frac{1}{8\pi^3} \int d\beta \frac{dk_{\perp}^2}{k^2} d\alpha \tilde{A}_q^{(\perp)}(q, k) \left(\frac{w\alpha\beta}{k^2} \right) T_q^{(\perp)}(k^2, 2pk). \end{aligned} \quad (30)$$

Taking the s -imaginary part of Eq. (26), we arrive at the totally unintegrated, or fully unintegrated as was used in Ref. [11], distribution of unpolarized quarks in the hadron $D_q^{unpol\ B}$ in the Born approximation:

$$\begin{aligned} D_q^{(unpol\ B)} &= \frac{1}{8\pi^2} \int d\beta \frac{dk_{\perp}^2}{k^2} d\alpha \left[g^2 C_F \delta(\beta - x - z) \right] \frac{k_{\perp}^2}{k^2} \Psi_q^{(1)}(k^2, 2pk) \\ &= \frac{1}{8\pi^2} \int \frac{d\beta}{\beta} \frac{dk_{\perp}^2}{k^2} d\alpha \left[g^2 C_F \delta(1 - x/\beta - z/\beta) \right] \frac{k_{\perp}^2}{k^2} \Psi_q^{(1)}(k^2, 2pk) \end{aligned} \quad (31)$$

where $x = -q^2/w$, $z = k_{\perp}^2/w$ and $\Psi_q^{(1)}$ is the primary quark distribution of unpolarized quarks in the hadron, $\Psi_q^{(1)} = (1/\pi) \Im T_q^{(U)}$. This object is altogether non-perturbative. Applying the Optical theorem to Eq. (30), we arrive at the parton distributions beyond the Born approximation:

$$\begin{aligned} D_q^{(unpol)}(x, q^2) &= \frac{1}{8\pi^2} \int \frac{d\beta}{\beta} \frac{dk_{\perp}^2}{k^2} d\alpha \tilde{D}_q^{(unpol)}(x/\beta, q^2/k^2) d\beta \left(\frac{k_{\perp}^2}{k^2} \right) \Psi_q^{(1)}(k^2, w\alpha), \\ D_q^{(\parallel)}(x, q^2) &= \frac{1}{8\pi^2} \int \frac{d\beta}{\beta} \frac{dk_{\perp}^2}{k^2} d\alpha \tilde{D}_q^{(\parallel)}(x/\beta, q^2/k^2) \left(\frac{k_{\perp}^2}{k^2} \right) \Psi_q^{(\parallel)}(k^2, w\alpha), \\ D_q^{(\perp)}(x, q^2) &= \frac{1}{8\pi^2} \int \frac{d\beta}{\beta} \frac{dk_{\perp}^2}{k^2} d\alpha \tilde{D}_q^{(\perp)}(x/\beta, q^2/k^2) \left(\frac{w\alpha\beta}{k^2} \right) \Psi_q^{(\perp)}(k^2, w\alpha). \end{aligned} \quad (32)$$

IV. REDUCTION OF BASIC FACTORIZATION TO CONVENTIONAL FACTORIZATIONS

Conventional forms of factorization are Collinear and k_T - factorizations. In Ref. [5] we described reduction of Basic factorization to K_T - and Collinear factorizations for the Compton scattering amplitudes and DIS structure functions without specifying the non-perturbative amplitudes T_q . In this Sect. we show that these results perfectly agree with our assumption in Eq. (25) concerning structure of T_q . We demonstrate that the parton distributions in both conventional factorizations can be obtained with step-by-step reductions of the expressions for $D_q^{(unpol)}$, $D_q^{(\parallel)}$, $D_q^{(\perp)}$ in Basic factorization. This reductions are the same for both the parton-hadron amplitudes and parton distributions, they are insensitive to spin and stands when the quarks are replaced by gluons. Because of that we consider such reductions for a generic parton-hadron distribution D in Basic factorization and skip unessential factors:

$$D(x, q_{\perp}^2) = \int d\beta \frac{dk_{\perp}^2}{k^2} d\alpha D^{(pert)}(x/\beta, q_{\perp}^2/k^2) \left(\frac{k_{\perp}^2}{k^2} \right) \Psi(w\alpha, k^2), \quad (33)$$

where $D^{(pert)}$ stands for a perturbative contribution and Ψ is the altogether non-perturbative (we address it as primary) parton-hadron distribution. Actually, $\Psi(w\alpha, k^2)$ is the starting point for the perturbative evolution. Integration in Eq. (33) runs over the whole phase space. Let us note that in the literature very often are considered purely transverse q : $q^2 \approx q_\perp^2$. Because of this reason we will use the notation q_\perp^2 instead of q^2 in what follows, though Eqs. (32,33) are also valid when $q^2 \neq q_\perp^2$.

A. Reduction to k_T -factorization

In order to reduce Eq. (33) to k_T -factorization, we have to perform integration with respect to α . However, this integration should not involve $D^{(pert)}$, which, strictly speaking, is impossible because $D^{(pert)}$ depends on k^2 and thereby it depends on α : $k^2 = -w\alpha\beta - k_\perp^2$. The only way out is to assume that the main contributions to Eq. (33) come from the region where

$$\alpha \ll \alpha_{\max} = k_\perp^2 / (w\beta), \quad (34)$$

i.e. $k^2 \approx -k_\perp^2$. Let us notice that approximating ladder partons virtualities k^2 by their transverse momenta is well-known. It is used in all available evolution equations, including DGLAP and BFKL, and now it allows us to convert Eq. (33) into an expression for the unintegrated (transverse momentum dependent[12]) parton distributions D_{KT} in k_T -factorization:

$$D_{KT}(x, q_\perp^2) \approx \int_x^1 \frac{d\beta}{\beta} \int_0^r \frac{dk_\perp^2}{k_\perp^2} D_{KT}^{(pert)}(x/\beta, q_\perp^2/k_\perp^2) \Phi(w\beta, k_\perp^2), \quad (35)$$

where $r = w\beta - q_\perp^2$. Obviously, $r \approx w \approx q_\perp^2$ for large x while at very small x one can use $r \approx w\beta$. Φ denotes the primary (i.e. non-perturbative) k_T -parton distribution. It is related to Ψ as follows:

$$\Phi(w\beta, k_\perp) = \int_0^{k_\perp^2/w\beta} d\alpha \Psi(w\alpha, k_\perp^2). \quad (36)$$

B. Reduction to Collinear factorization

In Ref. [5] we discussed how to reduce K_T -factorization to Collinear one, using DIS structure functions as an example. The same argumentation can be applied to the parton distributions. We briefly repeat it below. In order to reduce k_T -factorization to Collinear factorization, we should perform integration of Eq. (35) with respect to k_\perp without integrating $D^{(pert)}$. Of course it cannot be done straightforwardly, because $D^{(pert)}$ explicitly depends on k_\perp . However, we can do it approximately, assuming a sharp peaked dependence of $\Psi(w\alpha, k_\perp^2)$ on k_\perp^2 with maximum at $k_\perp^2 = \mu^2$ as shown in Fig. 7. The closer this dependence is to $\delta(k_\perp^2 - \mu^2)$, the higher is accuracy of the reduction. As discussed in Ref. [5], the number of such maximums can be unlimited. We remind that Φ is non-perturbative, so typical values of μ must be of non-perturbative range, $\mu \sim \Lambda_{QCD}$. After the integration of Φ we arrive at Collinear factorization convolution:

$$D^{(col)}(x, q_\perp^2/\mu^2) \approx \int_x^1 \frac{d\beta}{\beta} D_{col}^{(pert)}(x/\beta, q_\perp^2/\mu^2) \phi(\beta, \mu^2), \quad (37)$$

with μ being the intrinsic factorization scale and ϕ being the primary (non-perturbative) integrated parton distribution:

$$\phi(\beta, \mu^2) = \int_\Omega \frac{dk_\perp^2}{k_\perp^2} \Phi(w\beta, k_\perp^2), \quad (38)$$

where the integration region Ω is located around the maximum $k_\perp^2 = \mu^2$. At the first sight, the form of Collinear factorization presented in Eq. (37) contradicts to the conventional form. Indeed, the scale μ in Eq. (37) corresponds to the maximum in Fig. 7 and therefore its value is fixed. On the contrary, the conventional form of Collinear




FIG. 7. The peaked form of $\Phi(\beta, k_{\perp}^2)$ with one maximum

factorization operates with integrated parton distributions $\varphi(\beta, \tilde{\mu}^2)$, where the scale $\tilde{\mu}$ can have any arbitrary value. Then, we expect the value of μ to be of non-perturbative range whereas usually $\tilde{\mu} \sim \text{few GeV}$, i.e. typically $\tilde{\mu} \gg \mu$. However, this contradiction can easily be solved as was shown in Ref. [5]. The point is that transition from $\phi(\beta, \mu^2)$ to $\varphi(\beta, \tilde{\mu}^2)$ can be done, applying perturbative evolution in the μ^2 -space to $\phi(\beta, \mu^2)$ and keeping β fixed. It can be written symbolically as

$$\varphi(\beta, \tilde{\mu}^2) = E(\tilde{\mu}^2, \mu^2) \otimes \phi(\beta, \mu^2), \quad (39)$$

where E is the evolution operator in the μ^2 -space. Specific expressions for E are different in different perturbative approaches (see Appendix E for detail). Eq. (39) makes possible to arrive at the conventional unintegrated distribution $\varphi(\beta, \tilde{\mu}^2)$ fixed at an arbitrary scale $\tilde{\mu}^2$. In contrast to $\phi(\beta, \mu^2)$, the distribution $\varphi(\beta, \tilde{\mu}^2)$ accumulates both perturbative and non-perturbative contributions. It is easy to show that our reasoning remains true in the case when Φ has several maximums or an infinite series of them. This point was discussed in detail in Ref. [5], so we will not do it in the present paper. Instead, we focus on modeling invariant amplitudes $T^{(U,S)}$ introduced in Eq. (25).

V. MODELING THE INVARIANT QUARK-HADRON AMPLITUDES AND PRIMARY QUARK DISTRIBUTIONS

In this Sect. we suggest a model which mimics non-perturbative QCD contributions in the primary hadron-quark invariant amplitudes $T^{(U,S)}$ and in the primary quark distributions in all available forms of factorization. Once again we begin with consideration of the invariant amplitudes $T^{(U,S)}$ and then proceed to the quark distributions.

A. Resonance model for the primary quark-hadron invariant amplitudes

Amplitudes $T^{(U,S)}$ can be introduced in a model-dependent way only because QCD has not been solved in the non-perturbative region. All such models should satisfy several restrictions:

(i): The IR stability conditions in Eq. (12) and the UV stability conditions in Eqs. (20 - 23) should be respected because they guarantee integrability of the factorization convolutions. We remind that the UV stability conditions derived in Sect. II depend on UV-behavior of the factors regulating IR divergences. Namely, Eqs. (14,20,22) correspond to the case **(A)** while Eqs. (15,21,23) correspond to the case **(B)**. In the present paper we focus on the most UV-divergent case **(A)**, although our conclusions hold true for the case **(B)** as well.

(ii): the invariant amplitudes should respect the Optical Theorem, so they should have s -channel imaginary parts.

(iii): These amplitudes should guarantee the step-by-step reductions from Basic factorization to k_T -factorization and then to Collinear factorization described in Sect. IV.

The expressions for the unpolarized and spin-dependent amplitudes with the longitudinal spin in Eq. (30) are much alike while the expression for the transverse spin amplitude differs from them. Despite this difference, our model equally stands for all spin-dependent amplitudes and quark distributions regardless of the spin orientation. To describe the invariant primary quark-hadron amplitudes, we suggest a model of the resonance type for $T_q^{(U,S)}$:

$$\begin{aligned}
T_q^{(U)}(pk, k^2) &= \frac{R_U(k^2)}{((k-p)^2 - M_1^2 + i\Gamma_1)((k-p)^2 - M_2^2 + i\Gamma_2)} \\
T_q^{(S)}(pk, k^2) &= \frac{R_S(k^2)}{((k-p)^2 - M_3^2 + i\Gamma_3)((k-p)^2 - M_4^2 + i\Gamma_4)}
\end{aligned} \tag{40}$$

where $R_{U,S}(k^2)$ are supposed to behave as $R_{U,S}(k^2) \sim (k^2)^{1+\eta}$ at small k^2 . We need at least two resonances to satisfy the UV stability requirement of Eq. (14). Indeed, Eq. (40) leads to $\chi = 1$ while a model with one resonance corresponds to $\chi = 0$. Formally, Eq. (40) contains independent parameters $M_{1,2,3,4}$ and $\Gamma_{1,2,3,4}$ but we do not see a physical reason forbidding to identify $T_q^{(U)}$ and $T_q^{(S)}$, which would left us with the parameters $M_{1,2}$ and $\Gamma_{1,2}$ only. In terms of the Sudakov variables $T_q^{(U,S)}$ are:

$$\begin{aligned}
T_q^{(U)}(w\alpha, k^2) &= \frac{R_U(k^2)}{(w\alpha + k^2 - \mu_1^2 + i\Gamma_1)(w\alpha + k^2 - \mu_2^2 + i\Gamma_2)} \\
T_q^{(S)}(w\alpha, k^2) &= \frac{R_S(k^2)}{(w\alpha + k^2 - \mu_3^2 + i\Gamma_3)(w\alpha + k^2 - \mu_4^2 + i\Gamma_4)},
\end{aligned} \tag{41}$$

where $k^2 = -w\alpha\beta - k_\perp^2$ and

$$\mu_j^2 = M_j^2 - p^2. \tag{42}$$

We suggest that values of μ_j^2 and Γ_j should be within the non-perturbative scale domain, with $M_j^2 > \Gamma_j$. It is convenient to write $T_{U,S}$ as the sum of two resonances:

$$\begin{aligned}
T_q^{(U)}(w\alpha, k^2) &= \frac{R_U(k^2)}{(\mu_1^2 - \mu_2^2) - i(\Gamma_1 - \Gamma_2)} \left[\frac{1}{(w\alpha + k^2 - \mu_1^2 + i\Gamma_1)} - \frac{1}{(w\alpha + k^2 - \mu_2^2 + i\Gamma_2)} \right], \\
T_q^{(S)}(w\alpha, k^2) &= \frac{R_S(k^2)}{(\mu_3^2 - \mu_4^2) - i(\Gamma_3 - \Gamma_4)} \left[\frac{1}{(w\alpha + k^2 - \mu_3^2 + i\Gamma_3)} - \frac{1}{(w\alpha + k^2 - \mu_4^2 + i\Gamma_4)} \right].
\end{aligned} \tag{43}$$

It seems that specifying $R_{U,S}$ cannot be done unambiguously. We postpone investigating this problem to the future while in the present paper we use $R_{U,S}$ defined as follows:

$$R_U = \lambda_U \left(\frac{k^2}{k^2 + \mu_U^2} \right)^{1+\eta}, \quad R_S = \lambda_S \left(\frac{k^2}{k^2 + \mu_S^2} \right)^{1+\eta}, \tag{44}$$

where $\lambda_{U,S}$ and $\mu_{U,S}^2$, ($\mu_{U,S}^2 > 0$) are independent parameters, though we think that R_U and R_S could coincide which would diminish the number of free parameters. It is easy to check now that the expressions for $T_q^{(U)}, T_q^{(S)}$ introduced in Eq. (41) obey the condition of IR stability in Eq. (12) with arbitrary η . In contrast, the value of the UV parameter χ (introduced in Eqs. (14,20) to guarantee UV stability) is now fixed: $\chi = 1$ in Eq. (41). Eqs. (25,41) are suggested for invariant amplitudes $T_q^{(U,S)}$ in Basic factorization. Reducing Basic factorization to k_T -factorization converts $T_q^{(U,S)}$ into new amplitudes $\tilde{T}_q^{(U,S)}$. They are obtained from $T_q^{(U,S)}$ by integrating them with respect to α :

$$\tilde{T}_q^{(r)}(\beta, k_\perp^2) = \int_0^{\alpha_{\max}} d\alpha T^{(r)}(\alpha, k^2), \tag{45}$$

where $r = U, S$. The upper limit of integration, α_{\max} should obey Eq. (34), so we choose

$$\alpha_{\max} \approx k_\perp^2 / (w\beta). \tag{46}$$

According to Eq. (34), $k^2 \approx -k_\perp^2$. The integration leads to the following expression for $\tilde{T}_q^{(j)}$ (see Appendix D for detail):

$$\begin{aligned}\tilde{T}_q^{(U)}(\beta, k_\perp^2) &\approx \frac{1}{2} R_U(k_\perp^2) \left[\frac{1}{k_\perp^2(1-\beta)/\beta - \mu_1^2 + i\Gamma_1} + \frac{1}{k_\perp^2(1-\beta)/\beta - \mu_2^2 + i\Gamma_2} \right] + \Delta\tilde{T}_q^{(U)}, \\ \tilde{T}_q^{(S)}(\beta, k_\perp^2) &\approx \frac{1}{2} R_S(k_\perp^2) \left[\frac{1}{k_\perp^2(1-\beta)/\beta - \mu_3^2 + i\Gamma_3} + \frac{1}{k_\perp^2(1-\beta)/\beta - \mu_4^2 + i\Gamma_4} \right] + \Delta\tilde{T}_q^{(S)},\end{aligned}\quad (47)$$

where $\Delta\tilde{T}_q^{(U)}$ and $\Delta\tilde{T}_q^{(S)}$ are

$$\begin{aligned}\Delta\tilde{T}_q^{(U)} &= \frac{1}{(\mu_1^2 - \mu_2^2) + i(\Gamma_1 - \Gamma_2)} \ln \left(\frac{k_\perp^2 + \mu_1^2 - i\Gamma_1}{k_\perp^2 + \mu_2^2 - i\Gamma_2} \right), \\ \Delta\tilde{T}_q^{(S)} &= \frac{1}{(\mu_3^2 - \mu_4^2) + i(\Gamma_3 - \Gamma_4)} \ln \left(\frac{k_\perp^2 + \mu_3^2 - i\Gamma_3}{k_\perp^2 + \mu_4^2 - i\Gamma_4} \right).\end{aligned}\quad (48)$$

They depend on k_\perp very slowly and they can be neglected at large k_\perp^2 .

B. Primary quark distributions

The Optical theorem relates the s -channel imaginary parts of $T^{(U,S)}$ and $\tilde{T}^{(U,S)}$ to the primary quark distributions $\Psi_{U,S}$ in Basic factorization and to unintegrated (or) quark distributions $\Phi_{U,S}$ in k_T -factorization respectively. So applying the Optical theorem, we obtain the following expression for the primary quark distribution Ψ_r in Basic factorization:

$$\begin{aligned}\Psi_U(w\alpha, k^2) &= \frac{1}{\pi} \frac{R_U(k^2)}{(\mu_1^2 - \mu_2^2)} \left[\frac{\Gamma_1}{(w\alpha + k^2 - \mu_1^2)^2 + \Gamma_1^2} - \frac{\Gamma_2}{(w\alpha + k^2 - \mu_2^2)^2 + \Gamma_2^2} \right], \\ \Psi_S(w\alpha, k^2) &= \frac{1}{\pi} \frac{R_S(k^2)}{(\mu_3^2 - \mu_4^2)} \left[\frac{\Gamma_3}{(w\alpha + k^2 - \mu_3^2)^2 + \Gamma_3^2} - \frac{\Gamma_4}{(w\alpha + k^2 - \mu_4^2)^2 + \Gamma_4^2} \right].\end{aligned}\quad (49)$$

and a similar expression for the primary quark distribution $\Phi_{U,S}$ in k_T -factorization:

$$\begin{aligned}\Phi_U(\beta, k_\perp^2) &= \frac{1}{\pi} R_U(k_\perp^2) \left[\frac{\Gamma_1}{(k_\perp^2(1-\beta)/\beta - \mu_1^2)^2 + \Gamma_1^2} + \frac{\Gamma_2}{(k_\perp^2(1-\beta)/\beta - \mu_2^2)^2 + \Gamma_2^2} \right], \\ \Phi_S(\beta, k_\perp^2) &= \frac{1}{\pi} R_S(k_\perp^2) \left[\frac{\Gamma_3}{(k_\perp^2(1-\beta)/\beta - \mu_3^2)^2 + \Gamma_3^2} + \frac{\Gamma_4}{(k_\perp^2(1-\beta)/\beta - \mu_4^2)^2 + \Gamma_4^2} \right].\end{aligned}\quad (50)$$

Obviously, the expressions in Eqs. (49,50) are of the Breit-Wigner type. Substituting Eq. (50) in Eq. (35) and integrating over k_\perp^2 , we arrive at the quark parton distribution $D_j^{(col)}$ in k_T -factorization, where the non-perturbative contributions i.e. the unintegrated parton distributions are specified:

$$\begin{aligned}D_U^{(k_T)}(x, q_\perp^2) &= \frac{1}{\pi} \int_x^1 \frac{d\beta}{\beta} \int_0^r \frac{dk_\perp^2}{k_\perp^2} D_U^{(pert)}(x/\beta, k_\perp^2/q_\perp^2) R_U(k_\perp^2) \\ &\quad \left[\frac{\Gamma_1}{(k_\perp^2(1-\beta)/\beta - \mu_1^2)^2 + \Gamma_1^2} + \frac{\Gamma_2}{(k_\perp^2(1-\beta)/\beta - \mu_2^2)^2 + \Gamma_2^2} \right], \\ D_S^{(k_T)}(x, q_\perp^2) &= \frac{1}{\pi} \int_x^1 \frac{d\beta}{\beta} \int_0^r \frac{dk_\perp^2}{k_\perp^2} D_S^{(pert)}(x/\beta, k_\perp^2/q_\perp^2) \\ &\quad R_S(k_\perp^2) \left[\frac{\Gamma_3}{(k_\perp^2(1-\beta)/\beta - \mu_3^2)^2 + \Gamma_3^2} + \frac{\Gamma_4}{(k_\perp^2(1-\beta)/\beta - \mu_4^2)^2 + \Gamma_4^2} \right],\end{aligned}\quad (51)$$

where r is defined in Eq. (35). Let us consider the k_\perp -dependence in Eqs. (50,35) in more detail. Obviously, the structures of expressions for $D_U^{(k_T)}$ and $D_S^{(k_T)}$ (or Φ_u and Φ_S) are quite similar, so we consider $D_U^{(k_T)}$ only. Then, the expression in the squared brackets in Eq. (35), i.e. Φ_U of Eq. (50), is symmetric with respect to replacement $1 \rightleftharpoons 2$. Each term in the parentheses has a peaked form, with maximums at $k_\perp^2 = \mu_{1,2}^2$. The less $\Gamma_{1,2}$, the sharper the peaks are. We remind that $R_{U,S} \sim (k_\perp^2)^{1+\eta}$ at small k_\perp^2 . By definition, see Eq. (42), $\mu_{1,2}^2 = M_{1,2}^2 - p^2$, so they can be either positive or negative while k_\perp^2 cannot be negative. In any case the both terms in Φ_U and Φ_S contribute to $D_{U,S}^{(k_T)}$ but a result of interference of the two peaks depends on values of the parameters. There are possible three particular cases:

Case (i): both μ_1^2 and μ_2^2 are positive.

In this the both maximums are within the integration region of Eq. (35) and interference of the two peaks generates various forms of $\Phi_U(\beta, k_\perp^2)$ ranging from the picture with two isolated peaks to a kind of plateau, depending on values of $\Gamma_{1,2}$.

Case (ii): $\mu_1^2 > 0$ and $\mu_2^2 < 0$ or vice versa.

Here the peak from the first term in Eq. (50) combines with a tail of the contribution of the second term whose maximum is beyond the integration region of Eq. (35). The resulting picture has a resemblance to the dual model combining a resonant and a constant term.

Case (iii): both μ_1^2 and μ_2^2 are negative.

The both maximums now are out of the integration region, so tails of the peaks, taken by themselves, generate a form slow decreasing with growth of k_\perp^2 . However, this slope is affected by an impact of R_U . We remind that $R_U = 0$ at $k_\perp^2 = 0$.

C. Primary quark distributions in Collinear factorization

Performing integration over k_\perp^2 in Eq. (35), we arrive at the parton distributions $D_j^{(col)}$ in Collinear factorization. Presuming that parameter Γ_j is small, we write the result of the integration in the following form (see Appendix C for detail):

$$D_U^{(col)}(x, q_\perp^2) \approx \int_x^1 \frac{d\beta}{\beta} D_U^{(pert)}(x/\beta, q_\perp^2/\mu_1^2) \phi_U(\beta, \mu_1^2) + \int_x^1 \frac{d\beta}{\beta} D_U^{(pert)}(x/\beta, q_\perp^2/\mu_2^2) \phi_U(\beta, \mu_2^2), \quad (52)$$

with

$$\begin{aligned} \phi_U(\beta, \mu_1^2) &\approx \frac{1}{\pi} \int_{\Omega_1} \frac{dk_\perp^2}{k_\perp^2} \frac{R_U(k_\perp^2) \Gamma_1}{(k_\perp^2(1-\beta)/\beta - \mu_1^2)^2 + \Gamma_1^2}, \\ \phi_U(\beta, \mu_2^2) &\approx \frac{1}{\pi} \int_{\Omega_2} \frac{dk_\perp^2}{k_\perp^2} \frac{R_U(k_\perp^2) \Gamma_2}{(k_\perp^2(1-\beta)/\beta - \mu_2^2)^2 + \Gamma_2^2}, \end{aligned} \quad (53)$$

where the integration regions $\Omega_1 = \Omega'_1 \cap [0, w]$ and $\Omega_2 = \Omega'_2 \cap [0, w]$, with the subregions Ω'_1, Ω'_2 being located around the maximums of the peaks. Formally, the both terms in Eq. (53) contribute to ϕ_U at any signs of μ_1^2, μ_2^2 , but in the limit of sharp peaks these contributions have different weights: At $\mu_1^2 > 0, \mu_2^2 > 0$ the both terms contribute equally:

$$\phi_U \approx R_U(\mu_1^2\beta/(1-\beta))/\mu_1^2 + R_U(\mu_2^2\beta/(1-\beta))/\mu_2^2 + O(\Gamma_1, \Gamma_2). \quad (54)$$

Mostly the first term contributes, when $\mu_1^2 > 0, \mu_2^2 < 0$:

$$\phi_U \approx R_U(\mu_1^2\beta/(1-\beta))/\mu_1^2 + O(\Gamma_1). \quad (55)$$

and vice versa. Finally, at $\mu_1^2, \mu_2^2 < 0$ only tails of the both peaks contribute and therefore ϕ_U is small and flat compared to the previous cases:

$$\phi_U \approx const. \quad (56)$$

(see Appendix E for detail). When $\mu_1^2 > 0, \mu_2^2 > 0$

$$\varphi_U(\omega, \mu^2) = \int_x^1 \frac{d\beta}{\beta} \beta^\omega [E(\omega, \mu^2, \mu_1^2) R_U(\mu_1^2 \beta / (1 - \beta)) \mu_2^{-2} + E(\omega, \mu^2, \mu_2^2) R_U(\mu_2^2 \beta / (1 - \beta)) \mu_2^{-2}], \quad (57)$$

$$R_U(\mu_j^2 \beta / (1 - \beta)) \mu_j^{-2} = \lambda_U \beta. \quad (58)$$

Combining Eqs. (57) and (58), integrating over β and remembering that at small x essential values of ω are small leads to the following expression for $\varphi_U(\omega, \mu^2)$ (see Appendix E for detail):

$$\begin{aligned} \varphi_U(\omega, \mu^2) &= \int_x^1 \frac{d\beta}{\beta} \beta^{\omega+1} \left[\frac{\lambda_U}{\mu_1^2} E(\omega, \mu^2, \mu_1^2) + \frac{\lambda_U}{\mu_2^2} E(\omega, \mu^2, \mu_2^2) \right] \\ &\approx \frac{\lambda_U}{\mu_1^2} E(\omega, \mu^2, \mu_1^2) + \frac{\lambda_U}{\mu_2^2} E(\omega, \mu^2, \mu_2^2). \end{aligned} \quad (59)$$

VI. CONCLUSION

In the present paper we have considered the quark-hadron scattering amplitudes and distributions of polarized and unpolarized quarks in hadrons in the framework of the factorization concept where the both amplitudes and distributions are expressed through convolutions of the perturbative and non-perturbative components. We began with considering the quark-hadron amplitudes in Basic factorization where integration over momenta of connecting partons runs over the whole phase space and obtained the conditions for the factorization convolution to be stable both in IR and UV regions. Then we demonstrated how to reduce Basic factorization to K_T - and Collinear factorizations. We suggested a Resonance Model for non-perturbative contributions to the unpolarized and spin-dependent parton-hadron scattering amplitudes. This model is based on the simple argumentation: after emitting an active quark by a hadron, the remaining colored quark-gluon state cannot be stable and therefore it can be described by quasi-resonant expressions. We needed at least two resonances in Basic factorization and this remained true when Basic factorization was reduced to K_T -factorization. Applying the Optical theorem to the Resonance Model provided us first with the expressions of the Breit-Wigner type for non-perturbative (primary) contributions to the quark distributions in Basic and K_T -factorizations and then, after one more reduction, to the parton distributions in Collinear factorization. To conclude, let us notice that the Resonance Model can also be used for analysis of the non-singlet components of the DIS structure functions.

ACKNOWLEDGMENTS

We are grateful to B.L. Webber for interesting discussions. S.I. Troyan is supported by Russian Science Foundation, Grant No. 14-22-00281.

Appendix A: Amplitude for the forward Compton scattering off a gluon in the box approximation

$$M_{\mu\nu\lambda\rho} = (t_l t_r) \frac{e^2 \alpha_s}{8\pi^2} w \int d\beta' d\alpha' dk_\perp'^2 [M_{\mu\nu\lambda\rho}^{(1)} + M_{\mu\nu\lambda\rho}^{(2)} + M_{\mu\nu\lambda\rho}^{(3)}] \quad (A1)$$

$$\begin{aligned} M_{\mu\nu\lambda\rho}^{(1)} &= \frac{Tr [\gamma_\nu (\hat{q} + \hat{k}') \gamma_\mu \hat{k}' \gamma_\lambda (\hat{k}' - \hat{k}) \gamma_\rho \hat{k}']}{k'^2 k'^2 (q + k')^2 (k' - k)^2}, \\ M_{\mu\nu\lambda\rho}^{(2)} &= \frac{Tr [\gamma_\nu (\hat{q} + \hat{k}') \gamma_\mu \hat{k}' \gamma_\rho (\hat{k}' + \hat{k}) \gamma_\lambda \hat{k}']}{k'^2 k'^2 (q + k')^2 (k' + k)^2}, \\ M_{\mu\nu\lambda\rho}^{(3)} &= \frac{Tr [\gamma_\nu (\hat{q} + \hat{k}' - \hat{k}) \gamma_\rho (\hat{k}' - \hat{k}) \gamma_\mu \hat{k}' \gamma_\lambda (\hat{k}' + \hat{q})]}{k'^2 (q + k')^2 (k' - k)^2 (q + k' - k)^2} \end{aligned} \quad (A2)$$

Appendix B: Projection operators for forward Compton amplitudes

The conventional of dealing with the forward Compton scattering amplitude $A_{\mu\nu}$ is, in the first place, to simplify their tensor structure. To this end, $A_{\mu\nu}$ is represented as an expansion of $A_{\mu\nu}$ into the series of simpler tensors, each multiplied by an invariant amplitude. Such tensors are called projection operators. Through the Optical theorem the invariant amplitudes are related to the DIS structure functions.

In the case of the unpolarized Compton scattering such an expansion looks as follows:

$$A_{\mu\nu} = P_{\mu\nu}^{(1)} A_1 + P_{\mu\nu}^{(2)} A_2, \quad (\text{B1})$$

where

$$P_{\mu\nu}^{(1)} = -g_{\mu\nu} + q_\mu q_\nu / q^2, \quad P_{\mu\nu}^{(2)} = (1/pq) (p_\mu - q_\mu(pq/q^2)) (p_\nu - q_\nu(pq/q^2)) \quad (\text{B2})$$

are the projection operators and A_1 , A_2 are invariant amplitudes. According to the Optical theorem

$$F_1 = \frac{1}{\pi} \Im A_1, \quad F_2 = \frac{1}{\pi} \Im A_2. \quad (\text{B3})$$

Similarly, for the polarized Compton scattering

$$A_{\mu\nu} = P_{\mu\nu}^{(3)} A_3 + P_{\mu\nu}^{(4)} A_4, \quad (\text{B4})$$

where

$$P_{\mu\nu}^{(3)} = i\epsilon_{\mu\nu\lambda\rho} M q_\lambda S_\rho, \quad P_{\mu\nu}^{(4)} = i\epsilon_{\mu\nu\lambda\rho} M q_\lambda [S_\rho - p_\rho (qS/pq)], \quad (\text{B5})$$

with M and S being the hadron mass and spin respectively, and $A_{3,4}$ are spin-dependent invariant amplitudes. The Optical theorem states that

$$g_1 = \frac{1}{\pi} \Im A_3, \quad g_2 = \frac{1}{\pi} \Im A_4. \quad (\text{B6})$$

All operators $P_{\mu\nu}^{(n)}$ respect the electromagnetic current conservation: $q_\mu P_{\mu\nu}^{(n)} = q_\nu P_{\mu\nu}^{(n)} = 0$. It is convenient to introduce the longitudinal, S^\parallel and transverse, S^\perp components of the spin, so that $S^\perp p = S^\perp q = 0$ and $S_\rho^\parallel = p_\rho (qS/pq)$. In such terms Eq. (B4) can be written as follows:

$$A_{\mu\nu} = i\epsilon_{\mu\nu\lambda\rho} M q_\lambda \left[S_\rho^\parallel A_3 + S_\rho^\perp (A_3 + A_4) \right] \equiv i\epsilon_{\mu\nu\lambda\rho} M q_\lambda \left[S_\rho^\parallel A^\parallel + S_\rho^\perp A^\perp \right]. \quad (\text{B7})$$

This expression is useful for practical attributing different terms in the spin-dependent $A_{\mu\nu}$ to proper invariant amplitudes. In the unpolarized case one can use the simple rule: expressions $\sim g_{\mu\nu}$ contribute to A_1 while expressions $\sim p_\mu p_\nu / pq$ form A_2 . In contrast, the gauge invariance admits adding arbitrary terms $\sim q_\mu$, q_ν .

Appendix C: Convolutions involving the Breit-Wigner formula

Let us consider the following convolution:

$$F = \frac{1}{\pi} \int_{-\infty}^{\infty} dx f(x) \frac{\Gamma}{(x - x_0)^2 + \Gamma^2}. \quad (\text{C1})$$

Replacing x by t , with $t = (x - x_0)/\Gamma$, we convert Eq. (C1) into

$$F = \frac{1}{\pi} \int_{-\infty}^{\infty} dt f(t\Gamma + x_0) \frac{\Gamma}{t^2 + 1}. \quad (\text{C2})$$

At small Γ , we can expand $f(t\Gamma + x_0)$ in the power series and retain several terms:

$$f(t\Gamma + x_0) = f(x_0) + f'(x_0)t\Gamma + O(\Gamma^2). \quad (\text{C3})$$

Substituting Eq. (C3) in (C2) and integrating (C2) yields

$$F = f(x_0) + O(\Gamma). \quad (\text{C4})$$

The first term in Eq. (C4) corresponds to the well-known representation of the δ -function:

$$\frac{1}{\pi} \lim_{\epsilon \rightarrow 0} \frac{\epsilon}{x^2 + \epsilon^2} = \delta(x) \quad (\text{C5})$$

Appendix D: Integration in Eq. (45)

A generic expression to integrate can be written as

$$\tilde{T} = \int_0^{\alpha_{\max}} \frac{d\alpha}{(\alpha - A)(\alpha - B)} = \frac{1}{(A - B)} \int_0^{\alpha_{\max}} d\alpha \left[\frac{1}{\alpha - A} - \frac{1}{\alpha - B} \right] = \frac{1}{(A - B)} \left[\ln \left(\frac{\alpha_{\max} - A}{\alpha_{\max} - B} \right) - \ln \left(\frac{A}{B} \right) \right]. \quad (\text{D1})$$

Assuming that $\alpha_{\max} \gg A, B \gg |A - B|$ allows us to expand the logarithms into the power series and retain the first terms only:

$$\tilde{T} \approx \frac{1}{2} \left[\frac{1}{\alpha_{\max} - A} + \frac{1}{\alpha_{\max} - B} \right] - \frac{1}{(A - B)} \ln(A/B). \quad (\text{D2})$$

We have written Eq. (D2) in the symmetrical form with respect to A, B because Eq. (D1) has this feature.

Appendix E: Evolving the factorization scale in Collinear factorization

Using the Mellin transform, we can rewrite Eq. (37) as follows:

$$D^{(col)}(x, q_{\perp}^2) = \int_{-\imath\infty}^{\imath\infty} \frac{d\omega}{2\pi\imath} x^{-\omega} C_U(\omega) E(\omega, q^2, \mu^2) \phi(\omega, \mu^2), \quad (\text{E1})$$

where the primary quark distribution is fixed at the scale μ , with $\mu^2 < q_{\perp}^2$. $E(\omega, q_{\perp}^2, \mu^2)$ is a generic notation for an operator evolving the distribution $\phi(\omega, \mu^2)$ from the factorization scale μ^2 to q_{\perp}^2 , while $C(\omega)$ is responsible for the x -evolution. Choosing a scale $\tilde{\mu}$ such that $\mu^2 < \tilde{\mu} < q_{\perp}^2$ and representing $E(\omega, q_{\perp}^2, \mu^2)$ as

$$E(\omega, q_{\perp}^2, \mu^2) = E(\omega, q_{\perp}^2, \tilde{\mu}^2) E(\omega, \tilde{\mu}^2, \mu^2), \quad (\text{E2})$$

we bring $D^{(col)}$ to the conventional form

$$D^{(col)}(x, q_{\perp}^2) = \int_{-\imath\infty}^{\imath\infty} \frac{d\omega}{2\pi\imath} x^{-\omega} C_U(\omega) E(\omega, q^2, \tilde{\mu}^2) \varphi(\omega, \tilde{\mu}^2), \quad (\text{E3})$$

where

$$\varphi(\omega, \tilde{\mu}^2) = E(\omega, \tilde{\mu}^2, \mu^2) \phi(\beta, \mu^2), \quad (\text{E4})$$

which corresponds to Eq. (39). Actually, $\varphi(\omega, \tilde{\mu}^2)$ is the conventional parton distribution in the ω -space (momentum space). It is fixed at an arbitrary scale $\tilde{\mu}^2$ and related to the standard integrated distribution $\delta q(x, \tilde{\mu}^2)$ by the Mellin transform. The evolution operator $E(\omega, q^2, \mu^2)$ is expressed in different terms, depending on the perturbative approach in use. For instance, in LO DGLAP with fixed α_s it is given by

$$E = \exp [\alpha_s \gamma_0(\omega) \ln(q^2/\mu^2)], \quad (\text{E5})$$

with γ_0 being the LO DGLAP anomalous dimension and $C_F = 4/3$. When in LO DGLAP α_s is running and the standard parametrization $\alpha_s = \alpha_s(k_\perp^2)$ is used in the Feynman graphs, Eq. (E5) is changed by

$$E = \left(\frac{q^2}{\mu^2} \right)^{\gamma_0/b}, \quad (\text{E6})$$

with b being the first coefficient of the β -function. The parametrization $\alpha_s = \alpha_s(k_\perp^2)$ should not be used at small x (see Ref. [14]). When it is replaced by the appropriate parametrization and when the total resummation of the leading logarithms is done, Eq. (E6) is replaced by

$$E = \exp [h(\omega) \ln(q^2/\mu^2)], \quad (\text{E7})$$

where $h(\omega)$ is a new anomalous dimension. It accounts for the total resummation of the leading double-logarithmic contributions and running QCD coupling effects (see Ref. [13] and overview [14] for detail).

-
- [1] D. Amati, R. Petronzio, G. Veneziano. Nucl. Phys. B 140 (1978) 54; A.V. Efremov, A.V. Radyushkin. Teor.Mat.Fiz. 42 (1980) 147; Theor.Math.Phys.44 (1980)573; Teor.Mat.Fiz.44 (1980)17; Phys.Lett.B63 (1976) 449; Lett.Nuovo Cim.19 (1977)83; S. Libby, G. Sterman. Phys. Rev. D18 (1978) 3252. S.J. Brodsky and G.P. Lepage. Phys. Lett. B 87 (1979) 359; Phys. Rev. D 22 (1980) 2157; J.C. Collins and D.E. Soper. Nucl. Phys.B 193 (1981) 381; J.C. Collins and D.E. Soper. Nucl. Phys.B 194 (1982) 445; J.C. Collins, D.E. Soper and G. Sterman. Nucl. Phys.B 250 (1985) 199. A.V. Efremov and A.V. Radyushkin. Report JINR E2-80-521; Mod.Phys.Lett. A24 (2009) 2803.
- [2] G. Altarelli and G. Parisi, Nucl. Phys.B126 (1977) 297; V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. 15 (1972) 438; L.N.Lipatov, Sov. J. Nucl. Phys. 20 (1972) 95; Yu.L. Dokshitzer, Sov. Phys. JETP 46 (1977) 641.
- [3] E.A. Kuraev, L.N. Lipatov and V.S. Fadin, Sov. Phys. JETP 44, 443 (1976); E.A. Kuraev, L.N. Lipatov and V.S. Fadin, Sov. Phys. JETP 45, 199 (1977); I.I. Balitsky and L.N. Lipatov, Sov. J. Nucl. Phys. 28, 822 (1978).
- [4] S. Catani, M. Ciafaloni, F. Hautmann. Phys. Lett. B (1990) 97; Nucl.Phys.B366 (1991) 135.
- [5] B.I. Ermolaev, M. Greco, S.I. Troyan. Eur.Phys.J. C71 (2011) 1750; Eur.Phys.J. C72 (2012) 1953.
- [6] V.V. Sudakov. Sov. Phys. JETP 3(1956)65.
- [7] S.J. Brodsky, P. Hoyer, C. Peterson, N. Sakai. Phys.Lett. B93 (1980) 451; Stanley J. Brodsky, C. Peterson, N. Sakai. Phys.Rev. D23 (1981) 2745
- [8] J.C. Collins. PoS LC2008 (2008) 028.
- [9] I.O. Cherednikov, N.G. Stefanis. AIP Conf.Proc. 1105 (2009) 327; Int.J.Mod.Phys.Conf.Ser. 4 (2011) 135-145.
- [10] O. Erdogan, G. Sterman. Phys. Rev. D 91 (2015) 6, 065033.
- [11] J.C. Collins, T.C. Rogers, A.M. Stasto. Phys. Rev. D77 (2008) 085009.
- [12] J. Collins. Int.J.Mod.Phys.Conf.Ser. 4 (2011) 85.
- [13] B.I. Ermolaev, M. Greco, S.I. Troyan. Nucl.Phys. B571 (2000) 137; Nucl.Phys. B594 (2001) 71.
- [14] B.I. Ermolaev, M. Greco, S.I. Troyan. Riv.Nuovo Cimento 33 (2010) 37.