Froissart Bound on Inelastic Cross Section Without Unknown Constants

André Martin^{1, *} and S. M. $Roy^{2, \dagger}$

¹Theoretical Physics Division, CERN, Geneva ²HBCSE, Tata Institute of Fundamental Research, Mumbai

Assuming that axiomatic local field theory results hold for hadron scattering, André Martin and S. M. Roy recently obtained absolute bounds on the D-wave below threshold for pion-pion scattering and thereby determined the scale of the logarithm in the Froissart bound on total cross sections in terms of pion mass only. Previously, Martin proved a rigorous upper bound on the inelastic cross-section σ_{inel} which is one-fourth of the corresponding upper bound on σ_{tot} , and Wu, Martin,Roy and Singh improved the bound by adding the constraint of a given σ_{tot} .

PACS numbers: 03.67.-a, 03.65.Ud, 42.50.-p

I. Introduction. Recently we [1] have obtained bounds on energy averages of the total cross-section without any unknown constants such as an overall constant factor or the scale factor in the logarithm. The purpose of the present work is to obtain analogous bounds on the energy averaged inelastic cross section without any unknown constants. The background and the basic postulates are again summarized below to make this work self-contained.

The Froissart [2] bound on the total cross-section $\sigma_{tot}(s)$ for two particles at c.m. energy \sqrt{s} ,

$$\sigma_{tot}(s) \leq_{s \to \infty} C \left[\ln(s/s_0) \right]^2,\tag{1}$$

(where C, s_0 are unknown constants) was initially proved assuming the Mandelstam representation. This assumption might not be valid, for example, if there are rising Regge trajectories. Fortunately, [3] it was possible to prove this bound rigorously in the much more general frame work of Wightman's [4] axiomatic local field theory as applied to hadrons. Later, the needed analyticity properties , and polynomial boundedness at fixed momentum transfer squared t, were obtained by Epstein, Glaser and Martin [5] in the even more general framework of the theory of local observables of Haag, Kastler and Ruelle [6]. It has nevertheless been questioned [7] if these properties apply to hadrons made of quarks and gluons. Zimmermann [8] has shown that local fields can be associated to composite particles. We decide to believe that this proof applies to the present situation. We postulate that the analyticity and polynomial boundedness derived from local field theory holds for hadrons .

In the proof of the Froissart bound in [3] a crucial role is played by the use of unitarity to enlarge the Lehmann ellipse of analyticity [9] for the absorptive part A(s,t) to show that the right extremity t_0 of the enlarged ellipse in the *t*-plane stays non-zero and positive

^{*}Electronic address: martina@mail.cern.ch

 $^{^{\}dagger} Electronic address: smroy@hbcse.tifr.res.in$

when $s \to \infty$. For many processes, for example for $\pi\pi, KK, K\overline{K}, \pi K, \pi N, \pi \Lambda$ scattering it is known [10] that $t_0 = 4m_{\pi}^2, m_{\pi}$ being the pion-mass . (Except when especially necessary to show the dependence on pion-mass, we shall choose units $m_{\pi} = 1$). Using unitarity and validity of dispersion relations with a finite number of subtractions for $-T < t \leq 0$, Jin and Martin [11]) proved twice subtracted fixed-t dispersion relations for $|t| < t_0$. From this Lukaszuk and Martin [12] fixed the constant C in the Froissart bound to obtain,

$$\sigma_{tot}(s) \leq_{s \to \infty} \sigma_{max}(s), \tag{2}$$

where,

$$\sigma_{max}(s) \equiv 4\pi/(t_0 - \epsilon) \left[\ln(s/s_0)\right]^2 \tag{3}$$

where ϵ is an arbitrarily small positive constant. The Froissart-Martin bounds have inspired much work on high energy theorems (see e.g. [13], [14]) and on models of high energy scattering [15]. Further, Martin proved a bound on the total inelastic cross section $\sigma_{inel}(s)$ at high energy [16] which is one-fourth of the above bound $\sigma_{max}(s)$ on the total cross-section; and, Wu, Martin, Roy and Singh [17] obtained a bound on $\sigma_{inel}(s)$ which improves that bound if $\sigma_{tot}(s)$ is known,

$$\sigma_{inel}(s) \leq_{s \to \infty} \sigma_{tot}(s) \left(1 - \sigma_{tot}(s) / \sigma_{max}(s)\right).$$
(4)

The motivation for getting a bound on the inelastic cross section is the almost general belief [15] that at high energies hadron total cross sections cannot exceed twice the inelastic cross section. Hence, gaining a factor of 4 in the inelastic cross section gains a factor of 2 in the total cross section.

One shortcoming of these bounds from the standpoint of rigour is that [18] they are deduced by assuming that the absorptive part $A(s,t), 0 \leq t < t_0$ is bounded by $\text{Const.} s^2/\ln(s/s_0)$ for $s \to \infty$, whereas axiomatic field theory results only guarantee that

$$C(t) \equiv \int_{s_{th}}^{\infty} ds A(s,t) / s^3 < \infty, \ 0 \le t < t_0, \qquad (5)$$

where s_{th} is the *s*-channel threshold. From the practical point of view a more serious shortcoming is that they involve the unknown scale factor s_0 in the argument of the logarithm and the unknown arbitrarily small but non-zero constant ϵ .

In the case of the total cross section, both these shortcomings were removed recently [1]. Bounds on energy averages of the total cross-section were obtained in which the scale s_0 is determined in terms of C(t). C(t) can be bounded rigorously in terms of pion mass alone for $\pi^0 \pi^0$ scattering. Thus we obtained the absolute bound[1],

$$\bar{\sigma}_{tot}(s,\infty) \leq \pi(m_{\pi})^{-2} [\ln(s/s_0) + (1/2) \ln \ln(s/s_0) + 1]^2 + O(\ln \ln(s/s_0)), \ s_0^{-1} = 17\pi \sqrt{\pi/2} \ m_{\pi}^{-2}, \ (6)$$

where,

$$\bar{\sigma}_{tot}(s,\infty) \equiv s \int_{s}^{\infty} ds' \sigma_{tot}(s') / s'^{2}.$$
 (7)

We also obtained somewhat improved bounds by using the additional phenomenological inputs for the D-wave scattering length [19] for pion-pion scattering.

We prove here analogous bounds on energy averages of the inelastic cross section. We choose the same normalizations as in Martin-Roy [1]. F(s,t) denotes an $ab \rightarrow ab$ scattering amplitude at c.m. energy \sqrt{s} and momentum transfer squared t normalized for non-identical partcles a, b such that the differential cross-section $d\sigma/d\Omega(s,t)$ is given by $|4F(s,t)/\sqrt{s}|^2$, with t being given in terms of the c.m. momentum k and the scattering angle θ by the relation,

$$t = -2k^2(1 - \cos\theta); \ z \equiv \cos\theta = 1 + t/(2k^2).$$
 (8)

Then, for fixed s larger than the physical s-channel threshold, $F(s; \cos \theta) \equiv F(s, t)$ is analytic in the complex $\cos \theta$ -plane inside the Lehmann-Martin ellipse with foci -1 and +1 and semi-major axis $\cos \theta_0 = 1 + t_0/(2k^2)$. Within the ellipse ,in particular, for $|t| < t_0, F(s, t)$ has the convergent partial wave expansion,

$$F(s,t) = \frac{\sqrt{s}}{4k} \sum_{l=0}^{\infty} (2l+1)P_l(z)a_l(s), \qquad (9)$$

with the unitarity constraint,

$$Ima_l(s) \ge |a_l(s)|^2, \ s \ge s_{th} \ .$$
 (10)

Correspondingly, the optical theorem gives, for nonidentical partcles a, b,

$$\sigma_{tot}(s) = \frac{4\pi}{k} Im(4F(s,0)/\sqrt{s})$$

= $\frac{4\pi}{k^2} \sum_{l=0}^{\infty} (2l+1)Ima_l(s)$. (11)

For identical particles ,e.g. for $\pi^0 \pi^0$ scattering, or for pion-pion scattering with Iso-spin *I*, the partial waves $a_l(s) \rightarrow 2a_l^I(s)$ in the partial wave expansion, i.e.

$$F^{I}(s,t) = \frac{\sqrt{s}}{4k} \sum_{l=0}^{\infty} (2l+1)2a_{l}^{I}(s)P_{l}(z), \qquad (12)$$

$$\sigma_{tot}^{I}(s) = \frac{4\pi}{k^2} \sum_{l=0}^{\infty} (2l+1) 2Ima_l^{I}(s) , \qquad (13)$$

and we have the same formula for the differential crosssection in terms of F(s,t), and the same form of the unitarity constraint, $Ima_l^I(s) \ge |a_l^I(s)|^2$, $s \ge 4$, as for non-identical particles. At threshold, $F^I(4,0) = a_0^I$, the S-wave scattering length for Iso-spin I.

It will be seen that proofs of the bounds for inelastic cross sections are considerably more involved than those for total cross sections, but the basic principles are the same. We give detailed derivations for the case of non-identical particles $a \neq b$, and also quote the identical particle results when needed.

II. Convexity Properties of Lower Bound on Absorptive Part in terms of Total Inelastic Cross Section. We obtain a lower bound on the absorptive part of F(s,t) (for $s \ge s_{th}$ and $0 \le t < t_0$) in terms of the inelastic cross section $\sigma_{inel}(s)$. Following [20] we prove that the bound is a convex function of the inelastic cross section. The absorptive part has the partial wave expansion,

$$F_s(s,t) \equiv A(s,t) = \frac{\sqrt{s}}{4k} \sum_{l=0}^{\infty} (2l+1)P_l(z)Ima_l(s).$$
(14)

The corresponding expansion of the inelastic cross section is,

$$\sigma_{inel}(s) = \frac{4\pi}{k^2} \sum_{l=0}^{\infty} (2l+1) \left(Ima_l(s) - |a_l(s)|^2 \right).$$
(15)

Actually we shall vary $Ima_l(s)$ subject to the positivity restrictions (due to unitarity),

$$Ima_l(s) \ge 0 \tag{16}$$

to minimise A(s,t), given,

$$\sigma_{inel,im}(s) \equiv \frac{4\pi}{k^2} \sum_{l=0}^{\infty} (2l+1) \left(Ima_l(s) - (Ima_l(s))^2 \right).$$
(17)

The bound will be seen to be an increasing function of $\sigma_{inel.im}(s)$. Further,

$$\sigma_{inel,im}(s) \ge \sigma_{inel}(s); \tag{18}$$

therefore the bound will still hold when we replace $\sigma_{inel,im}(s)$ by the experimentally accessible $\sigma_{inel}(s)$. We work at a fixed s; so, unless specially needed, we shall suppress the s-dependence of $Ima_l(s), \ \sigma_{inel,im}(s)$ and $\sigma_{inel}(s)$. The Lagrange multiplier method with positivity constraints on partial waves gives the variational solution $(Ima_l)_0$,

$$(Ima_l)_0 = \frac{1}{2} \left(1 - \frac{P_l(z)}{P_\lambda(z)} \right), \ l \le L; L \le \lambda < L+1$$
$$(Ima_l)_0 = 0, \ l > L; \ (19)$$

where the integer L and the non-negative fraction $\lambda - L$, are to be determined so as to reproduce the given $\sigma_{inel,im}$; here $P_{\lambda}(z)$ for non-integer λ and $z \geq 1$ is defined by,

$$P_{\lambda}(z) = \frac{1}{\pi} \int_0^{\pi} (z + \cos\phi\sqrt{z^2 - 1})^{\lambda} d\phi .$$
 (20)

If $A_0(s,t)$ denotes the absorptive part with partial waves $(Ima_l)_0$ and A(s,t) that with arbitrary positive partial waves with the given $\sigma_{inel,im}(s)$, we obtain by direct subtraction,

$$4(k/\sqrt{s}) \left(A(s,t) - A_0(s,t)\right)$$

= $P_{\lambda}(z) \sum_{l=0}^{L} (2l+1) \left(Ima_l - (Ima_l)_0\right)^2 + P_{\lambda}(z) \sum_{l=L+1}^{\infty} (2l+1) \left((Ima_l)^2 + Ima_l \left(\frac{P_l(z)}{P_{\lambda}(z)} - 1\right)\right)$
 $\ge 0.$ (21)

The last inequality follows because for $z \ge 1$, and $\lambda \ge 0$, $P_{\lambda}(z)$ is an increasing function of λ . We then have,

$$4(k/\sqrt{s})A(s,t) \ge \sum_{l=0}^{L} (2l+1)P_l(z)\frac{1}{2} \left(1 - \frac{P_l(z)}{P_{\lambda}(z)}\right) \\ \equiv A(\lambda), \quad (22)$$

where,

$$\sigma_{inel.im} \frac{k^2}{4\pi} = \sum_{l=0}^{l=L} (2l+1)(1/4) \left(1 - \left(\frac{P_l(z)}{P_\lambda(z)}\right)^2\right)$$
$$\equiv \Sigma_I(\lambda). \quad (23)$$

Note that $\Sigma_I(\lambda)$ and $A(\lambda)$ are monotonically increasing continuous functions of λ ; hence λ and $A(\lambda)$ may be considered functions of Σ_I , and

$$dA/d\Sigma_I = \left((dA/d\lambda)/(d\Sigma_I/d\lambda) \right) = P_\lambda(z) ,$$
 (24)

which is always positive, and also continuous at integer λ although $(dA/d\lambda)$ and $(d\Sigma_I/d\lambda)$ are discontinuous there. Hence,

$$d^{2}A/d\Sigma_{I}^{2} = \left((dP_{\lambda}(z)/d\lambda)/(d\Sigma_{I}/d\lambda) \right) > 0, \qquad (25)$$

which is discontinuous at integer λ , but always positive. This completes the proof that $A(\lambda(\Sigma_I))$ is a convex function of Σ_I ; i.e. at a given s the lower bound on A(s,t) is an increasing and convex fuction of $\sigma_{inel.im}$, and hence of σ_{inel} .

III. Explicit evaluation of the bound. Explicitly,

$$A(\lambda(\Sigma_I)) = \int_0^{\Sigma_I} \frac{dA}{d\Sigma'_I} d\Sigma'_I = \int_0^{\Sigma_I} P_{\lambda'}(z) d\Sigma'_I, \quad (26)$$

where λ', L' corresponds to the value Σ'_I of $\sigma_{inel.im}k^2/(4\pi)$. When $0 < \Sigma'_I < (1-z^{-2})/4 \equiv \Sigma_I(1)$ we get L' = 0 and the corresponding part of the integral can be evaluated exactly. Hence,

$$A(\lambda(\Sigma_I)) = (1 - z^{-1})/2 + \int_{\Sigma_I(1)}^{\Sigma_I} P_{\lambda'}(z) d\Sigma'_I.$$
 (27)

In the remaining integral $L' \geq 1$ and we shall prove that for $\Sigma'_I \geq \Sigma_I(1)$,

$$(\lambda')^2 \ge 4\Sigma'_I. \tag{28}$$

Proof. From the partial wave expansion for $\Sigma_I(\lambda') \equiv \Sigma'_I$, we obtain

$$4\Sigma_I' \le (L')^2 + (2L'+1)\left(1 - \left(\frac{P_{L'}(z)}{P_{\lambda'}(z)}\right)^2\right).$$
(29)

The integral representation for $P_{\lambda}(z)$ given before yields, for z > 1,

$$\left(\frac{P_{L'}(z)}{P_{\lambda'}(z)}\right)^2 \ge \exp\left(-2(\lambda'-L')\ln(z+\sqrt{z^2-1})\right) \\\ge 1-2(\lambda'-L')\ln(z+\sqrt{z^2-1}), \quad (30)$$

where the last inequality uses $\exp(-x) \ge 1 - x$. At high energies $\ln(z + \sqrt{z^2 - 1})$ goes to zero, and we assume moderately high energies $(k > 6m_{\pi})$ such that, with $t < 4m_{\pi}^2$,

$$\ln(z + \sqrt{z^2 - 1}) < 1/3. \tag{31}$$

Then,

$$\begin{split} 4\Sigma'_{I} &\leq (L')^{2} + (2L'+1)2(\lambda'-L')\ln(z+\sqrt{z^{2}-1}) \\ &\leq (\lambda')^{2} - (\lambda'-L')(\lambda'+L'-(2/3)((2L'+1))) \\ &\leq (\lambda')^{2}, for \ L' \geq 1, \end{split}$$
(32)

which completes the proof.

Since $P_{\lambda'}(z)$ is an increasing function of λ' , we obtain,

$$A(\lambda(\Sigma_I)) = (1 - z^{-1})/2 + \int_{\Sigma_I(1)}^{\Sigma_I} P_{\sqrt{4\Sigma_I'}}(z) d\Sigma_I'.$$
 (33)

Using the integral representation for $P_{\lambda}(z)$ given before and the analogous representation

$$I_0(z) = \frac{1}{\pi} \int_0^\pi \exp(z\cos\phi)d\phi \tag{34}$$

for the modified Bessel function , and the elementary inequality

$$\ln((z + \cos\phi\sqrt{z^2 - 1}) \ge (\cos\phi)\ln((z + \sqrt{z^2 - 1}), z > 1)$$

we obtain [20],

$$P_{\lambda}(z) \ge I_0 \left(\lambda \ln((z + \sqrt{z^2 - 1})) \right)$$
(35)

$$\geq I_0 \left(\lambda \sqrt{z^2 - 1/z}\right), \ z > 1.$$
 (36)

Substituting the above inequalities , the integral over Σ'_I in the expression for the lower bound can be evaluated exactly, and we have the exact result (without any high energy approximation),

$$4(k/\sqrt{s})A(s,t) \ge A(\Sigma_I) > \frac{(1-z^{-1})}{2} + (1/2)\left(\ln((z+\sqrt{z^2-1}))\right)^{-2} [x'I_1(x')] \Big|_{x'=u_1}^{x'=u}, \quad (37)$$
$$u \equiv \ln((z+\sqrt{z^2-1}))\sqrt{k^2\sigma_{inel}/\pi};$$
$$u_1 \equiv \ln((z+\sqrt{z^2-1}))\sqrt{1-z^{-2}}, \quad (38)$$

and the slightly weaker but simpler result,

$$A(\Sigma_{I}) > \frac{z^{2}}{2(z^{2}-1)} \frac{\sqrt{4\Sigma_{I}(z^{2}-1)}}{z} I_{1}\left(\frac{\sqrt{4\Sigma_{I}(z^{2}-1)}}{z}\right) + \frac{(1-z^{-1})}{2} - \frac{1}{2} I_{1}\left(\frac{z^{2}-1}{z^{2}}\right).$$
(39)

Note that at high energies $z - 1 \rightarrow 0$, the last two terms give only a small positive contribution,

$$\frac{(1-z^{-1})}{2} - \frac{1}{2}I_1\left(\frac{z^2-1}{z^2}\right) \approx (z-1)^2/4, z-1 \to 0.$$
(40)

Hence, at sufficiently high energies, but without any high energy approximation, we have the bound given by Eqs. (37)-(38) and the slightly weaker but simpler bound,

$$A(s,t) > \frac{k\sqrt{s}}{4t} \frac{z^2}{z+1} x I_1(x),$$

$$x \equiv \sqrt{\frac{z+1}{2z^2}} \sqrt{\frac{t\sigma_{inel}(s)}{\pi}}.$$
 (41)

IV. Bound on energy averaged inelastic cross section.

Multiplying by s^{-3} and integrating over s, we obtain a lower bound on $C_{(s_1,s_2)}(t)$ which is the contribution from s_1 to s_2 to C(t),

$$C_{(s_1,s_2)}(t) \equiv \int_{s_1}^{s_2} A(s',t) \frac{ds'}{(s')^3} \\ \geq \int_{s_1}^{s_2} \frac{ds'}{(s')^2} \frac{\sqrt{(1-4/s_1)}}{16t} x'_1 I_1(x'_1), \quad (42)$$

where $s_2 > s_1$, and we used $2k'/\sqrt{s'} > \sqrt{(1-4/s_1)}$, and $2z_1^2/(z_1+1) \ge 2z^2/(z+1) \ge 1$ for s' in the interval (s_1, s_2) , and

$$z_1 \equiv \frac{s_1 - 4 + 2t}{s_1 - 4}, \ x'_1 \equiv \sqrt{\frac{z_1 + 1}{2z_1^2}} \sqrt{\frac{t\sigma_{inel}(s')}{\pi}}.$$
 (43)

The lower bound on $C(s_1, s_2)(t)$ is an average with the normalized weight function

$$\rho(s') = \frac{s_1 s_2}{(s_2 - s_1)(s')^2} \tag{44}$$

of an integrand which is a convex function of $\sigma_{inel}(s')$. The convexity is readily proved; using $(xI_1(x))' = xI_0(x)$, and denoting,

$$t_1 = t \frac{z_1 + 1}{2z_1^2} = t \frac{(s_1 - 4)(s_1 - 4 + t)}{(s_1 - 4 + 2t)^2} < t,$$
(45)

we have,

$$\frac{d(x_1'I_1(x_1'))}{d\sigma_{inel}(s')} = \frac{t_1}{2\pi} I_0(\sqrt{\frac{t_1\sigma_{inel}(s')}{\pi}}).$$
 (46)

Since the right-hand side is an increasing function of $\sigma_{inel}(s')$, we get the convexity property,

$$\frac{d^2(x_1'I_1(x_1'))}{d\sigma_{inel}(s')^2} > 0.$$
(47)

Since the average of a convex function is greater than the convex function of the average [21], we have the bound

$$C_{(s_1,s_2)}(t) \ge \frac{s_2 - s_1}{16ts_1s_2} \sqrt{(1 - 4/s_1)} x_1 I_1(x_1) , \qquad (48)$$

where,

$$x_1 \equiv \sqrt{\frac{t_1 \bar{\sigma}_{inel}(s_1, s_2)}{\pi}},\tag{49}$$

$$\bar{\sigma}_{inel}(s_1, s_2) \equiv \int_{s_1}^{s_2} ds' \rho(s') \sigma_{inel}(s').$$
(50)

To get bounds on $\bar{\sigma}_{inel}(s, \infty)$, and $\bar{\sigma}_{inel}(s, 2s)$, we just choose the corresponding values for (s_1, s_2) . Thus we obtain, without any asymptotic approximations in s,

$$C_{(s,\infty)}(t) \ge \frac{1}{16ts} \sqrt{(1-4/s)} x_1 I_1(x_1) ,$$
 (51)

$$x_1 \equiv \sqrt{\frac{t_1 \bar{\sigma}_{inel}(s, \infty)}{\pi}},\tag{52}$$

and,

$$C_{(s,2s)}(t) \ge \frac{1}{32ts} \sqrt{(1-4/s)} x_2 I_1(x_2) ,$$
 (53)

$$x_2 \equiv \sqrt{\frac{t_1 \bar{\sigma}_{inel}(s, 2s)}{\pi}},\tag{54}$$

where

$$t_1 = t \frac{(s-4)(s-4+t)}{(s-4+2t)^2} < t.$$
(55)

Asymptotic bounds Since we want asymptotic upper bounds on the energy averaged inelastic cross sections, we can assume without loss of generality that the arguments x_1, x_2 of the modified Bessel functions tend to infinity, and obtain,

$$16stC_{(s,\infty)}(t)\sqrt{2\pi} > \left(\sqrt{\xi_1}\exp\xi_1\right)(1+O(1/\xi_1))(56)$$

$$\xi_1 = \sqrt{\frac{t\bar{\sigma}_{inel}(s,\infty)}{\pi}},$$
(57)

and

$$32stC_{(s,2s)}(t)\sqrt{2\pi} > \left(\sqrt{\xi_2}\exp\xi_2\right)(1+O(1/\xi_2))(58)$$

$$\xi_2 = \sqrt{\frac{t\bar{\sigma}_{inel}(s,2s)}{\pi}}.$$
 (59)

We now use the elementary lemma proved in [20], [1],

Lemma. If $\xi > 1$, and $y \ge \sqrt{\xi} \exp \xi$, then,

$$\xi < f(y) \equiv \ln y - (1/2) \ln \left(\ln y - \frac{1}{2} \ln \ln y \right).$$
 (60)

With f(y) as defined above, we obtain the asymptotic bounds,

$$\bar{\sigma}_{inel}(s,\infty) \leq_{s \to \infty} \frac{\pi}{t} \left(f(4s/s_0) \right)^2, \tag{61}$$

where,

$$\frac{1}{s_0} = 4t C_{(s,\infty)}(t) \sqrt{2\pi}, \ t = 4m_\pi^2 - \epsilon, \tag{62}$$

and,

$$\bar{\sigma}_{inel}(s,2s) \leq_{s \to \infty} \frac{\pi}{t} \left(f(8s/s_0) \right)^2.$$
(63)

Notice that the coefficients of $(\ln s)^2$ in these bounds on the inelastic cross section are one-fourth of those in the corresponding bounds on the total cross section at high energies , and the scale factors in the inelastic case are also one-fourth of those in the corresponding total cross section bounds [1],

$$\bar{\sigma}_{tot}(s,\infty) \leq_{s \to \infty} \frac{4\pi}{t} (f(s/s_0))^2 ,$$

$$\bar{\sigma}_{tot}(s,2s) \leq_{s \to \infty} \frac{4\pi}{t} (f(2s/s_0))^2 .$$
(64)

In the case of pion-pion scattering, we may remove the unknown ϵ in $t = 4m_{\pi}^2 - \epsilon$ rigorously by using absolute bounds on the D-wave below threshold derived in [1], or use phenological inputs on the D-wave scattering length and set $\epsilon = 0$. The main qualitative difference from the case of non-identical particles is that only even partial waves occur in $\pi^0 \pi^0$ scattering, We first show that inspite of this difference, the bounds of this section at moderate energies as well as the asymptotic bounds on inelastic cross sections hold for $\pi^0 \pi^0$ scattering.

V. Bounds on pion-pion inelastic cross sections. We shall exploit iso-spin invariance.

$$F^{\pi^{0}\pi^{0}\to\pi^{0}\pi^{0}} = \frac{1}{3}F^{0} + \frac{2}{3}F^{2}$$
$$= \frac{\sqrt{s}}{4k} \sum_{l=0,2,}^{\infty} (2l+1)2a_{l}^{\pi^{0}\pi^{0}\to\pi^{0}\pi^{0}}(s)P_{l}(z),$$
$$F^{\pi^{0}\pi^{0}\to\pi^{+}\pi^{-}} = \frac{1}{3}F^{0} - \frac{1}{3}F^{2}$$
$$= \frac{\sqrt{s}}{4k} \sum_{l=0,2,}^{\infty} (2l+1)\sqrt{2}a_{l}^{\pi^{0}\pi^{0}\to\pi^{+}\pi^{-}}(s)P_{l}(z).$$
(65)

Unitarity then implies,

$$Ima_{l}^{\pi^{0}\pi^{0}\to\pi^{0}\pi^{0}} \geq |a_{l}^{\pi^{0}\pi^{0}\to\pi^{0}\pi^{0}}|^{2} + |a_{l}^{\pi^{0}\pi^{0}\to\pi^{+}\pi^{-}}|^{2}$$
$$= \frac{1}{3}|a_{l}^{0}(s)|^{2} + \frac{2}{3}|a_{l}^{2}(s)|^{2}.$$
(66)

Hence we define the inelastic cross section considering $\pi^0 \pi^0 \to \pi^+ \pi^-$ also as an elastic channel,

$$\sigma_{inel}^{\pi^0\pi^0} \equiv \sigma_{tot}^{\pi^0\pi^0} - \sigma^{\pi^0\pi^0 \to \pi^0\pi^0} - \sigma^{\pi^0\pi^0 \to \pi^+\pi^-} = \frac{8\pi}{k^2} \times \sum_{l=0,2,}^{\infty} (2l+1) \left(\frac{1}{3} (Ima_l^0 - |a_l^0|^2) + \frac{2}{3} (Ima_l^2 - |a_l^2|^2)\right).$$

Note that,

$$\sigma_{inel}^{\pi^0\pi^0} \le \sigma_{inel,im}^{\pi^0\pi^0} \equiv \frac{8\pi}{k^2} \sum_{l=0,2,}^{\infty} (2l+1) \times \left(\frac{1}{3}(Ima_l^0 - (Ima_l^0)^2) + \frac{2}{3}(Ima_l^2 - (Ima_l^2)^2)\right).$$
(67)

As before, we vary the Ima_l^I subject to positivity constraints , and the given $\sigma_{inel,im}^{\pi^0\pi^0}$ to minimise the aborptive part,

$$A^{\pi^{0}\pi^{0} \to \pi^{0}\pi^{0}}(s,t) = \frac{\sqrt{s}}{4k} \sum_{l=0,2}^{\infty} (2l+1) \times P_{l}(z) 2\left(\frac{1}{3}(Ima_{l}^{0}) + \frac{2}{3}(Ima_{l}^{2})\right).$$
(68)

The minimum is reached when

$$Ima_{l}^{0} = Ima_{l}^{2} = \frac{1}{2} \left(1 - \frac{P_{l}(z)}{P_{\lambda}(z)} \right), \ l \le L;$$

$$Ima_{l}^{0} = Ima_{l}^{2} = 0, \ l > L, L \le \lambda < L + 2.$$
(69)

The minimum is an increasing and a convex function of $\sigma_{inel,im}^{\pi^0\pi^0}$. The lower bound on the absorptive part therefore remains valid if we replace $\sigma_{inel,im}^{\pi^0\pi^0}$ by $\sigma_{inel}^{\pi^0\pi^0}$. Again, defining

$$4(k/\sqrt{s})A^{\pi^{0}\pi^{0}\to\pi^{0}\pi^{0}}(s,t) \equiv A^{\pi^{0}\pi^{0}}(\lambda),$$
(70)

$$\sigma_{inel}^{\pi^0\pi^0} \frac{k^2}{4\pi} \equiv \Sigma_I^{\pi^0\pi^0}(\lambda), \qquad (71)$$

we prove that if $ln(z + \sqrt{z^2 - 1}) < 1/6$, which holds at moderately high energies, $\Sigma_I^{\pi^0 \pi^0}(\lambda) < \lambda^2/4$ if $L \geq 2$. Finally, proceeding as for non-identical particles, we obtain a bound without any high energy approximations,

$$A^{\pi^{0}\pi^{0}}(\lambda(\Sigma_{I}^{\pi^{0}\pi^{0}})) \geq (1 - 1/P_{2}(z)) + \int_{\Sigma_{I}^{\pi^{0}\pi^{0}}(2)}^{\Sigma_{I}^{\pi^{0}\pi^{0}}} P_{\sqrt{4\Sigma_{I}'}}(z)d\Sigma_{I}'.$$
 (72)

This yields the exact bound,

$$4(k/\sqrt{s})A^{\pi^{0}\pi^{0}}(s,t) \ge (1-1/P_{2}(z)) + (1/2)\alpha(z)^{-2} [x'I_{1}(x')]|_{x'=v_{1}}^{x'=v},$$
(73)

$$\alpha(z) \equiv \ln((z+\sqrt{z^{2}-1})), v \equiv \alpha(z)\sqrt{\frac{k^{2}}{\pi}\sigma_{inel}^{\pi^{0}\pi^{0}}};$$

$$v_{1} \equiv \alpha(z)\sqrt{2(1-P_{2}(z)^{-2})};$$
(74)

a slightly weaker bound is obtained by replacing $\alpha(z)$ by the smaller quantity $\sqrt{z^2 - 1/z}$, and noting that $(1 - 1/P_2(z)) - (1/2)\alpha(z)^{-2}[x'I_1(x')]|_{x'=v_1}$ is then positive at moderately high energies. Thus we obtain a slightly weaker but rigorous bound,

$$A^{\pi^{0}\pi^{0}}(s,t) > \frac{k\sqrt{s}}{4t} \frac{z^{2}}{z+1} x I_{1}(x),$$
$$x \equiv \sqrt{\frac{z+1}{2z^{2}}} \sqrt{\frac{t\sigma_{inel}^{\pi^{0}\pi^{0}}(s)}{\pi}}$$
(75)

which is identical to the result given earlier for nonidentical particles. Therefore the asymptotic bounds of the last section on energy averages of inelastic cross sections also hold for $\pi^0\pi^0$ scattering.

VI. Absolute bounds on $\pi^0 \pi^{\tilde{0}}$ inelastic cross sections. In [1] we derived absolute bounds on $\pi^0 \pi^0$ D-waves below threshold and on C(t) in terms of pion-mass alone, for 0 < t < 4. In particular,

$$f_2(t) <_{t \to 4-} \frac{4-t}{120} (34+6.25(4-t)+O(4-t)^2).$$
 (76)

and

$$C^{\pi^0 \pi^0 \to \pi^0 \pi^0}(t) <_{t \to 4-} \frac{17\pi}{4(4-t)},$$
 (77)

where $f_l(s) = \frac{\sqrt{s}}{4k} 2a_l^{\pi^0 \pi^0 \to \pi^0 \pi^0}(s)$. We now use these rigorous bounds in conjunction with the asymptotic bounds on inelastic $\pi^0 \pi^0$ cross sections to remove the unknown $\epsilon = 4 - t$ in these bounds. The price to pay for the rigour is that we cannot choose t = 4. For the upper bound on $\bar{\sigma}_{inel}^{\pi^0 \pi^0}(s, \infty)$, the optimum choice is,

$$\epsilon = 4 - t = \frac{8}{\ln(s/s_1)},\tag{78}$$

which yields,

(

$$\bar{\sigma}_{inel}^{\pi^0 \pi^0}(s,\infty) \leq_{s \to \infty} (\pi/4)(m_{\pi})^{-2} \times \ln(s/s_1) + (1/2) \ln \ln(s/s_1) + 1 \right)^2 + O(\ln \ln(s/s_1)).(79)$$

where the scale factor s_1 is given by

$$1/s_1 = 34\pi\sqrt{2\pi} \ m_\pi^{-2},\tag{80}$$

and is one-fourth of that for the total cross section case[1]. For the upper bound on $\bar{\sigma}_{inel}^{\pi^0\pi^0}(s, 2s)$, the optimum choice is,

$$\epsilon = 4 - t = \frac{8}{\ln(s/s_2)},\tag{81}$$

which yields,

$$\bar{\sigma}_{inel}^{\pi^0 \pi^0}(s, 2s) \leq_{s \to \infty} (\pi/4)(m_{\pi})^{-2} \times \left(\ln(s/s_2) + (1/2)\ln\ln(s/s_2) + 1\right)^2 + O(\ln\ln(s/s_2)).(82)$$

where the scale factor s_2 is given by

$$1/s_2 = 2/s_1 = 68\pi\sqrt{2\pi} \ m_\pi^{-2}.$$
 (83)

These are bounds from first principles on a cross section fundamental in strong interaction physics. But for phenomenological comparisons it is more useful to use some crossed channel low energy data to get stronger bounds , particularly on the scale of the logarthm.

VII. Phenomenological comparisons for Pion-Pion scattering .

(i) First, the basic lower bound (from unitarity alone) on the absorptive part A(s,t) in terms of the inelastic cross section, given by Eq. (41), or in terms of the total cross section (Eq. (21) of [1]) can be compared directly with phenomenological estimates of the absorptive part at energies where such estimates are available [26], [27]. This can be done for the amplitudes

$$F^{\pi^{+}\pi^{0}\to\pi^{+}\pi^{0}(s,t)} = 1/2(F^{1}+F^{2})(s,t),$$

$$F^{\pi^{0}\pi^{0}\to\pi^{0}\pi^{0}}(s,t) = \frac{1}{3}F^{0} + \frac{2}{3}F^{2}(s,t),$$

which have positive absorptive parts for $s \ge 4$, 0 < t < 4. A violation of the bounds will indicate that the input absorptive part violates unitarity.

(ii) Secondly, bounds on energy averages of cross sections in the intervals $(s.\infty)$ and (s.2s) in terms of phenomenological inputs for c.m. energies less than \sqrt{s} follow from unitarity, analyticity and crossing. The crossing relations,

$$\frac{1}{2}(F^{1} + F^{2})(s,t) = \frac{1}{3}(F^{0} - F^{2})(t,s),$$

$$F^{\pi^{0}\pi^{0} \to \pi^{0}\pi^{0}}(s,t) = F^{\pi^{0}\pi^{0} \to \pi^{0}\pi^{0}}(t,s)$$
(84)

and the Froissart Gribov formula yield,

$$C_{(s.\infty)}^{\pi^+\pi^0 \to \pi^+\pi^0}(t=4) = \frac{5\pi}{16}m_{\pi}(a_2^0 - a_2^2) - C_{(4,s)}^{\pi^+\pi^0 \to \pi^+\pi^0}(t=4), \quad (85)$$

and

$$C_{(s,\infty)}^{\pi^0\pi^0\to\pi^0\pi^0}(t=4) = \frac{5\pi}{16}m_{\pi}(a_2^0+2a_2^2) - C_{(4,s)}^{\pi^0\pi^0\to\pi^0\pi^0}(t=4).$$
(86)

Here , as in [1], we defined the *l*-wave scattering lengths a_l^I as the $q \to 0$ limits of the phase shifts $\delta_l^I(q)$ divided by q^{2l+1} where q is the c.m. momentum . The Bern group [19] already has estimates of the *D*-wave scattering lengths, and has recently obtained [28] estimates of the absorptive part integrals upto $\sqrt{s} = 1, 6 GeV$,

$$\begin{split} C^{\pi^+\pi^0\to\pi^+\pi^0}_{(4m^2_\pi,s)}(t=4) &= 1.48\times 10^{-3},\\ C^{\pi^0\pi^0\to\pi^0\pi^0}_{(4m^2_\pi,s)}(t=4) &= 2.031\times 10^{-3}. \end{split}$$

Hence the bounds on energy averaged cross sections $\bar{\sigma}_{inel}^{\pi^0\pi^0}(s,2s)$ and $\bar{\sigma}_{inel}^{\pi^+\pi^0}(s,2s)$, as well as $\bar{\sigma}_{inel}^{\pi^0\pi^0}(s,\infty)$ and $\bar{\sigma}_{inel}^{\pi^+\pi^0}(s,\infty)$ implied by Eqs. (51)-(55) can be directly tested against the corresponding experimental values.

(iii) Thirdly, explicit asymptotic bounds on the averages of the inelastic cross section in the intervals (s, ∞) and (s, 2s) are given by Eqs. (61) and (63), and in terms of the corresponding averages of the total cross section , in terms of a scale parameter s_0 ; s_0 is given by Eq. (62) in terms of $C_{(s,\infty)}$, an integral over absorptive parts in the interval (s, ∞) . Substituting the values of the D-wave scattering lengths given by [19],

$$a_2^0 \approx 0.00175 m_\pi^{-5}; a_2^2 \approx 0.00017 m_\pi^{-5},$$
 (87)

we have, choosing for s a value up to which absorptive parts can be reliably estimated,

$$\pi^{0}\pi^{0}: \ s_{0}^{-1} = m_{\pi}^{-2} 16\sqrt{2\pi}$$
$$\times \left(2.05 \times 10^{-3} - C_{(4,s)}^{\pi^{0}\pi^{0} \to \pi^{0}\pi^{0}}(t=4)\right), \tag{88}$$

$$\pi^{+}\pi^{0}: s_{0}^{-1} = m_{\pi}^{-2} 16\sqrt{2\pi}$$
$$\times (1.55 \times 10^{-3} - C_{(4,s)}^{\pi^{+}\pi^{0}} \rightarrow \pi^{+}\pi^{0}}(t=4)).$$
(89)

These equations give much stronger bounds than the absolute bounds. E.g. using only positivity of $C_{(4,s)}^{\pi^0\pi^0\to\pi^0\pi^0}(t=4)$ we get,

$$s_0 \ge 12m_{\pi}^2$$

which is 800 times the absolute bound $s_0 \geq .015m_{\pi}^2$. As the absorptive part integrals in the *D*-wave scattering length sum rules are rapidly convergent, even for the moderate value of $\sqrt{s} = 1, 6GeV$, Colangelo et al [28] obtained a further big improvement in the values of the scale factor, when phenomenological values of absorptive parts upto $\sqrt{s} = 1, 6GeV$ are utilised,

$$\pi^0 \pi^0 : s_0 \ge 1312m_\pi^2 \pi^+ \pi^0 : s_0 \ge 356m_\pi^2,$$
(90)

which are not very far from the scale factors used in phenomenological fits [27]. We should remember that the phenomenological values may be dependent on the particular parametrisation used to fit experimental cross sections. The implicit bounds (51)-(55) discussed in (i) and (ii) are without asymptotic approximations, and therefore can be compared directly with experiment.

VIII. Concluding Remarks.

In this paper on inelastic cross sections and the previous one on total cross sections [1] we believe to have put the Froissart Bound on a solid ground, by using the notion of average cross sections which avoids completely the problem of the scale in the Froissart bound. These averages can be chosen rather arbitrarily but once you have chosen one you must stick to it. The simplest averages that we use are the ones from s to infinity and from s to 2s. The averaging interval must be sufficiently large if one wants to preserve the coefficients appearing in the Lukaszuk-Martin bound. The only unknown is the value of a certain integral on the absoptive part for some positive t. In the special case of pion-pion scattering all unknown constants are eliminated. The advantage of introducing the bound on the inelastic cross section is that, asymptotically, it is 4 times smaller than the one on the total cross section. So if you accept to believe that the elastic cross-section cannot be larger than the inelastic cross section, the limiting case being an expanding black disk you gain a factor 2 on the bound on the total cross section. However, not everybody agrees with this, for instance Troshin and Tyurin [29] believe that at high energy the scattering amplitude is dominantly elastic. It is tempting to make a rather daring and nonrigorous suggestion: if the amplitude is essentially elastic (a small inelastic part is unavoidable according to well known theorems) then the effective large Lehmann ellipse has a right extremity at $t = 16m_{\pi}^2$, and the Froissart bound is divided by 4.

Anyway a factor of 2 or 4 is not sufficient to bring the absolute bounds near the experimental values [30],[31],[32] including the most recent experiments at LHC [33], which indicate a definite increase of the cross-sections compatible with a $(\ln(s))^2$ behaviour. There is little doubt that this trend will continue when LHC reaches higher energies. Towards quantitative improve-

8

ment we may find unitarity bounds on the energy averages of the inelastic cross section given the total cross section as an input , in addition to absorptive part integrals at positive t [34].

However, as explained in Sec. VII above, if we are prepared to make phenomenological inputs such as the *D*wave scattering lengths and low energy absorptive parts , the situation with respect to experimental comparisons improves dramatically [27].

What can we do on the theoretical side? In the case of pion pion scattering, Kupsch[35] has constructed an amplitude, crossing-symmetric satisfying "inelastic" unitarity and saturating the Froissart bound [35], but he does not give numbers. The result of Gribov [36] shows the importance of satisfying elastic unitarity in the "elastic strips", [37]. This might help, but we dont know how, and there is the problem of finding people interested in working on this.

We acknowledge very stimulating correspondence with I. Caprini, G. Colangelo, J. Gasser and H. Leutwyler of the University of Bern , and thank them for sending us their unpublished results, Eq. (90), on phenomenological evaluation of the bounds on scale factors for pion-pion scattering. We also thank Maurice Haguenauer for giving us up to date references on the work of the Atlas Alpha collaboration. SMR would like to thank the Indian National Science Academy for an INSA senior scientist grant.

- A. Martin and S. M. Roy , Phys. Rev. D89, 045015 (2014).
- [2] M. Froissart, Phys. Rev. 123, 1053 (1961).
- [3] A. Martin, Nuov. Cimen. 42, 930 (1966).
- [4] See e.g. R. F. Streater and A. S. Wightman, "PCT. Spin and Statistics and all that" (2nd revised printing, Benjamin, New York (1978)) reprinted by Princeton University Press (2000).
- [5] H. Epstein, V. Glaser and A. Martin, Comm. Math. Phys., 13, 257 (1969).
- [6] R. Haag and D. Kastler, J. Math. Phys., 5 848 (1964); D. Ruelle, Helv. Phys. Acta 35, 147 (1962); H. Araki, "Mathematical Theory of Quantum Fields", Oxford Univ. Press, Oxford (2000).
- [7] Y. I. Azimov, Phys. Rev. D84,056012 (2011).
- [8] W. Zimmermann, Nuov. Cimen. 10, 597 (1958), 21,249(1961) and 21,268(1961).
- [9] H. Lehmann, Nuovo Cimen. 10, 579 (1958); Fortschr. Physik 6 159 (1959).
- [10] J. D. Bessis and V. Glaser, Nuov. Cimen. 50, 568 (1967);
 G. Sommer, Nuov. Cimen. A48, 92 (1967); for reviews, see G. Sommer, Fortschritte der Physik, 18,577 (1970);
 A. Martin, CERN-TH/99-110 (1999).
- [11] Y. S. Jin and A. Martin, Phys. Rev. **135B**, 1375(1964).
- [12] L. Lukaszuk and A. Martin, Nuov. Cimen. **52A**, 122 (1967).
- [13] V. Singh and S. M. Roy, Ann. Phys. 57, 461 (1970).
- [14] S. M. Roy, Phys. Reports, **5C**, 125 (1972).
- [15] H. Cheng and T. T. Wu, Phys. Rev. Letters 24,1456

(1970); C. Bourrely, J. Soffer, and T. T. Wu, Phys. Rev.
D19, 3249 (1979) and Nucl. Phys. B247, 15 (1984), M.
M. Block, Phys. Reports, 436, 71 (2006) and references therein.

- [16] A. Martin, Phys. Rev. **D80**, 065013 (2009).
- [17] T. T. Wu, A. Martin, S. M. Roy and V. Singh, Phys. Rev. **D84**, 025012 (2011).
- [18] F J. Yndurain, Phys. Letters **31B**,368 (1970); A. K. Common, Nuovo Cim. **69A** ,115(1970).
- [19] G. Colangelo, J. Gasser, and H. Leutwyler, Nuclear Physics B603, 125(2001); B. Ananthanarayan,G. Colangelo, J. Gasser, and H. Leutwyler, Physics Reports 353, 207 (2001).
- [20] A. Martin, Pomeranchuk prize lecture, ITEP, Moscow (2010), https://cds.cern.ch/record/1591290.
- [21] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequali*ties, p.74, Cambridge University Press (1952).
- [22] A. Martin, Nuovo Cimento 47,265 (1967).
- [23] A. Martin, High Energy Physics and Elementary Particles, p.155, IAEA Vienna (1965).
- [24] G. Auberson, L. Epele, G. Mahoux and F. R. A. Simao, Nucl. Phys. B73 ,314 (1974); Nucl. Phys. B94 ,311 (1975).
- [25] C. Lopez and G. Mennessier, Phys. Lett. B58,437(1975), and Nucl. Phys. B 118, 426(1977).
- [26] I. Caprini, G. Colangelo, J. Gasser and H. Leutwyler, Phys.Rev. D68, 074006 (2003); e-Print: hep-ph/0306122.
- [27] I. Caprini, G. Colangelo, and H. Leutwyler, Eur. Phys. J.

C72,1860,(2012) and arXiv:1111.7160v2 [hep-ph] 7 Aug 2012; I. Caprini,G. Colangelo, J. Gasser and H. Leutwyler, (private communication).

- [28] I. Caprini,G. Colangelo, and H. Leutwyler, private communication (to be published).
- [29] S. M. Troshin and N. E. Tyurin, Phys. Lett. 208 B, 517 (1988).
- [30] U. Amaldi et al, Phys. Lett. B44,112(1973);S. R. Amendolia et al, Phys. Lett. B44,119(1973).
- [31] M. Bozzo et al, Phys. Lett. B147,392(1984); C. Augier et al, Phys. Lett. B316,448(1993).
- [32] N. A. Amos et al, Phys. Rv. Lett. 63,2784 (1989), Phys.
 Lett. B243,158(1990), and Phys. Rev. Lett. 68 ,2433

(1992).

- [33] Totem Collaboration, EPL 101 ,21004 (2013); Atlas-Alpha collaboration, Nucl. Phys. B889, 486-548(2014)
- [34] A. Martin and S. M. Roy, manuscript in preparation.
- [35] J. Kupsch, Nuovo Cim. **71A**,85 (1982).
- [36] V. N. Gribov, Proc. International Conf. on High Energy Physics at Rochester, Edited by E. C. G. Sudershan, J. H. Tinlot and A. C. Melissos (Univ. of Rochester, Rochester, NY, 1960), p. 340.
- [37] A. Martin and J.-M. Richard, Proc. 'Forward Physics and Luminosity Determination at LHC', Helsinki 2000, K. Haiti et al Eds., p.27 (World Scientific 2001).