Larry Zamick

Department of Physics and Astronomy, Rutgers University, Piscataway, New Jersey 08854

October 31, 2019

Abstract

There has been a great deal of attention to the low lying energy spectum in a nucleus because of the abundance of experimetal data. Likewise ,perhaps to a lesser extent but still significant the high end for a given configuration has been examined. Here ,using single j shell calculations as a guide we examine the middle part of the spectum resulting form single j shell calculations. Seniority arguments are used to partially explain the midshell behaviours even though in general seniority is not a good quantum number for mixed systems of neutrons and protons.

1 Introduction

.

One of the first things the present author learned was that for a system of neutrons and protons the seniority is not a good quantum number.This is when the author joined a collaboration with John McCullen and Ben Bayman on wave functions in the $f_{7/2}$ shell[1]. Only for a system of particles of one kind e.g. neutrons in the Ca isotopes is seniority conserved but once one has mixed systems the neutron-proton interaction strongly mixes states of different seniority. Nevetheless in this work it will becontend that some remnants of the seniority conserving $J=0$ T=1 pairing interaction survive.

We have recently performed single j shell studies of a system of 3 protons and one neutron (or holes) e.g. ⁹⁶Ag as 3 g_{9/2} proton holes and one g_{9/2} neutron hole.[2,3] We focused on the yrast T=1 sttes and came up with a $(2j-1)$ rule, namey that states with total angular momentum I=2j-1 lay very low in energy sometimes being the ground state. This value of I corresponds to the middle of the calculated spectrum. The spectrum of ⁹⁶Ag is poorly known but the rule has been verified experimentally for lighter nuclei such as ⁴⁴Sc and $^{52}\mathrm{Mn}.$

Table I Experimental and calculated spectra of odd-odd nuclei

	T	EXP	TН
$\overline{^{44}Sc}$	5	1.513	1.276
	6	0.271	0.381
	7	0.968	1.272
$\overline{52}_{\rm Mn}$	5	1.254	1.404
	6	0	
	7	0.870	1.819
96 Ag			0.861
	8	0	
	9	$\overline{0.470}$	0.492
$\rm h_{11/2}$	9	?	1.30
Q.Q	10	?	0.21
	11	?	0.85

In the present work we extend the study to even-even nuclei such as ⁴⁴Ti and ⁹⁶ Cd. Our contention will be that there is a fairly wide gap that separates the lower part of the spectrum from the upper part.

2 Calculations.

In the following 4 tables (II,III, IV and V) we present the energies of calculated spectra (MeV) as well as the difference in energies of adjacent spectra (Diff).

TableII Calculated even I spectra of ⁴⁴Ti (a) and ⁵²Fe (b)

Table III Calculated even I spectrum of $\rm{^{96}Cd}$ –INTd

Τ	E(MeV)	Diff.
0	0.0000	
\mathfrak{D}	0.2791	0.2971
4	0.9434	0.6463
6	1.8344	0.8905
8	1.9276	0.0923
10	3.1649	1.2373
12	3.9119	0.7470
14	4.1382	0.2263
16	3.4830	-0.6552

.Table IV Calculated Spectrum of ⁹⁶Cd–Qi

.Table V Calculated odd I spectrum of ⁹⁶Cd –INTd

In a single j shell calculation for 2 protons and 2 neutrons in a single j shell the maximum angular momentum of the 2 protons is 2j-1 and likewise for the 2 neutrons. Hence for the 4 particle system the maximum anguar momentum I_{max} is equal to 4j-2 and the middle angular momentum is (2j-1).

We have recently performed single j shell studies of a system of 3 protons and one neutron (or holes) e.g. ⁹⁶Ag as 3 g_{9/2} proton holes and one g_{9/2} neutron hole. We focused on the yrast T=1 states and came up with a $(2j-1)$ rule, namey that states with total angular momentum I=2j-1 lay very low in energy sometimes being the ground state. This value of I corresponds to the middle of the calculated spectrum. The spectrum of ⁹⁶Ag is poorly known but the rule has been verified experimentally for lighter nuclei such as ⁴⁴Sc and ⁵²Mn. Results from ref [3] are shown in Table I.

In the present work we extend the study to even-even nuclei such as ^{44}Ti , $^{52}Fe^{96}Cd$. Our contention will be that there is a fairly wide gap that separates the lower part of the spectrum from the upper part. The spectra are shown in Tables II for the INTa $(44Ti)$ and INTb interactions (5^2Fe) [2,4] ;Table III for INTd1,4] and TableIV for the Qi interaction[5] (both for 96 Cd. All the states considered have isospin T=0. We focus only on even I in these tables. The lower half of the spectrum consists of states up to I=6 for the $f_{7/2}$ shell and up to I=8 for the $g_{9/2}$ shell. The midshell angular momenta are I=6 and I =8 respectively. These gaps, between I=2j+1 and I=2j=1($8 \text{ to } 6$ and $10 \text{ to } 8$ respectively) are larger than the neighboring ones. These effects persist for several different interactions. Besides the INT d interaction of ref [1] and Qi[3] there is the one of Coraggio et al. [6]. They all give qualitatively similar results.

We show more briefly the calculated odd I spectrum for 96 Cd with the INTd interaction. We see that the first few levels are spaced very close to each other but these is a sudden gap between $I=9^+$ and $I=11^+$ of 1.4384 MeV.

Let us make a brief digression to the highest energy levels. We note that in 44 Ti the 12^+ state is correctly predicted to be higher in energy than the 10^+ state but in 52 Fe the 12^+ is lower. This is due to the fact that the J=7 T=0 two-body input matrix element in ${}^{54}Co$ is smaller than in ${}^{42}Sc$. These results are in agreement with experiment. The consequences are that the 12^+ state in 52 Fe has a much longer half- life (45.9s) than the one in 44 Ti(2.1 ns).

In ⁹⁶Cd the calculated I=16⁺ state is lower than I=14⁺ (and also 15⁺). This implies that the 16⁺ state is isomeric. This is in agreement with the experiment of Nara Singh[7].

3 Experimental results compared with theory

There is not enough experimental data in the $g_{9/2}$ shell i.e. ⁹⁶Cd, to make a comparison of theory and experiment, but such a comparison can be made in the $f_{7/2}$ region. The interaction used for ⁴⁴Ti is INTa from the two-particle spectrum of ${}^{42}\text{Sc}$; for 52 Fe we use INTb from the two-hole spectrum of ${}^{54}\text{Co}$.

In ⁴⁴Ti we have : E(6)-E(4) = 1.561 MeV—E(8)-E(6) = 2.493 MeV— E(10)-E(8) = 1.163 MeV

In ⁵²Fe we have : E(6)-E(4)= 1.941 MeV — E(8) -E(6) = 2.035 MeV—E(10)- E(8) = 1.021 MeV. These empirical results are in qualitative agreement with the predictions of the single j shell model-that

there is indeed a midshell gap in energies of levels below midshell and those above midshell. The effect is not as pronounced in ${}^{52}Fe$ as it is in ${}^{44}Ti$ but it is there nevertheless. It would be of great interest to find more details of the spectra of ⁹⁶Cd and ⁹⁶Ag to see if indeed there is such a gap in these heavier nuclei.

44T _{ii}	MeV	EXPT.	INTa
INTa	$E(6)$ - $E(4)$	1.5611	1.272
	$E(8) - E(6)$	2.493	2.024
	$\overline{E(10)}$ -E(8)	1.163	1.307
$^{52}\mathrm{Fe}$		EXPT.	INTb
INTb	$E(6)$ - $E(4)$	1.941	1.403
	$E(8)$ - $E(6)$	2.035	1.783
	$E(10)$ - $E(8)$	1.021	1.072

. Table VI Comparison of experiment and theory (INT) for the gaps.

4 Explanation via Pairing interaction

We feel we can explain the above gap in the spectrum via the pairing interaction of B.H. Flowers [8] and A.R. Edmonds and B.H. Flowers [9]. The two body matrix elements in say the $g_{9/2}$ shell from J=0 to J=9 are $-A,0,0,0,0,0,0,0,0,0$, with A positive. The expression for the energies is :

 $E = C \int (n-v)/4 * (4j+8-n-v) - T(T+1+t(t+1)).$

with C negative. Here n is the number of valence nucleons, v is the seniority, T is the total isospin and t is the reduced isospin. The relation between A and C has ben discusssed by Harper and Zamick [10]. They note that for any j shell C=- $A/((2j+1)$. This can be obtained by noting that the quantum numbers (T,t,v) for the lowest I=0T=0 state are $(0,0,0)$ whilst for the unique I=0 T=2 state they are $(2,0,0)$.

In the N=Z nuclei we are dealing only with $T=0$ states whilst for the odd-odd nuclei in ref [1] all states had T=1.

The resulting yrast spectra for even I are as follows when we choose $A=1$: $:E(0)=0$, $E(I)=1$ for I=2,4,6and 8, $E(I) = 2.2$ for I=10, 12, 14and 16. So we see there is a break after I=8. We can understand this because the energy for this interaction does not depend explicitly on I. But it does depend on seniority. For I=2,4,6,and 8 we can have seniority $v=2$ states by coupling one pair of nucleons to J=0. But we cannot reach $I=10,12,14$ or 16 by coupling one pair to to $J=0$. Hence the latter states must have seniority $v=4$.

We can also look at the odd specra. For $I=1,3,5,7,9$ E(I) is equal to 1.4 whilst for $I=11, 13$ and 15 E(I)=2.2 Hence there is a predicted gap between $I=9$ and $I=11$. The same seniority argument applies for odd I. Such a gap also appears with more realistic interactions as seen in TableV.

It should be emphasized that for most interactions e.g. INT seniority is not a good quantum number for a system of both neutrons and protons. With the interactions of Edmonds and Flowers [7,8] seniority is a good quantum number . Some remnants of the senioriy behaviour in their simple interaction seem to have not been completely lost in the more complex INT and other interactions.

For completenss we show results with shematic interactions for yrast states in ⁹⁶Cd in Table IX and for yrast states in ⁹⁶Ag in Table X.

The iteractions are:

- $E(0)$ J=0 pairing -1.0, 0,0,0,0,0,0,0,0,0,0
- E(9) J_{max} pairing 0,0,0,0,0,0,0,0,0,0, -1.0
- $E(0,9)$ Sum of above -1.0, 0,0,0,0,0,0,0,0,0,-1.0

Note that with E(0) there is a high degeneracy–the low lying spectra do not spread out. Also note that with $E(9)$ the ground state does not have I=0. Rather it has I=16, the maximum I, as the grounsd state. Although both $E(0)$ and $E(9)$ yield terrible spectra when we mix them to form $E(0,9)$ we have the beginning of a reasonable spectrum, with $I=0$ as the ground state and some spreading out of the energy levels.

5 Appendix: Interactions used

The interactions used in the $f_{7/2}$ shell are shown in Table VII. The interactions used in the $g_{9/2}$ shell are shown in Table VIII. We also show the Q.Q interaction.In some but not all cases a constant has been added so that the J=0 matrix element is zero. This does not affect the spectra.

Table VII: $f_{7/2}$ matrix elements

Table VIII $g_{9/2}$ matrix elements

.
Table IX Calculated spectra with schematic interactions
 $^{96}\mathrm{Cd}$

T	E(0)	E(9)	E(0,9)
EVEN			
$\overline{0}$	0.0000	0.5294	0.0000
$\overline{2}$	$\mathbf{1}$	0.5294	0.6370
$\overline{4}$	$\mathbf 1$	0.5294	0.9292
$\overline{6}$	$\mathbf{1}$	0.5294	$\overline{1.}1965$
8	$\mathbf{1}$	0.5286	1.2562
10	2.2	0.5253	1.6071
12	2.2	0.4835	1.5674
14	2.2	0.3285	1.4124
16	2.2	0.0000	1.0839
ODD			
$\mathbf 1$	$\overline{1.4}$	1.5298	2.4323
3	1.4	1.5278	2.1862
5	1.4	1.5114	2.1059
7	1.4	1.4247	2.1113
9	1.4	1.0293	1.4943
11	2.2	1.0247	2.1086
13	2.2	0.9700	2.0539
15	2.2	0.7941	1.8787

.Table X Calculated Spectra with schematic interactions $96Ag$

References

- [1] J.D. McCullen, B.F. Bayman and L. Zamick, Phys. Rev. 134, 515B (1964).
- [2] L.Zamick and A.Escuderos, Nucl.Phys.A 889 (2012) 8
- [3] A. Escuderos and L. Zamick, Romanian Journal of Physics, Vol. 58, Nos. 9-10, pp 1064-1075, (2013)
- [4] L. Zamick and S.J.Q. Robinson, Phys. Atom. And Nucl.,65 ,740 (2002)
- [5] Chong Qi ,Phys.Rev.C83, 014307 (2011)
- [6] L. Coraggio, A. Covello, A. Gargano, and N. Itaco, Phys. Rev. C 85, 034335 (2012).
- [7] B.S. Nara Singh et al., Phys. Rev. Lett. 107 ,172502 (2011).
- [8] B.H. Flowers, Proc. Roy. Soc. (London) A212 ,248 (1951)
- [9] A.R. Edmonds and B.H. Flowers , Proc. Roy. Soc. (London) A214 ,515 (1952)
- [10] M. Harper and L.Zamick,Phys. Rev. C91,014304 (2015)