

Parametrizations of the $\gamma^*N \rightarrow \Delta(1232)$ quadrupole form factors and Siegert's theorem

G. Ramalho

*International Institute of Physics, Federal University of Rio Grande do Norte,
Campus Universitário - Lagoa Nova, CP. 1613, Natal-RN, 59078-970, Brazil*

(Dated: November 11, 2018)

The large N_c limit provides relations that can be used to calculate the $\gamma^*N \rightarrow \Delta(1232)$ quadrupole form factors at low and intermediate Q^2 under the assumption of the pion cloud dominance. There are two limitations in those parametrizations. First, the parametrization of the Coulomb quadrupole form factor underestimate the low Q^2 data. Second, when extrapolated for the timelike region, the form factors violate Siegert's theorem by terms of the order $1/N_c^2$. We propose here corrections to the parametrization of the electric quadrupole form factor, which violate Siegert's theorem only by terms of the order $1/N_c^4$. Combining the improved large N_c pion cloud parametrizations with the valence quark contributions based on a covariant quark model for the quadrupole transition form factors, we obtain an extrapolation to the timelike region consistent with Siegert's theorem, and accomplish also a very good description of the data.

I. INTRODUCTION

The $\gamma^*N \rightarrow \Delta(1232)$ transition is characterized by the dominant dipole form factor (G_M) and two sub-leading quadrupole form factors: the electric (G_E) and the Coulomb (G_C) form factors [1–4]. The nonzero results for the quadrupole form factors are a consequence of the deviation of the $\Delta(1232)$ from a spherical shape [2, 5, 6].

Calculations based on the limit of a large number of colors (N_c) and $SU(6)$ quark models with symmetry breaking suggest that, in the low Q^2 region the $\gamma^*N \rightarrow \Delta(1232)$ quadrupole form factors are dominated by pion cloud effects [2, 7–12]. Estimates based exclusively on valence quarks underestimate the data by about an order of magnitude [3, 4, 13–15]. Although small comparative to the leading order pion cloud contributions, the valence quark contributions can nevertheless help to improve the description of the data [15–17]. The large N_c parametrizations of the $\gamma^*N \rightarrow \Delta(1232)$ electric and Coulomb quadrupole form factors have, however, a problem: they are in conflict with Siegert's theorem [18–21].

In the form factors representation, Siegert's theorem is expressed by the identity at the pseudo-threshold: $G_E = \frac{M_\Delta - M}{2M_\Delta} G_C$ [1, 19, 20] (M and M_Δ are respectively the nucleon and the Δ masses). The pseudo-threshold is the limit where the magnitude of the photon three-momentum, $|\mathbf{q}|$, vanishes, and $Q^2 = Q_{pt}^2 = -(M_\Delta - M)^2$. A test for the validity of Siegert's theorem is the value of

$$\mathcal{R}_{pt} = G_E(Q_{pt}^2) - \kappa G_C(Q_{pt}^2), \quad (1)$$

where

$$\kappa = \frac{M_\Delta - M}{2M_\Delta}. \quad (2)$$

When $\mathcal{R}_{pt} = 0$, Siegert's theorem is verified. When

$\mathcal{R}_{pt} \neq 0$, the form factors are inconsistent with Siegert's theorem.

The combination of $SU(6)$ quark models with two-body exchange currents and the large N_c limit provides a connection between the neutron charge distribution and the $\gamma^*N \rightarrow \Delta(1232)$ quadrupole form factors [10, 11]. In an exact $SU(6)$ quark model, the neutron electric form factor vanishes and the electric and Coulomb quadrupole moments of the $\gamma^*N \rightarrow \Delta(1232)$ transition are both zero. When the $SU(6)$ symmetry is broken we can relate the quadrupole moments with the neutron square charge radius r_n^2 [7–12]. We can then conclude that the $SU(6)$ symmetry breaking induces an asymmetric distribution of charge in the nucleon, which generates nonzero results for the neutron electric form factor G_{En} , and for the $\gamma^*N \rightarrow \Delta(1232)$ quadrupole transition form factors, G_E and G_C [8, 9, 11].

Moreover, based on the low Q^2 expansion of the neutron electric form factor, $G_{En} \simeq -\frac{1}{6}r_n^2 Q^2$, we can extrapolate the Q^2 dependence of the quadrupole form factors to [7, 9–11]

$$G_E(Q^2) = \left(\frac{M}{M_\Delta}\right)^{3/2} \frac{M_\Delta^2 - M^2}{2\sqrt{2}} \tilde{G}_{En}(Q^2) \quad (3)$$

$$G_C(Q^2) = \left(\frac{M}{M_\Delta}\right)^{1/2} \sqrt{2} M M_\Delta \tilde{G}_{En}(Q^2), \quad (4)$$

where $\tilde{G}_{En} = G_{En}/Q^2$. Hereafter we refer those results as large N_c relations, since they can be derived exclusively in the large N_c limit [7].

In this work we present improved large N_c pion cloud parametrizations for the quadrupole form factors in order to obtain a parametrization consistent simultaneously with Siegert's theorem and with the empirical data of the $\gamma^*N \rightarrow \Delta(1232)$ quadrupole form factors.

We conclude first that the relations (3)–(4) implies that

Siebert's theorem is violated by terms $\mathcal{R}_{pt} = \mathcal{O}(1/N_c^2)$, which may be a sizable error in the case $N_c = 3$. Since the relations (3)-(4) are extrapolated from large N_c they can have relative deviations of the order $1/N_c^2$. We then use the constraints of Siebert's theorem to modify the relation for G_E . We obtain parametrizations for the quadrupole form factors that violate Siebert's theorem only by terms $\mathcal{R}_{pt} = \mathcal{O}(1/N_c^4)$. This result is thus compatible with Siebert's theorem apart from higher-order corrections in $1/N_c^2$.

We look also for additional contributions for the transition form factors G_E and G_C , namely the contributions from the valence quarks from the nucleon and $\Delta(1232)$ systems. As mentioned, those contributions are small in the context of quark models but combined with the parametrizations of the pion cloud contributions they can reduce the gap between theory and data. An interesting propriety of the valence quark contributions for the electromagnetic form factors is that they vanish in the pseudo-threshold limit, as consequence of the orthogonality between the nucleon and $\Delta(1232)$ wave functions. As a consequence, the test of Siebert's theorem condition $\mathcal{R}_{pt} = 0$ needs to be tested only for the pion cloud contribution of the transitions form factors.

At the end, we combine valence and pion cloud contributions using a model compatible with Siebert's theorem apart from higher-order corrections in $1/N_c^2$. The results are then compared with the empirical data for G_E and G_C , showing a fair description of the overall data.

II. PION CLOUD CONTRIBUTIONS

We can test the quality of the relations (3)-(4) comparing those functions with the data based on some parametrization for G_{En} . To represent the electric form factor of the neutron, we consider the Galster parametrization [22]

$$G_{En}(Q^2) = -\mu_n \frac{a\tau_N}{1 + d\tau_N} G_D, \quad (5)$$

where $\mu_n = -1.913$ is the neutron magnetic moment, $\tau_N = \frac{Q^2}{4M^2}$, $G_D = 1/(1 + Q^2/0.71)^2$ is the dipole factor, and a, d are two dimensionless parameters.

The quadrupole form factors obtained with the parameters $a = 0.9$ and $d = 2.8$ [11] are presented in Fig. 1. For a better test of Siebert's theorem we multiply the function G_C and the data for G_C by κ . The calculations are compared with the data from Mainz [13], MIT-Bates [23] and Jefferson Lab [24] for finite square momentum transfer, Q^2 , and the world average from the Particle Data Group for $Q^2 = 0$ [25]. The data are compiled in Ref. [26].

It is clear in Fig. 1 that, the difference between the parametrizations for G_E and κG_C is not zero in the pseudo-threshold limit, when $Q^2 \simeq -0.1 \text{ GeV}^2$. This result implies that Siebert's theorem is violated, because $\mathcal{R}_{pt} \neq 0$. The explicit calculation of the deviation using




FIG. 1: $\gamma^*N \rightarrow \Delta$ quadrupole form factors estimated by the pion cloud parametrization of Eqs. (3)-(4). Data from Ref. [26].

$G_{En}(Q_{pt}^2) \simeq -\frac{1}{6}r_n^2 Q_{pt}^2$, gives

$$\mathcal{R}_{pt} \simeq -\left(\frac{M}{M_\Delta}\right)^{3/2} \frac{r_n^2}{12\sqrt{2}} Q_{pt}^2. \quad (6)$$

Since $Q_{pt}^2 = -(M_\Delta - M)^2$ and $M_\Delta - M = \mathcal{O}(1/N_c)$ we can conclude that $\mathcal{R}_{pt} = \mathcal{O}(1/N_c^2)$. Although a result $\mathcal{O}(1/N_c^2)$ may be seen as a small quantity, the numerical value is still sizable, as we can see in the graph for $\mathcal{R} = G_E - \kappa G_C$ at the pseudo-threshold (\mathcal{R}_{pt}).

III. VALENCE QUARK CONTRIBUTIONS

Before discussing how to improve the pion cloud parametrization of the quadrupole form factors G_E and G_C , we may question if Siebert's theorem can in fact be verified for the valence quark sector.

We look then for the results obtained within valence quark models. We consider, in particular, the covariant spectator quark model developed in Refs. [3, 16, 27-29] for the nucleon and $\Delta(1232)$ systems. The basic assumptions of the model are that: (i) in the electromagnetic interaction the photon couples with the single quark (impulse approximation) while the other two quarks can be interpreted as an effective diquark, (ii) the quarks have their own internal structure (dressed by gluons and quark-antiquark states), and (iii) the radial quark-diquark wave functions are calibrated in terms of momentum range parameters that can be estimated by physical or lattice QCD data.

Concerning the nucleon and $\Delta(1232)$ systems the model is quite successful in the description of the data. The parameters associated with the quark structure (quark electromagnetic form factors) were first fixed by the nucleon elastic form factor data [28]. After that the model was used to estimate the valence quark contribution for the $\gamma^*N \rightarrow \Delta$ magnetic dipole form factor [27]. The results are compatible simultaneously with estimates from dynamical reactions models [15] and with lattice QCD simulations [3, 16, 27, 30]. The model for




FIG. 2: Valence quark contributions for the $\gamma^* N \rightarrow \Delta$ quadrupole form factors estimated by the covariant spectator quark model [16].

the $\Delta(1232)$ was then extended with the inclusion of D states in the wave function [3, 16].

We consider, in particular, the parametrization from Ref. [16]. In that work the $\Delta(1232)$ system is described by a combination a S -wave and two D -wave quark-diquark states. The D states can be labeled as $D3$ (quark total spin 3/2) and $D1$ (quark total spin 1/2) [3, 27]. The nucleon system is represented just by a S state [28]. In the limit where the D -state mixtures vanish, we obtain $G_E \equiv 0$ and $G_C \equiv 0$. The D -state mixtures and the D -radial wave functions are determined by a fit to the lattice QCD data from Ref. [31] and then extrapolated to the physical point. The admixture of D states is about 0.72% for both states. The extrapolation from lattice QCD for the physical regime is performed using a vector meson parametrization of the quark current (quark electromagnetic form factors). More details can be found in Refs. [16, 29, 30].

The valence quark contributions to the quadrupole form factors G_E and G_C from Ref. [16] are presented in Fig. 2. In the figure, we again compare G_E with κG_C . The more interesting aspect of the figure is that Siegert's theorem is exactly verified, as we can see from the result $\mathcal{R} = G_E - \kappa G_C = 0$ at the pseudo-threshold. This result is a consequence of $G_E = G_C = 0$ at the same point.

In the covariant spectator quark model, the quadrupole form factors G_E and G_C are calculated in terms of angular integrals of a function $b(k, q)$, where k and q are, respectively, the diquark and the photon momenta. In the pseudo-threshold limit ($|\mathbf{q}| = 0$), the function $b(k, q)$ reduces to the spherical harmonic $Y_{20}(\hat{\mathbf{k}})$ [3]. The presence of the function $Y_{20}(\hat{\mathbf{k}})$ is then the consequence of the overlap between the $\Delta(1232)$ D states and the nucleon S state. Since in the pseudo-threshold limit there is no dependence in the photon momentum $|\mathbf{q}|$, the angular integrals are reduced to the angular integration of $Y_{20}(\hat{\mathbf{k}})$, which vanishes. Consequently the form factors vanish too. The result $G_E = G_C = 0$ at the pseudo-threshold is then the corollary of the orthogonality between S and D states.

In Fig. 2, we can notice the turning of the functions

G_E and G_C near $Q^2 = 0.15 \text{ GeV}^2$, just above the photon point $Q^2 = 0$ and a soft convergence to zero at the pseudo-threshold. The reduction of the quadrupole form factors near $Q^2 = 0$ can also be seen in the lattice QCD data [16, 31]. As we will see next, the presence of the maximum for G_C near $Q^2 = 0.15 \text{ GeV}^2$, instead at $Q^2 = 0$, has implications in the values of G_C at small Q^2 .

Another interesting aspect concerning Fig. 2 is the function $\mathcal{R} = G_E - \kappa G_C$ for finite Q^2 . We can see that the function \mathcal{R} is very small compared with G_E or κG_C . We then concludes, that in the covariant spectator quark model, the results G_E and κG_C are very similar. It is possible that the relation $G_E \simeq \kappa G_C$ is also valid for other quark models. We note, in particular, that the estimate of the quark core contribution used in the Sato-Lee model assumes $G_E = \kappa \frac{M_\Delta + M}{2M_\Delta} G_C \simeq \kappa G_C$ [15].

IV. SIEGERT'S THEOREM

We can now discuss if, as for the valence quark sector, the pion cloud parametrization is also consistent with Siegert's theorem.

In the large N_c limit the baryons are infinitely heavy and can be treated as static. For this reason large N_c is mostly used to calculate static proprieties of the baryon and transition between baryon states [32, 33]. In particular, the relations (3)-(4) are extensions of large N_c relations for G_E and G_C at $Q^2 = 0$ for finite Q^2 [7]. Those relations are not derived explicitly from large N_c , but, since they derivation for $Q^2 = 0$ is based on large N_c , one can infer that they are also limited by the accuracy from the large N_c analysis, and can therefore be affected by relative corrections of the order $1/N_c^2$ [7, 9, 11, 12, 32].

Instead of using the $1/N_c$ expansion to calculate possible $1/N_c^2$ relative corrections, one can use Siegert's theorem to check if there are corrections that are consistent with the theorem. One notes that the use of constraints external to the $SU(6)$ and large N_c formalisms were used already in the calculation of coefficients associated with physical quantities as the charge radius, quadrupole moments and others [12, 34–36].

In a first attempt we checked if we can have an exact description of Siegert's theorem, modifying the functions G_E or G_C keeping at the same time the results for $Q^2 = 0$, derived from large N_c . Considering the replacement $G_E \rightarrow (1 + \alpha Q^2)G_E$, we can preserve the result from large N_c at $Q^2 = 0$, if we calculate α in order to obtain $\mathcal{R}_{pt} = 0$ at the pseudo-threshold. The solution for this condition is $\alpha = -\frac{1}{M_\Delta^2 - M^2}$. We obtain the same effect if we replace $G_C \rightarrow G_C/(1 + \alpha Q^2)$. The problem of the new form for G_E is that it vanishes when $Q^2 = M_\Delta^2 - M^2$, in conflict with the data.

We then try an approximated solution, replacing $G_E \rightarrow G_E/(1 + Q^2/(M_\Delta^2 - M^2))$, which induces no zeros

for $Q^2 > 0$. One obtains in this case

$$G_E(Q^2) = \left(\frac{M}{M_\Delta}\right)^{3/2} \frac{M_\Delta^2 - M^2}{2\sqrt{2}} \frac{\tilde{G}_{En}(Q^2)}{1 + \frac{Q^2}{M_\Delta^2 - M^2}}. \quad (7)$$

The previous relation differs from Eq. (3) at the pseudo-threshold only by a term $\mathcal{O}(1/N_c^2)$.

With the new form for G_E , one obtains

$$\mathcal{R}_{pt} \simeq \left(\frac{M}{M_\Delta}\right)^{3/2} \frac{M_\Delta - M}{2M} \frac{r_n^2}{12\sqrt{2}} Q_{pt}^2, \quad (8)$$

which is now a term $\mathcal{O}(1/N_c^4)$, since $M_\Delta - M = \mathcal{O}(1/N_c)$ and $M = \mathcal{O}(N_c)$.

The expected falloff for large Q^2 of the pion cloud contributions for the form factors G_E and G_C given by Eqs. (4) and (7) are $G_E \propto 1/Q^8$ and $G_C \propto 1/Q^6$ respectively. One recalls however those contributions are derived from the low Q^2 expansion of the neutron electric form factor and its application is in principle limited to intermediate values of Q^2 [7]. The high Q^2 region is expected to be dominated by the valence quark degrees of freedom [37] as discussed later.

To summarize, we use Siegert's theorem to find a correction for the form factor G_E that minimizes the violation of Siegert's theorem. The solution proposed, given by Eq. (7) is not exact, but reduces the violation of Siegert's theorem to a term of the order $1/N_c^4$. The proposed function preserves the result for $G_E(0)$ obtained in the large N_c framework, and corresponds to a relative correction of $1/N_c^2$ to the result of G_E at the pseudo-threshold.

V. COMBINATION OF PION CLOUD AND VALENCE QUARK CONTRIBUTIONS

Since, as discussed, the pion cloud component represents only the leading order effect in G_E and G_C , we combine the new pion cloud parametrizations with the valence quark contributions of the covariant spectator quark model (consistent with Siegert's theorem). The sum of the two contributions is presented in Fig. 3.

From Fig. 3, we conclude that, apart from the results for G_C below 0.2 GeV^2 , to be discussed later, we obtain a very good description of the overall data. This represents a considerable improvement comparative to the previous pion cloud parametrizations (see Fig. 1). In addition, the form factors are now compatible with Siegert's theorem, within an error of the order $1/N_c^4$. The smallness of the error can be visualized in the figure if we look for $\mathcal{R} = G_E - \kappa G_C$ at the pseudo-threshold.

For future reference we call the attention for the fact that, the nonzero results for the form factors G_E and G_C are a direct consequence of the pion cloud contributions, since as discussed, the valence quark contribution vanishes at the pseudo-threshold.




FIG. 3: $\gamma^*N \rightarrow \Delta$ quadrupole form factors estimated as a combination of the valence quark and pion cloud contributions. Data from Ref. [26].

For a final discussion of the results, we need to take into account that the pion cloud contributions for G_E and G_C can have relative deviations of the order $1/N_c^2$ from Eqs. (4) and (7), as discussed previously. To represent those deviations we include a band of $\pm 10\%$ in the graphs for G_E and κG_C to represent the possible relative deviation (term of order $1/N_c^2$) from the estimate of the pion cloud contribution. Note that the $\pm 10\%$ deviation is more relevant for the discussion of the space-like region, where the data are available, than near the pseudo-threshold. The final results for G_E and G_C including the variation band are presented in Fig. 4. In the figure we compare also the results with the MAID-SG2 parametrization from Ref. [19], in order to better visualize the difference between our model and the data at low Q^2 . The MAID-SG2 gives a high quality description of the data, and it is compatible with Siegert's theorem.

From the graph for G_C , we can conclude that, the gap between the present model and the data for $Q^2 < 0.2 \text{ GeV}^2$ may not be explained by the pion cloud contribution, since the data are out of the band. We tested unsuccessfully if the quality of the description at low Q^2 could be improved considering a parametrization of the G_{En} data more complex than the Galster parametrization (5). Those results are an indication that the gap between model and data may be a consequence of the valence quark contributions. Parametrizations of the quark core contributions closer to a dipole form as in the Sato-Lee and DMT models from Ref. [15, 17] are more appropriate to describe the data measured at low Q^2 . Those parametrizations, however, differ in shape, from the estimates presented in Fig. 2, and are not compatible with the shape required by Siegert's theorem and the soft convergence to zero at the pseudo-threshold. A comparison between the results from the covariant spectator quark model and the parametrization from Ref. [15] can be found in Ref. [16]. As discussed in the context of the covariant spectator quark model, the convergence to zero at the pseudo-threshold is a consequence of the orthogonality between the nucleon and the $\Delta(1232)$ states.




FIG. 4: Results for the $\gamma^*N \rightarrow \Delta$ quadrupole form factors in comparison with the MAID-SG2 parametrization from Ref. [19]. The bands indicate an estimate of the pion cloud error of 10%, equivalent to the order of $1/N_c^2$. Data from Ref. [26].

The shape of the valence quark contributions in the region $Q^2 = 0-0.2 \text{ GeV}^2$ can in principle be tested with the help of lattice QCD simulations. With the advent of the lattice QCD simulations near the physical point, it is expected that in the near future the lattice QCD simulations approaches the physical point. In those conditions, lattice simulations in quenched QCD and partially quenched QCD approximations may be compared with our estimate of the valence quark contributions. Also, the pion cloud contributions can be estimated from the comparison between full QCD and quenched QCD. An indication that the low Q^2 data for G_C are reliable comes from the chiral effective-field theory, which connects the lattice QCD data with large pion masses with the physical regime [38].

Independent of the source of the discrepancy of G_C for $Q^2 < 0.2 \text{ GeV}^2$, the shape of the form factor G_C has implications in the Coulomb quadrupole square radius. Some authors argue that the low Q^2 behavior of G_C is a consequence of the long extension of the pion cloud [11]. Other works suggest instead, that, as a consequence of Siegert’s theorem, G_C becomes smoother near $Q^2 = 0$ [19], which according with the present analysis may be a consequence of the valence quark contributions, since the pion cloud contribution is sharper near $Q^2 = 0$.

In the range of the data presented $Q^2 < 2 \text{ GeV}^2$ the pion cloud contributions are still sizable (see Fig. 1). As discussed the pion cloud parametrization are in principle

valid for low Q^2 and may be modified for larger values of Q^2 . For G_E the valence quark component falls as $1/Q^4$, as expected from a quark model [3, 16], and dominates over the pion cloud component ($1/Q^6$). As for G_C the valence quark component falls as $1/Q^6$, as the pion cloud component. The final falloff is then $G_E \propto 1/Q^4$ and $G_C \propto 1/Q^6$, respectively, consistent with pQCD [37]. We conclude then that, for very large Q^2 , the present model is compatible with the expected pQCD falloff. Recall that the pQCD falloff is the consequence of the dominance of the valence quark contributions [37]. Another pQCD prediction is that, $G_E \simeq -G_M$, for very large Q^2 [37, 39]. Experimentally we are nowadays far away from this result [4, 37].

Overall, our calculations support the idea that the physics associated with the $\gamma^*N \rightarrow \Delta(1232)$ transition can be described by adding pionic degrees of freedom to the quark models [4, 40, 41]. Dynamical reaction models such as the Sato-Lee model [15] and the DMT model [17, 18], which calculate the pion cloud dynamically, are also in qualitative agreement with the data. In those models the bare core contributions are estimated phenomenologically as already discussed for the case of Ref. [15].

VI. SUMMARY AND CONCLUSIONS

In conclusion, we have proposed a new pion cloud parametrization for the $\gamma^*N \rightarrow \Delta(1232)$ electric quadrupole form factor. The new form for G_E is consistent with Siegert’s theorem, $G_E = \kappa G_C$, at the pseudo-threshold, within an error of $1/N_c^4$.

We have also discussed the implications of Siegert’s theorem to the bare core contribution of the quadrupole form factors. Contrary to the pion cloud contributions, the valence quark contributions vanish at the pseudo-threshold.

Combining the new parametrizations of pion cloud contributions with the valence quark contributions for the same form factors, we have obtained a very accurate description of the quadrupole form factors data, apart from a small underestimation of the G_C data in the region $Q^2 = 0-0.15 \text{ GeV}^2$.

Future developments in lattice QCD may help to clarify if the the gap between theory and data at low Q^2 is a consequence of the underestimation of the valence quark component or of the pion cloud component.

Acknowledgments

The author thanks João Pacheco B. C. de Melo and Kazuo Tsushima for the hospitality at Universidade Cruzeiro do Sul, where this work started, and Pulak Giri for useful suggestions. This work is supported by the Brazilian Ministry of Science, Technology and Innovation and by the project “Science without Borders” from

-
- [1] H. F. Jones and M. D. Scadron, *Annals Phys.* **81**, 1 (1973).
- [2] V. Pascalutsa, M. Vanderhaeghen and S. N. Yang, *Phys. Rept.* **437**, 125 (2007) [hep-ph/0609004].
- [3] G. Ramalho, M. T. Peña and F. Gross, *Phys. Rev. D* **78**, 114017 (2008) [arXiv:0810.4126 [hep-ph]].
- [4] I. G. Aznauryan *et al.*, *Int. J. Mod. Phys. E* **22**, 1330015 (2013) [arXiv:1212.4891 [nucl-th]].
- [5] G. Ramalho, M. T. Peña and A. Stadler, *Phys. Rev. D* **86**, 093022 (2012) [arXiv:1207.4392 [nucl-th]].
- [6] G. Ramalho, M. T. Peña and F. Gross, *Phys. Lett. B* **678**, 355 (2009) [arXiv:0902.4212 [hep-ph]]. G. Ramalho, M. T. Peña and F. Gross, *Phys. Rev. D* **81**, 113011 (2010) [arXiv:1002.4170 [hep-ph]].
- [7] V. Pascalutsa and M. Vanderhaeghen, *Phys. Rev. D* **76**, 111501 (2007) [arXiv:0711.0147 [hep-ph]].
- [8] A. J. Buchmann, E. Hernandez and A. Faessler, *Phys. Rev. C* **55**, 448 (1997) [nucl-th/9610040].
- [9] P. Grabmayr and A. J. Buchmann, *Phys. Rev. Lett.* **86**, 2237 (2001) [hep-ph/0104203].
- [10] A. J. Buchmann, *Phys. Rev. Lett.* **93**, 212301 (2004) [hep-ph/0412421].
- [11] A. J. Buchmann, *Can. J. Phys.* **87**, 773 (2009) [arXiv:0910.4747 [physics.atom-ph]].
- [12] A. J. Buchmann, J. A. Hester and R. F. Lebed, *Phys. Rev. D* **66**, 056002 (2002) [hep-ph/0205108].
- [13] S. Stave *et al.* [A1 Collaboration], *Phys. Rev. C* **78**, 025209 (2008) [arXiv:0803.2476 [hep-ex]].
- [14] S. Capstick and G. Karl, *Phys. Rev. D* **41**, 2767 (1990).
- [15] B. Julia-Diaz, T.-S. H. Lee, T. Sato and L. C. Smith, *Phys. Rev. C* **75**, 015205 (2007) [nucl-th/0611033].
- [16] G. Ramalho and M. T. Peña, *Phys. Rev. D* **80**, 013008 (2009) [arXiv:0901.4310 [hep-ph]].
- [17] S. S. Kamalov and S. N. Yang, *Phys. Rev. Lett.* **83**, 4494 (1999) [nucl-th/9904072].
- [18] D. Drechsel, S. S. Kamalov and L. Tiator, *Eur. Phys. J. A* **34**, 69 (2007) [arXiv:0710.0306 [nucl-th]].
- [19] G. Ramalho, *Phys. Rev. D* **93**, 113012 (2016) [arXiv:1602.03832 [hep-ph]].
- [20] G. Ramalho, *Phys. Lett. B* **759**, 126 (2016) [arXiv:1602.03444 [hep-ph]].
- [21] A. J. Buchmann, E. Hernandez, U. Meyer, and A. Faessler, *Phys. Rev. C* **58**, 2478 (1998).
- [22] S. Galster, H. Klein, J. Moritz, K. H. Schmidt, D. We-
gener and J. Bleckwenn, *Nucl. Phys. B* **32**, 221 (1971).
- [23] N. F. Sparveris *et al.* [OOPS Collaboration], *Phys. Rev. Lett.* **94**, 022003 (2005) [nucl-ex/0408003].
- [24] J. J. Kelly *et al.*, *Phys. Rev. C* **75**, 025201 (2007) [nucl-ex/0509004]; I. G. Aznauryan *et al.* [CLAS Collaboration], *Phys. Rev. C* **80**, 055203 (2009) [arXiv:0909.2349 [nucl-ex]].
- [25] K. A. Olive *et al.* [Particle Data Group Collaboration], *Chin. Phys. C* **38**, 090001 (2014).
- [26] V. I. Mokeev, https://userweb.jlab.org/~mokeev/resonance_electrocouplings/
- [27] G. Ramalho, M. T. Peña and F. Gross, *Eur. Phys. J. A* **36**, 329 (2008) [arXiv:0803.3034 [hep-ph]].
- [28] F. Gross, G. Ramalho and M. T. Peña, *Phys. Rev. C* **77**, 015202 (2008) [nucl-th/0606029].
- [29] G. Ramalho, K. Tsushima and F. Gross, *Phys. Rev. D* **80**, 033004 (2009) [arXiv:0907.1060 [hep-ph]].
- [30] G. Ramalho and M. T. Peña, *J. Phys. G* **36**, 115011 (2009) [arXiv:0812.0187 [hep-ph]].
- [31] C. Alexandrou, G. Koutsou, H. Neff, J. W. Negele, W. Schroers and A. Tsapalis, *Phys. Rev. D* **77**, 085012 (2008) [arXiv:0710.4621 [hep-lat]].
- [32] E. E. Jenkins, X. Ji and A. V. Manohar, *Phys. Rev. Lett.* **89**, 242001 (2002) [hep-ph/0207092].
- [33] R. F. Dashen, E. E. Jenkins and A. V. Manohar, *Phys. Rev. D* **49**, 4713 (1994) Erratum: [*Phys. Rev. D* **51**, 2489 (1995)] [hep-ph/9310379].
- [34] G. Dillon and G. Morpurgo, *Phys. Lett. B* **448**, 107 (1999).
- [35] A. J. Buchmann and R. F. Lebed, *Phys. Rev. D* **62**, 096005 (2000) [hep-ph/0003167].
- [36] A. J. Buchmann and E. M. Henley, *Phys. Rev. D* **65**, 073017 (2002).
- [37] C. E. Carlson and N. C. Mukhopadhyay, *Phys. Rev. Lett.* **81**, 2646 (1998) [hep-ph/9804356].
- [38] V. Pascalutsa and M. Vanderhaeghen, *Phys. Rev. Lett.* **95**, 232001 (2005) [hep-ph/0508060].
- [39] C. E. Carlson, *Phys. Rev. D* **34**, 2704 (1986).
- [40] M. Fiolhais, B. Golli and S. Širca, *Phys. Lett. B* **373**, 229 (1996).
- [41] D. H. Lu, A. W. Thomas and A. G. Williams, *Phys. Rev. C* **55**, 3108 (1997).