

Random local complex dynamics.

Lorenzo Guerini and Han Peters

March 19, 2018

Abstract

The study of the dynamics of an holomorphic map near a fixed point is a central topic in complex dynamical systems. In this paper we will consider the corresponding random setting: given a probability measure ν with compact support on the space of germs of holomorphic maps fixing the origin, we study the compositions $f_n \circ \dots \circ f_1$, where each f_i is chosen independently with probability ν . As in the deterministic case, the stability of the family of the random iterates is mostly determined by the linear part of the germs in the support of the measure. A particularly interesting case occurs when all Lyapunov indices vanish, in which case stability implies simultaneous linearizability of all germs in $\text{supp}(\nu)$.

1. INTRODUCTION

An elementary but fundamental result in the theory of local complex dynamical systems is the following:

Let $f : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^m, 0)$ be a neutral germ, i.e. all eigenvalues of $df(0)$ have norm 1. Then $\{f^n\}$ is normal in a neighborhood of the origin if and only if f is locally linearizable and $df(0)$ is diagonalizable.

Our goal in this paper is to generalize this statement to the random setting, i.e. when studying compositions of maps that are chosen i.i.d.. Our result is the following:

Theorem 1.1. *Suppose that ν is a neutral probability measure with compact support on $\mathcal{O}(\mathbb{C}^m, 0)$. Then the origin lies in the random Fatou set if and only if all the germs in $\text{supp}(\nu)$ are simultaneously linearizable, and the semigroup of differentials*

$$dS_\nu := \{df_\omega^n(0) \mid \omega \in \text{supp}(\nu)^\mathbb{N}, n \in \mathbb{N}\},$$

is conjugate to a sub-semigroup of $U(m)$.

As an illustration, consider the case where all maps in the support of ν are of the form $z \mapsto \lambda z + z^2$, with $|\lambda| = 1$. The fact that all the maps must be simultaneously linearizable implies that ν is supported at a single point: a germ with a Siegel disk.

Let us be more precise about our setting. Instead of considering normality of a family of iterates $\{f^n\}$ we will consider the family $\mathcal{F}_\omega = \{f_\omega^n\}$. Here

$\omega = (f_n)$ is a sequence of germs each chosen independently with a probability ν and $f_\omega^n := f_n \circ \cdots \circ f_1$.

The investigation on the random dynamics of holomorphic maps began with the work of Fornæss and Sibony. In [FS91] they showed that a generic rational map, with attracting cycles, admits a neighborhood W such that for almost every sequence of functions, chosen i.i.d. with respect to an absolutely continuous probability measure supported in W , the Julia set of the family \mathcal{F}_ω has zero measure. Furthermore for every $z \in \widehat{\mathbb{C}}$ the point z belongs almost certainly to the Fatou set of the family \mathcal{F}_ω . We refer to the paper of Sumi [Su11] for a generalization of this result.

In this paper we will consider the special case in which all functions in the support of a probability measure ν share a common fixed point. In particular we will be interested in the comparison of the *random Fatou set*, i.e. the points z that lie in the Fatou set of \mathcal{F}_ω for almost every ω , and the Fatou set of the semigroup generated by $\text{supp}(\nu)$. Our main motivation is to find analogies between this type of random systems and the study of local fixed point theory.

The local dynamics of a holomorphic germ f fixing the origin is one of the earliest problems studied in complex dynamics. A rough first description of the local behavior of f is depends on the eigenvalues of $df(0)$. Depending on their absolute value we distinguish between *hyperbolic* germs, where all eigenvalues have norm unequal to 1, *neutral* germs, where all eigenvalues have norm 1, and *semi-neutral* germs.

Determining the linearizability of neutral and semi-neutral germs is a subtle problem. In one dimension a complete description of this phenomenon was given in the works of Cremer [C38], Siegel [Si42], Brjuno [Brj71] and Yoccoz [Y95].

The local dynamics of germs in several complex variables has many analogies with the one dimensional case, but yet many differences. We refer to the survey of Abate [A03] for more details. In this setting the presence of resonances constitute an obstacle to linearizability, as studied by Poincaré [P28] for attracting germs. The description of attracting germs was completed in dimension 2 by Lattés [La11] and in arbitrary dimensions independently by Sternberg [St57, St58] and Rosay–Rudin [RR88].

Neutral and semi-neutral germs in several complex variables have also been intensively studied, particularly in the parabolic case, where all eigenvalues of norm 1 roots of unity. See for instance the works of Écalle [Ec85], Hakim [H98], and Abate [A01] for germs tangent to the identity, and Ueda [U86, U91] and Rivi [Ri01] for a description of semi-parabolic germs. For the non-parabolic case consider for example the papers of Bracci–Molino [BM04] and Bracci–Zaitsev [BZ13] for neutral germs, and recent papers of Firsova–Lyubich–Radu–Tanase [LRT16, FLRT16] for semi-neutral germs.

In the random setting we may introduce Lyapunov indices $\kappa_1 > \cdots > \kappa_s$, which play the same role as the (logarithms of absolute values of the) eigenvalues of $df(0)$, see [O68] or [GM89] for more information on Lyapunov indices. In analogy with the deterministic case, we say that the measure ν is *attracting* if $\kappa_1 < 0$, it is *repelling* if $\kappa_1 > 0$, is *neutral* if the only Lyapunov index is $\kappa_1 = 0$, and is *semi-neutral* if $\kappa_1 = 0$ and $\kappa_2 < 0$.

In the neutral case, the measure ν plays a marginal role: the origin lies in the random Fatou set if and only if it lies in the Fatou set of the semigroup generated by $\text{supp}(\nu)$. Let us present the proof of this fact in the simpler one-dimensional case when ν is assumed to be supported on a finite set. The general statement follows from Lemma 4.5, whose proof follows same ideas.

Proof. Suppose that the origin lies in the random Fatou set. Then $|f'(0)| = 1$ for every $f \in \text{supp}(\nu)$, otherwise the sequence $(f_\omega^n)'(0)$ is almost surely unbounded.

Choose a sequence $\omega_0 \in \text{supp}(\nu)^\mathbb{N}$, such that \mathcal{F}_{ω_0} is normal and given any finite word (g_1, g_2, \dots, g_n) , with $g_i \in \text{supp}(\nu)$, there exists $N > 0$ for which

$$f_{i+N} = g_i, \quad 1 \leq i \leq n.$$

If T denotes the left shift operator, this is equivalent to saying that for every $g \in S_\nu$ there exists $N, n > 0$ for which $f_{T^N \omega_0}^n = g$.

Given $\varepsilon > 0$ small enough, we may assume that every $f \in \text{supp}(\nu)$ is holomorphic and univalent on \mathbb{B}_ε . Furthermore by Ascoli–Arzelá Theorem, there exists $0 < \delta_0 < \varepsilon$ such that $f_{\omega_0}^n(\mathbb{B}_{\delta_0}) \subset \mathbb{B}_\varepsilon$, which implies that every $f_{\omega_0}^n$ is univalent on \mathbb{B}_{δ_0} . By Koebe quarter Theorem, given $g \in S_\nu$ we have

$$g(\mathbb{B}_{\delta_0/4}) \subset f_{T^N \omega_0}^n \circ f_{\omega_0}^N(\mathbb{B}_{\delta_0}) \subset \mathbb{B}_\varepsilon.$$

Weak Montel Theorem implies normality of the semigroup S_ν . Proving that normality of S_ν implies that the origin lies in the random Fatou set is trivial. \square

We note that the semi-group need not be normal in the attracting (or similarly, semi-neutral) case, as we will see in section 4.1.

The behavior of semi-neutral measures turns out to be substantially more complicated than the other cases, and we are unable to give a satisfying description of when the origin belongs to the random Fatou set. In section 3 we do show that it is not possible to decide whether the origin lies in the random Fatou set just by looking at the Lyapunov indices or at linearizability. In the two-dimensional setting, the fact that the origin is in the random Fatou set implies the existence of stable manifolds, analogous to their deterministic setting.

Theorem 1.2. *Let ν be a semi-neutral measure on $\mathcal{O}(\mathbb{C}^2, 0)$ with compact support. If the origin lies in the random Fatou set, then almost surely every limit germ $g = \lim_{k \rightarrow \infty} f_\omega^{n_k}$ has rank one and given z sufficiently close to the origin, its stable set $W_\omega^s(z)$ is locally a one-dimensional complex manifold.*

The paper is organized as follows. In section 2 we review background on random matrices, and prove that a neutral measure of linear maps is stable if and only if there is a conjugation to a sub-semigroup of $U(m)$. In section 3 we give a more precise formulation of the problem, and treat several examples showing that our assumptions are necessary. We will study normality of the family \mathcal{F}_ω in section 4.1 when $\kappa_1 \neq 0$, and in section 4.2 for neutral measures. Semi-neutral measures will be considered in section 5

Throughout the paper we use the inductive limit topology on $\mathcal{O}(\mathbb{C}^m, 0)$. Its construction and properties are discussed in the appendix.

Acknowledgement. We thank László Lempert and Jasmin Raissy for several useful discussions.

2. PRODUCTS OF RANDOM MATRICES

Let $(\Omega, \mathcal{B}, \mu)$ be a probability space and $T : \Omega \rightarrow \Omega$ be an ergodic transformation. Given a measurable function $M : \Omega \rightarrow \text{Mat}(m, \mathbb{C})$, the (linear) cocycle defined by M over T is the skew-product transformation

$$F : \Omega \times \mathbb{C}^m \rightarrow \Omega \times \mathbb{C}^m, \quad (\omega, v) \mapsto (T\omega, M_\omega v).$$

Observe that $F^n(\omega, v) = (T^n\omega, M_\omega^n v)$, where

$$M_\omega^n := M_{T^{n-1}\omega} \cdots M_\omega.$$

By elementary properties of measurable functions, the function $\omega \mapsto \sup_n \|M_\omega^n\|$ is also measurable. Therefore the set

$$\Omega_b = \{\omega \in \Omega \mid M_\omega^n \text{ is bounded}\},$$

is measurable and it satisfies $T^{-1}\Omega_b = \Omega_b$. By ergodicity of T we either have $\mu(\Omega_b) = 0$ or $\mu(\Omega_b) = 1$.

Problem For which cocycles is the family $\mathcal{M}_\omega = \{M_\omega^n\}$ almost surely bounded?

A cocycle of particular importance for our work is the following

Definition (i.i.d. cocycle) Let ν be a probability measure $\text{Mat}(m, \mathbb{C})$. Let Ω be the space of all sequences in $\text{supp}(\nu)$, T be the left shift operator and $M : \Omega \rightarrow \text{Mat}(m, \mathbb{C})$ be the function which returns the first element of the sequence ω .

Let U_1, \dots, U_n be open subsets of $\text{supp}(\nu)$, we define the set

$$\mathcal{C}_{U_1, \dots, U_n} := \{\omega \in \Omega \mid M_\omega \in U_1, \dots, M_{T^{n-1}\omega} \in U_n\}.$$

This collection of sets forms a basis for the product topology of Ω . With \mathcal{B} we will denote the corresponding Borel σ -algebra and with μ the measure for which $\mu(\mathcal{C}_{U_1, \dots, U_n}) = \nu(U_1) \cdots \nu(U_n)$. It is a well known fact that T is ergodic on $(\Omega, \mathcal{B}, \mu)$.

The function M is continuous and therefore measurable. We notice that $M_{T^{n-1}\omega}$ coincides with the n -th element of the initial sequence and that $X_i(\omega) := M_{T^{i-1}\omega}$ is an i.i.d. sequence of random variables with values in $\text{Mat}(m, \mathbb{C})$, each chosen with probability ν . In this case, M_ω^n is given by the product of the first n elements of the sequence ω .

The problem of the iteration of random matrices was first studied by Furstenberg and Kesten in [FK60]. An important generalization of their result is the multiplicative ergodic Theorem of Oseledec [O68]. The following version of the theorem can be found in [Rue79].

Theorem 2.1 (Multiplicative ergodic Theorem). *Let $T : \Omega \rightarrow \Omega$ be an ergodic transformation and $M : \Omega \rightarrow \text{Mat}(m, \mathbb{C})$ be a measurable function such that*

$$\log^+ \|M_\omega\| \in L^1(\Omega, \mu).$$

*There exist numbers $+\infty > \kappa_1 > \dots > \kappa_s \geq -\infty$ called **Lyapunov indices** and natural numbers $\alpha_1, \dots, \alpha_s$ called **Lyapunov multiplicities** satisfying the equation $\alpha_1 + \dots + \alpha_s = m$, such that, for almost every $\omega \in \Omega$, we have*

- a. *There exists a Hermitian matrix Λ_ω with eigenvalues $e^{\kappa_1} > \dots > e^{\kappa_s}$, with respective multiplicities α_i , such that*

$$\lim_{n \rightarrow \infty} ((M_\omega^n)^* M_\omega^n)^{\frac{1}{2n}} = \Lambda_\omega. \quad (1)$$

- b. *Suppose that $\mathcal{U}_1(\omega), \dots, \mathcal{U}_s(\omega)$ are the eigenspaces of Λ_ω . Let $\mathcal{V}_{s+1}(\omega) = \{0\}$ and $\mathcal{V}_i(\omega) = \mathcal{U}_i(\omega) \oplus \dots \oplus \mathcal{U}_s(\omega)$. Then, given $v \in \mathcal{V}_i(\omega) \setminus \mathcal{V}_{i+1}(\omega)$, we have*

$$\lim_{n \rightarrow \infty} n^{-1} \log \|M_\omega^n v\| = \kappa_i. \quad (2)$$

*The set $\sigma = \{(\kappa_1, \alpha_1), \dots, (\kappa_s, \alpha_s)\}$ is called the **Lyapunov spectrum**.*

Definition Suppose that a given cocycle satisfies $\log^+ \|M_\omega\| \in L^1(\Omega, \mu)$ and let $\kappa_1 > \dots > \kappa_s$ be the Lyapunov indices. We say that the linear cocycle is

Attracting if the maximal Lyapunov index $\kappa_1 < 0$,

Repelling if the maximal Lyapunov index $\kappa_1 > 0$,

Neutral if the Lyapunov spectrum $\sigma_\mu = \{(0, m)\}$,

Semi-neutral if the maximal Lyapunov index $\kappa_1 = 0$ and $\alpha_1 \neq m$.

Remark In the work of Furstenberg and Kesten [FK60] it is proved that, for every cocycle, we may define a number κ_μ , called *Lyapunov exponent*, as

$$\kappa_\mu = \lim_{n \rightarrow \infty} n^{-1} \mathbf{E} \log \|M_\omega^n\|. \quad (3)$$

Furthermore under the assumption that $\log^+ \|M_\omega\| \in L^1(\Omega, \mu)$, for almost every $\omega \in \Omega$ we have

$$\lim_{n \rightarrow \infty} n^{-1} \log \|M_\omega^n\| = \kappa_\mu. \quad (4)$$

In the same hypothesis of Theorem 2.1 we can describe the connection between Lyapunov indices and Lyapunov exponent as follows

Lemma 2.2. *The Lyapunov exponent κ_μ coincides with the maximal Lyapunov index κ_1 . Furthermore the following equality holds*

$$\alpha_1 \kappa_1 + \dots + \alpha_s \kappa_s = \mathbf{E} \log |\det(M_\omega)|.$$

Proof. Let κ_1 be the maximal Lyapunov index and choose $\omega \in \Omega$ such that (1), (2) and (4) hold. It is not hard to prove that $\kappa_\mu \geq \kappa_1$, thus it remains to prove that $\kappa_\mu \leq \kappa_1$. Let v_n be a unit vectors such that $\|M_\omega^n\| = \|M_\omega^n v_n\|$. We take n_k such that $v_{n_k} \rightarrow v$. By (2), for some $i = 1, \dots, s$, we have

$$\begin{aligned} \kappa_i &= \lim_{k \rightarrow \infty} n_k^{-1} \log \|M_\omega^{n_k} v\| \\ &\geq \lim_{k \rightarrow \infty} n_k^{-1} \log (\|M_\omega^{n_k} v_{n_k}\| - \|M_\omega^{n_k} (v_{n_k} - v)\|) \\ &\geq \lim_{k \rightarrow \infty} n_k^{-1} \log \|M_\omega^{n_k}\| + \lim_{k \rightarrow \infty} n_k^{-1} \log (1 - \|v_{n_k} - v\|) \\ &\geq \kappa_\mu, \end{aligned}$$

since $\kappa_i \leq \kappa_1$, the equality $\kappa_\mu = \kappa_1$ follows.

If we apply $\log |\cdot|$ to both side of (1), we obtain that

$$\begin{aligned} \alpha_1 \kappa_1 + \dots + \alpha_s \kappa_s &= \log |\det(\Lambda_\omega)| \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log |\det(M_\omega^n)| \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \log |\det(M_{T^i \omega})|. \end{aligned}$$

By Birkhoff ergodic theorem the last term of this equality converge almost surely to $\mathbf{E} \log |\det(M_\omega)|$. By choosing an appropriate ω , we obtain the desired equality. \square

Corollary 2.3. *Suppose that $\log^+ \|M_\omega\| \in L^1(\Omega, \mu)$. We have the following list of equivalences:*

- a. *the cocycle is attracting if and only if $\kappa_\mu < 0$,*
- b. *the cocycle is repelling if and only if $\kappa_\mu > 0$,*
- c. *the cocycle is neutral if and only if $\kappa_\mu = 0$ and $\mathbf{E} \log |\det(M_\omega)| = 0$,*
- d. *the cocycle is semi-neutral if and only if $\kappa_\mu = 0$ and $\mathbf{E} \log |\det(M_\omega)| < 0$.*

When the cocycle is attracting or repelling, (4) gives an answer to the problem of the boundedness of the family $\mathcal{M}_\omega = \{M_\omega^n\}$.

Corollary 2.4. *If a cocycle is attracting then the family \mathcal{M}_ω is almost certainly bounded in $\text{Mat}(m, \mathbb{C})$. If a cocycle is repelling \mathcal{M}_ω is almost certainly unbounded.*

2.1 Neutral measures

The stability problem is considerably more complicated in the neutral setting. However, for i.i.d. cocycles arising from probability measures with compact support, we can give a precise description of the stable systems in Proposition 2.7 below. In the two examples that follow we show that no such statement can hold for general neutral cocycles.

Given a probability measure ν on $\text{Mat}(m, \mathbb{C})$ with compact support, by Tychonoff Theorem, the set Ω is compact, thus $\log^+ \|M_\omega\| \in L^1(\Omega, \mu)$.

Definition Let ν be a measure on $Mat(m, \mathbb{C})$ with compact support. We say that ν is attracting (respectively repelling, neutral and semi-neutral) if the corresponding i.i.d. cocycle is attracting (respectively repelling, neutral and semi-neutral).

Lemma 2.5. *Let ν be a probability measure on $Mat(m, \mathbb{C})$ with compact support. Then the set*

$$\Omega_a := \left\{ \omega \in \Omega \mid \forall n \in \mathbb{N} \text{ and } \forall \alpha \in \Omega, \exists k_j : M_{T^{k_j} \omega}^n \rightarrow M_\alpha^n \right\}$$

is a full measure subset of Ω .

Proof. Let $n \in \mathbb{N}$ and choose a sequence $\varepsilon_j \rightarrow 0$. We define

$$\Omega_{n,j} := \{ \omega \in \Omega \mid \forall \alpha \in \Omega, \exists k : \|M_{T^k \omega}^n - M_\alpha^n\| < \varepsilon_j \}.$$

Is it clear that $\bigcap_{n,j} \Omega_{n,j} = \Omega_a$. Since the sets $\Omega_{n,j}$ are countably many it suffices to prove that, for every $n, j \in \mathbb{N}$, the set $\Omega_{n,j}$ has full measure.

Suppose that n and j are fixed. Given $\alpha \in \Omega$, using the continuity of T and M , we may find an open neighborhood $U_\alpha \ni \alpha$ so that, given $\beta \in U_\alpha$, we have

$$\|M_\alpha^n - M_\beta^n\| < \varepsilon_j/2.$$

By compactness of $supp(\nu)$, the set Ω is also compact, therefore we may find $\alpha_1, \dots, \alpha_N$ such that $\Omega = \cup_i U_{\alpha_i}$. For simplicity we will write U_i for U_{α_i} . Notice that $supp(\mu) = \Omega$, therefore all the sets U_i have positive measure. Using standard results from ergodic theory and the fact that the α_i are a finite number, we find that

$$\Omega'_{n,j} = \{ \omega \in \Omega \mid \forall i = 1, \dots, N, \exists k_i : T^{k_i} \omega \in U_i \}$$

has full measure in Ω .

Finally, given $\omega \in \Omega'_{n,j}$ and $\alpha \in \Omega$, there exists U_i so that $\alpha \in U_i$ and k_i so that $T^{k_i} \omega \in U_i$. By the definition of the U_i -s we obtain that

$$\|M_{T^{k_i} \omega}^n - M_\alpha^n\| \leq \|M_{T^{k_i} \omega}^n - M_{\alpha_i}^n\| + \|M_{\alpha_i}^n - M_\alpha^n\| < \varepsilon_j,$$

which proves that $\Omega'_{n,j} \subset \Omega_{n,j}$ and thus that $\Omega_{n,j}$ has full measure. \square

Remark The previous lemma is valid in a much more general context. As a matter of fact it holds, with the same exact proof, for continuous cocycles, i.e. where both T and M are continuous functions, over a compact space.

Lemma 2.6. *Let \mathcal{G} be a compact subgroup of $GL(m, \mathbb{C})$. Then \mathcal{G} is conjugated to a subgroup of the standard unitary group $U(m)$.*

Proof. Let $\nu_{\mathcal{G}}$ the Haar measure on \mathcal{G} normalized in such a way that $\nu_{\mathcal{G}}(\mathcal{G}) = 1$. We define a Hermitian inner product on \mathbb{C}^m as

$$\langle u, v \rangle_\mu = \int_{\mathcal{G}} \langle Mu, Mv \rangle_E d\nu_{\mathcal{G}}(M),$$

where $\langle \cdot, \cdot \rangle_E$ is the standard Hermitian inner product on \mathbb{C}^m . Thank to the properties of the measure $\nu_{\mathcal{G}}$, for every $M \in \mathcal{G}$ and $u, v \in \mathbb{C}^m$, we have

$$\langle Mu, Mv \rangle_{\mathcal{G}} = \langle u, v \rangle_{\mathcal{G}}.$$

Let $\beta_E = \{e_E^i\}$ and $\beta_{\mathcal{G}} = \{e_{\mathcal{G}}^i\}$ be two orthonormal basis of $\langle \cdot, \cdot \rangle_E$ and $\langle \cdot, \cdot \rangle_{\mathcal{G}}$ respectively. Let $H \in GL(m, \mathbb{C})$ be the matrix such that $He_{\mathcal{G}}^i = e_E^i$. Given $u, v \in \mathbb{C}^m$ we can easily prove that

$$\langle u, v \rangle_{\mathcal{G}} = \langle Hu, Hv \rangle_E.$$

Finally given $M \in \mathcal{G}$, than for every $u, v \in \mathbb{C}^m$ we have

$$\begin{aligned} \langle H^{-1}MHu, H^{-1}MHv \rangle_E &= \langle MH^{-1}u, MH^{-1}v \rangle_{\mathcal{G}} \\ &= \langle H^{-1}u, H^{-1}v \rangle_{\mathcal{G}} \\ &= \langle u, v \rangle_E \end{aligned}$$

which implies that $H^{-1}\mathcal{G}H$ is a subgroup of $U(m)$. □

We define the set

$$S_{\nu} = \{M_{\omega}^n \mid \omega \in \Omega, n \in \mathbb{N}\}.$$

In this case $S_{\nu} \subset Mat(m, \mathbb{C})$ has a semigroup structure, which is not necessarily true for general neutral cocycles.

Proposition 2.7. *let ν be a neutral measure on $Mat(m, \mathbb{C})$ with compact support, and consider the corresponding i.i.d. cocycle. The following are equivalent*

1. *the family \mathcal{M}_{ω} is almost certainly bounded,*
2. *the semigroup S_{ν} is relatively compact in $GL(m, \mathbb{C})$ and is conjugated to a sub-semigroup of the unitary group $U(m)$.*

Before proceeding to the proof, we want to discuss why there is no hope that a (similar) proposition holds for a generic neutral cocycle or even for a continuous neutral cocycle over a compact space. We will show this by constructing two cocycles for which \mathcal{M}_{ω} is almost certainly bounded but for which $S_{\Omega} = \{M_{\omega}^n \mid \omega \in \Omega, n \in \mathbb{N}\}$ is not relatively compact in $GL(m, \mathbb{C})$.

Example Let $X = [0, 1)$ with the Borel σ -algebra \mathcal{B} and Lebesgue measure λ . Let $\theta \in \mathbb{R} \setminus \mathbb{Q}$ and define $T : X \rightarrow X$ by $Tx = x + \theta \pmod{1}$. It is well known that the transformation T is ergodic. Given $x_1, x_2 \in X$ we write

$$d_X(x_1, x_2) = \inf_{m \in \mathbb{Z}} |x_1 + m - x_2|.$$

Let $\varphi \in L^1(X, \lambda)$ be a nonnegative and unbounded function. The function $M : X \rightarrow \mathbb{R}^+$, defined as

$$\log(M_x) = f(x) := \varphi(x) - \varphi(Tx),$$

is measurable and defines a neutral cocycle over T .

Notice that $M_x^n = e^{\varphi(x) - \varphi(T^n x)} \leq e^{\varphi(x)}$. Since $\varphi(x) < \infty$ for almost every $x \in X$, it follows that M_x^n is almost surely a bounded sequence. On the other hand we can find $K_0 > 0$ so that the set $X_{K_0} = \{x \in X \mid \varphi(x) < K_0\}$ has positive measure. Since T is ergodic, for almost every $x \in X$ there exists n_0 such that $T^{n_0}x \in X_{K_0}$. We conclude that $M_x^{n_0} > e^{\varphi(x) - K_0}$. Since φ is unbounded, it follows that S_X is unbounded in \mathbb{R}^+ .

Example In the previous example the function f may be unbounded, in which case the unboundedness of S_X is not particularly surprising. In this second example we will construct a function φ for which f is continuous, and therefore bounded.

Let $\omega_n = (n-1)\theta \bmod 1 = T^{n-1}0$ and let $a_k = 2^k k$. For every $k > 0$ we may choose $\varepsilon_k > 0$ such that

- a. $\varepsilon_k \leq 1/a_k^2$;
- b. the sets $[\omega_1 - \varepsilon_k, \omega_1 + \varepsilon_k], \dots, [\omega_{2a_k} - \varepsilon_k, \omega_{2a_k} + \varepsilon_k]$ are pairwise disjoint.

Write $I_{j,k} = [\omega_j - \varepsilon_k, \omega_j + \varepsilon_k]$ and define f_k as

$$f_k(x) := \begin{cases} -\frac{1}{2^k} \left(1 - \frac{d_X(x, \omega_j)}{\varepsilon_k}\right) & \text{if } x \in I_{j,k} \text{ for } j = 1, \dots, a_k; \\ +\frac{1}{2^k} \left(1 - \frac{d_X(x, \omega_j)}{\varepsilon_k}\right) & \text{if } x \in I_{j,k} \text{ for } j = a_k + 1, \dots, 2a_k; \\ 0 & \text{elsewhere.} \end{cases}$$

We define a corresponding function φ_k as

$$\varphi_k(x) := \begin{cases} -\sum_{i=1}^{j-1} f_k(T^{-i}x) & \text{if } x \in I_{j,k} \text{ for } j = 2, \dots, 2a_k; \\ 0 & \text{elsewhere.} \end{cases}$$

With some elementary calculations, one can verify that $f_k(x) = \varphi_k(x) - \varphi_k(Tx)$ and that

$$\int_X \varphi_k d\lambda = a_k^2 \varepsilon_k \frac{1}{2^k} \leq \frac{1}{2^k}.$$

Let $f(x) = \sum_{k=1}^{\infty} f_k(x)$ and $\varphi(x) = \sum_{k=1}^{\infty} \varphi_k(x)$. Since $|f_k(x)| \leq (1/2)^k$, the function $f(x)$ is continuous. On the other hand, since all the φ_k are nonnegative functions, also φ is nonnegative. Furthermore the above estimate on the integral of φ_k implies that $\varphi \in L^1(X, \lambda)$. Finally we notice that $\varphi_k(\omega_{a_k+1}) = k$, and therefore that φ is unbounded. The function φ we constructed satisfies all the hypotheses of the previous example and $f(x) = \varphi(x) - \varphi(Tx)$ is a continuous function.

Proof of Proposition 2.7. The implication (2) \Rightarrow (1) is trivial. Suppose on the other hand that for almost every $\omega \in \Omega$ the family \mathcal{M}_ω is bounded.

Given $\omega \in \Omega$, we define $f(\omega) = \log |\det(M_\omega)|$. By elementary properties of the determinant, we obtain that

$$\log |\det(M_\omega^n)| = \sum_{i=0}^{n-1} f(T^i \omega).$$




Figure 1: the graphs of the functions f_1 and φ_1

Let $Y_i := f(T^{i-1}\omega)$ and $X_n = \sum_{i=1}^n Y_i$. The Y_i -s form a sequence of i.i.d. random variables with expected value $\mathbf{E}(Y_i) = 0$. If $\sqrt{\text{Var}(Y_i)} > 0$, almost surely there exists a sequence n_k so that $X^{n_k} \rightarrow \infty$, contradicting the fact that \mathcal{M}_ω is almost certainly bounded. It follows that $\text{Var}(Y_i) = 0$, which implies that $|\det(M_\omega^n)| = 1$ for all $\omega \in \Omega$ and $n \in \mathbb{N}$.

Let Ω_a as in the Lemma 2.5 and choose $\omega_0 \in \Omega_a$ for which \mathcal{M}_{ω_0} is bounded. We write $M_{\omega_0}^{-n} = (M_{\omega_0}^n)^{-1}$. Since $|\det(M_\omega^n)| = 1$ for every n , the extended family $\widehat{\mathcal{M}}_{\omega_0} := \{(M_{\omega_0}^n)\}_{n \in \mathbb{Z}}$ is also bounded. Let C be a bound on the norms of the matrices of this family.

Let $\omega \in \Omega$ and $n \in \mathbb{N}$. Since $\omega_0 \in \Omega_a$, there exists k_j such that $M_{T^{k_j}\omega_0}^n \rightarrow M_\omega^n$. Now

$$\|M_{T^{k_j}\omega_0}^n\| = \|M_{\omega_0}^{n+k_j} \cdot M_{\omega_0}^{-k_j}\| < C^2.$$

This proves that the set S_ν is bounded in $GL(m, \mathbb{C})$. Its closure $G_\nu = \overline{S_\nu}$ is compact in $GL(m, \mathbb{C})$.

Given $M \in G_\nu$, its orbit is also contained in G_ν , thus it is bounded. Since $|\det(M)| = 1$ the matrix M is diagonalizable and every eigenvalue of M has norm 1. In this case there exists n_k such that $M^{n_k} \rightarrow \text{id}$. Furthermore M^{n_k-1} is a convergent sequence and its limit coincide with M^{-1} , hence G_ν is a compact subgroup of $GL(m, \mathbb{C})$. By the previous lemma it follows that G_ν is conjugated

to a subgroup of $U(m)$. □

2.2 Semi-neutral measures.

Throughout the rest of this section we assume that ν is semi-neutral and with compact support, and consider the induced i.i.d. cocycle. Almost surely $M_\omega^n \neq 0$ for every n . Let $\mathbb{P}\Omega = \Omega \times \mathbb{P}^{m-1}$ and $\mathbb{P}F : \mathbb{P}\Omega \rightarrow \mathbb{P}\Omega$ be the map defined by

$$\mathbb{P}F : (\omega, [v]) \rightarrow (T\omega, [M_\omega v]).$$

Finally let $\Phi : \mathbb{P}\Omega \rightarrow \mathbb{R}$ be the map $\Phi(\omega, [v]) = \log(\|M_\omega v\|/\|v\|)$. For every $\omega \in \Omega$ and $v \in \mathbb{C}^m$ we have

$$\sum_{k=0}^{n-1} \Phi \circ \mathbb{P}F^k(\omega, [v]) = \log \frac{\|M_\omega^n v\|}{\|v\|} \leq \log \|M_\omega^n\|. \quad (5)$$

Let $0 = \kappa_1 > \dots > \kappa_s$ be the Lyapunov indices of the measure ν and $\mathbb{P}^{m-1} = \mathcal{V}_1(\omega) \supset \dots \supset \mathcal{V}_s(\omega)$ be the collection of vector subspaces introduced in Theorem 2.1, defined for almost every $\omega \in \Omega$. We write $\mathbb{P}\Omega_1, \dots, \mathbb{P}\Omega_s$ for the family of disjoint subsets of $\mathbb{P}\Omega$ given by

$$\mathbb{P}\Omega_j = \{(\omega, [v]) : v \in \mathcal{V}_j(\omega) \setminus \mathcal{V}_{j-1}(\omega)\}.$$

We recall the following statement:

Theorem 2.8 ([Le84],[V14]). *Given any $\mathbb{P}F$ -invariant ergodic probability measure m on $\mathbb{P}\Omega$ that projects down to $\mu = \nu^\infty$, there exists $j \in \{1, \dots, s\}$ such that*

$$\int \Phi dm = \kappa_j, \quad \text{and} \quad m(\mathbb{P}\Omega_j) = 1. \quad (6)$$

Conversely, given $j \in \{1, \dots, s\}$ there exists a $\mathbb{P}F$ -invariant ergodic probability measure m_j projecting to μ and satisfying (6).

The following lemma closely resembles Lemma 3.6 of [GR85], we provide a proof for the sake of completeness.

Lemma 2.9. *Let (X, T, μ) be an ergodic dynamical system and $f : X \rightarrow \mathbb{R}$ an integrable function. Suppose that $\int_X f d\mu = 0$, then almost surely there exists a sequence n_k such that*

$$\lim_{k \rightarrow \infty} \sum_{i=0}^{n_k-1} f \circ T^i x = 0$$

Proof. By replacing (X, T, μ) with its natural extension if necessary, we may assume that T is invertible. We consider the product $Y = X \times \mathbb{R}$ and the map $S : Y \rightarrow Y$ defined as

$$S(x, r) := (Tx, r + f(x)).$$

The transformation S preserves the measure $\nu = \mu \times \lambda$, where λ is the Lebesgue measure of \mathbb{R} . We can write

$$S^n(x, r) = \left(T^n x, r + \sum_{i=0}^{n-1} f \circ T^i x \right) = (T^n x, s_n(x, r)).$$

Suppose that there exists a *wandering* subset $A \subset Y$, i.e. $\nu(A) > 0$ and for every couple $i \neq j$ the sets $S^i(A)$ and $S^j(A)$ are disjoint, up to a set of measure 0. By ergodicity of μ , for almost every $(x, r) \in Y$ we have

$$\lim_{n \rightarrow \infty} \frac{s_n(x, r)}{n} = 0.$$

We can choose $B \subset A$ of finite positive measure and such that the limit above holds on B uniformly. Since $\nu(\bigcup_{0 \leq i \leq n} S^i(B)) \leq \lambda(\bigcup_{0 \leq i \leq n} s_i(B))$, it follows that $\lim_n \nu(\bigcup_{0 \leq i \leq n} S^i(B))/n = 0$. On the other hand, since A is wandering and ν is S -invariant we find that $\nu(\bigcup_{0 \leq i \leq n} S^i(B))/n = \nu(B) > 0$; which contradicts the assumption on A .

Let $\varepsilon > 0$ and $Y_\varepsilon = X \times [-\varepsilon, +\varepsilon]$. We write C_ε for the set of all points of Y_ε which do not return to Y_ε , under iterations of the map S . The map S is invertible, therefore if $\nu(C_\varepsilon) > 0$ then the set C_ε is wandering. Since there are no wandering sets, it follows that $\nu(C_\varepsilon) = 0$.

Let $X_\varepsilon \subset X$ be the set of all points x for which there exists $n > 0$ such that $-\varepsilon \leq s_n(x, 0) \leq \varepsilon$. We note that $(X \setminus X_\varepsilon) \times [-\varepsilon/2, \varepsilon/2] \subset C_{\varepsilon/2}$, thus X_ε is a full measure subset of X . The set

$$X_0 = \bigcap_n T^{-n} \left(\bigcap_k X_{1/k} \right)$$

is the intersection of countable many full measure sets, thus it has full measure. Take $x_0 \in X_0$, since $X_0 \subset \bigcap_k X_{1/k}$, then for every k there exists a minimal $n_k > 0$ such that $-1/k \leq s_{n_k}(x_0, 0) \leq 1/k$. We notice that $n_{k+1} \geq n_k$, thus either $n_m \rightarrow \infty$, in which case the lemma is proved, or there exists n'_1 such that $s_{n'_1}(x_0, 0) = 0$. If the second case occurs, we let $x_1 = T^{n'_1}x_0 \in X_0$ and we repeat the argument for x_1 . It follows that one can always construct a sequence which satisfies the requirements of the Lemma. \square

Proposition 2.10. *Let ν be a semi-neutral measure with compact support. Suppose that \mathcal{M}_ω is almost surely bounded, then for every $M_\omega^n \in S_\nu$ we have*

$$\|M_\omega^n\| \geq 1.$$

Proof. Given $K > 0$, let

$$\Omega_K := \left\{ \omega \in \Omega \mid \sup_n \log \|M_\omega^n\| \leq K \right\}.$$

The set $\Omega_b = \bigcup_{K>0} \Omega_K$ coincides with the set of all ω for which \mathcal{M}_ω is bounded. It then follows that $\lim_{K \rightarrow \infty} \nu(\Omega) = 1$ and that we can choose $K_0 > 0$ so that $\nu(\Omega_{K_0}) > 0$.

Suppose that there exists $M_{\omega_0}^N \in S_\nu$ with $\|M_{\omega_0}^N\| < 1$. By the semigroup structure of S_ν we may further assume that $\|M_{\omega_0}^N\| \leq 1/(3K_0)$.

For $i = 1, \dots, N$ we choose $U_i \subset \text{Mat}(m, \mathbb{C})$ such that the set $\mathcal{C}_{U_1, \dots, U_N}$ has positive μ -measure and $\|M_\omega^N\| \leq 1/(2K_0)$ for every $\omega \in \mathcal{C}_{U_1, \dots, U_N}$.

We define the set

$$\mathcal{U} := \mathcal{C}_{U_1, \dots, U_N} \cap T^{-N}(\Omega_{K_0}) = U_1 \times \dots \times U_N \times \Omega_{K_0}.$$

It is clear that this set has positive μ -measure.

Let m_1 be the $\mathbb{P}F$ -invariant ergodic measure on $\mathbb{P}\Omega$ defined in Theorem 2.8. Given $(\omega, [v]) \in \mathcal{U} \times \mathbb{P}^{m-1}$, for every $n \geq N$ we have

$$\begin{aligned} \sum_{k=0}^{n-1} \Phi \circ \mathbb{P}F^k(\omega, [v]) &\leq \log \|M_\omega^n\| \\ &\leq \log \|M_\omega^N\| \|M_{\sigma^N(\omega)}^{n-N}\| \\ &\leq -\log 2. \end{aligned}$$

By (6) we have $\int_\Omega \Phi dm_1 \neq 0$ and $m_1(\mathcal{U} \times \mathbb{P}^{m-1}) = \mu(\mathcal{U}) > 0$, contradicting the previous Lemma. It follows that for every $M_\omega^n \in S_v$ we have $\|M_\omega^n\| \geq 1$. \square

3. COMPOSITIONS OF RANDOM GERMS

Let $\mathcal{O}(\mathbb{C}^m, 0)$ be the space of all germs of holomorphic functions fixing the origin. This set can be endowed with the so-called inductive limit topology τ_{ind} , see the appendix of this paper for more details. Let ν be a Borel probability measure on the set $\mathcal{O}(\mathbb{C}^m, 0)$. Let Ω be the space of all sequences in $\text{supp}(\nu)$ and $\mu = \nu^\infty$. Let $T : \Omega \rightarrow \Omega$ denote the left shift function and $f : \Omega \rightarrow \mathcal{O}(\mathbb{C}^m, 0)$ be the function which returns the first element of the sequence. As we will see later, the function f defines a (non linear) cocycle over T . In analogy with the previous section we will write

$$f_\omega^n = f_{T^{n-1}\omega} \circ \cdots \circ f_\omega,$$

and $\mathcal{F}_\omega = \{f_\omega^n\}$. Finally we define

$$\Omega_{nor} = \{\omega \in \Omega \mid \mathcal{F}_\omega \text{ is a normal family near the origin}\}.$$

Proposition 3.1. *The set Ω_{nor} is measurable.*

Proof. Normality near the origin is equivalent to relatively compactness of the set \mathcal{F}_ω with respect to τ_{ind} . Furthermore by Corollary A.9 the set \mathcal{F}_ω is relatively compact in $\mathcal{O}(\mathbb{C}^m, 0)$ if and only if $\mathcal{F}_\omega \subset \overline{B}_n(0, r)$ for some natural number n and $r > 0$. The terminology $\overline{B}_n(0, r)$ is introduced in the appendix.

Let r_n be an increasing sequence of positive real numbers such that $r_n \rightarrow \infty$. Since $\overline{B}_n(0, r) \subset \overline{B}_{n+1}(0, r)$, it follows that

$$\begin{aligned} \Omega_{nor} &= \bigcup_n \{\omega \in \Omega \mid \mathcal{F}_\omega \subset \overline{B}_n(0, r_n)\} \\ &= \bigcup_n \bigcap_m \{\omega \in \Omega \mid f_\omega^m \in \overline{B}_n(0, r_n)\}. \end{aligned}$$

The function $\omega \mapsto f_\omega^m$ is measurable for every m . By Lemma A.5 and Corollary A.9, $\overline{B}_n(0, r_n)$ is a closed set in the inductive limit topology. This proves that Ω_{nor} is the intersection of countably many measurable sets, and thus that it is a measurable set. \square

Problem Let $\omega \in \Omega$ be a random sequence of germs. What is the probability that \mathcal{F}_ω is a normal family?

We notice that the set Ω_{nor} is backward invariant. By ergodicity of T it follows that

Proposition 3.2. *Normality of \mathcal{F}_ω occurs either with probability 1 or 0, depending on the probability measure μ .*

Random Fatou set We say that the origin belongs to the *(local) random Fatou set* if Ω_{nor} has full measure. If this is not the case we say that the origin belongs to the *(local) random Julia set*.

Suppose now that the measure ν has compact support. By Corollary A.9, there exists $R > 0$ such that each $f \in \text{supp}(\nu)$ is holomorphic and bounded on \mathbb{B}_R , the open ball of \mathbb{C}^m with radius R and center the origin. Given $\omega \in \Omega$ we may define the set

$$D_\omega := \bigcap_{n=0}^{\infty} (f_\omega^n)^{-1}(\mathbb{B}_R) \quad (7)$$

Clearly $0 \in D_\omega$, thus D_ω is not empty. The following invariance relation holds

$$f_\omega^n(D_\omega) = D_{T^n(\omega)}.$$

Given $z \in D_\omega$ its orbit $\{f_\omega^n(z) \mid n \in \mathbb{N}\}$ is well defined and bounded. By the weak Montel Theorem the family \mathcal{F}_ω is normal on $\text{int}(D_\omega)$. We will call this set the *(local) Fatou set* of the sequence ω and we will denote it as F_ω . It is clear that the origin lies in the random Fatou set if and only if it almost certainly lies in the Fatou set of ω .

We define

$$D := \{(\omega, z) \in \Omega \times \mathbb{C}^m \mid z \in D_\omega\}.$$

The *(non linear) cocycle* defined by f over T is the skew-product transformation $F : D \rightarrow D$

$$F(\omega, z) = (T\omega, f_\omega(z)), \quad F^n(\omega, z) = (T^n\omega, f_\omega^n(z)).$$

By Lemma A.1 the map $d_0 : f \mapsto df(0)$ is a continuous map. Following the previous section, we introduce the following classification of probability measures on $\mathcal{O}(\mathbb{C}^m, 0)$:

Definition Let ν be a probability measure on $\mathcal{O}(\mathbb{C}^m, 0)$ so that $(d_0)_*\nu$ has compact support. We say that the measure ν is *attracting, repelling, neutral or semi-neutral* if $(d_0)_*\nu$ is respectively attracting, repelling, neutral or semi-neutral.

We are now ready to state the central theorem of this paper. Notice that the neutral case coincides with Theorem 1.1 of the introduction .

Theorem 3.3. *Let ν be a probability measure on $\mathcal{O}(\mathbb{C}^m, 0)$ with compact support, and suppose that ν is either attracting, repelling or neutral.*

Attracting: the origin lies in the random Fatou set, and it is almost surely an attracting point for the system $\{f_\omega^n|_{D_\omega}\}$.

Repelling: the origin lies in the random Julia set.

Neutral: the origin lies in the random Fatou set if and only if all the germs in $\text{supp}(\nu)$ are simultaneously linearizable, and the semigroup of the differentials

$$dS_\nu := \{df_\omega^n(0) \mid \omega \in \Omega, n \in \mathbb{N}\},$$

is conjugate to a sub-semigroup of $U(m)$.

If we drop the compactness hypothesis on $\text{supp}(\nu)$, then this theorem is false in general. We will show a counterexample, for an attracting measure ν , with non compact support.

Example Let $0 < \lambda_1 < 1$ and let $a_i = \lambda_1^{-3 \cdot 2^i}$. We consider the family

$$\mathcal{A} := \{f_i(z) = \lambda_1 z + a_i z^2\}.$$

Let ν be the probability measure on $\mathcal{O}(\mathbb{C}, 0)$ such that $\nu(f_i) = \frac{1}{2^i}$. The support of ν coincides with \mathcal{A} and is not compact. On the other hand we have $\text{supp}((d_0)_* \nu) = \{\lambda\}$, which implies that the measure ν is attracting. In this case the (unique) Lyapunov index is $\kappa_1 = \log \lambda_1 < 0$.

Let $\omega = (f_{i_1}, f_{i_2}, \dots)$ be a sequence of i.i.d. random germs of \mathcal{A} . We write $\lambda_2^{(n)}$ for the coefficient relative to z^2 of f_ω^n . It follows that

$$\lambda_2^{(n)} = \lambda_1^n \sum_{k=1}^n \lambda_1^{k-1} a_{i_k} \geq \lambda_1^{2n} a_{i_n}.$$

Write p_n for the probability that $\lambda_1^{2n} a_{i_n} \geq \lambda_1^{-n}$. The inequality holds if and only if $i_n \geq \log n$, thus

$$p_n = \sum_{k \geq \log n} \frac{1}{2^k} \sim \frac{2}{n}.$$

Given ω and $m > 0$, by the independence assumption, the probability that for every $n \geq m$ the inequality $\lambda_1^{2n} a_{i_n} < \lambda_1^{-n}$ holds is equal to

$$P_m = \prod_{n \geq m} (1 - p_n) = \text{Exp} \left(\sum_{n \geq m} \log(1 - p_n) \right).$$

Since the general term of the series above is asymptotic to $-2/n$ we see that $\sum_{n \geq m} \log(1 - p_n) = -\infty$ and therefore $P_m = 0$. It follows that almost surely there exists $n_k \rightarrow \infty$ such that $\lambda_2^{(n_k)} \rightarrow \infty$, thus that almost surely \mathcal{F}_ω is not a normal family.

We conclude this section with the discussion of a particular semi-neutral measure. The example shows that, in the semi-neutral case, $\text{supp}(\nu)$ may contain germs which are not linearizable but at the same time the origin can be in the random Fatou set. Furthermore the semigroup S_ν generated by $\text{supp}(\nu)$ may not be bounded in $\mathcal{O}(\mathbb{C}^m, 0)$. Thus the methods developed in section 4.2 for the neutral case, do not work for the semi-neutral one.

Example Suppose that f and g are given by

$$\begin{aligned} f(z, w) &= (z, w/2), \\ g(z, w) &= (z + zw, w). \end{aligned}$$

The n -th iterate of g has the form $g^n(z, w) = (z + nzw + O(|z|^3), w)$. Therefore $\{g^n\}$ is not normal on any neighborhood of the origin, thus it cannot be linearized.

Let $\nu := \frac{1}{2}\delta_f + \frac{1}{2}\delta_g$ be a probability measure. A short computation gives $\kappa_1 = 0$ and $\kappa_2 = \frac{1}{2} \log \frac{1}{2} < 0$.

Let $\omega \in \Omega$ be a sequence of i.i.d. random germs and write $f_\omega^n = (f_{\omega,1}^n, f_{\omega,2}^n)$, then

$$\begin{aligned} |f_{\omega,1}^{n+1}(z, w)| &\leq \begin{cases} |f_{\omega,1}^n(z, w)| & \text{if } f_n = f; \\ |f_{\omega,1}^n(z, w)| \left(1 + |f_{\omega,2}^n(z, w)|\right) & \text{if } f_n = g; \end{cases} \\ |f_{\omega,2}^{n+1}(z, w)| &= \begin{cases} \frac{1}{2}|f_{\omega,2}^n(z, w)| & \text{if } f_n = f; \\ |f_{\omega,2}^n(z, w)| & \text{if } f_n = g. \end{cases} \end{aligned}$$

Let $\alpha(n) = \#\{k \in \{1, \dots, n\} \mid f_k = f\}$. Using the estimates above we find that

$$\begin{aligned} |f_{\omega,2}^n(z, w)| &= \frac{|w|}{2^{\alpha(n)}}, \\ |f_{\omega,1}^n(z, w)| &\leq |z| \prod_{i=1}^n \left(1 + \frac{|w|}{2^{\alpha(i)}}\right). \end{aligned}$$

It is a well known fact that almost surely we have $\alpha(n) \sim n/2$. It follows that

$$\log |f_{\omega,1}^n(z, w)| \leq \log |z| + \sum_{i=1}^n \log \left(1 + \frac{|w|}{2^{\alpha(i)}}\right).$$

As $n \rightarrow \infty$, almost certainly $a_i := \log \left(1 + \frac{|w|}{2^{\alpha(i)}}\right) \sim \frac{|w|}{2^{i/2}}$, therefore $\sum_{i=1}^{\infty} a_i < \infty$ and $f_\omega^n(z, w)$ is uniformly bounded for every $(z, w) \in \mathbb{C}^2$. We conclude that in this case \mathcal{F}_ω is normal for almost every sequence ω .

4. PROOF OF THEOREM 3.3

We start this section with the simpler attracting and repelling cases, and will cover the neutral case afterwards.

4.1 Attracting and repelling measures.

If the measure ν is repelling, then the sequence of the differentials $df_\omega^n(0)$ diverges almost surely. If this is the case, the origin lies in the random Julia set.

The following statement implies the attracting case. It is likely well known, we provide a proof for the sake of completeness.

Proposition 4.1. *Let ν be an attracting measure on $\mathcal{O}(\mathbb{C}^m, 0)$ with compact support, then the origin lies in the random Fatou set. Furthermore, there exists almost surely a neighborhood of the origin on which all orbits converge to the origin.*

Proof. By Lemma 2.2 and (4), there exists $n_0 > 0$ such that $\mathbf{E} \log \|df_\omega^{n_0}(0)\| < 0$. Without loss of generality we may assume that $n_0 = 1$.

Given $\delta > 0$ we write

$$\begin{aligned} \mathbf{E} \log \|df(0)\| &= \int_{\|df(0)\| \leq \delta} \log \|df(0)\| d\nu + \int_{\|df(0)\| > \delta} \log \|df(0)\| d\nu \\ &= L_\delta + U_\delta. \end{aligned}$$

Suppose that $U_\delta < 0$ for some $\delta > 0$. In this case we can find $\varepsilon > 0$ sufficiently small so that

$$\mathbf{E} \log (\|df(0)\| + \varepsilon) < 0. \quad (8)$$

On the other hand, suppose that $U_\delta \geq 0$ for every $\delta > 0$. Let δ_n be a decreasing sequence of positive numbers converging to 0 and such that $\delta_0 > \sup_{f \in \text{supp}(\nu)} \|df(0)\|$. It then follows that

$$\begin{aligned} \int_{df(0) \neq 0} \log \|df(0)\| d\nu &= \sum_{n=1}^{\infty} \int_{\delta_n < \|df(0)\| \leq \delta_{n-1}} \log \|df(0)\| d\nu \\ &= \lim_{N \rightarrow \infty} U_{\delta_N} \\ &\geq 0. \end{aligned}$$

Since $\mathbf{E} \log \|df(0)\| < 0$ we must have $M_0 := \nu\{df(0) = 0\} > 0$. We conclude that

$$\begin{aligned} \mathbf{E} \log (\|df(0)\| + \varepsilon) &= M_0 \log \varepsilon + \int_{\|df(0)\| \neq 0} (\log \|df(0)\| + \varepsilon) d\mu(f) \\ &\leq M_0 \log \varepsilon + \sup_{f \in \text{supp}(\mu)} \log (\|df(0)\| + \varepsilon). \end{aligned}$$

Therefore also in this second case we can choose $\varepsilon > 0$ sufficiently small so that (8) is satisfied.

Let $\alpha_f := \|df(0)\| + \varepsilon$. By the compactness of $\text{supp}(\nu)$ there exists $0 < r < R$ such that every $f \in \text{supp}(\nu)$ is holomorphic on \mathbb{B}_R and

$$\|f(z)\| \leq \alpha_f \|z\|, \quad \forall z \in \mathbb{B}_r.$$

Since $\mathbf{E} \log \alpha_f < 0$, then for almost every sequence $\omega \in \Omega$ we have $\alpha_\omega^n := \alpha_{f_{T^{n-1}\omega}} \cdots \alpha_{f_\omega} \rightarrow 0$. In particular there exists $0 < \delta \leq r$, which depends on ω , such that if $\|z\| < \delta$ then $\alpha_\omega^n \|z\| < r$ for every $n > 0$.

Let $\|z\| < \delta$, a simple induction argument shows that

$$\|f_\omega^n(z)\| \leq \alpha_\omega^n \|z\| < r, \quad \forall n \in \mathbb{N}.$$

This shows that $\mathbb{B}_\delta \subset D_\omega$ and that the family \mathcal{F}_ω is normal at the origin. This is true for almost every $\omega \in \Omega$, therefore the origin lies in the random Fatou set. Furthermore since $\alpha_\omega^n \rightarrow 0$ as $n \rightarrow \infty$, the orbits of all points in \mathbb{B}_δ converge to the origin. \square

4.2 Neutral measures

The following lemma is proved along the lines of Lemma 2.5.

Lemma 4.2. *Let ν be a probability measure on $\mathcal{O}(\mathbb{C}^m, 0)$ with compact support. Then the set*

$$\Omega_a := \left\{ \omega \in \Omega \mid \forall n \in \mathbb{N} \text{ and } \forall \alpha \in \Omega, \exists k_j : f_{T^{k_j} \omega}^n \rightarrow f_\alpha^n \right\}$$

is a full measure subset of Ω .

Proof. Unlike the standard topology of $\text{Mat}(m, \mathbb{C})$, the inductive limit topology is not metrizable. However, given $n \in \mathbb{N}$, the image of Ω under the continuous map $\omega \mapsto f_\omega^n$ is a compact set in the inductive limit topology. By Corollary A.9, we can find a natural number N_n so that, for every $\omega \in \Omega$, the germ f_ω^n belongs to X_{N_n} .

We choose a sequence $\varepsilon_j \rightarrow 0$ and we define

$$\Omega_{n,j} = \{ \omega \in \Omega \mid \forall \alpha \in \Omega, \exists k : d_{N_n}(f_\omega^n, f_\alpha^n) < \varepsilon_j \}.$$

By Theorem A.6, we have $\Omega_a = \cup_{n,j} \Omega_{n,j}$. From this point on we follow the proof of Lemma 2.5, replacing the standard metric of $\text{Mat}(m, \mathbb{C})$ with the metric d_{N_n} . \square

Theorem 4.3 (Hurwitz Theorem). *Suppose that $(f_n)_n$ is a sequence of injective holomorphic maps on a domain $U \subset \mathbb{C}^m$ converging uniformly on compact subsets to a function g . Then g is either injective or degenerate.*

Given a probability measure ν with compact support, choose $R > 0$ and define D_ω as explained in section 3.

Lemma 4.4. *Let ν be a neutral measure with compact support. The origin lies in the random Fatou set if and only if given $\varepsilon > 0$ there exists $\rho > 0$ such that, for every $\omega \in \Omega$, we have $\mathbb{B}_\rho \subset D_\omega$ and*

$$f_\omega^n(\mathbb{B}_\rho) \subset \mathbb{B}_\varepsilon, \quad \forall n \in \mathbb{N}. \quad (9)$$

Proof. Suppose that for every $\varepsilon > 0$ there exists such ρ , then by the weak Montel Theorem the origin lies in the random Fatou set.

Suppose on the other hand that the origin lies in the random Fatou set and let $\omega_0 \in \Omega_a \cap \Omega_{nor}$. By Ascoli–Arzelá theorem, given $\varepsilon > 0$ there exists $\delta_0 > 0$ such that $\mathbb{B}_{\delta_0} \subset D_{\omega_0}$ and

$$f_{\omega_0}^n(\mathbb{B}_{\delta_0}) \subset \mathbb{B}_\varepsilon, \quad \forall n \in \mathbb{N}.$$

By the compactness of $\text{supp}(\nu)$, by choosing $\varepsilon > 0$ small enough, we may assume that every $f \in \text{supp}(\nu)$ is holomorphic and injective on \mathbb{B}_ε . Therefore $f_{\omega_0}^n : \mathbb{B}_{\delta_0} \rightarrow \mathbb{B}_\varepsilon$ is injective for all values of n .

Suppose that there exists n_k so that

$$\mathbb{B}_{1/k} \not\subset f_{\omega_0}^{n_k}(\mathbb{B}_{\delta_0}). \quad (10)$$

By taking a subsequence if necessary, we may assume that $f_{\omega_0}^{n_k} \rightarrow g$ uniformly on compact subsets of \mathbb{B}_{δ_0} . By Proposition 2.7 we have $|\det(df_{\omega_0}^{n_k}(0))| = 1$ for

every k , thus $|\det(dg(0))| = 1$. Thanks to Hurwitz Theorem we can conclude that g is injective on \mathbb{B}_{δ_0} .

Since g is an open function, we can choose $\rho' > 0$ such that $\mathbb{B}_{\rho'} \subset g(\mathbb{B}_{\delta_0})$ and thus such that $\mathbb{B}_{\rho'} \subset f_{\omega_0}^{nk}(\mathbb{B}_{\delta_0})$ for k big enough, which contradicts (10).

It follows that there exists $\rho_0 > 0$ so that $\mathbb{B}_{\rho_0} \subset f_{\omega_0}^n(\mathbb{B}_{\delta_0})$ for every n . By the invariance relation (7), given $n, k \in \mathbb{N}$, we find that $\mathbb{B}_{\rho_0} \subset D_{T^k \omega_0}$ and

$$\begin{aligned} f_{T^k \omega_0}^n(\mathbb{B}_{\rho_0}) &\subset f_{T^k \omega_0}^n \circ f_{\omega_0}^k(\mathbb{B}_{\delta_0}) \\ &\subset \mathbb{B}_{\varepsilon}. \end{aligned}$$

Let $\rho < \rho_0$ be fixed. Given $\omega \in \Omega$ and $n \in \mathbb{N}$, there exists a sequence k_j so that $f_{T^{k_j} \omega_0}^n$ converges to f_{ω}^n in the inductive limit topology. On the other hand, by the weak Montel Theorem, we may also assume that $f_{T^{k_j} \omega_0}^n$ converges to some g , uniformly on compact subsets of \mathbb{B}_{ρ_0} . By Theorem A.6 the maps g and f_{ω}^n agree on a small ball containing the origin, therefore, by the identity principle, they coincide as germs. We conclude that f_{ω}^n is holomorphic on \mathbb{B}_{ρ} and, since ρ is independent from n and ω , that $\mathbb{B}_{\rho} \subset D_{\omega}$ for every $\omega \in \Omega$. Finally uniform convergence on compact subsets of \mathbb{B}_{ρ_0} implies (9). \square

We write S_v for the semigroup

$$S_v := \{f_{\omega}^n \mid \omega \in \Omega, n \in \mathbb{N}\}.$$

The following corollary is an immediate consequence of the previous lemma.

Corollary 4.5. *Let v be a neutral measure with compact support. Then the origin lies in the random Fatou set if and only if S_v is relatively compact in $\mathcal{O}(\mathbb{C}^m, 0)$.*

Definition Given $f \in \mathcal{O}(\mathbb{C}^m, 0)$ we say that f is *linearizable*, if there exists $\varphi \in \mathcal{O}(\mathbb{C}^m, 0)$, locally invertible at the origin, such that

$$\varphi \circ f(z) = df(0) \cdot \varphi.$$

If this is the case, we say that f is *linearized* by φ .

Corollary 4.6. *Let v be a neutral measure with compact support. If the origin lies in the random Fatou set, then every element of S_v is linearizable.*

Proof. If the semigroup S_v is relatively compact in $\mathcal{O}(\mathbb{C}^m, 0)$ then, given $f \in S_v$, the family $\{f^n\}$ is also relatively compact. It follows that the origin lies in the Fatou set of f , and therefore that the germ is linearizable. \square

We notice that linearizability of every element of S_v does not imply that the origin lies in the random Fatou set. Consider the following example.

Example Let $\lambda = e^{2\pi i \alpha}$ be an irrational rotation, with α Brjuno number. We define the maps

$$f_1(z) = \lambda(z + z^2), \quad \text{and} \quad f_2(z) = \lambda(z - z^2),$$

and consider a measure such that $\text{supp}(v) = \{f_1, f_2\}$.

In this case we see immediately that every function in the semigroup $S_\nu = \langle f_1, f_2 \rangle$ is linearizable. Nevertheless f_1 and f_2 are not simultaneously linearizable.

Let $z_0 \neq 0$ and define integers $n_i \in \{1, 2\}$ recursively as follows: if $\operatorname{Re}(z_i) \geq 0$ then $n_{i+1} = 1$, else $n_{i+1} = 2$. Let $\omega := (f_{n_1}, f_{n_2}, \dots)$, and $z_i = f_\omega^i(z_0)$.

We claim that the orbit of z_0 converges to infinity. To see this suppose that $\operatorname{Re}(z_i) \geq 0$. Then it follows that the angle between the vectors z_i and z_i^2 is at most $\pi/2$, and hence

$$|z_{i+1}| = |z_i + z_i^2| \geq |z_i|.$$

The irrational rotation guarantees that the norm increases often enough to converge to infinity, proving the claim.

The conclusion is that in this case S_ν is not relatively compact, therefore the origin does not lie in the random Fatou set.

Suppose that ν is a neutral measure with compact support for which the origin lies in the random Fatou set. We write G_ν for the closure of S_ν in $\mathcal{O}(\mathbb{C}^m, 0)$.

Lemma 4.7. *The set G_ν is a compact topological group. Moreover there exists an open subset $M \subset \mathbb{C}^m$ such that every $f \in G_\nu$ belongs to $\operatorname{Aut}(M)$.*

Proof. By Lemma 4.4 the set G_ν is compact. Furthermore, by Corollary 4.6, every $f \in S_\nu$ is linearizable and, by Proposition 2.7 the differential $df(0)$ is conjugated to an element of $U(m)$. It follows that for some sequence n_j we have $f^{n_j} \rightarrow \operatorname{id}$. This proves that G_ν contains the identity element. Furthermore the germ $g = \lim_{j \rightarrow \infty} f^{n_j^{-1}}$ is the inverse of f in $\mathcal{O}(\mathbb{C}^m, 0)$, thus G_ν contains also the inverse of f .

Let $f \in G_\nu$ and $f_n \in S_\nu$ such that $f_n \rightarrow f$. Every f_n has an inverse $f_n^{-1} \in G_\nu$. By taking a subsequence if necessary, we may assume that $f_n^{-1} \rightarrow g \in G_\nu$. It is not difficult to prove that g is the inverse element of f in $\mathcal{O}(\mathbb{C}^m, 0)$, which finally proves that G_ν is a compact group.

Let $\varepsilon > 0$ and $\rho > 0$ as in Lemma 4.4. If we choose ε small enough we may further assume that all the elements of $\operatorname{supp}(\nu)$ are univalent on \mathbb{B}_ε . It follows that every $f \in S_\nu$ is univalent on \mathbb{B}_ρ . Since every $g \in G_\nu$ is invertible, by Hurwitz's theorem, it is also univalent on \mathbb{B}_ρ .

By taking a smaller ε if necessary, we may further assume that every $g \in G_\nu$ is univalent on \mathbb{B}_ε . We define the open set

$$M = \bigcup_{g \in G_\nu} g(\mathbb{B}_\rho) \subset \mathbb{B}_\varepsilon.$$

Given $g \in G_\nu$ and $x \in M$, then $x = \hat{g}(z)$ for some $\hat{g} \in G_\nu$ and $z \in \mathbb{B}_\rho$. It follows that $g(M) \subset M$. Furthermore, since $g^{-1} \in G_\nu$, we obtain the equality $g(M) = M$. Which proves that $g \in \operatorname{Aut}(M)$. \square

The following result is known as Bochner's Linearization Theorem. A proof of the theorem, valid for C^k diffeomorphism, can be found in [DK00]. The same proof is valid also in the holomorphic case, up to small modifications.

Theorem 4.8 (Bochner's Linearization Theorem). *Let M be a complex manifold and $x_0 \in M$. Let A be a continuous homomorphism from a compact group K to $\text{Aut}(M)$ such that $A_k(x_0) = x_0$, for all $k \in K$. Then there exists a K invariant open neighborhood U of x_0 in M and biholomorphism φ from U onto an open neighborhood $V \subset T_{x_0}M$ of 0 , such that:*

$$\varphi(x_0) = 0, \quad d\varphi(x_0) = \text{id} : T_{x_0}M \rightarrow T_{x_0}M$$

and

$$\varphi \circ A_k(x) = dA_k(x_0) \cdot \varphi(x) \quad \forall k \in K, x \in U).$$

Definition We say that two or more germs are *simultaneously linearizable* if there exists a locally invertible $\varphi \in \mathcal{O}(\mathbb{C}^m, 0)$ such that all the germs are linearized by φ .

Lemma 4.9. *Let ν be a neutral measure on $\mathcal{O}(\mathbb{C}^m, 0)$ with compact support. If the origin lies in the random Fatou set then all the elements $f \in \text{supp}(\nu)$ are simultaneously linearizable.*

Proof. Let G_ν be the closure of S_ν and M be the G_ν -invariant open set described in Lemma 4.7. The compact group G_ν induces an action on M that satisfies the hypothesis of Bochner's Linearization Theorem. It follows that all the germs $f \in G_\nu$ can be simultaneously linearized. \square

We are finally ready to conclude the proof of Theorem 3.3.

Proof of Theorem 3.3. Suppose that the origin lies in the random Fatou set. By the previous lemma it follows that all the germs in $\text{supp}(\nu)$ are simultaneously linearizable, which implies that the semigroup S_ν is conjugated to the semigroup of the differentials dS_ν . Furthermore by Proposition 2.7, the semigroup dS_ν is itself conjugate to a sub-semigroup of $U(m)$. The other implication of the theorem is trivial. \square

5. DISCUSSION OF SEMI-NEUTRAL MEASURES

Recall from section 3 that for every $\omega \in \Omega_{\text{nor}}$ the origin belongs to the Fatou set $F_\omega := \text{int}(D_\omega)$. Let U_ω be the connected component of F_ω containing the origin.

Proposition 5.1. *Let ν be a semi-neutral measure on $\mathcal{O}(\mathbb{C}^m, 0)$ with compact support, so that the origin lies in the random Fatou set. Then almost surely, every limit map $g = \lim_{k \rightarrow \infty} f_\omega^{n_k}$ is degenerate on U_ω .*

Proof. By Corollary 2.3 we have $\mathbf{E} \log |\det df(0)| < 0$. Along the lines of Proposition 4.1, we can choose $\varepsilon > 0$ such that $\mathbf{E} \log (|\det df(0)| + \varepsilon) < 0$. In particular for almost every $\omega \in \Omega$ we have

$$\prod_{k=1}^{\infty} (|\det df_{T^{k-1}\omega}(0)| + \varepsilon) = 0. \quad (11)$$

By compactness of $\text{supp}(\nu)$, we may choose $r > 0$ so that for every $\|z\| < r$ we have

$$|\det df(z)| < |\det df(0)| + \varepsilon, \quad \forall f \in \text{supp}(\mu).$$

Since the origin belongs to the random Fatou set, for almost every $\omega \in \Omega$ there exists $\delta_\omega > 0$ such that $\|f_\omega^n(z)\| < r$ for every $z \in \mathbb{B}_{\delta_\omega}$ and $n \geq 0$. We conclude that for $z \in \mathbb{B}_{\delta_\omega}$

$$\begin{aligned} |\det df_\omega^n(z)| &= \prod_{k=1}^n |\det df_{T^{k-1}\omega}(f_\omega^{k-1}(p))| \\ &\leq \prod_{k=1}^n (|\det df_{T^{k-1}\omega}(0)| + \varepsilon). \end{aligned}$$

Let n_k be a sequence such that $f_\omega^{n_k} \rightarrow g$ locally uniformly on a neighborhood U of the origin, then on a possibly smaller neighborhood we have $\det dg \equiv 0$. By the identity principle we conclude that the same is true on U_ω . \square

Given $z \in U_\omega$ we define the *stable set*

$$\mathbb{W}_\omega^s(z) := \{w \in U_\omega \mid \|f_\omega^n(z) - f_\omega^n(w)\| \rightarrow 0\}.$$

When $m = 2$, we will prove that, given ν semi-neutral with compact support and such that the origin lies in the random Fatou set, the stable set through every point z sufficiently close to the origin is locally a complex submanifold. It is a natural question whether the same is true when $m > 2$.

Before proceeding to the proof of this result we will present two examples of this phenomenon in the case the germs in $\text{supp}(\nu)$ are linear maps.

Example Suppose $\nu = \frac{1}{2}\delta_g + \frac{1}{2}\delta_h$, where

$$g(z, w) = \begin{pmatrix} 1/2 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix}, \quad h(z, w) = \begin{pmatrix} 1/2 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix}.$$

We notice that given $\omega \in \Omega$

$$f_\omega^{n+1}(z, w) = \begin{pmatrix} 1/2^n & \alpha_n \\ 0 & 1 \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix}, \quad \text{with } \alpha_n = \begin{cases} \frac{\alpha_n}{2} & \text{if } f_{T^n\omega} = g \\ \frac{\alpha_n}{2} + 1 & \text{if } f_{T^n\omega} = h. \end{cases}$$

The two Lyapunov indices of the measure ν are $\kappa_1 = 0$ and $\kappa_2 = -\log 2$, therefore the measure is semi-neutral. Furthermore we have $0 \leq \alpha_n < 2$ for every n and every ω , which implies that the origin lies in the random Fatou set. Given $(z_0, w_0) \in U_\omega = \mathbb{C}^2$, we can write its stable set as

$$\mathbb{W}_\omega^s(z_0, w_0) = \{(z, w) \in \mathbb{C}^2 \mid w = w_0\},$$

which is a one-dimensional manifold. We notice that in this case the stable manifolds are independent from the sequence $\omega \in \Omega$, but that this is not true in general.

Consider for example the measure $\tilde{\nu}$, obtained by taking instead the maps

$$g(z, w) = \begin{pmatrix} 1/2 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix}, \quad h(z, w) = \begin{pmatrix} 1/2 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix}.$$

This measure has the same Lyapunov indices and the origin lies in the random Fatou set. However the sequence f_ω^n converges, for every ω , to a map of the form $\tilde{g}(z, w) = (z + \beta_\omega w, 0)$. Hence the stable manifolds are again parallel complex planes, but now depend on ω .

Remark The last two examples are helpful models in order to understand the random dynamics of semi-neutral measures. We notice that in both cases $df(0)$ and $dg(0)$ share an eigenvalue. In the theory of linear cocycles one says that a cocycle is *strongly irreducible* if there is no finite family of proper subspaces invariant by M_ω for ν -almost every ω . This raises the following question

Does there exist a semi-neutral measure, such that M_ω^n is almost surely bounded, but the corresponding cocycle is strongly irreducible?

Lemma 5.2. *There exists $r > 0$ and, for almost every $\omega \in \Omega$, a sequence n_k dependent by ω , so that $f_\omega^{n_k} \rightarrow g$ locally uniformly on U_ω and so that*

$$\mathbb{W}_\omega^s(z) = \{w \in U_\omega \mid g(w) = g(z)\}$$

for every $z \in U_\omega$ which satisfies $\|g(z)\| \leq r$.

Proof. Given $r > 0$ we define the measurable sets

$$\mathcal{A}_r := \{\omega \in \Omega \mid \mathbb{B}_r \subseteq U_\omega\}.$$

It is clear that $\mathcal{A}_{r_1} \supset \mathcal{A}_{r_2}$, when $r_1 < r_2$, and that $\bigcup_{r>0} \mathcal{A}_r = \Omega_{nor}$. Since the latter is a full measure set, we may choose $r > 0$ so that $\mu(\mathcal{A}_r) > 1/2$.

Let $\varepsilon_1 > 0$, by the Ascoli–Arzelà Theorem the collection of measurable sets

$$\mathcal{B}_\delta^{(1)} := \left\{ \omega \in \mathcal{A}_r \mid \begin{array}{l} \forall z, w \in \mathbb{B}_r \text{ with } \|z - w\| < \delta : \\ \sup_n \|f_\omega^n(z) - f_\omega^n(w)\| < \varepsilon_1 \end{array} \right\},$$

defined for $\delta > 0$, covers \mathcal{A}_r . Furthermore it satisfies $\mathcal{B}_\delta^{(1)} \supset \mathcal{B}_{\delta'}^{(1)}$, for $\delta < \delta'$, thus we may choose $\delta_1 > 0$ so that $\mu(\mathcal{B}_{\delta_1}^{(1)}) > 1/2$. We will write $\mathcal{B}^{(1)}$ for the set $\mathcal{B}_{\delta_1}^{(1)}$.

Given a sequence of positive real numbers ε_k with $\lim_{k \rightarrow \infty} \varepsilon_k = 0$, we may recursively define a sequence δ_k of positive real numbers, in such a way that

$$\mathcal{B}^{(k+1)} := \left\{ \omega \in \mathcal{B}^{(k)} \mid \begin{array}{l} \forall z, w \in \mathbb{B}_r \text{ with } \|z - w\| < \delta_{k+1} : \\ \sup_n \|f_\omega^n(z) - f_\omega^n(w)\| < \varepsilon_{k+1} \end{array} \right\}$$

has measure strictly greater than $1/2$.

Since the nested sets $\mathcal{B}^{(k)}$ all have measure at least $1/2$, the intersection, which we will simply denote as \mathcal{B} satisfies $\mu(\mathcal{B}) \geq 1/2$.

By ergodicity of the transformation T there exists, for almost every $\omega \in \Omega_{nor}$ a sequence n_k so that $T^{n_k}(\omega) \in \mathcal{B}$. By taking a subsequence if necessary, we find that $f_\omega^{n_k}$ converges locally uniform on U_ω to a function g .

Given $z \in U_\omega$ such that $\|g(z)\| < r$, it is clear that $g(w) = g(z)$ for every $w \in \mathbb{W}_\omega^s(z)$. On the other hand, given $w \in U_\omega$ which satisfies $g(w) = g(z)$,

we can find k_0 so that $f_\omega^{n_k}(z), f_\omega^{n_k}(w) \in \mathbb{B}_r$ for every $k \geq k_0$. Furthermore we may also find a sequence $k_0 \leq k_1 \leq k_2 \leq \dots$, so that whenever $k \geq k_i$ we have $\|f_\omega^{n_k}(z) - f_\omega^{n_k}(w)\| < \delta_i$.

Since $T^{n_k}\omega \in \mathcal{B}$ for every k , we conclude that, given $n \geq n_{k_i}$,

$$\|f_\omega^n(z) - f_\omega^n(w)\| = \left\| f_{T^{n_{k_i}}\omega}^{n-n_{k_i}}(f_\omega^{n_{k_i}}(z)) - f_{T^{n_{k_i}}\omega}^{n-n_{k_i}}(f_\omega^{n_{k_i}}(w)) \right\| < \varepsilon_i.$$

This shows that $w \in \mathbb{W}_\omega^s(z)$, which proves the desired equality. \square

When $m = 2$ and the measure is semi-neutral we obtain the following (local) version of the Stable Manifold theorem

Corollary 5.3 (Stable Manifold Theorem). *Let μ be a semi-neutral measure on $\mathcal{O}(\mathbb{C}^2, 0)$ with compact support, for which the origin lies in the random Fatou set. Then there exists, for almost every $\omega \in \Omega$, a constant $\rho > 0$ sufficiently small so that, given $z \in \mathbb{B}_\rho$, the set $\mathbb{W}_\omega^s(z)$ is locally a one-dimensional complex manifold.*

Proof. Let $\omega \in \Omega$ be such that the map g described in the previous lemma exists. Since $g(0) = 0$ we can choose $\rho > 0$ sufficiently small so that $\mathbb{B}_\rho \subset U_\omega$ and $g(\mathbb{B}_\rho) \subset \mathbb{B}_r$. Here the value of r is again determined by the previous lemma.

By Proposition 5.1 the function g is degenerate. Furthermore, by Proposition 2.10, we must have $\|dg(0)\| \geq 1$, thus we may further assume, by shrinking ρ is necessary, that g has rank 1 on every point of \mathbb{B}_ρ .

By the constant rank Theorem $g(\mathbb{B}_\rho)$ is a one-dimensional submanifold and every point in it is a regular value for the holomorphic map $g : \mathbb{B}_\rho \rightarrow g(\mathbb{B}_\rho)$. By the Implicit Function Theorem, given $z \in \mathbb{B}_\rho$, the level set $\{w \in \mathbb{B}_\rho \mid g(w) = g(z)\}$ is a one dimension complex manifold. The level sets of the map g coincide with the stable sets of the sequence ω . It follows that the latter are locally one-dimensional manifolds. \square

A. APPENDIX: HOLOMORPHIC GERMS AND THEIR TOPOLOGY

We write $\mathcal{O}(\mathbb{C}^m, 0)$ for the space of germs at 0 of holomorphic maps from \mathbb{C}^m to itself. Note that in the main text we required that the germs fix the origin, but this requirement plays no role here.

Our goal in this appendix is to define a topology on $\mathcal{O}(\mathbb{C}^m, 0)$ with the following property:

Local uniform convergence. *A sequence of germs (f_k) converges to a germ f if there exists $r > 0$ such that f and all the f_n admit a representative which is bounded and holomorphic on \mathbb{B}_r and*

$$\sup_{\|z\| \leq r} \|f_k(z) - f(z)\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

The construction of such a topology, given in Theorem A.6 below, resembles very closely the so-called inductive limit topology of a sequence of nested

Fréchet spaces $X_1 \subset X_2 \subset \dots \subset X_n \subset \dots$. See [Rud91] or [Gr09] for an example of this construction in the case of $C_0^\infty(\Omega)$. For a discussion of inductive limit topology for holomorphic germs, see for instance [M79], where germs in possibly infinite dimensional spaces are considered. While the results presented here are undoubtedly known to experts on the topic, we include them for the sake of completeness.

Suppose that $\varepsilon_n \rightarrow 0$ is a strictly decreasing sequence. Let X_n be the space of all bounded and holomorphic functions on $\mathbb{B}_{\varepsilon_n}$ equipped with the distance

$$d_n(f, g) := \sup_{\|z\| \leq \varepsilon_n} \|f(z) - g(z)\|.$$

There exists a natural injection of X_n into $\mathcal{O}(\mathbb{C}^m, 0)$, which we will denote by π_n . Furthermore it is clear that $X_n \subset X_{n+1}$.

We write τ_n for the standard metric topology on X_n . We note that τ_n is not equivalent to the compact-open topology. However, it will be clear that the inductive limit topology on $\mathcal{O}(\mathbb{C}^m, 0)$ obtained using either sequence of topologies is identical.

Inductive limit topology The *inductive limit topology*, which we will denote by τ_{ind} , is the finest topology on $\mathcal{O}(\mathbb{C}^m, 0)$ such that each injection $\pi_n : X_n \rightarrow \mathcal{O}(\mathbb{C}^m, 0)$ is a continuous function.

In other words a set U is open in the inductive limit topology if and only if, for each n , $X_n \cap \pi_n^{-1}(U)$ is open with respect to the topology τ_n .

Since each map π_n is injective, from now on we will avoid writing the map π_n and we will consider each X_n as a subset of $\mathcal{O}(\mathbb{C}^m, 0)$.

Remark In the classical construction of the inductive limit topology of a nested sequence of Fréchet spaces, $X_1 \subset X_2 \subset \dots$, it is assumed that the topology on X_n is induced by the topology on X_{n+1} , which fails in our setting. Consider for example the germs $f_n = n \left(\frac{z}{\varepsilon_1}\right)^n$. This sequence converges uniformly to 0 on \mathbb{B}_r for $r < \varepsilon_1$, in particular we have $d_2(f_n, 0) \rightarrow 0$. On the other hand we see that $d_1(f_n, 0) \rightarrow \infty$. This shows that τ_1 is different from $\tau_2|_{X_1}$.

Lemma A.1. *Let $\alpha \in \mathbb{N}^m$. The derivation $D_\alpha : f \mapsto \partial_\alpha f(0)$ is a continuous map from $\mathcal{O}(\mathbb{C}^m, 0)$, endowed with the inductive limit topology, to \mathbb{C} .*

Proof. If D_α was not continuous, there would exist $U \subset \mathbb{C}$ open and a natural number n such that $V := D_\alpha^{-1}(U) \cap X_n$ is not open with respect to the metric topology of X_n . If so, there exists $(f_k) \subset X_n \setminus V$ such that $f_k \rightarrow g \in V$ with respect to the metric d_n . By Weierstrass we have $D_\alpha f_k \rightarrow D_\alpha g$, which is not possible, since $f_k \notin V$. \square

Corollary A.2. *The inductive limit topology on $\mathcal{O}(\mathbb{C}^m, 0)$ is Hausdorff.*

Proof. Given two germs $f_1 \neq f_2$ then there exists $\alpha \in \mathbb{N}^m$ such that $D_\alpha f_1 \neq D_\alpha f_2$. The Hausdorff property of τ_{ind} follows from the continuity of D_α . \square

Recall that a subset U of a topological space X is *sequentially open* if each sequence (x_k) in X converging to a point of U is eventually in U . We say that X is a *sequential space* if every sequentially open subset of X is open.

Lemma A.3. $\mathcal{O}(\mathbb{C}^m, 0)$, endowed with the inductive limit topology, is a sequential space.

Proof. First of all we notice that if $f_k \rightarrow f$ with respect to the metric topology of X_n , then the convergence is valid in τ_{ind} . If this was not the case, we could find an open set $U \ni f$ in the inductive limit topology such that (f_k) is not eventually contained in U . However this is not possible since $U \cap X_n$ is open, with respect to the metric topology of X_n , and f_k converges to f in this topology.

As a consequence, given $U \subset \mathcal{O}(\mathbb{C}^m, 0)$ sequentially open, the set $U \cap X_n$ is sequentially open in the metric topology of X_n . Metric spaces are sequential, thus $U \cap X_n$ is open in X_n . By the definition of τ_{ind} it follows that U is open, which concludes the proof. \square

A proof of the following proposition can be found in [En89, pg. 209].

Proposition A.4. *Sequential compactness and countable compactness are equivalent in the class of sequential Hausdorff spaces.*

Given $r > 0$ and $f \in X_n$, we write $B_n(f, r)$ (respectively $\overline{B}_n(f, r)$) for the open (respectively closed) ball of radius r and center f with respect to the metric d_n .

Lemma A.5. *The closed ball $\overline{B}_n(f, r)$ is compact in (X_{n+1}, d_{n+1}) .*

Proof. It is sufficient to prove the lemma for the case $f = 0$. Suppose (f_k) is a sequence in $\overline{B}_n(0, r)$, then by the weak Montel Theorem there exists a subsequence k_m such that $f_{k_m} \rightarrow f_\infty$ uniformly on every compact subset $K \subset \mathbb{B}_{\varepsilon_n}$. In particular we have $d_{n+1}(f_{k_m}, f_\infty) \rightarrow 0$.

The function f_∞ is holomorphic on $\mathbb{B}_{\varepsilon_n}$. Furthermore, by uniform convergence on compact subsets of $\mathbb{B}_{\varepsilon_n}$, we have $\|f_\infty(z)\| \leq r$ for every $z \in B_{\varepsilon_n}(0)$. This proves that $f_\infty \in \overline{B}_n(0, r)$, which concludes the proof of the lemma. \square

Theorem A.6. *A sequence (f_k) is a convergent sequence with respect to τ_{ind} if and only if there exists N such that $(f_k) \subset X_N$ and (f_k) is convergent with respect to the metric d_N .*

Proof. We already proved the only part in Lemma A.3. Suppose now that (f_k) is a convergent sequence in the inductive limit topology. Write $f_\infty = \lim_{k \rightarrow \infty} f_k$ and let n_0 so that $f_\infty \in X_{n_0}$. Firstly we prove that there exists N_0 such that $(f_k) \subset X_{N_0}$. If this was not the case, by taking a subsequence if necessary, we may assume that for every n , $X_n \cap (f_k)$ is a finite set, and that the intersection is empty when $n \leq n_0$.

Let $U = \mathcal{O}(\mathbb{C}^m, 0) \setminus (f_k)$. The set $X_n \setminus U$ is finite for every $n \in \mathbb{N}$, therefore U is open in the inductive limit topology. It is clear that $f_\infty \in U$ but that the sequence (f_k) is not eventually contained in U , contradicting $f_\infty = \lim_{k \rightarrow \infty} f_k$. The existence of N_0 follows.

Suppose now that for every $N \geq N_0$ we have $d_N(f_k, f_\infty) \not\rightarrow 0$. If $\varepsilon > 0$ is fixed, then the sequence f_k is not eventually contained in $B_N(f_\infty, \varepsilon)$. This follows from the fact that $B_N(f_\infty, \varepsilon)$ are relatively compact in X_{N+1} , plus the fact that convergence in X_{N+1} implies convergence with respect to τ_{ind} .

Let $(k_n^N) \subset \mathbb{N}$ be the subsequence obtained by removing all the indices k for which $f_k \in B_N(f_\infty, \varepsilon)$. Since $B_N(f_\infty, \varepsilon) \subset B_{N+1}(f_\infty, \varepsilon)$ it follows that

$$(k_n^{N+1}) \subset (k_n^N) \subset \dots \subset (k_n^{N_0}).$$

Let $k_n := k_n^{n+N_0}$. Then, for every $n \geq N - N_0$, we have that $f_{k_n} \notin B_N(f_\infty, \varepsilon)$. Furthermore we notice that

$$d_N(f_{k_n}, f_\infty) \rightarrow \infty, \quad \forall N \geq N_0.$$

Suppose on the contrary that there exists $(k'_n) \subset (k_n)$ such that $d_N(f_{k'_n}, f_\infty) \rightarrow M$. By the previous lemma, by taking a subsequence of k'_n if necessary, we may assume that $f_{k'_n} \rightarrow g$ in X_{N+1} with $g \neq f_\infty$. Convergence in X_{N+1} implies convergence in τ_{ind} , therefore this is not possible.

The set $U = \mathcal{O}(\mathbb{C}^m, 0) \setminus (f_{k_n})$ contains $f_\infty \in U$. Since $(f_k) \subset X_{N_0}$, given $N \geq N_0$ we have that $U \cap X_N = X_N \setminus (f_{k_n})$. The sequence (f_{k_n}) is divergent in X_N , thus $U \cap X_N$ is open in X_N for $N \geq N_0$. Finally if V is open in X_{N+1} then $V \cap X_N$ is open in X_N . This proves that U is an open neighborhood of f_∞ in the inductive limit topology, which gives a contradiction. It follows that there exists N_1 such that $d_{N_1}(f_k, f_\infty) \rightarrow 0$. \square

Corollary A.7. *A set $K \subset \mathcal{O}(\mathbb{C}^m, 0)$ is sequentially compact in τ_{ind} if and only if there exists N such that $K \subset X_N$ and is compact with respect to the metric topology of X_N .*

Proof. Suppose that K is sequentially compact in τ_{ind} . First of all there exists N_0 such that $K \subset X_{N_0}$. If this was not the case, we could find a sequence $(f_k) \subset K$ such that $f_k \notin X_k$. By the previous theorem, such sequence does not have any convergent subsequence with respect to τ_{ind} , which contradicts sequential compactness.

Suppose that K is unbounded in X_N for every $k \geq N_0$. We can find $f_k \in K$ such that $d_k(f_k, 0) > N$. The previous theorem proves that this sequence does not have convergent subsequence, which contradicts sequential compactness. This proves that K is bounded in X_{N_1} for some $N_1 \geq N_0$. By Lemma A.5 the set K is relatively compact in X_{N_1+1} . Since the inductive limit topology is sequential, it follows that K is a closed set in τ_{ind} . This shows that K is a compact set in X_{N_1+1} .

On the other hand since convergence in X_N implies convergence in τ_{ind} , a compact set $K \subset X_N$ is sequentially compact in τ_{ind} . \square

Corollary A.8. *Compactness and sequential compactness are equivalent in $\mathcal{O}(\mathbb{C}^m, 0)$.*

Proof. Since $\mathcal{O}(\mathbb{C}^m, 0)$ is a Hausdorff sequential space, we only have to prove that every sequentially compact set is compact. By the previous corollary if K is sequentially compact then there exists N such that $K \subset X_N$ and K is compact in X_N . Let $\{U_\alpha\}$ be an open cover of K . By the definition of the inductive limit

topology it follows that $\{U_\alpha \cap X_N\}$ is an open cover of K in X_N . Using the compactness of K in X_N we can extract a finite open subcover $\{U_1, \dots, U_n\}$. This proves that K is compact with respect to the topology τ_{ind} . \square

The following corollary is an immediate consequence of the previous ones.

Corollary A.9. *A set $K \subset \mathcal{O}(\mathbb{C}^m, 0)$ is compact in τ_{ind} if and only if there exists N such that $K \subset X_N$ is compact with respect to the metric topology of X_N .*

We conclude this appendix by showing that the inductive limit topology is not metrizable. Notice that a similar proof also shows that this topology is not even first countable.

Proposition A.10. *The inductive limit topology is not metrizable.*

Proof. Suppose on the contrary that there exists a metric d_{ind} on $\mathcal{O}(\mathbb{C}^m, 0)$ such that τ_{ind} coincides with the metric topology of d_{ind} . We will write $B_{ind}(f, r)$ for the open ball of center f and radius r with respect of this metric.

We note that for every n we can construct a sequence $(f_k^n)_{k \in \mathbb{N}} \subset X_n \setminus X_{n-1}$, such that $\lim_{k \rightarrow \infty} d_n(f_k^n, 0) = 0$. Since convergence in X_n implies convergence in τ_{ind} , for every n we can find k_n such that

$$f_{k_n}^n \in B_{ind}(0, 1/n).$$

Now let $g_n = f_{k_n}^n$. It is clear that $d_{ind}(g_n, 0) \rightarrow 0$ as $n \rightarrow \infty$. By Theorem A.6, it follows that (g_n) is contained in some X_N , which is not possible by the definition of the f_k^n -s. This contradicts the fact that τ_{ind} is metrizable. \square

REFERENCES

- [A03] Marco Abate. “Discrete local holomorphic dynamics”. In: *Proceedings of 13th Seminar on Analysis and its Applications*. Isfahan Univ. Press, Isfahan, 2003, pp. 1–31.
- [A01] Marco Abate. “The residual index and the dynamics of holomorphic maps tangent to the identity”. In: *Duke Math. J.* 107.1 (Mar. 2001), pp. 173–207.
- [BM04] Filippo Bracci and Laura Molino. “The Dynamics near Quasi-Parabolic Fixed Points of Holomorphic Diffeomorphisms in \mathbb{C}^2 ”. In: *American Journal of Mathematics* 126.3 (2004), pp. 671–686.
- [BZ13] Filippo Bracci and Dmitri Zaitsev. “Dynamics of one-resonant bi-holomorphisms”. In: *J. Eur. Math. Soc. (JEMS)* 15.1 (2013), pp. 179–200.
- [Brj71] A. D. Brjuno. “Analytic form of differential equations. I, II”. In: *Trudy Moskov. Mat. Obšč.* 25 (1971), 131–288; *ibid.* 26 (1972), 199–239.
- [C38] H. Cremer. “Über die Häufigkeit der Nichtzentren.” In: *Mathematische Annalen* 115 (1938), pp. 573–580.

- [DK00] J.J. Duistermaat and Johan A.C. Kolk. *Lie Groups*. 1st ed. Springer-Verlag Berlin Heidelberg, 2000, p. 96.
- [Ec85] J. Écalle. “Les fonctions résurgentes, Tome III: L’équation du pont et la classification analytiques des objets locaux”. In: *Publ. Math. Orsay* 49 (1985).
- [En89] Ryszard Engelking. *General topology*. Second. Vol. 6. Sigma Series in Pure Mathematics. Heldermann Verlag, Berlin, 1989.
- [FLRT16] T. Firsova et al. “Hedgehogs for neutral dissipative germs of holomorphic diffeomorphisms of $(\mathbb{C}^2, 0)$ ”. In: *ArXiv e-prints* (Nov. 2016).
- [FS91] John Erik Fornæss and Nessim Sibony. “Random iterations of rational functions”. In: *Ergodic Theory and Dynamical Systems* 11.4 (1991), pp. 687–708.
- [FK60] H. Furstenberg and H. Kesten. “Products of Random Matrices”. In: *Ann. Math. Statist.* 31.2 (June 1960), pp. 457–469.
- [GM89] I Ya Gol’dsheid and G A Margulis. “Lyapunov indices of a product of random matrices”. In: *Russian Mathematical Surveys* 44.5 (1989), p. 11.
- [Gr09] Gerd Grubb. *Distribution and Operators*. First. Graduate Texts in Mathematics. Springer-Verlag New York, 2009, pp. 9–15.
- [GR85] Y. Guivarc’h and A. Raugi. “Frontière de Furstenberg, propriétés de contraction et théorèmes de convergence”. In: *Z. Wahrsch. Verw. Gebiete* 69.2 (1985), pp. 187–242.
- [H98] Monique Hakim. “Analytic transformations of $(\mathbb{C}^p, 0)$ tangent to the identity”. In: *Duke Math. J.* 92.2 (Apr. 1998), pp. 403–428.
- [La11] Samuel Lattès. “Sur les formes réduits des transformations ponctuelles à deux variables”. In: *Bulletin de la S.M.F.* 39 (1911), pp. 309–345.
- [Le84] F. Ledrappier. “Quelques propriétés des exposants caractéristiques”. In: *École d’été de probabilités de Saint-Flour, XII—1982*. Vol. 1097. Lecture Notes in Math. Springer, Berlin, 1984, pp. 305–396.
- [LRT16] M. Lyubich, R. Radu, and R. Tanase. “Hedgehogs in higher dimensions and their applications”. In: *ArXiv e-prints* (Nov. 2016).
- [M79] Jorge Mujica. “Spaces of germs of holomorphic functions”. In: *Studies in analysis*. Vol. 4. Adv. in Math. Suppl. Stud. Academic Press, New York-London, 1979, pp. 1–41.
- [O68] V. I. Oseledec. “A multiplicative ergodic theorem. Characteristic Ljapunov, exponents of dynamical systems”. In: *Trudy Moskov. Mat. Obšč.* 19 (1968), pp. 179–210.
- [P28] Henri Poincaré. *Oeuvres, Tome I*. Gauthier-Villars, Paris, 1928.
- [Ri01] Marzia Rivi. “Parabolic manifolds for semi-attractive holomorphic germs”. In: *Michigan Math. J.* 49.2 (Oct. 2001), pp. 211–241.
- [RR88] Jean-Pierre Rosay and Walter Rudin. “Holomorphic Maps from \mathbb{C}^n to \mathbb{C}^n ”. In: *Transactions of the American Mathematical Society* 310.1 (1988), pp. 47–86.

- [Rud91] Walter Rudin. *Functional analysis*. Second. International Series in Pure and Applied Mathematics. New York: McGraw-Hill Inc., 1991, pp. 149–156.
- [Rue79] David Ruelle. “Ergodic theory of differentiable dynamical systems”. In: *Inst. Hautes Études Sci. Publ. Math.* 50 (1979), pp. 27–58.
- [Si42] Carl Ludwig Siegel. “Iteration of analytic functions”. In: *Ann. of Math. (2)* 43 (1942), pp. 607–612.
- [St57] Shlomo Sternberg. “Local Contractions and a Theorem of Poincaré”. In: *American Journal of Mathematics* 79.4 (1957), pp. 809–824.
- [St58] Shlomo Sternberg. “On the Structure of Local Homeomorphisms of Euclidean n -Space, II”. In: *American Journal of Mathematics* 80.3 (1958), pp. 623–631.
- [Su11] Hiroki Sumi. “Random complex dynamics and semigroups of holomorphic maps”. In: *Proceedings of the London Mathematical Society* 102.1 (2011), pp. 50–112.
- [U86] Tetsuo Ueda. “Local structure of analytic transformations of two complex variables, I”. In: *J. Math. Kyoto Univ.* 26.2 (1986), pp. 233–261.
- [U91] Tetsuo Ueda. “Local structure of analytic transformations of two complex variables, II”. In: *J. Math. Kyoto Univ.* 31.3 (1991), pp. 695–711.
- [V14] Marcelo Viana. *Lectures on Lyapunov exponents*. Vol. 145. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2014, pp. xiv+202.
- [Y95] Jean-Christophe Yoccoz. “Théorème de Siegel, nombres de Bruno et polynômes quadratiques”. In: *Astérisque* 231 (1995), pp. 3–88.