

# Randomized subspace iteration: Analysis of canonical angles and unitarily invariant norms

Arvind K. Saibaba\*

November 13, 2018

## Abstract

This paper analyzes the randomized subspace iteration for the computation of low-rank approximations. We present three different kinds of bounds. First, we derive both bounds for the canonical angles between the exact and the approximate singular subspaces. Second, we derive bounds for the low-rank approximation in any unitarily invariant norm (including the Schatten- $p$  norm). This generalizes the bounds for Spectral and Frobenius norms found in the literature. Third, we present bounds for the accuracy of the singular values. The bounds are structural in that they are applicable to any starting guess, be it random or deterministic, that satisfies some minimal assumptions. Specialized bounds are provided when a Gaussian random matrix is used as the starting guess. Numerical experiments demonstrate the effectiveness of the proposed bounds.

## 1 Introduction

The computation of low-rank approximations of large-scale matrices is a vital step in many applications in data analysis and scientific computing. These applications include principal component analysis, facial recognition, spectral clustering, model reduction techniques such as proper orthogonal decomposition (POD) and discrete empirical interpolation method (DEIM), approximation algorithms for partial differential and integral equations. The celebrated Eckart-Young theorem [10] says that the optimal low-rank approximation can be obtained by means of the Singular Value Decomposition (SVD); however, computing the full or truncated SVD can be computationally challenging, or even prohibitively expensive for many applications of interest.

Randomized algorithms for computing low-rank approximations have become increasingly popular in the last two decades. For example, see the survey papers [12, 17]. Randomized methods have gained in popularity since they are easy to implement, computationally efficient, and numerically robust. Although randomized algorithms tend to have the same asymptotic cost compared to classical methods, they have several advantages that make them suitable for large-scale computing. Specifically, for datasets that are too large to fit in memory, randomized algorithms are able to exploit parallel computing efficiently and are efficient in the number of times they access the data. Randomized algorithms also have excellent numerical robustness and are very reliable in practical applications.

We focus on a specific randomized algorithm known as *randomized subspace iteration*. The main idea of this method is to use random sampling to identify a subspace that approximately captures the range of the matrix. A low-rank approximation to the matrix is then obtained by projecting the matrix onto this subspace. A post-processing step is then performed to compress the low-rank representation to achieve a desired target rank, and a conversion step to obtain an equivalent representation in the desired format (typically, a truncated SVD representation)—both these steps are deterministic.

---

\*Department of Mathematics, North Carolina State University asaibab@ncsu.edu. This work was funded, in part, by NSF DMS 1720398, OP: Collaborative Research: Novel Feature-Based, Randomized Methods for Large-Scale Inversion

Many advances have been made in the analysis of randomized algorithms for low-rank approximations. The analysis typically has two stages: a *structural, or deterministic stage*, in which minimal assumption about the distribution of the random matrix is made, and a *probabilistic stage*, in which the distribution of the random matrix is taken into account to derive bounds for expected and tail bounds of the error distribution. As mentioned earlier, existing literature only targets the error in the low-rank representation [11, 12]. When the low-rank representation is in the SVD format, it is desirable to understand the quality of the approximate subspaces and the individual singular triplets. This paper aims to fill in some of the missing gaps in the literature by a rigorous analysis of the accuracy of approximate singular values, vectors and subspaces obtained using randomized subspace iteration. This analysis will be beneficial in applications where an analysis beyond the low-rank approximation is desired. Examples include Model Reduction techniques [2, 9], Leverage Score computation [14], Spectral Clustering [6], FEAST eigensolvers [23], Canonical Correlation Analysis [1].

## 1.1 Contributions and overview of paper

We survey the contents and the main contributions of this paper.

**Canonical angles.** We have developed bounds for *all* the canonical angles between the spaces spanned by the exact and the approximate singular vectors. Several different flavors of bounds are provided:

1. The bounds in Section 3.1 relate the canonical angles between the exact and the approximate singular subspaces. Analysis is also provided for unitarily invariant norms of the canonical angles.
2. In applications where lower dimensional subspaces are extracted from the approximate singular subspaces, the bounds in Section 3.2 quantifies the accuracy in the extraction process.
3. Section 3.2 also presents bounds for the angles between the individual exact and approximate singular vectors, extracted from the appropriate subspaces.

Our bounds suggest that the accuracy of the singular values and vectors, in addition to the low-rank approximations, is high provided (1) singular values decay rapidly beyond the target rank  $k$ , and (2) the larger the singular value gaps, the higher is the accuracy to be expected. Furthermore, the truncation step to extract the  $k$  dimensional subspaces does not significantly lower the accuracy of the subspaces.

**Low-rank approximation.** This paper provides the first known analysis of the randomized subspace iteration for an arbitrary unitarily invariant norm, with stronger, specialized results for Schatten-p norms. Bounds for the special cases of the Schatten-p norm, namely the spectral and Frobenius norms, have already appeared in the literature—our result for the Schatten-p norm recovers these results as special cases.

**Singular values.** We derive upper and lower bounds on the approximate singular values obtained by the randomized subspace iteration. Similar bounds also appear in [11]; however, our proof technique is different. We also present Hoffman-Wielandt type bounds for the accuracy of the singular values.

The conclusion of the bounds for the low-rank approximations and the singular values are similar to those of the conclusions for the canonical angles.

**Generalization of sin theta theorem** The sin theta theorem [25] is a well known result in numerical analysis and relates the canonical angles between the true and approximate singular subspaces in the unitarily invariant norms. We derive a generalization of the sin theta theorem that derives bounds for the individual canonical angles between the two subspaces. The sin theta theorem is recovered as a special case. This result maybe of independent interest beyond the study of randomized algorithms.

## 2 Background and preliminaries

### 2.1 Notation

Denote the target rank by  $k$  and let  $1 \leq k \leq \text{rank}(A)$ . Let the matrix  $A \in \mathbb{C}^{m \times n}$ , have the SVD

$$A = [U_k \quad U_\perp] \begin{bmatrix} \Sigma_k & \\ & \Sigma_\perp \end{bmatrix} \begin{bmatrix} V_k^* \\ V_\perp^* \end{bmatrix}.$$

Here,  $\Sigma_k \in \mathbb{C}^{k \times k}$  and  $\Sigma_\perp \in \mathbb{C}^{(m-k) \times (n-k)}$ ; the columns of  $U_k$  and  $U_\perp$  are the corresponding left singular vectors, and columns of  $V_k$  and  $V_\perp$  are the corresponding right singular vectors. We also denote by  $A_k = U_k \Sigma_k V_k^*$  as the best rank- $k$  approximation to the matrix  $A$ , in any unitarily invariant norm (for a definition, see below). We also define  $A_\perp = U_\perp \Sigma_\perp V_\perp^*$  and observe that

$$A = A_k + A_\perp.$$

**Singular values and ratios** Let  $\|\cdot\|_2$  denote the spectral norm, so that  $\|\Sigma_\perp\|_2 = \sigma_{k+1}$  and  $\|\Sigma_k^{-1}\|_2 = \frac{1}{\sigma_k}$ . The singular values of  $A$  can be arranged in decreasing order as

$$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_k \geq \sigma_{k+1} \geq \dots \geq \sigma_n.$$

For later use, we define the singular value ratios

$$\gamma_j = \frac{\sigma_{k+1}}{\sigma_j} \quad j = 1, \dots, k. \quad (1)$$

Since the singular values are monotonically decreasing, the singular value ratios are monotonically increasing, i.e.,  $\gamma_1 \leq \dots \leq \gamma_k \leq 1$ .

**Norms** We have already defined the spectral norm. The Frobenius norm of a matrix is  $\|A\|_F = \sqrt{\text{trace}(A^*A)}$ . We use the symbol  $\|\cdot\|$  to denote any unitarily invariant norm, i.e., a norm that satisfies  $\|QAZ\| = \|A\|$  for unitary matrices  $Q, Z$ . An example of the unitarily invariant norms is Schatten- $p$  class of norms, defined as the vector  $\ell_p$  norm of the singular values of  $A$ , i.e.,

$$\|A\|_p = \left( \sum_{j=1}^{\min\{m,n\}} \sigma_j^p \right)^{1/p}.$$

With this definition, it can be readily seen that  $\|A\|_2 = \|A\|_\infty$  and  $\|A\|_F = \|A\|_2$ . Another example is the Ky-Fan- $k$  class of norms defined  $\|A\|_{(k)} = \sum_{j=1}^k \sigma_j$  for every  $k = 1, \dots, \min\{m, n\}$ . Associated with every unitarily invariant norm is a symmetric gauge function acting on the singular values of the matrix that it acts on.

**Projection matrices** Suppose the matrix  $Z$  has full column rank with column space  $\mathcal{R}(Z)$ ;  $Z^\dagger$  is a left multiplicative inverse and where  $^\dagger$  represents the Moore-Penrose inverse. We define the (orthogonal) projection matrix  $\mathcal{P}_Z = ZZ^\dagger$ . An orthogonal projection matrix is uniquely defined by its range, and  $\mathcal{R}(\mathcal{P}_Z) = \mathcal{R}(Z)$ . For a matrix  $Q$  with orthonormal columns, the formula simplifies and  $\mathcal{P}_Q = QQ^*$ .

**Canonical angles** The separation between subspaces can be measured by the principal or canonical angles. Let  $\mathcal{M}$  and  $\mathcal{N}$  be two subspaces of  $\mathbb{C}^n$ , such that  $\dim \mathcal{M} = \ell$ ,  $\dim \mathcal{N} = k$  and  $\ell \geq k$ . Then the principal angles between the subspaces  $\mathcal{M}$  and  $\mathcal{N}$  are recursively defined to be the numbers  $0 \leq \theta_i \leq \pi/2$  such that

$$\cos \theta_i = \max_{u \in \mathcal{M}, v \in \mathcal{N} \|u\|_2 = \|v\|_2 = 1} v^* u = v_i^* u_i, \quad i = 1, \dots, k$$

subject to the constraints  $\|u_i\|_2 = \|v_i\|_2 = 1$ , and

$$u_j^* u, \quad v_j^* v = 0, \quad j = 1, \dots, i-1.$$

The canonical angles are arranged in increasing order as

$$0 \leq \theta_1 \leq \dots \leq \theta_k \leq \pi/2.$$

It can also be shown that  $\sin \theta_i$  are also the singular values of  $\mathcal{P}_M - \mathcal{P}_N$ .

We denote  $\angle(\mathcal{M}, \mathcal{N})$  to be the canonical angles between subspaces  $\mathcal{M}$  and  $\mathcal{N}$ . Let  $M$  and  $N$  be matrices with orthonormal columns, which form bases for subspaces  $\mathcal{M}$  and  $\mathcal{N}$  respectively. Then, the singular values of singular values of  $(I - MM^*)N$  can be used to compute  $\sin \angle(\mathcal{M}, \mathcal{N})$  and the singular values of  $M^*N$  can be used to compute  $\cos \angle(\mathcal{M}, \mathcal{N})$  [5, Section 3]. For ease of notation, in the rest of this paper, we write  $\angle(M, N)$  instead of  $\angle(\mathcal{M}, \mathcal{N})$ .

## 2.2 Randomized subspace iteration

The basic version of the randomized subspace iteration is summarized in Algorithm 1. Given a starting guess, denoted by  $\Omega \in \mathbb{C}^{n \times (k+\rho)}$ , the algorithm performs  $q$  steps of the randomized subspace iteration to obtain the matrix  $Y$ , also known as the “sketch.” A thin-QR factorization of  $Y$  is performed to obtain  $Q$  whose columns form an orthonormal basis for the range of  $Y$ . The main idea is that, under suitable conditions, the range of  $Q$  is a good approximation for the range of  $A$ . We obtain a low-rank approximation to  $A$  by the projection  $\hat{A} = QQ^*A$ . The rest of the algorithm involves converting this low-rank approximation into the SVD format. The algorithm to compute

---

### Algorithm 1 Idealized version of Subspace iteration for Singular Value Decomposition

---

**Require:** Matrix  $A$ , Starting guess  $\Omega \in \mathbb{C}^{n \times (k+\rho)}$ , an integer  $q \geq 0$ .

- 1: Compute  $Y = (AA^*)^q A \Omega$
  - 2: Compute thin QR factorization of  $Y$ , so that  $Y = QR$ .
  - 3: Compute  $B = Q^*A$  and its SVD  $B = U_B \hat{\Sigma} \hat{V}^*$ .
  - 4: Compute  $\hat{U} = QU_B$ .
  - 5: **return** Matrices  $\hat{U}, \hat{\Sigma}, \hat{V}$  that define  $\hat{A} \equiv \hat{U} \hat{\Sigma} \hat{V}^*$ .
- 

an approximate singular value decomposition, given starting guess  $\Omega \in \mathbb{C}^{n \times (k+\rho)}$  is summarized in Algorithm 1. We say that this is an idealized version, since the algorithm can behave poorly in the presence of round-off errors. A practical implementation of this algorithm alternates the QR factorization with matrix-vector products (matvecs) involving  $A$ ; for more details regarding the implementation, the reader is referred to [12, 20]. In Algorithm 1, the output

$$\hat{A} \equiv QQ^*A = \hat{U} \hat{\Sigma} \hat{V}^*$$

may have a larger rank than (or equal to)  $k$ . If a rank- $k$  approximation to  $A$  is desired, then it can be obtained by discarding the  $\rho$  smallest singular values of  $\hat{A}$ . We denote this low-rank representation by

$$\hat{A}_k = \hat{U}_k \hat{\Sigma}_k \hat{V}_k^*.$$

This is summarized in Algorithm 2.

Before we state the assumptions needed for our analysis, we introduce the following notation. The matrix  $V^* \Omega$  captures the influence of the starting guess on the right singular matrix  $V$ . Partition this matrix as

$$V^* \Omega = \begin{bmatrix} V_k^* \Omega \\ V_{\perp}^* \Omega \end{bmatrix} = \begin{bmatrix} \Omega_1 \\ \Omega_2 \end{bmatrix}, \quad (2)$$

where  $\Omega_1 = V_k^* \Omega \in \mathbb{C}^{k \times (k+\rho)}$  and  $\Omega_2 = V_{\perp}^* \Omega \in \mathbb{C}^{(n-k) \times (k+\rho)}$ . As was mentioned earlier, we assume that the target rank  $k$  satisfies  $1 \leq k \leq \text{rank}(A)$ . Additionally, the following assumptions will be required for our analysis.

---

**Algorithm 2** Truncated SVD of  $\widehat{A} = QQ^*A$ 


---

**Require:** Matrix  $A \in \mathbb{C}^{m \times n}$  and  $Q \in \mathbb{C}^{m \times (k+\rho)}$ . Target rank  $1 \leq k \leq \text{rank}(A)$ .

- 1: Form matrix  $B = Q^*A$ .
  - 2: Compute the truncated SVD representation  $B_k = \widehat{U}_{B,k} \widehat{\Sigma}_k \widehat{V}_k^*$ .
  - 3: Form  $\widehat{U}_k = Q \widehat{U}_{B,k}$
  - 4: **return** Matrices  $\widehat{U}_k, \widehat{\Sigma}_k, \widehat{V}_k$  such that  $\widehat{A}_k = \widehat{U}_k \widehat{\Sigma}_k \widehat{V}_k^*$ .
- 

**Assumption 1.** Let  $\Omega_1 \in \mathbb{C}^{k \times (k+\rho)}$  be defined as above. We assume that

$$\text{rank}(\Omega_1) = k. \quad (3)$$

The singular value gap at index  $k$  is inversely proportional to the singular value ratio

$$\gamma_k = \|\Sigma_{\perp}\|_2 \|\Sigma_k^{-1}\|_2 = \frac{\sigma_{k+1}}{\sigma_k} < 1. \quad (4)$$

The first assumption guarantees that the starting guess  $\Omega$  has a significant influence over the right singular vectors, whereas the second assumption ensures that the  $k$  dimensional subspace  $\mathcal{R}(U_k)$  is well defined. In practice, it is highly desirable that  $\gamma_k \ll 1$ , which ensures that there is a large singular value gap.

### 3 Accuracy of singular vectors

We want to understand how well  $\mathcal{R}(\widehat{U})$  approximates  $\mathcal{R}(U_k)$ , measured in terms of the canonical angles between the subspaces. To this end, abbreviate the subspace angles between  $\widehat{U} \in \mathbb{C}^{m \times \ell}$  and  $U_k \in \mathbb{C}^{m \times k}$  as  $\theta_1, \dots, \theta_k$ . Similarly, denote the angles between  $\widehat{V} \in \mathbb{C}^{n \times \ell}$  and  $V_k \in \mathbb{C}^{n \times k}$  by  $\nu_1, \dots, \nu_k$ . We are also interested in obtaining bounds for the canonical angles  $\angle(U_k, \widehat{U}_k)$  and  $\angle(V_k, \widehat{V}_k)$ . To distinguish these angles from  $\angle(U_k, \widehat{U})$  and  $\angle(V_k, \widehat{V})$ , we call them  $\theta'_j$  and  $\nu'_j$  for  $j = 1, \dots, k$ .

#### 3.1 Bounds for canonical angles

Our first result derives bounds for the canonical angles  $\angle(U_k, \widehat{U}_k)$ . The analysis is based on the perturbation of projectors and the tools used here are similar to [12].

**Theorem 1.** Let  $\widehat{U}$  and  $\widehat{V}$  be obtained from Algorithm 1. With Assumption 1, the canonical angles  $\theta_j$  and  $\nu_j$  satisfy

$$\sin \theta_j \leq \frac{\gamma_j^{2q+1} \|\Omega_2 \Omega_1^\dagger\|_2}{\sqrt{1 + \gamma_j^{4q+2} \|\Omega_2 \Omega_1^\dagger\|_2^2}} \quad \sin \nu_j \leq \frac{\gamma_j^{2q+2} \|\Omega_2 \Omega_1^\dagger\|_2}{\sqrt{1 + \gamma_j^{4q+4} \|\Omega_2 \Omega_1^\dagger\|_2^2}}$$

for  $j = 1, \dots, k$ .

This theorem has several interesting features worth pointing out. First, if the matrix has exact rank  $k$ , then all of the canonical angles are uniformly equal to zero; that is, the randomized subspace iteration identifies the subspace exactly. On the other hand, when  $\gamma_k$  is very close to 1, the subspaces may not be well-defined and may be difficult to identify. In practice, it is highly desirable that  $\gamma_k \ll 1$ , so that the angles are captured accurately.

Second, the bounds for the canonical angles show explicit dependence on the singular value ratios  $\gamma_j$ . In particular, the canonical angles  $\theta_j$  and  $\nu_j$  converge to zero quadratically but at different rates depending on the singular value ratios. Specifically, the smaller the singular value ratio, smaller the canonical angles.

Third, the term  $\left\| \Omega_2 \Omega_1^\dagger \right\|_2$  can be written in terms of the right singular vector matrix  $V$  and the starting guess  $\Omega$  as

$$\left\| \Omega_2 \Omega_1^\dagger \right\|_2 = \left\| (V_\perp^* \Omega) (V_k^* \Omega)^\dagger \right\|_2.$$

When the columns of  $\Omega$  is linearly independent, this quantity is nothing but the tangent of the largest canonical angle between  $\mathcal{R}(V_k)$  and  $\mathcal{R}(\Omega)$ . This term appears frequently in randomized linear algebra and can be interpreted as a measure of the subspace overlap between the starting guess and the right singular vectors. In the ideal case,  $\Omega$  contains the singular vectors in  $V_k$ . A discussion of the meaning and interpretation of this term, is provided in [8, Section 2.5]. In particular, when  $\Omega$  is a Gaussian random matrix,  $\left\| \Omega_2 \Omega_1^\dagger \right\|_2$  is roughly on the order of  $\sqrt{(n-k)k}$ .

Fourth, the influence of  $\left\| \Omega_2 \Omega_1^\dagger \right\|_2$  is subdued by the singular value ratios  $\gamma_j^{2q+1}$ . With sufficiently large number of iterations  $q$ , the canonical angles are smaller than a user-defined tolerance. Rigorous bounds for the requisite number of iterations are provided in Section 3.4.

Lastly, the bounds for the canonical angles  $\theta_j$  are smaller than  $\nu_j$  because the latter contains an additional power of  $\gamma_j$ . The reason for this higher accuracy is as follows: the columns of  $\widehat{V}$  are the right singular vectors of  $Q^*A$ . Therefore, the multiplication step with  $Q$  amounts to an additional step of subspace iteration and gives the extra factor.

**Remark 1.** *Theorem 1 gives the sine of the canonical angles; these bounds can also be used to obtain upper bounds for the tangents and lower bounds for the cosines. With the same assumptions and notation as in Theorem 1, the relationship between the tangent and sine implies*

$$\tan \theta_j \leq \gamma_j^{2q+1} \left\| \Omega_2 \Omega_1^\dagger \right\|_2 \quad \tan \nu_j \leq \gamma_j^{2q+2} \left\| \Omega_2 \Omega_1^\dagger \right\|_2$$

for  $j = 1, \dots, k$ . Lower bounds for cosine of the canonical angles follow similarly.

**Unitarily invariant norms** The following result derives bounds for the canonical angles in any unitarily invariant norm, in contrast to Theorem 1 which bounds the individual canonical angles.

**Theorem 2.** *Let the approximate singular vectors  $\widehat{U}$  and  $\widehat{V}$  for a matrix  $A$  be computed according to Algorithm 1. Under Assumption 1, for every unitarily invariant norm,*

$$\begin{aligned} \left\| \sin \angle(U_k, \widehat{U}) \right\| &\leq \gamma_k^{2q} \frac{\left\| \Sigma_\perp \right\|}{\sigma_k} \left\| \Omega_2 \Omega_1^\dagger \right\|_2, \\ \left\| \sin \angle(V_k, \widehat{V}) \right\| &\leq \gamma_k^{2q+1} \frac{\left\| \Sigma_\perp \right\|}{\sigma_k} \left\| \Omega_2 \Omega_1^\dagger \right\|_2. \end{aligned} \tag{5}$$

The interpretation of this theorem is similar to that of Theorem 1. The connection between the two theorems follows from the identity  $\sin \theta_k = \left\| \sin \angle(U_k, \widehat{U}) \right\|_2$ . If we specialize the result in Theorem 2 to the spectral norm, then it is clear that this result weaker than the bound in Theorem 1.

### 3.2 Extraction of $k$ -dimensional subspaces

In the previous subsection, the columns of  $\widehat{U}$  and  $\widehat{V}$  spanned  $\ell = k + \rho$  dimensional subspaces. Many applications, however, require the extraction of  $k$  dimensional singular subspaces from the low-rank approximation  $\widehat{A} \equiv QQ^*A$ . One way to extract the appropriate subspaces is to first compute the optimal rank- $k$  truncation of  $\widehat{A}$ , denoted by  $\widehat{A}_k$ . The singular vectors of  $\widehat{A}_k$ , denoted by  $\widehat{U}_k$  and  $\widehat{V}_k$ , are then used instead of  $\widehat{U}$  and  $\widehat{V}$ . See Algorithm 2, for details regarding implementation. The bounds derived in the previous subsection are not directly applicable since [26, Corollary 10] says

$$\theta_j \leq \theta'_j \quad \nu_j \leq \nu'_j \quad j = 1, \dots, k.$$

To understand how much additional error is incurred during this extraction process, we present several results. The important conclusion of all these results is that the accuracy of the extracted

subspaces of dimension  $k$  is comparable to the accuracy of the  $k + \rho$  dimensional subspace provided the singular values are sufficiently well separated.

The approach we take is different from that of the previous section. The starting point of our analysis is the well-known sin theta theorem for singular subspaces [25]. Let  $A, \hat{A}$  be two matrices of conformal dimensions. Assuming that

$$\zeta \equiv \sigma_k(A) - \sigma_{k+1}(\hat{A}) > 0, \quad (6)$$

we have

$$\max \left\{ \left\| \left\| \sin \angle(U_k, \hat{U}_k) \right\| \right\|, \left\| \left\| \sin \angle(V_k, \hat{V}_k) \right\| \right\| \right\} \leq \frac{\max\{\|E_{12}\|, \|E_{21}\|\}}{\zeta}, \quad (7)$$

where the two matrices  $E_{12}$  and  $E_{21}$  are

$$\begin{aligned} E_{12} &= (I - \mathcal{P}_{\hat{U}_k})(A - \hat{A})\mathcal{P}_{V_k} \\ E_{21} &= \mathcal{P}_{U_k}(A - \hat{A})(I - \mathcal{P}_{\hat{V}_k}). \end{aligned} \quad (8)$$

However, this version of the sin theta theorem does not provide us with a way to obtain bounds for the individual canonical angles. To this end, we first present a new generalization of the sin theta theorem.

**Theorem 3.** *Let  $A \in \mathbb{C}^{m \times n}$  with  $\text{rank}(A) \geq k$  and let  $\hat{A}$  be the perturbed matrix with same dimensions. Suppose the singular value gap satisfies Equation (6). Let  $\hat{A}_k = \hat{U}_k \hat{\Sigma}_k \hat{V}_k^*$  be the truncated SVD of  $\hat{A}$ . Then*

$$\max\{\sin \theta'_j, \sin \nu'_j\} \leq \frac{\sigma_k(A)}{\sigma_j(A)} \max\{\sin \theta'_k, \sin \nu'_k\} \quad j = 1, \dots, k.$$

This theorem states that the sine of the canonical angles  $\sin \theta'_j$  are bounded by  $\sin \theta'_k$  up to a multiplicative factor, which is at most 1.

Our main result provides the following bounds for canonical angles between the exact and the approximate singular subspaces, when both the subspaces have the same dimension. The proof involves simplifying every term in Equation (8).

**Theorem 4.** *Let  $\hat{U}$  and  $\hat{V}$  be obtained from Algorithm 1, and matrices  $\hat{U}_k$  and  $\hat{V}_k$  from Algorithm 2. Under Assumption 1,*

- for every unitarily invariant norm

$$\max \left\{ \left\| \left\| \sin \angle(U_k, \hat{U}_k) \right\| \right\|, \left\| \left\| \sin \angle(V_k, \hat{V}_k) \right\| \right\| \right\} \leq \phi \frac{\gamma_k^{2q}}{1 - \gamma_k} \frac{\|\Sigma_{\perp}\|}{\sigma_k} \left\| \Omega_2 \Omega_1^{\dagger} \right\|_2.$$

The factor  $\phi$  takes different values depending on the specific norm used. For an arbitrary unitarily invariant norm, we have  $\phi = \sqrt{2}$ , whereas for the spectral and Frobenius norms, we have  $\phi = 1$ .

- canonical angles  $\theta'_j$  and  $\nu'_j$  satisfy

$$\max\{\sin \theta'_j, \sin \nu'_j\} \leq \gamma_j \frac{\gamma_k^{2q}}{1 - \gamma_k} \left\| \Omega_2 \Omega_1^{\dagger} \right\|_2 \quad j = 1, \dots, k.$$

The interpretation of this theorem is: (1) as the number of iterations  $q$  increase, the largest canonical angle converges to 0 quadratically, and (2) a larger singular value gap means that the subspace is computed more accurately. Comparing this result with Theorem 2, we see that the upper bound in Theorem 4 has additional factors which depend on the specific norm used. For an arbitrary unitarily invariant norm, there is an additional factor  $\max\{1, \sqrt{2}\gamma_k\}/(1 - \gamma_k)$ . For the spectral and Frobenius norms, the additional factor is  $1/(1 - \gamma_k)$ . Both factors are greater than 1, suggesting that the truncation process can introduce additional error. The additional factor is also independent of the number of iterations  $q$ , suggesting that it is a one-time price to be paid for the extraction process. The bound is devastating when  $\gamma_k \approx 1$ , but this also means that the subspaces may not be well-defined.

**Individual singular vectors** The previous results give insight into the accuracy measured using the canonical angles between the exact and approximate singular subspaces. When individual singular vectors need to be extracted, does the extraction process introduce additional error? The following result quantifies the accuracy of the extraction process.

**Theorem 5.** *Let the approximate singular vectors  $\widehat{U}$  and  $\widehat{V}$  be computed according to Algorithm 1. With Assumption 1, we have the following inequalities*

$$\sin \angle(u_j, \widehat{U}) \leq \gamma_j^{2q+1} \left\| \Omega_2 \Omega_1^\dagger \right\|_2 \quad \sin \angle(v_j, \widehat{V}) \leq \gamma_j^{2q+2} \left\| \Omega_2 \Omega_1^\dagger \right\|_2 \quad (9)$$

for  $j = 1, \dots, k$ . Denote the approximate singular triplets  $(\widehat{\sigma}_j, \widehat{u}_j, \widehat{v}_j)$  for  $j = 1, \dots, k$ . Under Assumption 1

$$\max \{ \sin \angle(u_j, \widehat{u}_j), \sin \angle(v_j, \widehat{v}_j) \} \leq \sqrt{1 + 2 \frac{\tilde{\gamma}^2}{\delta^2} \gamma_j^{2q+1} \left\| \Omega_2 \Omega_1^\dagger \right\|_2}. \quad (10)$$

Here,  $\tilde{\gamma}^2 \equiv \|\Sigma_\perp\|_2^2 + \left\| \Sigma_\perp \Omega_2 \Omega_1^\dagger \right\|_2^2$  and  $\tilde{\delta} \equiv \min\{\min_{\tilde{\sigma}_i \neq \tilde{\sigma}_j} \{|\sigma_j - \tilde{\sigma}_i|, \sigma_j\}\}$ .

The first result bounds the angles between the exact singular vector and the corresponding approximate singular subspaces. The second result compares the angles of the exact and the approximate singular vectors. This result also says that the extraction process does not adversely increase the error in the singular subspaces, provided the singular values are well-separated.

The convergence of the individual singular vectors tell a similar story to that of Theorem 1. The singular vectors corresponding to the largest singular values converge earlier than the singular vectors corresponding to the smaller singular vectors. This is a consequence of the fact that the singular value ratios are non-decreasing.

### 3.3 Comparison with other bounds

The subspace iteration dates to a 1957 paper by Bauer [3] for eigenvalue problems. The analysis of the subspace iteration has also been well-established, for example, we refer to [19, Chapter 14]. Randomized subspace iteration has attracted a lot of attention in the last two decades, with a special emphasis on quantifying the influence of the starting guess  $\Omega$ . In particular, recent research has focused on the choice of the distribution and the effect of the oversampling parameter  $\rho$ . The effect of randomized subspace iteration on the accuracy of singular vectors was studied in the context of spectral clustering in [6]. However, the authors made the rather strong assumption that  $\Omega \in \mathbb{R}^{n \times k}$ , which amounts to setting the oversampling parameter  $\rho = 0$ . This is a strong requirement since Assumption 1 now requires  $\Omega_1$  to be invertible. The authors were able to show (in our notation)

$$\left\| \sin \angle(U_k, \widehat{U}_k) \right\|_2 \leq \frac{\gamma_k^{2q+1} \left\| \Omega_2 \Omega_1^{-1} \right\|_2}{\sqrt{1 + \gamma_k^{4q+2} \left\| \Omega_2 \Omega_1^{-1} \right\|_2^2}}.$$

Notice that this bound coincides with Theorem 1 (for  $\sin \theta_k$ ) when  $\rho = 0$ . Our results provide bounds for the right singular vectors as well as all the canonical angles.

Let us return to this assumption that  $\text{rank}(\Omega_1)$ . When  $\Omega$  is standard Gaussian matrix, [21, Theorem 3.3] says

$$\left\| \Omega_1^{-1} \right\|_2 \leq \frac{2.35\sqrt{k}}{\delta}$$

with probability at least  $1 - \delta$ . For a small probability of failure  $0 < \delta < 1$ , this bound can be devastating. By contrast, if we let  $\Omega_1 \in \mathbb{C}^{k \times (k+\rho)}$  with  $\rho \geq 2$ , and still suppose that  $\Omega$  is a Gaussian random matrix. Then, with probability at least  $1 - \delta$  [12, Proposition 10.4] says

$$\left\| \Omega_1^\dagger \right\|_2 \leq e \frac{\sqrt{k+\rho}}{\rho} \left( \frac{1}{\delta} \right)^{1/(\rho+1)}.$$



It is clear that when the random matrix is Gaussian, oversampling has an impact on the accuracy of the randomized subspace iteration. Specifically, larger the oversampling, the more accurate is the subspace.

Oversampling plays a bigger role for random matrices that have different distributions than Gaussian. When  $\Omega$  is generated from the subsampled randomized Hadamard transform (SRHT), or Rademacher distributions, a more aggressive form of oversampling  $\ell \sim k \log k$  is necessary to ensure that  $\text{rank}(\Omega_1) = k$ . Therefore, by allowing for oversampling, our bounds are applicable to starting guesses that are not restricted to Gaussian random matrices. Not only that, our bounds are also informative for matrices with decaying singular values and significant singular value gap.

A recent paper by Nakatsukasa [18] considered the issue of accuracy of extracting singular subspaces for general projection-based approximation methods. In our notation, these refer to relating bounds for  $\angle(U_k, \hat{U})$  to  $\angle(U_k, \tilde{U}_k)$ . Our bounds for the canonical angles appear to be tighter than the result implied by [18, Corollary 1]. This may be because the analysis was applicable to arbitrary subspace projections, whereas ours is specialized to randomized subspace iteration; we do not go into a detailed comparison here. Furthermore, our analysis is able to bound the individual canonical angles which is missing in [18].

### 3.4 Probabilistic bounds

Thus far, we have not made specific assumptions on the matrix  $\Omega$ , as long as it satisfies  $\text{rank}(\Omega_1) = k$ . In particular,  $\Omega$  need not be even be random, and may be deterministic. However, more can be said about the bounds when  $\Omega$  is random is drawn from a specific distribution.

In many applications, the matrix  $\Omega \in \mathbb{R}^{n \times (k+\rho)}$  is taken to be the standard Gaussian random matrix. That is, the entries of  $\Omega$  are i.i.d.  $\mathcal{N}(0, 1)$  random variables. Here we derive a few probabilistic results that provide insight into the accuracy of the subspaces. Let  $\rho \geq 2$  and define the constant

$$C_e = \sqrt{\frac{k}{\rho-1}} + \frac{e\sqrt{(k+\rho)(n-k)}}{\rho} \quad (11)$$

and for  $0 < \delta < 1$  define the constant

$$C_d = \frac{e\sqrt{k+\rho}}{\rho+1} \left(\frac{2}{\delta}\right)^{1/(\rho+1)} \left(\sqrt{n-k} + \sqrt{k+\rho} + \sqrt{2 \log \frac{2}{\delta}}\right). \quad (12)$$

**Theorem 6** (Probabilistic bounds). *Let  $\Omega \in \mathbb{R}^{n \times (k+\rho)}$  be a standard Gaussian random matrix with  $\rho \geq 2$ . Assume that the singular value ratio  $\gamma_k < 1$ . Let  $\hat{U}$  and  $\hat{V}$  be obtained from Algorithm 1. For  $j = 1, \dots, k$ , the expected value of the canonical angles satisfy*

$$\mathbb{E} [\sin \theta_j] \leq \frac{\gamma_j^{2q+1} C_e}{\sqrt{1 + \gamma_j^{4q+2} C_e^2}} \quad \mathbb{E} [\sin \nu_j] \leq \frac{\gamma_j^{2q+2} C_e}{\sqrt{1 + \gamma_j^{4q+4} C_e^2}}.$$

Let  $0 < \delta < 1$  be a user defined failure tolerance. With probability, at least  $1 - \delta$ , the following inequalities hold independently for  $j = 1, \dots, k$

$$\sin \theta_j \leq \frac{\gamma_{j+1}^{2q+1} C_d}{\sqrt{1 + \gamma_j^{4q+2} C_d^2}} \quad \sin \nu_j \leq \frac{\gamma_j^{2q+2} C_d}{\sqrt{1 + \gamma_j^{4q+4} C_d^2}}.$$

The main message of theorem can be seen from the following bound on the number of subspace iterations  $q$ . Specifically, suppose  $0 < \epsilon < 1$ , and the number of subspace iterations  $q$  we take satisfies

$$q \geq \frac{1}{2} \left( \frac{\log \epsilon / C_e}{\log \gamma_k} - 1 \right),$$

then  $\mathbb{E} \sin \theta_j \leq \mathcal{O}(\epsilon^2)$  for  $j = 1, \dots, k$ .

Several extensions of these results are possible. First, following the proof technique of Theorem 6, we can extend the probabilistic analysis to Theorems 2 and 5 as well. Second, following the strategy in [12], the probabilistic results can be extended to other distributions. However, we will not pursue these extensions here.

## 4 Low-rank approximation and Singular values

In this section, we provide several structural bounds for the accuracy of the low-rank approximation and the accuracy of the singular values.

### 4.1 Low-rank approximation

Several results are available for estimating the error in the low-rank approximation  $A \approx QQ^*A$  in the spectral and Frobenius norms, when the matrix  $Q$  is obtained from the randomized subspace iteration [11, 12, 27]. As was mentioned earlier, the spectral and Frobenius norms are special cases of the Schatten-p norm, which are examples of unitarily invariant norms.

Here we present the first known analysis of randomized subspace iteration in a unitarily invariant norm.

**Theorem 7.** *Let  $\hat{A} \in \mathbb{C}^{m \times n}$  be computed using Algorithm 1. Under Assumption 1, the following inequalities hold in every unitarily invariant norm*

$$\| (I - QQ^*)A \| \leq \| \Sigma_{\perp} \| + \gamma_k^{2q} \| \Sigma_{\perp} \Omega_2 \Omega_1^{\dagger} \| \quad (13)$$

$$\| (I - QQ^*)A_k \| \leq \gamma_k^{2q} \| \Sigma_{\perp} \Omega_2 \Omega_1^{\dagger} \|. \quad (14)$$

Let  $B = Q^*A$ , and let  $B_k$  be its best rank- $k$  approximation. If  $A$  is approximated using  $QB_k$ , then the error in the low-rank approximation is

$$\| A - QB_k \| \leq \left( 1 + \frac{\sigma_1}{\sigma_k} \frac{\phi \gamma_k^{2q}}{1 - \gamma_k} \| \Omega_2 \Omega_1^{\dagger} \|_2 \right) \| \Sigma_{\perp} \|. \quad (15)$$

As in Theorem 4,  $\phi = 1$  for spectral and Frobenius norms, and  $\sqrt{2}$  for an arbitrary unitarily invariant norm.

In this theorem, as the number of iterations  $q \rightarrow \infty$ , the error in the low-rank approximation goes to zero.

We present a variant of the error in the low-rank approximation for the special case that a Schatten-p norm is used. The proof for the special case of the Frobenius norm was provided in [27].

**Theorem 8.** *Let  $\hat{A}$  be computed using Algorithm 1. Under Assumption 1, we have*

$$\| (I - QQ^*)A \|_p^2 \leq \| \Sigma_{\perp} \|_p^2 + \gamma_k^{4q} \| \Sigma_{\perp} \Omega_2 \Omega_1^{\dagger} \|_p^2. \quad (16)$$

The error bound in Theorem 7 is weaker than Theorem 8 for the Schatten-p norm since for  $\alpha, \beta \geq 0$ , we have  $\sqrt{\alpha^2 + \beta^2} \leq \alpha + \beta$ . More generally, Theorem 8 is applicable to any unitarily invariant norm that is also a Q-norm [4, Definition IV.2.9]. A unitarily invariant norm  $\| \cdot \|_Q$  is a Q-norm, if there exists another unitarily invariant norm  $\| \cdot \|_a$  such that  $\| A \|_Q^2 = \| A^* A \|_a$ . Note that the Schatten-p norms satisfy this property for  $p \geq 2$ , since  $\| A \|_p^2 = \| A^* A \|_{p/2}$ .

### 4.2 Accuracy of singular values

How are the singular values of  $A$  related to the singular values of  $\hat{A}$ ? We now present a result that quantifies the accuracy of the individual singular values. This result is similar to [11, Theorem 4.3]. Our proof techniques are substantially different. We make extensive use of the Cauchy interlacing theorem and the multiplicative singular value inequalities Equation (20).

**Theorem 9.** Let  $\widehat{A} = \widehat{U}\widehat{\Sigma}\widehat{V}^*$  be computed using Algorithm 1. Under Assumption 1, the approximate singular values  $\sigma_j(\widehat{A})$  satisfy for  $j = 1, \dots, k$

$$\sigma_j(A) \geq \sigma_j(\widehat{A}) \geq \frac{\sigma_j(A)}{\sqrt{1 + \gamma_j^{4q+2} \|\Omega_2 \Omega_1^\dagger\|_2^2}}.$$

It can be readily seen that the large singular values are computed more accurately since the singular value ratio corresponding to larger singular values is smaller.

Rather than quantify the accuracy of the individual singular values, the next results are of the Hoffman-Wielandt type and account for all the singular values together. Define the two matrices of conformal sizes

$$\Sigma = \begin{bmatrix} \Sigma_k & \\ & \Sigma_\perp \end{bmatrix} \quad \Sigma' = \begin{bmatrix} \widehat{\Sigma} & \\ & 0 \end{bmatrix}.$$

Under Assumption 1, the error in the singular values satisfies

$$\|\|\Sigma - \Sigma'\|\| \leq \|\|\Sigma_\perp\|\| + \gamma_k^{2q} \|\|\Sigma_\perp \Omega_2 \Omega_1^\dagger\|\|. \quad (17)$$

The proof combines [4, III.6.13] with Theorem 7. For the Schatten- $p$  norm, with  $p \geq 2$ , we can derive the bound

$$\|\|\Sigma - \Sigma'\|_p \leq \sqrt{\|\|\Sigma_\perp\|_p^2 + \gamma_k^{4q} \|\|\Sigma_\perp \Omega_2 \Omega_1^\dagger\|_p^2}. \quad (18)$$

The proof is similar, and is therefore omitted.

## 5 Proofs

We recall some results here that will be useful in our analysis, see [15, Section 7.7] for proofs. Let  $M, N$  be Hermitian positive definite. The notation  $M \preceq N$  means  $N - M$  is positive semi-definite and it defines a partial ordering on the set of Hermitian matrices. Clearly, this also implies  $I - N \preceq I - M$ . The partial order is preserved under the conjugation rule. That is

$$SMS^* \preceq SNS^* \quad \forall S \in \mathbb{C}^{m \times n}.$$

Weyl's theorem implies that the eigenvalues satisfy  $\lambda_j(M) \leq \lambda_j(N)$  for all  $j = 1, \dots, n$ . If additionally,  $M, N$  are both positive semidefinite then  $M^{1/2} \preceq N^{1/2}$  [4, Proposition V.1.8] and  $(I + N)^{-1} \preceq (I + M)^{-1}$ .

**Singular value inequalities** Let  $A, B \in \mathbb{C}^{m \times n}$ . For all  $i, j$  such that  $1 \leq i, j \leq \min\{m, n\}$  and  $i + j - 1 \leq \min\{m, n\}$ , the following singular value inequalities hold for the sum  $A + B$  [15, Equation 7.3.13]

$$\sigma_{i+j-1}(A + B) \leq \sigma_i(A) + \sigma_j(B), \quad (19)$$

and product  $AB^*$  [15, Equation (7.3.14)]

$$\sigma_{i+j-1}(AB^*) \leq \sigma_i(A)\sigma_j(B). \quad (20)$$

A useful corollary of these results is that  $\sigma_i(A + B) \leq \sigma_i(A) + \sigma_1(B)$  and  $\sigma_i(AB^*) \leq \sigma_i(A)\sigma_1(B)$  for  $i = 1, \dots, \min\{m, n\}$ .

**Unitarily invariant norms** It is useful to recall some properties of the unitarily invariant norms. Every unitarily invariant norm  $\|\|\cdot\|\|$  on  $\mathbb{C}^n$  is associated with a symmetric gauge function on  $\mathbb{R}^n$ . The  $\|\|\cdot\|\|$  satisfies  $\|\|M\|\| = \|\|(M^*M)^{1/2}\|\|$ , since both matrices have the same nonzero singular values. The following inequality for unitarily invariant norms, also known as strong sub-multiplicativity, will be useful [4, (IV.40)]

$$\|\|ABC\|\| \leq \|A\|_2 \|C\|_2 \|\|B\|\|.$$

We will need the following lemma

**Lemma 1.** Let  $A, B, D \in \mathbb{C}^{n \times n}$  such that  $A, B$  Hermitian and  $0 \preceq A \preceq B$ , then

$$\left\| \left\| (D^*AD)^{1/2} \right\| \right\| \leq \left\| \left\| (D^*BD)^{1/2} \right\| \right\|.$$

*Proof.* Combining the properties of the partial ordering, the eigenvalues of the scaled matrices satisfy  $\lambda_j(D^*AD)^{1/2} \leq \lambda_j(D^*BD)^{1/2}$  for all  $j = 1, \dots, n$ . Since the matrices are positive semidefinite, the eigenvalues are the singular values and  $\|(D^*AD)^{1/2}\|_{(k)} \leq \|(D^*BD)^{1/2}\|_{(k)}$  for every Ky-Fan- $k$  norm  $k = 1, \dots, n$ . By the Fan dominance theorem [4, Theorem IV.2.2], the advertised inequality is true for every unitarily invariant norm.  $\square$

## 5.1 Proofs of Section 3.1 Theorems

*Theorem 1.* We tackle each case separately.

**Bounds for  $\sin \theta_j$ :** The proof is lengthy and proceeds in four steps. We give a great level of detail here, since the proof technique will be applicable to the subsequent proofs.

**1. Converting an SVD to an EVD** We compute the thin SVD of  $(I - \mathcal{P}_{\widehat{U}})U_k = K S U G^*$ . The matrix

$$S_U = \text{diag}(\sin \theta_k, \dots, \sin \theta_1) \in \mathbb{R}^{k \times k}$$

contains the sine of the canonical angles between the subspaces spanned by the columns of  $\widehat{U}$  and  $U_k$  [5, Equation (13)]. It is readily seen that

$$G S_U^2 G^* = U_k^* (I - \mathcal{P}_{\widehat{U}}) U_k. \quad (21)$$

**2. Shrinking space** In Algorithm 1, we had defined  $Y = (AA^*)^q A \Omega$ . It follows that

$$U^* Y = \begin{bmatrix} \Sigma_k^{2q+1} & \\ & (\Sigma_{\perp} \Sigma_{\perp}^{\top})^q \Sigma_{\perp} \end{bmatrix} (V^* \Omega) = \begin{bmatrix} \Sigma_k^{2q+1} \Omega_1 \\ (\Sigma_{\perp} \Sigma_{\perp}^{\top})^q \Sigma_{\perp} \Omega_2 \end{bmatrix},$$

where from Equation (2),  $\Omega_1 = V_k^* \Omega$  and  $\Omega_2 = V_{\perp}^* \Omega$ . Next, by Assumption 1,  $\Omega_1$  has full row rank and therefore it has a right multiplicative inverse. Define

$$Z \equiv U^* Y \Omega_1^{\dagger} \Sigma_k^{-(2q+1)} = \begin{bmatrix} I \\ F \end{bmatrix} \quad F \equiv (\Sigma_{\perp} \Sigma_{\perp}^{\top})^q \Sigma_{\perp} \Omega_2 \Omega_1^{\dagger} \Sigma_k^{-(2q+1)}.$$

Recall that  $Y = QR$  is the thin-QR factorization of  $Y$ . Let  $Q_1 R_1$  be the thin-QR factorization of  $R \Omega_1^{\dagger} \Sigma_k^{-(2q+1)}$ ; here,  $Q_1 \in \mathbb{C}^{(k+\rho) \times k}$ ,  $R_1 \in \mathbb{C}^{k \times k}$ .

From  $Q_1 Q_1^* \preceq I$ , the conjugation rule implies

$$\mathcal{P}_Z = U^* Q Q_1 Q_1^* Q^* U \preceq U^* Q Q^* U = \mathcal{P}_{U^* Q}.$$

Since  $\mathcal{R}(U^* Y) = \mathcal{R}(U^* Q) = \mathcal{R}(U^* \widehat{U})$ , they have the same projectors, so

$$\mathcal{P}_Z \preceq \mathcal{P}_{U^* \widehat{U}} \quad I - \mathcal{P}_{U^* \widehat{U}} \preceq I - \mathcal{P}_Z. \quad (22)$$

Plug in  $U U^* = I$  into (21), and use (22) to obtain

$$U_k^* (I - \mathcal{P}_{\widehat{U}}) U_k = U_k^* U (I - \mathcal{P}_{U^* \widehat{U}}) U^* U_k \preceq \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} (I - \mathcal{P}_Z) \begin{bmatrix} I \\ 0 \end{bmatrix}.$$

**3. Simplifying  $\mathcal{P}_Z$**  Since  $\mathcal{P}_Z = ZZ^\dagger$ , we have

$$\mathcal{P}_Z = \begin{bmatrix} I \\ F \end{bmatrix} (I + F^*F)^{-1} \begin{bmatrix} I & F^* \end{bmatrix},$$

from which, it can be readily seen that

$$\begin{aligned} \begin{bmatrix} I & 0 \end{bmatrix} (I - \mathcal{P}_Z) \begin{bmatrix} I \\ 0 \end{bmatrix} &= I - (I + F^*F)^{-1} \\ &= F^*F(I + F^*F)^{-1} \equiv H. \end{aligned} \quad (23)$$

Note that  $H$  is positive semidefinite. To summarize the story so far,  $GS_U^2G^* \preceq H$ .

**4. Applying singular value inequalities** A straightforward SVD argument shows that the  $j$ -th singular value of  $H$  satisfies

$$\sigma_j(H) = \sigma_j^2(F)/(1 + \sigma_j^2(F)) \quad j = 1, \dots, k.$$

The singular value inequalities Equation (20) imply

$$\sigma_j(F) \leq \sigma_1(\Sigma_\perp \Sigma_\perp^\top)^q \Sigma_\perp \Omega_2 \Omega_1^\dagger \sigma_j(\Sigma_k^{-2q-1}) \leq \left( \frac{\sigma_{k+1}}{\sigma_{k-j+1}} \right)^{2q+1} \left\| \Omega_2 \Omega_1^\dagger \right\|_2.$$

Plugging this inequality into  $\sigma_j(H)$

$$\sigma_j^2(H) \leq \frac{\gamma_{k-j+1}^{4q+2} \left\| \Omega_2 \Omega_1^\dagger \right\|_2^2}{1 + \gamma_{k-j+1}^{4q+2} \left\| \Omega_2 \Omega_1^\dagger \right\|_2^2} \quad j = 1, \dots, k.$$

Since  $GS_U^2G^* \preceq H$ , Weyl's theorem implies  $\sin^2 \theta_{k-j+1} \leq \sigma_j^2(H)$ . Take square roots on both sides and rename  $j \leftarrow k - j + 1$  to get the desired result.

**Bounds for  $\sin \nu_j$ :** Let  $GS_V^2G^*$  be the eigenvalue decomposition of  $V_k^*(I - \mathcal{P}_{\widehat{V}})V_k$ . Note that the diagonals of  $S_V$  are the sine of the canonical angles  $\angle(V_k, \widehat{V})$ . Since  $\widehat{V}$  is obtained from the thin SVD of  $A^*Q$ ,  $\mathcal{R}(A^*Q) = \mathcal{R}(\widehat{V})$  and  $\mathcal{P}_{\widehat{V}} = \mathcal{P}_{A^*Q}$ , since an orthogonal projection matrix is uniquely determined by the range. Next, consider  $\widehat{Z}$  defined as

$$\widehat{Z} \equiv \Sigma^\top U^* Y \Omega_1^\dagger \Sigma_k^{-2q-2} = \begin{bmatrix} I \\ \widehat{F} \end{bmatrix} \quad \widehat{F} \equiv (\Sigma_\perp^\top \Sigma_\perp)^{q+1} \Omega_2 \Omega_1^\dagger \Sigma_k^{-2q-2}, \quad (24)$$

from  $(AV)^*Q = \Sigma^*U^*Q$ , it can be verified that

$$\mathcal{R}(\widehat{Z}) \subset \mathcal{R}(\Sigma^\top U^* Y) = \mathcal{R}(\Sigma^\top U^* Q) = \mathcal{R}((AV)^*Q).$$

Using an argument similar to Equation (22), we obtain

$$V_k^* V (I - \mathcal{P}_{\widehat{V}}) V^* V_k \preceq V_k^* V (I - \mathcal{P}_{\widehat{Z}}) V^* V_k = \begin{bmatrix} I & 0 \end{bmatrix} (I - \mathcal{P}_{\widehat{Z}}) \begin{bmatrix} I \\ 0 \end{bmatrix}.$$

The right hand side simplifies to  $I - (I + \widehat{F}^* \widehat{F})^{-1}$ . The rest of the proof is similar to that of the proof for  $\sin \theta_j$ .  $\square$

*Theorem 2.* With the notation of Theorem 1, we follow steps 1-3 of the proof to obtain

$$GS_U^2G^* \preceq H \preceq F^*F.$$

Since the square root preserves partial ordering, implies  $GSUG^* \preceq (F^*F)^{1/2}$ . Note that  $(F^*F)^{1/2}$  and  $F$  have the same nonzero singular values. Therefore,

$$\left\| \left\| \sin \angle(U_k, \widehat{U}) \right\| \right\| \leq \left\| \left\| (F^*F)^{1/2} \right\| \right\| = \left\| \left\| F \right\| \right\|.$$

By using strong sub-multiplicativity of the unitarily invariant norm, we have

$$\left\| \left\| \sin \angle(U_k, \widehat{U}) \right\| \right\| \leq \gamma_k^{2q} \left\| \left\| \Omega_2 \Omega_1^\dagger \right\|_2 \right\| \frac{\left\| \left\| \Sigma_\perp \right\| \right\|}{\sigma_k}.$$

□

## 5.2 Proofs of Section 3.2 Theorems

*Theorem 3.* Let  $X = (I - \mathcal{P}_{\widehat{U}_k})\mathcal{P}_{U_k}$  and  $Y = (I - \mathcal{P}_{\widehat{V}_k})\mathcal{P}_{V_k}$ . In decreasing order, the singular values of  $X$  and  $Y$  are  $\{\sin \theta'_j\}_{j=1}^k$  and  $\{\sin \nu'_j\}_{j=1}^k$  respectively. Let  $B \equiv \widehat{A} - \widehat{A}_k$ . First, we observe that

$$\begin{aligned} E_{12} &= (I - \mathcal{P}_{\widehat{U}_k})(A - \widehat{A})\mathcal{P}_{V_k} \\ &= (I - \mathcal{P}_{\widehat{U}_k})A_k - (I - \mathcal{P}_{\widehat{U}_k})\widehat{A}\mathcal{P}_{V_k} \\ &= (I - \mathcal{P}_{\widehat{U}_k})\mathcal{P}_{U_k}A_k - (\widehat{A} - \widehat{A}_k)\mathcal{P}_{V_k} \\ &= XA_k - B(I - \mathcal{P}_{\widehat{V}_k})\mathcal{P}_{V_k} = XA_k - BY. \end{aligned}$$

A similar calculation shows that  $E_{21} = X^*B - A_kY^*$ . From the first relation, since  $\text{rank}(A) \geq k$ , we have

$$XA_kA_k^\dagger = (E_{12} + BY)A_k^\dagger.$$

But  $A_kA_k^\dagger = \mathcal{P}_{U_k}$  and  $X\mathcal{P}_{U_k} = X$ . Applying Equation (20), we have

$$\sigma_j(X) \leq (\|E_{12}\|_2 + \|B\|_2\|Y\|_2)/\sigma_{k-j+1}(A) \quad j = 1, \dots, k.$$

A similar argument gives

$$\sigma_j(Y) \leq (\|E_{21}\|_2 + \|B\|_2\|X\|_2)/\sigma_{k-j+1}(A) \quad j = 1, \dots, k.$$

Combining these relations

$$\max\{\sigma_j(X), \sigma_j(Y)\} \leq \frac{\max\{\|E_{21}\|_2, \|E_{12}\|_2\}}{\sigma_{k-j+1}(A)} + \frac{\|B\|_2}{\sigma_{k-j+1}(A)} \max\{\|X\|_2, \|Y\|_2\}.$$

Recognize that  $\|B\|_2 = \sigma_{k+1}(\widehat{A})$ . Applying Equation (7) in the spectral norm simplifies the expression since

$$\frac{1}{\sigma_{k-j+1}(A)} \left( 1 + \frac{\sigma_{k+1}(\widehat{A})}{\sigma_k(A) - \sigma_{k+1}(\widehat{A})} \right) = \frac{\sigma_k(A)}{\sigma_{k-j+1}(A)(\sigma_k(A) - \sigma_{k+1}(\widehat{A}))}.$$

Therefore,

$$\max\{\sigma_j(X), \sigma_j(Y)\} \leq \frac{\sigma_k(A)}{\sigma_{k-j+1}(A)} \frac{\max\{\|E_{21}\|_2, \|E_{12}\|_2\}}{\zeta}.$$

Now  $\sigma_j(X) = \sin \theta'_{k-j+1}$  and  $\sigma_j(Y) = \sin \nu'_{k-j+1}$ . Rename  $j \leftarrow k - j + 1$  to finish. □

*Theorem 4.* We tackle each case independently.

**Unitarily invariant norms:** Our proof involves simplifying each term in Equation (7), and Equation (8) and has several steps.

**1. Simplifying the gap** Recall  $\zeta = \sigma_k(A) - \sigma_{k+1}(\widehat{A})$  and  $\widehat{A} = QQ^*A$ . From the first part of Theorem 9

$$\zeta = \sigma_k(A) - \sigma_{k+1}(\widehat{A}) \geq \sigma_k(A) - \sigma_{k+1}(A).$$

**2. Simplifying**  $\|E_{12}\|$  First observe that  $AP_{V_k} = A_k$ . So

$$E_{12} = (I - \mathcal{P}_{\widehat{U}_k})(I - QQ^*)AP_{V_k} = (I - \mathcal{P}_{\widehat{U}_k})(I - QQ^*)A_k.$$

Then applying Equation (14) along with sub-multiplicativity gives

$$\|E_{12}\| \leq \|(I - QQ^*)A_k\| \leq \gamma_k^{2q} \|\Sigma_{\perp} \Omega_2 \Omega_1^{\dagger}\| \leq \gamma_k^{2q} \|\Sigma_{\perp}\| \|\Omega_2 \Omega_1^{\dagger}\|_2.$$

**3. Simplifying**  $\|E_{21}\|$  First,  $E_{21} = \mathcal{P}_{U_k}(I - QQ^*)AP_{\widehat{V}_k}$ , and since  $\|\mathcal{P}_{U_k}(I - QQ^*)\|_2 = \|\sin \angle(U_k, \widehat{U})\|_2$ ,

$$\|E_{21}\| \leq \|\sin \angle(U_k, \widehat{U})\|_2 \|(I - QQ^*)A\|,$$

because of strong sub-multiplicativity. Applying Theorem 1 and Equation (14)

$$\|E_{21}\| \leq \frac{\gamma_k^{2q+1} \|\Omega_2 \Omega_1^{\dagger}\|_2}{\sqrt{1 + \gamma_k^{4q+2} \|\Omega_2 \Omega_1^{\dagger}\|_2^2}} \left(1 + \gamma_k^{2q} \|\Omega_2 \Omega_1^{\dagger}\|_2\right) \|\Sigma_{\perp}\|.$$

Let  $\beta = \gamma_k^{2q} \|\Omega_2 \Omega_1^{\dagger}\|_2$ . Then for  $\beta \geq 0$ , since  $\gamma_k < 1$

$$\frac{\gamma_k(1 + \beta)}{\sqrt{1 + \gamma_k^2 \beta^2}} \leq \frac{1 + \gamma_k \beta}{\sqrt{1 + \gamma_k^2 \beta^2}} \leq \sqrt{2}.$$

Therefore,  $\|E_{21}\| \leq \sqrt{2} \gamma_k^{2q} \|\Sigma_{\perp}\| \|\Omega_2 \Omega_1^{\dagger}\|_2$ .

**4. Putting everything together** Plugging in the intermediate quantities into Equation (7), we have

$$\max \left\{ \|\sin \angle(U_k, \widehat{U}_k)\|, \|\sin \angle(V_k, \widehat{V}_k)\| \right\} \leq \sqrt{2} \gamma_k^{2q} \|\Omega_2 \Omega_1^{\dagger}\|_2 \frac{\|\Sigma_{\perp}\|}{\sigma_k - \sigma_{k+1}}.$$

Dividing the numerator and denominator by  $\sigma_k$  proves the stated result for unitarily invariant norms.

**Spectral/Frobenius norms:** Let  $\|\cdot\|_{\xi}$  denote the spectral and Frobenius norms. The first two steps are identical to the proof for unitarily invariant norms. For the third step, using Equation (14)

$$\|E_{21}\|_{\xi} \leq \frac{\gamma_k^{2q+1} \|\Omega_2 \Omega_1^{\dagger}\|_2}{\sqrt{1 + \gamma_k^{4q+2} \|\Omega_2 \Omega_1^{\dagger}\|_2^2}} \|\Sigma_{\perp}\|_{\xi} \sqrt{1 + \gamma_k^{4q} \|\Omega_2 \Omega_1^{\dagger}\|_2^2}.$$

With  $\beta$  defined as before, since  $\gamma_k < 1$ ,  $\sqrt{\gamma_k^2 + \gamma_k^2 \beta^2} / \sqrt{1 + \gamma_k^2 \beta^2} \leq 1$ . Therefore,

$$\|E_{21}\|_{\xi} \leq \gamma_k^{2q} \|\Sigma_{\perp}\|_{\xi} \|\Omega_2 \Omega_1^{\dagger}\|_2.$$

The rest of the proof is the same.

**Canonical angles:** The proof combines Theorem 3 with the above analysis for the spectral norm. The right hand side contains the term

$$\max \left\{ \|\sin \angle(U_k, \widehat{U}_k)\|_2, \|\sin \angle(V_k, \widehat{V}_k)\|_2 \right\}.$$

The rest of the proof involves some simple manipulations.  $\square$

*Theorem 5.* We first address Equation (9). Following the steps of the proof of Theorem 1, we have

$$\sin^2 \angle(u_j, \widehat{U}) = u_j^* U (I - \mathcal{P}_{U^* Q}) U^* u_j \preceq [e_j^{\top} \quad 0] (I - \mathcal{P}_Z) \begin{bmatrix} e_j \\ 0 \end{bmatrix},$$

where  $e_j$  is the  $j$ -th column of the  $k \times k$  identity matrix. Therefore, we have  $\sin^2 \angle(u_j, \widehat{U}) \leq e_j^\top H e_j$ , where  $H$  was defined in Equation (23). The inequality  $H \preceq F^* F$  implies

$$\begin{aligned} \sin^2 \angle(u_j, \widehat{U}) &\leq \sigma_j^{-4q-2} \left\| (\Sigma_\perp \Sigma_\perp^\top)^q \Sigma_\perp (\Omega_2 \Omega_1^\dagger) e_j \right\|_2^2 \\ &\leq \gamma_j^{4q+2} \left\| \Omega_2 \Omega_1^\dagger \right\|_2^2. \end{aligned}$$

Taking square-roots on both sides gives the desired results. The strategy for bounding the canonical angles  $\sin \angle(v_j, \widehat{V})$  is very similar and will be omitted.

We now address Equation (10), which is a straightforward application of [13, Theorem 2.5]. Let  $\mathcal{P}_U = QQ^*$  and  $\mathcal{P}_V = I$ . Then, in our notation, this result takes the form

$$\max \{ \sin \angle(u_j, \widehat{u}_j), \sin \angle(v_j, \widehat{v}_j) \} \leq \sqrt{1 + 2 \frac{\tilde{\gamma}'^2}{\tilde{\delta}^2}} \max \{ \sin \angle(u_j, \widehat{U}), \sin \angle(v_j, I) \}.$$

where  $\tilde{\gamma}' = \max\{0, \|(I - QQ^*)A\|_2\}$  and  $\tilde{\delta}$  is as defined in the statement of the theorem. Theorem 8 for the spectral norm implies  $\tilde{\gamma}' \leq \tilde{\gamma}$ , whereas Theorem 5 implies

$$\max \{ \sin \angle(u_j, \widehat{U}), \sin \angle(v_j, I) \} \leq \gamma_j^{2q} \left\| \Omega_2 \Omega_1^\dagger \right\|_2.$$

Plug in the intermediate steps to obtain the desired bound.  $\square$

*Theorem 6.* In Theorem 1, bounds for  $\left\| \Omega_2 \Omega_1^\dagger \right\|_2$  are available in the literature. From the proof of [12, Theorem 10.6] we find the inequality

$$\mathbb{E} \left\| \Omega_2 \Omega_1^\dagger \right\|_2 \leq C_e,$$

where the constant  $C_e$  was defined in Equation (11). Let  $\alpha > 0$  be a constant. The map  $x \mapsto x/\sqrt{1 + \alpha x^2}$  is convex. Therefore, by Jensen's inequality the results in expectation follow.

For the concentration inequalities, [11, Theorem 5.8] showed that  $\left\| \Omega_2 \Omega_1^\dagger \right\|_2 \leq C_d$  with a probability at least  $1 - \delta$ . Here,  $C_d$  was defined in Equation (12). Plug into Theorem 1 to obtain the desired bounds.  $\square$

### 5.3 Proofs of Section 4 Theorems

*Theorem 7. Proof of Equation (13):* Using the unitary invariance of the norms

$$\| (I - \mathcal{P}_Q)A \| = \| (I - \mathcal{P}_{U^*Q})\Sigma \| = \left\| (\Sigma^\top (I - \mathcal{P}_{U^*Q})\Sigma)^{1/2} \right\|.$$

We use Equation (22) combined with Lemma 1 to obtain

$$\left\| (\Sigma^\top (I - \mathcal{P}_{U^*Q})\Sigma)^{1/2} \right\| \leq \left\| (\Sigma^\top (I - \mathcal{P}_Z)\Sigma)^{1/2} \right\|.$$

With  $M_1 \equiv I - (I + F^*F)^{-1}$  and  $M_2 \equiv I - F(I + F^*F)^{-1}F^*$ , then  $\Sigma^\top (I - \mathcal{P}_Z)\Sigma$  simplifies as

$$\Sigma^\top (I - \mathcal{P}_Z)\Sigma = \begin{bmatrix} \Sigma_k M_1 \Sigma_k & * \\ * & \Sigma_\perp^\top M_2 \Sigma_\perp \end{bmatrix}. \quad (25)$$

The square root function is concave on  $[0, \infty)$  and  $\Sigma^\top (I - \mathcal{P}_Z)\Sigma$  is positive semidefinite. Therefore, an extension to Rotfel'd's theorem says [16, Theorem 2.1]

$$\left\| (\Sigma^\top (I - \mathcal{P}_Z)\Sigma)^{1/2} \right\| \leq \left\| (\Sigma_k^\top M_1 \Sigma_k)^{1/2} \right\| + \left\| (\Sigma_\perp^\top M_2 \Sigma_\perp)^{1/2} \right\|.$$



Use the inequalities  $M_1 \preceq F^*F$  and  $M_2 \preceq I$ , along with Lemma 1 gives

$$\begin{aligned} \|(I - \mathcal{P}_Q)A\| &\leq \left\| \left\| (\Sigma_k F^* F \Sigma_k)^{1/2} \right\| \right\| + \left\| \left\| (\Sigma_\perp \Sigma_\perp)^{1/2} \right\| \right\| \\ &\leq \|F \Sigma_k\| + \|\Sigma_\perp\|. \end{aligned} \quad (26)$$

Use  $F \Sigma_k = (\Sigma_\perp \Sigma_\perp)^q \Sigma_\perp \Omega_2 \Omega_1^\dagger \Sigma_k^{-2q}$  and the sub-multiplicativity to obtain the advertised bounds.

**Proof of Equation (14):** The proof for Equation (14) is similar and is omitted. The main observation is that  $A_k$  has only  $k$  nonzero singular values.

**Proof of Equation (15):** We follow the strategy in [7, Section 3.3]. Recall that  $B_k$  is the best rank- $k$  approximation to  $B = Q^*A$ . With the notation in Algorithm 2, note that

$$QB_k = Q\widehat{U}_{B,k}\widehat{U}_{B,k}^*B = \widehat{U}_k\widehat{U}_k^*A = \mathcal{P}_{\widehat{U}_k}A,$$

the triangle inequality gives

$$\left\| \left\| (I - \mathcal{P}_{\widehat{U}_k})A \right\| \right\| \leq \left\| \left\| (I - \mathcal{P}_{\widehat{U}_k})A_k \right\| \right\| + \left\| \left\| (I - \mathcal{P}_{\widehat{U}_k})A_\perp \right\| \right\|.$$

Since  $A_k = \mathcal{P}_{U_k}A_k$ , applying strong sub-multiplicativity

$$\left\| \left\| (I - \mathcal{P}_{\widehat{U}_k})A \right\| \right\| \leq \left\| \left\| (I - \mathcal{P}_{\widehat{U}_k})\mathcal{P}_{U_k} \right\| \right\| \|A_k\|_2 + \|\Sigma_\perp\|.$$

We recognize that  $\left\| \left\| (I - \mathcal{P}_{\widehat{U}_k})\mathcal{P}_{U_k} \right\| \right\| = \left\| \left\| \sin \angle(U_k, \widehat{U}_k) \right\| \right\|$ , apply Theorem 4 to complete the proof.  $\square$

*Theorem 8.* The proof is similar to that of the proof of Theorem 7. Consider the term of interest  $\|(I - QQ^*)A\|_p^2$ , which can be simplified to

$$\|(I - QQ^*)A\|_p^2 = \|A^*(I - QQ^*)A\|_{p/2} = \|\Sigma^\top(I - \mathcal{P}_{U^*Q})\Sigma\|_{p/2}.$$

The first equality holds only for  $p \geq 2$ , whereas the last equality follows because of the unitary invariance. As in the proof of Theorem 1, we have

$$Z = U^*Y\Omega_1^\dagger\Sigma_k^{-(2q+1)} \quad F = (\Sigma_\perp\Sigma_\perp^\top)^q\Sigma_\perp\Omega_2\Omega_1^\dagger\Sigma_k^{-(2q+1)}.$$

The use of Equation (22) and Lemma 1 ensures

$$\|\Sigma^\top(I - \mathcal{P}_{U^*Q})\Sigma\|_{p/2} \leq \|\Sigma^\top(I - \mathcal{P}_Z)\Sigma\|_{p/2}.$$

We apply [16, Theorem 2.1] to Equation (25) with  $f(t) = t$  to obtain

$$\begin{aligned} \|\Sigma^\top(I - \mathcal{P}_Z)\Sigma\|_{p/2} &\leq \|\Sigma_k M_1 \Sigma_k\|_{p/2} + \|\Sigma_\perp^\top M_2 \Sigma_\perp\|_{p/2} \\ &\leq \|\Sigma_k F^* F \Sigma_k\|_{p/2} + \|\Sigma_\perp^\top \Sigma_\perp\|_{p/2} \\ &= \|F \Sigma_k\|_p^2 + \|\Sigma_\perp\|_p^2. \end{aligned}$$

We have used  $M_1 \preceq F^*F$  and  $M_2 \preceq I$ . The rest of the proof is similar to that of Theorem 7.  $\square$

*Theorem 9.* The proof makes heavy use of the partial ordering which was reviewed in the start of Section 5. From the inequality  $I \succee QQ^*$ , the conjugation rule gives

$$A^*A \succee A^*QQ^*A.$$

Then, Weyl's theorem implies  $\lambda_j(A^*A) \geq \lambda_j(A^*QQ^*A)$  for  $j = 1, \dots, k$ . Relating the eigenvalues to the singular values proves the first inequality.

For the second inequality consider again  $A^*QQ^*A$ . With the aid of Equation (22)

$$A^*QQ^*A = V\Sigma^\top\mathcal{P}_{U^*Q}\Sigma V^* \succee V\Sigma^\top\mathcal{P}_Z\Sigma V^*. \quad (27)$$

Therefore,  $\lambda_j(A^*QQ^*A) \geq \lambda_j(V\Sigma^\top \mathcal{P}_Z \Sigma V^*)$  for  $j = 1, \dots, k$ . Since  $V\Sigma^\top \mathcal{P}_Z \Sigma V^*$  and  $\Sigma^\top \mathcal{P}_Z \Sigma$  are similar, they share the same eigenvalues. It can be readily shown that

$$\Sigma^\top \mathcal{P}_Z \Sigma = \begin{bmatrix} \Sigma_k(I + F^*F)^{-1}\Sigma_k & * \\ * & * \end{bmatrix}.$$

For  $j = 1, \dots, k$ , the eigenvalues of  $A^*QQ^*A$  satisfy

$$\lambda_j(A^*QQ^*A) \geq \lambda_j(V\Sigma^\top \mathcal{P}_Z \Sigma V^*) \geq \lambda_j(\Sigma_k(I + F^*F)^{-1}\Sigma_k). \quad (28)$$

The second inequality follows from the Cauchy interlacing theorem [19, Section 10-1]. Applying the properties of partial ordering, we obtain

$$F^*F \preceq \sigma_{k+1}^{4q+2} \left\| \Omega_2 \Omega_1^\dagger \right\|_2^2 \Sigma_k^{-(4q+2)} = \left\| \Omega_2 \Omega_1^\dagger \right\|_2^2 \Gamma_k^{4q+2},$$

where  $\Gamma_k = \text{diag}(\gamma_1, \dots, \gamma_k)$  is a diagonal matrix with the singular value gaps. Furthermore,

$$\Sigma_k(I + F^*F)^{-1}\Sigma_k \succeq \Sigma_k(I + \left\| \Omega_2 \Omega_1^\dagger \right\|_2^2 \Gamma_k^{4q+2})^{-1}\Sigma_k.$$

Since the diagonal matrix on the right hand side has its singular values on the diagonals; this fact, combined with Equation (28) gives for  $j = 1, \dots, k$

$$\sigma_j^2(Q^*A) = \lambda_j(A^*QQ^*A) \geq \lambda_j(\Sigma_k(I + F^*F)^{-1}\Sigma_k) \geq \frac{\sigma_j^2(A)}{1 + \left\| \Omega_2 \Omega_1^\dagger \right\|_2^2 \gamma_j^{4q+2}}.$$

Taking square-roots, we obtain the desired result.  $\square$

## 6 Numerical Results

### 6.1 Test matrices

To demonstrate the performance of the bounds, we use the following test matrices

1. **Controlled gap** The first set of test matrices  $A \in \mathbb{R}^{3000 \times 3000}$  are constructed using the formula

$$A = \sum_{j=1}^r \frac{\text{gap}}{j} x_j y_j^\top + \sum_{j=r+1}^{300} \frac{1}{j} x_j y_j^\top,$$

where  $x_j \in \mathbb{R}^{3000}$  and  $y_j \in \mathbb{R}^{300}$  are sparse random vectors with non-negative entries generated using the MATLAB commands `sprand(3000,1,0.025)` and `sprand(300,1,0.025)` respectively. The formula above is not an SVD, since the vectors do not form an orthonormal set. Nonetheless, the singular values decay like  $1/j$  and the gap between the singular values between 15 and 16 is controlled by the parameter gap. We consider three cases:

- (a) Small gap (GapSmall) gap = 1,
- (b) Medium gap (GapMedium) gap = 2,
- (c) Large gap (GapLarge) gap = 10.

2. **Low-rank plus noise** The matrices are of the form

$$A = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} + \sqrt{\frac{\gamma n^r}{2n^2}}(G + G^\top),$$

where  $G \in \mathbb{R}^{n \times n}$  is a random Gaussian matrix. We consider three cases:




Figure 1: Singular value of the matrices from the (left) ‘Controlled Gap’ example, (right) ‘Low-rank plus noise’ example, (below) ‘Low-rank plus decay’ example.

- (a) Small noise (NoiseSmall)  $\gamma_n = 10^{-2}$ ,
- (b) Medium noise (NoiseMedium)  $\gamma_n = 10^{-1}$ ,
- (c) Large noise (NoiseLarge)  $\gamma_n = 1$ .

3. **Low-rank plus decay** The matrices take the form

$$A = U \text{diag}(\underbrace{1, 1, \dots, 1}_r, 2^{-d}, 3^{-d}, \dots, (n-r+1)^{-d}) V^*.$$

The unitary matrices  $U, V$  are obtained by drawing a random Gaussian matrix, and taking its QR factorization. We distinguish between the following cases

- (a) Slow decay (DecaySlow):  $d = 0.5$ ,
- (b) Medium decay (DecayMedium):  $d = 1.0$ ,
- (c) Fast decay (DecayFast):  $d = 2.0$ .

The first example is adapted from [22], whereas the second and third examples are drawn from [24]. In all the examples, the random matrices were fixed by setting the random seed and we set the parameter  $r = 15$ . The singular values of all the test matrices are plotted in Figure 1.

## 6.2 Canonical angles

For the first numerical example, we use the 9 test matrices in Section 6.1. For each matrix, we chose an oversampling parameter  $\rho = 20$  and the target rank  $k$  was chosen to be 25. The starting guess  $\Omega$  was taken to be a random Gaussian matrix.

### 6.2.1 No extraction

We plot the canonical angles  $\sin \angle(U_k, \widehat{U})$  in solid lines, the corresponding bounds from Theorem 1 are also plotted in dashed lines. The results are displayed in Figure 2. We make the following general observations:

- The influence of the subspace iterations on the canonical angles is clear: the angles become smaller as the number of iterations  $q$  increases. This implies that the subspace is becoming more accurate.
- If there is a large singular value gap in the spectrum, this means that all the canonical angles below that index are captured accurately. This is prominently seen in Figure 2(c), in which there is a large gap between singular values 15 and 16. Similar observations can be made in the other figures.
- As the decay rate of the singular values increases, the corresponding canonical angles become smaller.




Figure 2: Plots of  $\sin \theta_j$  for  $j = 1, \dots, k$ . The test matrices were described in Section 6.1. The target rank  $k = 25$  and an oversampling parameter of 20 was chosen for all the experiments. The solid lines correspond to the computed values, the dashed lines correspond to bounds obtained using Theorem 1. The parameter  $q$  corresponds to the number of subspace iterations.

- In most figures the bounds are qualitatively informative, but in some figures, the bounds are also quantitatively accurate (e.g., GapLarge).
- Similar results were observed for  $\sin \angle(V_k, \hat{V})$  and, therefore, omitted.

We now make observations specific to the test examples:

- 1. Gap examples** The computed canonical angles decrease as the gap increases, and with more iterations. The test matrices (GapMedium and GapLarge) have both a decay in the singular values and a prominent singular value gap between indices 15 and 16. These matrices satisfy the assumptions of our analysis, and therefore the bounds can be expected to be good. We see that as the size of the gap increases, the bounds become more accurate in accordance with Theorem 1. GapSmall has decay in the singular values but no special singular value gap. Even in this case, the bounds are qualitatively good.
- 2. Noise examples** NoiseSmall is close to a low-rank matrix and there is a large singular value gap at index 15. For this example, the bounds are qualitatively good. As the level of noise increases, the gap decreases and therefore, the computed angles increase, as predicted by Theorem 1. The bounds are uninformative for  $q = 0$ , but qualitatively good for  $q = 1$  and 2. Compared to

the Gap examples, the bounds are not as sharp since there is very little decay in the singular values.

- 3. Decay examples** In these examples, the singular values decay beyond index 15 but there is no prominent gap. As the rate of decay increases, in general, the canonical angles decrease. It is also seen that the bounds are qualitatively accurate (except for  $q = 0$ ).

### 6.2.2 Extraction step

Our next experiment tests the effect of the extraction step on the accuracy of the canonical angles. We now compute  $\sin \theta'_j$  and  $\sin \nu'_j$  for the test matrices described in Section 6.1. We plot the quantities  $\max\{\sin \theta'_j, \sin \nu'_j\}$  for  $j = 1, \dots, k$  in solid lines. The corresponding bounds from Theorem 4 are plotted in dashed lines. Here, the target rank was chosen to be  $k = 15$ , to exploit the singular value gap in the matrices. We make the following general observations:

- The extraction step did not significantly affect the canonical angles and the accuracy is comparable to Figure 2. The subspaces are more accurate as the number of iterations increase, and if there is a large singular value gap at index  $j$ , then the canonical angles with index  $j' < j$  are captured accurately.
- Although the canonical angles are small, compared to Theorem 1, the bounds in Theorem 4 are not as accurate. One reason is that the upper bounds in Theorem 1 are at most 1, but the bounds in Theorem 4 are allowed to be greater than 1. Furthermore, the bound in Theorem 4 has the factor  $1/(1 - \gamma_k)$  in the denominator, which can be quite large when there is a small singular value gap. It may be possible to derive better bounds, but we could not immediately see how to derive them.
- We also compared the accuracy of the individual singular vectors (not shown here). The results and the conclusions are similar.

We now make observations specific to the test examples:

- 1. Gap examples** The behavior of the computed canonical angles is very similar to that without the extraction step. In general, the angles decrease as the gap increases. When the parameter gap is small, the singular value ratio  $\gamma_k$  is large, and  $(1 - \gamma_k)^{-1}$  is small. This explains why the bounds are bad for GapSmall and GapMedium, and show little improvement with more subspace iterations. Only for the GapLarge example with  $q = 0$ , the bounds are qualitatively good.
- 2. Noise examples** The computed canonical angles decrease as the noise decreases. In all three examples, the bounds are qualitatively good. The bounds are better for NoiseSmall and NoiseMedium because the singular value gap between indices 15 and 16 is bigger than that for NoiseLarge.
- 3. Decay examples** The computed canonical angles become smaller as the decay of the singular values increases. In these examples, there is no prominent gap, so the bounds don't capture the behavior well. However, the computed angles are small, and the subspace is accurate.

## 6.3 Singular Values

We now consider the accuracy of the singular values. We use the same test matrices and the remaining parameters are kept fixed. The computed singular values are plotted against the upper and lower bounds. We make the following general observations:

- For the large singular values, both the upper and lower bounds are qualitatively good for all the examples that we tested.




Figure 3: Plots of  $\max\{\sin \theta'_j, \sin \nu'_j\}$  for  $j = 1, \dots, k$ . The test matrices were described in Section 6.1. The target rank  $k = 15$  and an oversampling parameter of 20 was chosen for all the experiments. The solid lines correspond to the computed values, the dashed lines correspond to bounds obtained using Theorem 4. The parameter  $q$  corresponds to the number of subspace iterations.

- As the number of iterations increase, the singular values are computed more accurately and are close to the upper bounds (the exact singular values). However, for indices close to the target rank, the lower bounds are not tight. The bounds get tighter as the number of iterations  $q$  increase.
- The bounds for the singular values quantitatively better than the bounds for the canonical angles.

We now make observations specific to the test examples:

- 1. GapSmall** In these examples, the large singular values are captured accurately. As the number of iterations increase, both the lower bound and the approximate singular values approach the true singular values (upper bound). For GapMedium and GapLarge, the bounds were much more accurate.
- 2. NoiseMedium** There is a qualitatively different behavior before and after indices 15 – 16. The upper and lower bounds are tight before index 15, but only the upper bound is tight after index 16. The lower bound significantly under-predicts the singular values.




Figure 4: Plots of the singular values. The test matrices were described in Section 6.1. The target rank  $k = 25$  and an oversampling parameter of  $\rho = 20$  was chosen for all the experiments. The solid lines black and blue lines correspond to the upper and lower bounds respectively, the dashed red lines correspond to bounds obtained using Theorem 9. The parameter  $q$  corresponds to the number of subspace iterations.

**3. DecayMedium** Similar to the previous example, the lower bounds are good before index 15, and improve with number of iterations after index 15.

## 7 Acknowledgments

The author would like to thank Ilse C.F. Ipsen and Andreas Stathopoulos for helpful conversations. He would also like to acknowledge Ivy Huang for her help with the figures.

## References

- [1] H. Avron, C. Boutsidis, S. Toledo, and A. Zouzias. Efficient dimensionality reduction for canonical correlation analysis. In *International Conference on Machine Learning*, pages 347–355, 2013.
- [2] O. Balabanov and A. Nouy. Randomized linear algebra for model reduction. Part I: Galerkin methods and error estimation. *arXiv preprint arXiv:1803.02602*, 2018.

- [3] F. L. Bauer. Das verfahren der treppeniteration und verwandte verfahren zur lösung algebraischer eigenwertprobleme. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 8(3):214–235, May 1957.
- [4] R. Bhatia. *Matrix analysis*, volume 169 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997.
- [5] A. Björck and G. H. Golub. Numerical methods for computing angles between linear subspaces. *Mathematics of computation*, 27(123):579–594, 1973.
- [6] C. Boutsidis, P. Kambadur, and A. Gittens. Spectral clustering via the power method-provably. In *International Conference on Machine Learning*, pages 40–48, 2015.
- [7] P. Drineas and I. C. F. Ipsen. Low-rank matrix approximations do not need a singular value gap. *arXiv preprint arXiv:1801.00670*, 2018.
- [8] P. Drineas, I. C. F. Ipsen, E. M. Kontopoulou, and M. Magdon-Ismael. Structural convergence results for approximation of dominant subspaces from block Krylov spaces. *SIAM Journal on Matrix Analysis and Applications*, 39(2):567–586, 2018.
- [9] N. B. Erichson, S. L. Brunton, and J. N. Kutz. Randomized dynamic mode decomposition. *arXiv preprint arXiv:1702.02912*, 2017.
- [10] G. H. Golub and C. F. Van Loan. *Matrix Computations*. The Johns Hopkins University Press, Baltimore, fourth edition, 2013.
- [11] M. Gu. Subspace iteration randomization and singular value problems. *SIAM Journal on Scientific Computing*, 37(3):A1139–A1173, 2015.
- [12] N. Halko, P. G. Martinsson, and J. A. Tropp. Finding Structure with Randomness: Probabilistic Algorithms for Constructing Approximate Matrix Decompositions. *SIAM Rev.*, 53(2):217–288, 2011.
- [13] M. E. Hochstenbach. Harmonic and refined extraction methods for the singular value problem, with applications in least squares problems. *BIT Numerical Mathematics*, 44(4):721–754, 2004.
- [14] J. T. Holodnak, I. C. F. Ipsen, and T. Wentworth. Conditioning of leverage scores and computation by QR decomposition. *SIAM Journal on Matrix Analysis and Applications*, 36(3):1143–1163, 2015.
- [15] R. A. Horn and C. R. Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, second edition, 2013.
- [16] E. Y. Lee. Extension of Rotfel’d theorem. *Linear Algebra and its Applications*, 435(4):735–741, 2011.
- [17] M. W. Mahoney. Randomized algorithms for matrices and data. *Foundations and Trends® in Machine Learning*, 3(2):123–224, 2011.
- [18] Y. Nakatsukasa. Accuracy of singular vectors obtained by projection-based SVD methods. *BIT Numerical Mathematics*, pages 1–16, 2017.
- [19] B. N. Parlett. *The Symmetric Eigenvalue Problem*. Prentice Hall, Englewood Cliffs, 1980.
- [20] Y. Saad. *Numerical methods for large eigenvalue problems*, volume 66 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011. Revised edition of the 1992 original [ 1177405].
- [21] A. Sankar, D. A. Spielman, and S.-H. Teng. Smoothed analysis of the condition numbers and growth factors of matrices. *SIAM Journal on Matrix Analysis and Applications*, 28(2):446–476, 2006.



- [22] D. C. Sorensen and M. Embree. A DEIM induced CUR factorization. *SIAM Journal on Scientific Computing*, 38(3):A1454–A1482, 2016.
- [23] P. T. K. Tang and E. Polizzi. FEAST as a subspace iteration eigensolver accelerated by approximate spectral projection. *SIAM Journal on Matrix Analysis and Applications*, 35(2):354–390, 2014.
- [24] J. A. Tropp, A. Yurtsever, M. Udell, and V. Cevher. Practical sketching algorithms for low-rank matrix approximation. *SIAM Journal on Matrix Analysis and Applications*, 38(4):1454–1485, 2017.
- [25] P. Å. Wedin. On angles between subspaces of a finite dimensional inner product space. In B. Kågström and A. Ruhe, editors, *Matrix Pencils*, pages 263–285, Berlin, Heidelberg, 1983. Springer Berlin Heidelberg.
- [26] K. Ye and L. H. Lim. Schubert varieties and distances between subspaces of different dimensions. *SIAM Journal on Matrix Analysis and Applications*, 37(3):1176–1197, 2016.
- [27] J. Zhang, A. K. Saibaba, M. E. Kilmer, and S. Aeron. A randomized tensor singular value decomposition based on the t-product. *Numerical Linear Algebra with Applications*, page e2179, 2018.