

Self-Adjointness in Klein-Gordon Theory on Globally Hyperbolic Spacetimes

Albert Much^{*1} and Robert Oeckl^{†2}

¹Institut für Theoretische Physik, Universität Leipzig, D-04103 Leipzig

^{1,2}Centro de Ciencias Matemáticas, Universidad Nacional Autónoma de México,
C.P. 58190, Morelia, Michoacán, Mexico

20 April 2018

12 April 2019 (v2)

17 October 2019 (v3)

26 January 2021 (v4)

UNAM-CCM-2018-2

Abstract

We prove essential self-adjointness of the spatial part of the linear Klein-Gordon operator with external potential for a large class of globally hyperbolic manifolds. The proof is conducted by a fusion of new results concerning globally hyperbolic manifolds, the theory of weighted Hilbert spaces and related functional analytic advances.

1 Introduction

Quantum field theory (QFT) in curved spacetime studies the behavior of quantum fields that propagate in the presence of a classical gravitational field, where the quantum behavior of the gravitational field is neglected. It is seen as an intermediate (and mostly rigorous) step towards a complete theory of quantum gravity (see [1, 2, 3, 4, 5] for excellent reviews).

One particularly fruitful context arises from the restriction to globally hyperbolic spacetimes. The advantage of this class of spacetimes is the existence of a (non-canonical) choice of time, or equivalently the existence of a global Cauchy surface. Then the equations of motion given by the Klein Gordon equation have a well posed initial value formulation [2, Theorem 4.1.2], [1].

A well established path towards constructing a free QFT starts with the phase space of initial data of a classical field theory on a spacelike hypersurface. This comes naturally equipped with a symplectic structure. Provided one has a complex structure, which is compatible (tame) with the symplectic structure, a “one particle structure” is constructed by using the aforementioned objects to induce a positive definite inner product. This Hilbert space is then second quantized into a Fock space [6, 7]. For the case of stationary spacetimes this construction has been done in [8] and (more rigorously) in [9]. More recently, the question of finding a complex structure that

*much@itp.uni-leipzig.de

†robert@matmor.unam.mx

is compatible with a unitary time-evolution for non-stationary spacetimes has been addressed e.g. in [10, 11] (and references therein). In [12, 13] for example such a structure was constructed for the Gowdy cosmology (a time dependent globally hyperbolic spacetime).

In the present article we focus on the Klein-Gordon theory. An important ingredient in the construction of complex structures or *Hadamard states* (for the case of FRW-spacetime see [14, Appendix A]) in QFT is an essentially self-adjoint (possibly time-dependent) operator w^2 . This operator encodes the spatial part of the Klein Gordon equation (see Equation 2.4). In general this operator appears as a component in the Hamiltonian, that is represented as a 2×2 matrix, see [9, Equation 2.5], [15, Equation 1.17] and [14, Appendix A]. Therefore, in order for the Hamiltonian to be self-adjoint (which is of essence for a unitary time-evolution) it is essential for the operator w^2 that appears as one of its components to be self-adjoint. For a relation of the construction of a complex structure for a time dependent globally hyperbolic spacetime with the Hamiltonian approach see [16, Section VI.B]. Moreover, for globally hyperbolic manifolds it was recently realized, see [17], that essential self-adjointness of the Klein-Gordon operator is a requirement for the construction of various kinds of propagators needed in quantum field theory. Propagators are essential for the construction of states and thus for the formulation of QFT in such manifolds by the GNS-construction. A fundamental assumption [17, Assumption 1.a.] in the proof of essential self-adjointness of the Klein-Gordon operator is precisely the essential self-adjointness of the spatial part of the Klein-Gordon equation.

In this work we prove that in the case of a globally hyperbolic spacetime, the operator w^2 will take the form of a *weighted Laplace-Beltrami operator* (see Equation (3.1), Proposition 3.1, Lemma 4.1 and Theorem 4.1) plus a potential V (multiplied by a positive smooth function, see Equation 2.4).¹ The Laplace-Beltrami operator for a metric \mathbf{h} is essentially self-adjoint on $C_0^\infty(\Sigma)$ if the Riemannian manifold (Σ, \mathbf{h}) , where Σ is a Cauchy surface, is geodesically complete², see [19, 20]. This result was extended in [21] for the case of weighted Laplace-Beltrami operators and furthermore sufficient conditions for the potential were given in [21], in order for the (weighted) Laplace-Beltrami operator plus a potential to be essentially self-adjoint on $C_0^\infty(\Sigma)$. In these mentioned works geodesic completeness plays a fundamental role and it will be a necessary condition also for our main result, Theorem 4.1, to hold.

In addition to the essential self-adjointness of the operator w^2 we give a condition to guarantee strict positivity thereof. The positivity requirement is imposed in order to take the (unique) square-root and the inverse of the (closure of the) operator which is used in the construction of complex structures. Proving strict positivity and essential self-adjointness in a rather general setting is the goal of the present paper.

Kay proved the essential self-adjointness for the important class of *static* spacetimes, under certain additional boundedness requirements [9, Theorem 7.2]. Our results extend this by showing essential self-adjointness for a considerably larger class of globally hyperbolic spacetimes, while, in some instances, dropping at the same time the additional boundedness requirements. To this end we use, as already mentioned, the theory of weighted Hilbert spaces (see [22]) and related functional analytic advances [21, 23].

Recent results [24] have also shown an important connection between the existence of an essentially self-adjoint quantum Hamiltonian (that guarantees unitary evolution) and the requirement for those operators to be bounded from below. Hence, in addition to essential self-adjointness, positivity plays again an important role.

Throughout this work we use Greek letters $\mu, \nu = 0, \dots, 3$ for spacetime indices and we use Latin letters i, j, k, \dots for spatial components which run from $1, \dots, 3$.

¹The potential in the Klein-Gordon theory is usually given by $V = m^2 + \xi R$, where m denotes the mass of the scalar field and R the scalar curvature of the metric g .

²A complete Riemannian manifold is a Riemannian manifold for which every maximal (inextendible) geodesic is defined on \mathbb{R} , see [18, Definition 1.4.6].

2 Klein-Gordon Theory on Globally Hyperbolic Spacetimes

We start with basic definitions that are needed for the subsequent results.

DEFINITION 2.1 (Globally hyperbolic spacetime). *We denote a spacetime by (M, g) where M is a smooth, four-dimensional manifold and g is a Lorentzian metric on M with signature $(-1, +1, +1, +1)$. In addition we assume time-orientability of the manifold. This means that there exists a C^∞ -vector-field u on M that is everywhere timelike, i.e. $g(u, u) < 0$. A smooth curve $\gamma : I \rightarrow M$, I being a connected subset of \mathbb{R} , is called causal if $g(\dot{\gamma}, \dot{\gamma}) \leq 0$ where $\dot{\gamma}$ denotes the tangent vector of γ . A causal curve is called future directed if $g(\dot{\gamma}, u) < 0$ and past directed if $g(\dot{\gamma}, u) > 0$ all along γ and for a global timelike vector-field u . For any point $x \in M$, $J^\pm(x)$ denotes the set of all points in M which can be connected to x by a future(+)/past(-)-directed causal curve. An time-orientable spacetime is called globally hyperbolic if for each pair of points $x, y \in M$ the set $J^-(x) \cap J^+(y)$ is compact whenever it is non-empty. This definition is equivalent to the existence of a smooth foliation of M in Cauchy surfaces, where a smooth hypersurface of M is called a Cauchy surface if it is intersected exactly once by each inextendible causal curve.*

The advantages of requiring a spacetime to be globally hyperbolic are best displayed by the following theorem [15].

THEOREM 2.1. *Given a spacetime (M, g) the following statements are equivalent:*

- (M, g) is globally hyperbolic
- There exists a (global) Cauchy surface in (M, g)
- There exists a choice of time³ on (M, g)

Hence, an effective way of thinking of a globally hyperbolic spacetimes is to think that these spacetimes admit a choice of time or that they are topologically equivalent to $\mathbb{R} \times \Sigma$ for any Cauchy surface Σ (strictly, $\{t\} \times \Sigma$ for any $t \in \mathbb{R}$) [25]. The authors in [26] solved a long-standing conjecture by proving that any globally hyperbolic spacetime admits a *smooth* foliation into Cauchy surfaces [26, Theorem 1.1]. Moreover, the induced metric of such a globally hyperbolic spacetime admits a specific form [27, Theorem 1.1].

THEOREM 2.2. *Let (M, g) be a globally hyperbolic spacetime. Then, it is isometric to the smooth product manifold $\mathbb{R} \times \Sigma$ with a metric g , i.e.,*

$$g = -N^2 dt^2 + h_{ij} dx^i dx^j, \quad (2.1)$$

where Σ is a smooth 3-manifold, $t : \mathbb{R} \times \Sigma \mapsto \mathbb{R}$ is the natural projection, $N : \mathbb{R} \times \Sigma \mapsto (0, \infty)$ a smooth function, and \mathbf{h} a 2-covariant symmetric tensor field on $\mathbb{R} \times \Sigma$, satisfying the following condition: Each hypersurface $\Sigma_t \subset M$ at constant t is a Cauchy surface, and the restriction $\mathbf{h}(t)$ of \mathbf{h} to such a Σ_t is a Riemannian metric (i.e. Σ_t is spacelike).

In general N and \mathbf{h} depend on the time and space coordinates. By abuse of notation we also write $N = N(t)$ and $\mathbf{h} = \mathbf{h}(t)$ when denoting the corresponding objects pulled back to Σ for a fixed time t .

We proceed to consider the *Klein-Gordon equation* in (M, g) with an external potential V .⁴ A solution ϕ satisfies,

$$(\square_g - V)\phi = 0, \quad (2.2)$$

³See [15] for definition.

⁴The potential is usually given by $V = m^2 + \xi R$, where m denotes the mass of the scalar field and R the scalar curvature w.r.t. the metric g . The regularity of the potential will be discussed in more detail for the upcoming results.

where \square_g is the wave operator with respect to the metric g . That is,

$$\square_g = (\sqrt{|g|})^{-1} \partial_\mu (\sqrt{|g|} g^{\mu\nu} \partial_\nu),$$

with $|g|$ denoting the absolute value of the determinant of the metric g . We denote for each Cauchy surface Σ the space of smooth Cauchy data of compact support by

$$\mathcal{S}_\Sigma := C_0^\infty(\Sigma) \oplus C_0^\infty(\Sigma).$$

Moreover, due to Leray's Theorem [28], the Cauchy data $\Phi \in \mathcal{S}_\Sigma$ given by

$$\Phi = \begin{pmatrix} \varphi \\ \pi \end{pmatrix} = \begin{pmatrix} \phi \\ n^\mu \nabla_\mu \phi \end{pmatrix} \Big|_\Sigma = \begin{pmatrix} \phi \\ N^{-1} \partial_t \phi \end{pmatrix} \Big|_\Sigma \quad (2.3)$$

define a unique solution ϕ in $C^\infty(M)$, where $N^{-1} \partial_t = n^\mu \nabla_\mu$ and the vector n^μ is the future pointing unit vector normal to the surface Σ . Let us further define the corresponding symplectic form $\Omega_\Sigma : \mathcal{S}_\Sigma \times \mathcal{S}_\Sigma \mapsto \mathbb{R}$ on \mathcal{S}_Σ as

$$\Omega_\Sigma(\Phi_1, \Phi_2) = \int_\Sigma (\pi_1 \varphi_2 - \pi_2 \varphi_1) \sqrt{|\mathbf{h}|} d^3x,$$

where the integral exists since the functions Φ_1 and Φ_2 are compactly supported on Σ . Moreover, for given global solutions of the Klein-Gordon equation (2.2) the integral is independent of the choice of the Cauchy surface. The symplectic structure Ω_Σ makes \mathcal{S}_Σ into a symplectic vector space (a fact that relies on the requirement that the spacetime is globally hyperbolic, see [29]). Next, let the operator w^2 be given by

$$\begin{aligned} w^2 &= -\frac{N}{\sqrt{|\mathbf{h}|}} \partial_i (\sqrt{|\mathbf{h}|} N h^{ij} \partial_j) + N^2 V \\ &= -N^2 (\Delta_{\mathbf{h}} - V) - N h^{ij} \partial_i N \partial_j, \end{aligned} \quad (2.4)$$

where $\Delta_{\mathbf{h}}$ is the Laplace-Beltrami operator with respect to the associated spatial metric \mathbf{h} . The operator w^2 is defined such that the Klein-Gordon equation takes the form,

$$(\partial_t^2 + f(t, x) \partial_t + w^2) \phi = 0,$$

where $f(t, x) = -N^{-1} \partial_t N + (\sqrt{|\mathbf{h}|})^{-1} \partial_t \sqrt{|\mathbf{h}|}$.

3 Weighted Manifolds and Essential Self-Adjointness

We proceed by introducing the notion of weighted manifolds and weighted Hilbert spaces. For further details we direct the reader to the excellent reference [22]. We begin this section with the following definition, [22, Chapter 3.6, Definition 3.17].

DEFINITION 3.1. *A triple $(\Sigma, \mathbf{h}, \mu)$ is called a weighted manifold, if (Σ, \mathbf{h}) is a Riemannian manifold and μ is a measure on Σ with a smooth and everywhere positive density function ρ , i.e., $d\mu = \rho \sqrt{|\mathbf{h}|} d^3x$. A weighted Hilbert space, denoted by $L^2(\Sigma, \mu)$, is given as the space of all square-integrable functions on the manifold Σ with respect to the measure μ . The corresponding weighted Laplace-Beltrami operator (also called the Dirichlet-Laplace operator), denoted by Δ_μ is,*

$$\Delta_\mu = \frac{1}{\rho \sqrt{|\mathbf{h}|}} \partial_i (\rho \sqrt{|\mathbf{h}|} h^{ij} \partial_j). \quad (3.1)$$

We use the following proposition about weighted manifolds in the subsequent discussion [22, Chapter 3, Exercise 3.11].

PROPOSITION 3.1. *Let a, b be smooth and everywhere positive functions on a weighted manifold $(\Sigma, \mathbf{h}, \mu)$ and define a new metric $\tilde{\mathbf{h}}$ and measure $\tilde{\mu}$ by*

$$\tilde{\mathbf{h}} = a \mathbf{h}, \quad \text{and} \quad d\tilde{\mu} = b d\mu.$$

Then, the weighted Laplace-Beltrami operator $\tilde{\Delta}_{\tilde{\mu}}$ of the weighted manifold $(\Sigma, \tilde{\mathbf{h}}, \tilde{\mu})$ is given by

$$\tilde{\Delta}_{\tilde{\mu}} = \frac{1}{b} \operatorname{div}_{\mu} \left(\frac{b}{a} \nabla \right),$$

where in local coordinates the divergence of a vector field v is given by

$$\operatorname{div}_{\mu} v = \frac{1}{\rho} \frac{\partial}{\partial x^i} (\rho v^i).$$

In particular, if $a = b$ then

$$\tilde{\Delta}_{\tilde{\mu}} = \frac{1}{a} \Delta_{\mu}.$$

Proof. The proof is conducted by using functions of compact support and the Green formula [22, Chapter 3, Theorem 3.16],

$$\begin{aligned} \int_{\Sigma} u(\tilde{\Delta}_{\tilde{\mu}} v) d\tilde{\mu} &= - \int_{\Sigma} (\partial_i u) \tilde{h}^{ij} (\partial_j v) d\tilde{\mu} \\ &= - \int_{\Sigma} (\partial_i u) \frac{b}{a} h^{ij} (\partial_j v) d\mu = \int_{\Sigma} u \operatorname{div}_{\mu} \left(\frac{b}{a} \nabla v \right) d\mu \\ &= \int_{\Sigma} u \frac{1}{b} \operatorname{div}_{\mu} \left(\frac{b}{a} \nabla v \right) d\tilde{\mu}, \end{aligned}$$

for all $u, v \in C_0^{\infty}(\Sigma)$. For $b = a$ we have,

$$\int_{\Sigma} u(\tilde{\Delta}_{\tilde{\mu}} v) d\tilde{\mu} = \int_{\Sigma} u \frac{1}{a} \operatorname{div}_{\mu} (\nabla v) d\tilde{\mu} = \int_{\Sigma} u \left(\frac{1}{a} \Delta_{\mu} v \right) d\tilde{\mu}.$$

□

Remark 3.1. *From the Green Formula for a weighted Laplace-Beltrami operator it follows that the weighted Laplace-Beltrami operator is a positive operator (see [22, Lemma 4.4 and Equation 4.14]).*

Before proceeding to our general result we need to mention another theorem that we use. First, define a local $L^2(\Sigma, \mu)$ function f as a function that is square integrable (with respect to the scalar product of the weighted Hilbert space $L^2(\Sigma, \mu)$) on every compact subset of the manifold Σ and we write $f \in L_{loc}^2(\Sigma, \mu)$. Moreover, a symmetric operator H is semi-bounded on $C_0^{\infty}(\Sigma)$ if there exists a constant $C \in \mathbb{R}$ such that,

$$\langle \Psi, H \Psi \rangle \geq -C \langle \Psi, \Psi \rangle, \quad \Psi \in C_0^{\infty}(\Sigma).$$

Then, the theorem of Shubin states the following [21, Theorem 1.1] (see also [23] for an extension of this result to singular potentials).

THEOREM 3.1. *Let the Riemannian manifold (Σ, \mathbf{h}) be complete and let the potential $V \in L_{loc}^2(\Sigma, \mu)$ be such that we can write $V = V_+ + V_-$, where $V_+ \in L_{loc}^2(\Sigma, \mu) \geq 0$ and $V_- \in L_{loc}^2(\Sigma, \mu) \leq 0$ point-wise. Furthermore, let the operator $H_V = -\Delta_{\mu} + V$ be semi-bounded from below. Then, the operator H_V is an essentially self-adjoint operator on $C_0^{\infty}(\Sigma)$.*

This theorem is in particular an extension of the result that the (unweighted) Laplace-Beltrami operator for a metric \mathbf{h} is essentially self-adjoint on $C_0^{\infty}(\Sigma)$ if the Riemannian manifold (Σ, \mathbf{h}) is geodesically complete, see [19, 20], to the case of a weighted Laplace-Beltrami operator ([21, Theorem 6.1]).

4 Main Result on Self-Adjointness

In this section we present our main result, i.e., proving essential self-adjointness and positivity of the operator w^2 that was given by formula (2.4). In contrast to previous works, our result also covers the case of globally hyperbolic spacetimes with *unbounded* N (see next section).

We consider the weighted manifolds $(\Sigma, \mathbf{h}, \mu)$ and $(\Sigma, \tilde{\mathbf{h}}, \tilde{\mu})$, where $d\tilde{\mu} = N^{-2}d\mu$ and $\tilde{\mathbf{h}} = N^{-2}\mathbf{h}$. The measure $d\tilde{\mu} = N^{-1}\sqrt{|\tilde{\mathbf{h}}|}d^3x$ represents the one usually considered in field theory in curved spacetimes to define the symplectic structure or a real inner product (see [2, Equation 4.2.6] or [8]). Hence, one is in general interested in proving the essential self-adjointness of the operator w^2 with respect to the measure $d\tilde{\mu}$.

LEMMA 4.1. *Let the weighted manifold $(\Sigma, \mathbf{h}, \mu)$ be given by the Riemannian manifold (Σ, \mathbf{h}) and the corresponding measure μ with the smooth strictly positive density function N . Moreover, let the weighted manifold $(\Sigma, \tilde{\mathbf{h}}, \tilde{\mu})$ be given by the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$, with Riemannian metric $\tilde{\mathbf{h}} = N^{-2}\mathbf{h}$ and corresponding measure $\tilde{\mu}$ with the smooth strictly positive density function N^{-1} . Then, the operator w^2 (from Equation 2.4) is given by the sum of a weighted Laplace-Beltrami operator and a potential term,*

$$\begin{aligned} w^2 &= -N^2\Delta_\mu + N^2V \\ &= -\tilde{\Delta}_{\tilde{\mu}} + N^2V, \end{aligned}$$

where Δ_μ and $\tilde{\Delta}_{\tilde{\mu}}$ are the weighted Laplace-Beltrami operators of the weighted manifolds $(\Sigma, \mathbf{h}, \mu)$ and $(\Sigma, \tilde{\mathbf{h}}, \tilde{\mu})$, respectively.

Proof. We study the Laplace-Beltrami part of the operator w^2 , i.e., the operator $N^{-2}w^2 - V$. We write it as a weighted Laplace-Beltrami operator of the weighted manifold $(\Sigma, \mathbf{h}, \mu)$, where the measure is $d\mu = N\sqrt{|\mathbf{h}|}d^3x$. Positivity and smoothness of the function N follow from global hyperbolicity (see Theorem 2.2) and hence we have

$$N^{-2}w^2 - V = -\Delta_\mu = -\frac{1}{N\sqrt{|\mathbf{h}}|}\partial_i(N\sqrt{|\mathbf{h}|}h^{ij}\partial_j).$$

Next, we use Proposition 3.1 and make the following transformations

$$\tilde{\mathbf{h}} = N^{-2}\mathbf{h}, \quad d\tilde{\mu} = N^{-2}d\mu = N^{-1}\sqrt{\tilde{h}}d^3x.$$

Note that the operator N^{-2} satisfies the conditions of Proposition 3.1 (the conditions on a) as a multiplication operator since global hyperbolicity demands from the operator N to be smooth, positive and invertible. After applying the transformations we obtain the weighted Laplace-Beltrami operator of the weighted manifold $(\Sigma, \tilde{\mathbf{h}}, \tilde{\mu})$,

$$\tilde{\Delta}_{\tilde{\mu}} = N^2\Delta_\mu,$$

which is exactly the Laplace-Beltrami term (i.e., the term without the potential) of the operator w^2 . \square

By using the previous lemma we are able to obtain the following general result.

THEOREM 4.1. *Let the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$ (from Lemma 4.1) be complete and let the scaled potential $N^2V = V_+ + V_-$, where $V_+ \in L^2_{loc}(\Sigma, \tilde{\mu}) \geq 0$ and $V_- \in L^2_{loc}(\Sigma, \tilde{\mu}) \leq 0$ point-wise. Furthermore, let the operator $w^2 = -\tilde{\Delta}_{\tilde{\mu}} + N^2V$ be semi-bounded from below. Then, the operator w^2 is essentially self-adjoint on $C_0^\infty(\Sigma) \subset L^2(\Sigma, \tilde{\mu})$.*

Proof. By Lemma 4.1 we know that the operator w^2 takes the form of a weighted Laplace-Beltrami operator of the weighted manifold $(\Sigma, \tilde{\mathbf{h}}, \tilde{\mu})$ with the addition of a scaled potential term, i.e.

$$w^2 = -\tilde{\Delta}_{\tilde{\mu}} + N^2 V.$$

By demanding geodesic completeness of $(\Sigma, \tilde{\mathbf{h}})$, by imposing the requirements of local integrability on the scaled potential term $N^2 V$ and by the semi-boundedness condition on the operator w^2 , essential self-adjointness follows from Theorem 3.1. \square

PROPOSITION 4.1. *Let the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$ (from Lemma 4.1) be complete and let the potential V be strictly positive, i.e. $V > \epsilon$ for some $\epsilon > 0$. Moreover, let the scaled potential $N^2 V \in L^2_{loc}(\Sigma, \tilde{\mu})$ be locally square integrable. Then, the operator w^2 is essentially self-adjoint on $C_0^\infty(\Sigma) \subset L^2(\Sigma, \tilde{\mu})$ and the closure of the operator is strictly positive and invertible.*

Proof. By Theorem 2.2 we know that the function N is positive and hence the condition of strict positivity on the potential guarantees the semi-boundedness from below by a positive constant $c > 0$, i.e.

$$\langle \Phi, w^2 \Phi \rangle = \langle \Phi, (-\tilde{\Delta}_{\tilde{\mu}} + N^2 V) \Phi \rangle \geq c \|\Phi\|^2, \quad (4.1)$$

which holds $\forall \Phi \in C_0^\infty(\Sigma)$. Hence, by the completeness of the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$, by the local integrability condition of the scaled potential and by the semi-boundedness of the operator w^2 essential self-adjointness follows from Theorem 3.1. Since the operator w^2 is a strictly positive essentially self-adjoint operator, it has only one semi-bounded self-adjoint extension which coincides with the Friedrich extension ([30, Theorem X.23, Theorem X.26]) that is bounded by the same constant c (see Equation 4.1). \square

It follows from this proof that the square root (for $V > \epsilon$ for some $\epsilon > 0$) and its inverses are well-defined and given as a unique self-adjoint operator. The self-adjointness of the respective operators is proven by using the spectral theorem ([31, Chapter V.III.3]) or by using [32, Theorem 3.35], [33] or for a new shorter proof see [34].

Remark 4.1. *Since the operator w^2 is a positive symmetric operator (with the additional requirement on the potential) we could have used the Friedrich extension (as suggested in [8]) to prove that there exists a self-adjoint extension. However, the problem with this approach is lack of uniqueness. In particular, the extension given by the Friedrich theorem is, although self-adjoint, not necessarily unique. Proving essential self-adjointness for a strictly positive operator, on the other hand, implies the uniqueness of the Friedrich extension.*

5 Complete Riemannian Manifolds

In the last section we made the assumption that the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$ is complete in order to use Theorem 3.1 to prove essential self-adjointness. This section gives concrete results (and examples) for this condition.

Consider first the case where the hypersurface Σ used in the foliation of Theorem 2.2 is compact. From compactness it follows that the Riemann manifold (Σ, \mathbf{h}) is complete (see [18, Theorem 1.4.7] or [9, Corollary 5.4]). But the same is also true for the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$ with the conformally transformed metric $\tilde{\mathbf{h}} = N^{-2} \mathbf{h}$. Hence, in the case of compact Cauchy surfaces our proof applies without restriction.

We proceed to consider the case where Σ is non-compact.

5.1 Static Globally Hyperbolic Spacetimes

First, we investigate the case of static globally hyperbolic spacetimes, i.e., spacetimes (of the form given in Theorem 2.2) where N and all components of \mathbf{h} are time independent. The proposition that we use in the subsequent discussion is the following [9, Proposition 5.3].

PROPOSITION 5.1. *The Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$ is Cauchy with respect to the metric tensor*

$$g = -dt^2 + \tilde{h}_{ij}(\vec{x})dx^i dx^j \quad (5.1)$$

if and only if the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$ is complete.

By using the previous proposition we have the following result.

COROLLARY 5.1. *Let $(M = \mathbb{R} \times \Sigma, g)$ be a static globally hyperbolic spacetime given by*

$$g = -N^2(\vec{x})dt^2 + h_{ij}(\vec{x})dx^i dx^j.$$

Then, the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}} = N^{-2}\mathbf{h})$ is Cauchy (with respect to the metric $\tilde{g} = N^{-2}g$) and therefore geodesically complete.

Proof. Let us consider the conformally equivalent spacetime (M, \tilde{g}) that is ultra-static with metric

$$g = -dt^2 + N^{-2}h_{ij}dx^i dx^j. \quad (5.2)$$

The manifold (M, \tilde{g}) is globally hyperbolic since conformal transformations (that are smooth and strictly positive) preserve the causal structure, see [9, 35, 36] and [37, Appendix D].⁵ Hence, by the previous proposition (and the definition of globally hyperbolic manifolds) the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$ is Cauchy (for the metric \tilde{g}) and therefore necessarily complete. \square

Hence, our proof of essential self-adjointness of w^2 (see Theorem 4.1) holds for *all* static globally hyperbolic spacetimes (of the form given in Theorem 2.2 and a potential that is semi-bounded from below that in addition satisfies the required conditions in [21]).⁶

5.2 Globally Hyperbolic Spacetimes

Next, we provide a sufficient condition for a (not necessarily stationary) globally hyperbolic spacetime $(M = \mathbb{R} \times \Sigma, g)$, with Cauchy surface Σ (strictly, $\{t\} \times \Sigma$ for any $t \in \mathbb{R}$) and induced metric \mathbf{h} , to admit a conformally transformed geodesically complete Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$. In this context we use a fundamental theorem, [38, Theorem 2.1] (see also [39, 40]), for general spacetime manifolds (M, g) .⁷ Let the spacetime M be given by the product $M = \mathbb{R} \times \Sigma$, where Σ is an n -dimensional smooth manifold. Equip the manifold M with a $n + 1$ -dimensional Lorentz metric g of the form

$$g = -N^2(\vec{x}, t)dt^2 + h_{ij}(\vec{x}, t)dx^i dx^j, \quad (5.3)$$

where $N(\vec{x}, t)$ is the lapse function and $\Sigma_t = \{t\} \times \Sigma$, spatial slices of M , are space-like sub-manifolds equipped with the time-dependent metric $\mathbf{h}_t = h_{ij}(\vec{x}, t)dx^i dx^j$.⁸ Moreover, let the following Assumption be met.

⁵In fact, if the spacetimes (M, g) and (M, \tilde{g}) have identical causal structures, the respective metrics must be related by a conformal transformation.

⁶It can be easily seen that our result contains [14, Theorem A.1.] as a special case.

⁷The authors give the theorem for a metric with shift vector. However, if the shift vector is equal to zero the condition of the Theorem concerning the shift vector is trivially satisfied.

⁸Such a product space M is called a *sliced space*. The spacetime is time-oriented by increasing t .

Assumption 5.1. *With respect to the components of the metric g we have the following bounds.*

1. *The lapse function is bounded from above and below for all t by $\alpha_B, \alpha_C \in \mathbb{R}$, i.e.,*

$$0 < \alpha_B \leq N(\vec{x}, t) \leq \alpha_C.$$

2. *The metric $\mathbf{h}(\vec{x}, t)$ is uniformly bounded by the metric $\mathbf{h}(\vec{x}, 0)$ for all $t \in \mathbb{R}$ and tangent vectors $u \in T\Sigma$. That is, there exists universal constants $A, D \in \mathbb{R} > 0$ such that,*

$$A h_{ij}(\vec{x}, 0) u^i u^j \leq h_{ij}(\vec{x}, t) u^i u^j \leq D h_{ij}(\vec{x}, 0) u^i u^j. \quad (5.4)$$

THEOREM 5.1 ([38, Theorem 2.1]). *For the spacetime manifold (M, g) , with metric g that satisfies Assumption 5.1, the two following statements are equivalent.*

1. *(Σ, \mathbf{h}) is a complete Riemannian manifold.*
2. *The spacetime (M, g) is globally hyperbolic.*

Since with regards to our proof of essential self-adjointness we are interested in the conformally transformed metric $\tilde{\mathbf{h}}$ we obtain the following result.

THEOREM 5.2. *Let the globally hyperbolic spacetime M be given by the product $M = \mathbb{R} \times \Sigma$, with a 4-dimensional Lorentz metric g of the Form (5.3). Moreover, let Inequality (5.4) be fulfilled for the conformally transformed metric $\tilde{\mathbf{h}} := N^{-2}\mathbf{h}$. Then, the Riemannian manifold $(\Sigma, \tilde{\mathbf{h}})$ is geodesically complete.*

Proof. Since conformal transformations do not change the causal structure, the spacetime (M, \tilde{g}) with metric,

$$\tilde{g} = -dt^2 + \tilde{h}_{ij}(\vec{x}, t) dx^i dx^j,$$

is globally hyperbolic. Therefore, the first bound of Assumption 5.1 is trivially satisfied and the second bound is the demanded condition. Hence, the proof is concluded by the globally hyperbolicity of (M, \tilde{g}) and Theorem 5.1. \square

6 Discussion

Essential self-adjointness of the spatial part w^2 of the Klein Gordon operator has been proven for the case that the conformally transformed Cauchy surface under consideration is complete, i.e. that $(\Sigma, N^{-2}\mathbf{h})$ is a complete metric space. In addition we gave a necessary condition on the conformal factor for the manifold $(\Sigma, N^{-2}\mathbf{h})$ to be complete without necessarily having a complete Riemannian manifold (Σ, \mathbf{h}) .

This generalizes a corresponding result [9, Theorem 7.2] of Kay, who had to assume that (Σ, \mathbf{h}) is complete and that N is bounded as a multiplication operator. In particular, if N is bounded by positive constants from above and below, it follows (by strong equivalence) that $(\Sigma, N^{-2}\mathbf{h})$ is a complete metric space. However, the restriction on N and more specifically the assumption that (Σ, \mathbf{h}) is complete⁹ is not necessary as long as the conformally transformed metric becomes complete. Hence, given a globally hyperbolic manifold that is smoothly foliated by non-complete Cauchy surfaces, the proof of essential self-adjointness of the operator w^2 is still possible (by standard

⁹Kay was aware of this observation and gave explicit examples in [9] for which a conformal transformation completes an incomplete metric space.

methods) if the conformal factor N^{-2} acts as a completion. In the case of stationary globally hyperbolic spacetimes of the form given in Theorem 2.2 and a potential that is semi-bounded from below that in addition satisfies the required conditions in [21], our main result, Theorem 4.1, holds without restriction to generality.

Although we require global hyperbolicity of spacetime, a possible application of our theorem to QFT on certain spacetimes that are not globally hyperbolic might be possible. The reason therefore lies in the manner how QFT is studied on such spacetimes. In particular, in [41] a generalization was proposed in the context of the *algebraic approach* to quantum field theory to study the case of non-globally hyperbolic spacetimes. Basically one requires that every point in a given spacetime M should have a globally hyperbolic neighborhood N . Moreover, the algebra of observables of the spacetime M restricted to N should be equal to the algebra obtained by regarding the neighborhood N as a globally hyperbolic spacetime in its own right (with some choice of time). With regards to our result this means that we can define a Klein-Gordon equation on each such globally hyperbolic neighborhood and use Theorem 4.1 to prove the essential self-adjointness of the spatial part of the Klein-Gordon operator in that neighborhood.

Another approach for doing QFT on not necessarily globally hyperbolic spacetimes is *general boundary quantum field theory* [42, 43]. There, Hilbert spaces of states are associated to hypersurfaces that need not even be spacelike. However, in many contexts the restriction to spacelike hypersurfaces is sufficient and our result is useful there.

Acknowledgments

The authors would like to thank Suzanne Lanéry for many useful discussions and one of the authors (AM) would like to thank Bernard Kay for enlightening and interesting discussions on quantum field theory in curved spacetimes and related functional analytical questions. The authors acknowledge partial support from CONACYT project 259258.

References

- [1] B. S. Kay, R. M. Wald, *Theorems on the Uniqueness and Thermal Properties of Stationary, Nonsingular, Quasifree States on Space-Times with a Bifurcate Killing Horizon*, Phys. Rept. **207** (1991) 49–136.
- [2] R. Wald, *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*, Chicago Lectures in Physics, University of Chicago Press, 1994.
- [3] L. H. Ford, *Quantum field theory in curved space-time*, Particles and fields. Proceedings, 9th Jorge Andre Swieca Summer School, Campos do Jordao, Brazil, February 16-28, 1997, 1997, pp. 345–388.
- [4] B. S. Kay, *Quantum Field Theory in Curved Spacetime*, Mathematical Physics X, (ed. K. Schmüdgen), Springer Berlin Heidelberg, Berlin, Heidelberg, 1992, pp. 383–387.
- [5] S. Hollands, R. M. Wald, *Quantum fields in curved spacetime*, Phys. Rept. **574** (2015) 1–35.
- [6] I. E. Segal, *Foundations of the theory of dynamical systems of infinitely many degrees of freedom.*, Mat.-Fys. Medd. Danske Vid. Selsk. **31** (1959), no. 12 1–39.
- [7] I. E. Segal, *Foundations of the theory of dynamical systems of infinitely many degrees of freedom. II*, Canad. J. Math. **13** (1961) 1–18.
- [8] A. Ashtekar, A. Magnon, *Quantum Fields in Curved Space-Times*, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences **346** (1975), no. 1646 375–394.

- [9] B. S. Kay, *Linear Spin 0 Quantum Fields in External Gravitational and Scalar Fields. 1. A One Particle Structure for the Stationary Case*, Commun. Math. Phys. **62** (1978) 55–70.
- [10] A. Corichi, J. Cortez, H. Quevedo, *On the Schrodinger representation for a scalar field on curved space-time*, Phys. Rev. **D66** (2002) 085025.
- [11] I. Agullo, A. Ashtekar, *Unitarity and ultraviolet regularity in cosmology*, Phys. Rev. **D91** (2015), no. 12 124010.
- [12] C. G. Torre, *Quantum dynamics of the polarized Gowdy T^{*3} model*, Phys. Rev. **D66** (2002) 084017, [gr-qc/0206083](#).
- [13] A. Corichi, J. Cortez, G. A. Mena Marugán, J. M. Velhinho, *Quantum Gowdy T^3 model: Schrödinger representation with unitary dynamics*, Phys. Rev. D. **76** (2007), no. 12 124031.
- [14] M. Brum, K. Fredenhagen, *'Vacuum-like' Hadamard states for quantum fields on curved spacetimes*, Class. Quant. Grav. **31** (2014), no. 2 025024.
- [15] B. S. Kay, *Generally Covariant Perturbation Theory: Linear Spin 0 Quantum Fields in External Gravitational and Scalar Fields. 2.*, Commun. Math. Phys. **71** (1980) 29.
- [16] A. Corichi, J. Cortez, H. Quevedo, *On the relation between Fock and Schrodinger representations for a scalar field*, Annals Phys. **313** (2004) 446–478, [hep-th/0202070](#).
- [17] J. Dereziński, D. Siemssen, *An Evolution Equation Approach to the Klein-Gordon Operator on Curved Spacetime*, Pure Appl. Analysis **1** (2019) 215–261, [1709.03911](#).
- [18] J. Jost, *Riemannian Geometry and Geometric Analysis*, 4th ed ed., Universitext, Springer, 2005.
- [19] R. S. Strichartz, *Analysis of the Laplacian on the complete Riemannian manifold*, J. Funct. Anal. **52** (1983), no. 1 48 – 79.
- [20] P. R. Chernoff, *Essential self-adjointness of powers of generators of hyperbolic equations*, J. Funct. Anal. **12** (1973), no. 4 401 – 414.
- [21] M. Shubin, *Essential Self-Adjointness for Semi-bounded Magnetic Schroedinger Operators on Non-compact Manifolds*, J. Funct. Anal. **186** (2001), no. 1 92 – 116.
- [22] A. Grigor'yan, *Heat Kernel and Analysis on Manifolds*, AMS/IP Studies in Advanced Mathematics 47, American Mathematical Society, 2009.
- [23] M. Braverman, O. Milatovic, M. Shubin, *Essential self-adjointness of Schrödinger-type operators on manifolds*, Russ. Math. Surv. **57** (2002) 641–692.
- [24] A. D. Helfer, *The Hamiltonians of linear quantum fields. 2. Classically positive Hamiltonians* (1999), [hep-th/9908012](#).
- [25] R. Geroch, *Domain of Dependence*, J. Math. Phys. **11** (1970), no. 2 437–449.
- [26] A. N. Bernal, M. Sanchez, *On Smooth Cauchy hypersurfaces and Geroch's splitting theorem*, Commun. Math. Phys. **243** (2003) 461–470, [gr-qc/0306108](#).
- [27] A. N. Bernal, M. Sánchez, *Smoothness of Time Functions and the Metric Splitting of Globally Hyperbolic Spacetimes*, Commun. Math. Phys. **257** (2005), no. 1 43–50.
- [28] J. Leray, *Hyperbolic partial differential equations*, Princeton lecture notes., Princeton University, Princeton, 1952.
- [29] U. Yurtsever, *Algebraic approach to quantum field theory on nonglobally hyperbolic space-times*, Class. Quant. Grav. **11** (1994) 999–1012.
- [30] M. Reed, B. Simon, *Methods of Modern Mathematical Physics. 2. Fourier Analysis, Selfadjointness*, Gulf Professional Publishing, 1975.

-
- [31] M. Reed, B. Simon, *Methods of Modern Mathematical Physics. 1. Functional Analysis*, Gulf Professional Publishing, 1975.
- [32] T. Kato, *Perturbation theory for linear operators; 2nd ed.*, Grundlehren Math. Wiss., Springer, Berlin, 1976.
- [33] A. Wouk, *A Note on Square Roots of Positive Operators*, SIAM Review **8** (1966), no. 1 100–102.
- [34] Z. Sebestyén, Z. Tarcsay, *On the square root of a positive selfadjoint operator*, Period Math. Hung. **75** (2017), no. 2 268–272.
- [35] S. W. Hawking, A. R. King, P. J. McCarthy, *A New Topology for Curved Space-Time Which Incorporates the Causal, Differential, and Conformal Structures*, J. Math. Phys. **17** (1976) 174–181.
- [36] D.-H. Kim, *A Note on Non-compact Cauchy surface*, Class. Quant. Grav. **25** (2008) 238002, [0801.2671](#).
- [37] R. Wald, *General Relativity*, University of Chicago Press, 2010.
- [38] S. Cotsakis, *Global hyperbolicity of sliced spaces*, Gen. Rel. Grav. **36** (2004) 1183–1188, [gr-qc/0309048](#).
- [39] Y. Choquet-Bruhat, S. Cotsakis, *Global hyperbolicity and completeness*, J. Geom. Phys. **43** (2002) 345–350, [gr-qc/0201057](#).
- [40] K. Papadopoulos, N. Kurt, B. K. Papadopoulos, *On sliced spaces: Global Hyperbolicity revisited*, arXiv e-prints (2019) arXiv:1901.07381, [1901.07381](#).
- [41] B. S. Kay, *The Principle of locality and quantum field theory on (nonglobally hyperbolic) curved space-times*, Rev. Math. Phys. **4** (1992) 167–195.
- [42] R. Oeckl, *General boundary quantum field theory: Foundations and probability interpretation*, Adv. Theor. Math. Phys. **12** (2008) 319–352, [hep-th/0509122](#).
- [43] R. Oeckl, *Schrödinger-Feynman quantization and composition of observables in general boundary quantum field theory*, Adv. Theor. Math. Phys. **19** (2015) 451–506, [1201.1877](#).