

PBZ*-Lattices: Structure Theory and Subvarieties

Roberto Giuntini, Claudia Mureşan, Francesco Paoli
Dept. of Pedagogy, Psychology, Philosophy, University of Cagliari

May 16, 2019

Abstract

We investigate the structure theory of the variety of *PBZ*-lattices* and some of its proper subvarieties. These lattices with additional structure originate in the foundations of quantum mechanics and can be viewed as a common generalisation of orthomodular lattices and Kleene algebras expanded by an extra unary operation. We lay down the basics of the theories of ideals and of central elements in PBZ*-lattices, we prove some structure theorems, and we explore some connections with the theories of subtractive and binary discriminator varieties.

1 Introduction

The papers [14] and [15] contain the beginnings of an algebraic investigation of a variety of lattices with additional structure, the variety \mathbb{PBZL}^* of *PBZ*-lattices*. The key motivation for the introduction of this class of algebras comes from the foundations of quantum mechanics. Consider the structure

$$\mathbf{E}(\mathbf{H}) = (\mathcal{E}(\mathbf{H}), \wedge_s, \vee_s, ', \sim, \mathbb{O}, \mathbb{I}),$$

where:

- $\mathcal{E}(\mathbf{H})$ is the set of all effects of a given complex separable Hilbert space \mathbf{H} , i.e., positive linear operators of \mathbf{H} that are bounded by the identity operator \mathbb{I} ;
- \wedge_s and \vee_s are the meet and the join, respectively, of the *spectral ordering* \leq_s so defined for all $E, F \in \mathcal{E}(\mathbf{H})$:

$$E \leq_s F \text{ iff } \forall \lambda \in \mathbb{R} : M^F(\lambda) \leq M^E(\lambda),$$

where for any effect E , M^E is the unique spectral family [19, Ch. 7] such that $E = \int_{-\infty}^{\infty} \lambda dM^E(\lambda)$ (the integral is here meant in the sense of norm-converging Riemann-Stieltjes sums [22, Ch. 1]);

- \mathbb{O} and \mathbb{I} are the null and identity operators, respectively;
- $E' = \mathbb{I} - E$ and $E^\sim = P_{\ker(E)}$ (the projection onto the kernel of E).

The operations in $\mathbf{E}(\mathbf{H})$ are well-defined. The spectral ordering is indeed a lattice ordering [20, 18] that coincides with the usual ordering of effects induced via the trace functional when both orderings are restricted to the set of projection operators of the same Hilbert space.

A PBZ^* -lattice can be viewed as an abstraction from this concrete physical model, much in the same way as an orthomodular lattice can be viewed as an abstraction from a certain structure of projection operators in a complex separable Hilbert space. The faithfulness of PBZ^* -lattices to the physical model whence they stem is further underscored by the fact that they reproduce at an abstract level the “collapse” of several notions of *sharp physical property* that can be observed in $\mathbf{E}(\mathbf{H})$.

Further motivation for the study of PBZL^* comes from its emerging relationships with many related algebraic structures (orthomodular lattices, Kleene algebras, Stone algebras). In particular, PBZ^* -lattices can be seen as a common generalisation of orthomodular lattices and of Kleene algebras with an additional unary operation.

This paper is devoted to laying down the basics of the structure theory of the variety PBZL^* and of some of its subvarieties; let us briefly summarise its contents. In Section 2 we dispatch a number of preliminaries in order to keep the paper reasonably self-contained, including a short résumé of the results in [14] and [15]. In Section 3, we study decompositions of PBZ^* -lattices. As it happens for orthomodular lattices, and more generally for members of all Church varieties [21], direct decompositions in a PBZ^* -lattice \mathbf{L} are induced by certain members of L (the so-called *central elements*) that form a Boolean algebra and that can be conveniently described. In particular, we show that the central elements in a PBZ^* -lattice \mathbf{L} are those elements that “commute” with any $a \in L$, and that this “commuting” relation generalises the analogous relation of decisive importance in the context of orthomodular lattices. In Section 4, we introduce the notion of a p -ideal (ideal closed under perspectivity), mimicking the corresponding definition available for orthomodular lattices. Although in the general case p -ideals lack many of the strong properties one would expect from a reasonable notion of an ideal, as soon as we zoom in on the subvariety SDM satisfying the strong De Morgan law $(x \wedge y)^\sim \approx x^\sim \vee y^\sim$, we can show that such ideals coincide with the SDM -ideals in the sense of Ursini (whence also with 0 -classes of congruences, since PBZL^* and all its subvarieties are 0 -subtractive). We also prove that the 0 -assertional logic of SDM is strongly algebraisable and we characterise its equivalent variety semantics. Finally, we observe that the variety $V(\text{AOL})$ generated by antiortholattices — that is, PBZ^* -lattices with no nontrivial sharp element — is a binary discriminator variety and we further simplify the description of ideals in that case. In the concluding Section 5, after streamlining the known equational basis for $V(\text{AOL})$, we axiomatise the varietal join of orthomodular lattices and the variety generated by antiortholattices in the lattice of subvarieties of PBZL^* .

2 Preliminaries

2.1 Universal Algebra and Lattice Theory

For basic information on universal algebra, the reader is referred to [6, 17].

Throughout this paper, all algebras will be nonempty; by a *trivial algebra* we will mean a one-element algebra, and a *trivial variety* will be a variety consisting solely of trivial algebras. If \mathbf{A} is an algebra, then A will be the universe of \mathbf{A} ; in some cases, such as those of congruence lattices, lattices will be designated by their set reducts. If \mathbb{V} is a variety of algebras of similarity type ν and \mathbf{A} or a reduct of \mathbf{A} is a member of \mathbb{V} , then $(\text{Con}_{\mathbb{V}}(\mathbf{A}), \cap, \vee, \Delta_A, \nabla_A)$ will be the bounded lattice of the congruences of \mathbf{A} with respect to ν ; when \mathbb{V} is the variety of lattices, $\text{Con}_{\mathbb{V}}(\mathbf{A})$ will be denoted by $\text{Con}(\mathbf{A})$. With \mathbb{V} assumed implicit, the congruence of \mathbf{A} generated by an $S \subseteq A \times A$ will be denoted by $Cg(S)$; for all $a, b \in A$, the principal congruence $Cg(\{(a, b)\})$ will be denoted by $Cg(a, b)$.

For any lattice \mathbf{L} and any $x, y \in L$, the principal filter (resp. ideal) of \mathbf{L} generated by x will be denoted by $[x]$ (resp. $(x]$), and, if $x \leq y$, then $[x, y] = [x] \cap (y]$ will be the interval of \mathbf{L} bounded by x and y . The dual of any (bounded) lattice \mathbf{M} will be denoted by \mathbf{M}^d . If \mathbf{A} is an algebra with a bounded lattice reduct, then such a reduct will be indicated by \mathbf{A}_l . In this case, a congruence θ of \mathbf{A} (or any of its reducts) is said to be *pseudo-identical* iff $0^{\mathbf{A}}/\theta = \{0^{\mathbf{A}}\}$ and $1^{\mathbf{A}}/\theta = \{1^{\mathbf{A}}\}$.

2.2 PBZ* –lattices

We recap in this section some definitions and results on PBZ* –lattices (the latter mostly from [14] and [15], except when explicitly noted) that will be needed in the following.

Definition 1 *A bounded involution lattice is an algebra $\mathbf{L} = (L, \wedge, \vee, ', 0, 1)$ of type $(2, 2, 1, 0, 0)$ such that $(L, \wedge, \vee, 0, 1)$ is a bounded lattice with partial order \leq and the following conditions are satisfied for all $a, b \in L$:*

- $a'' = a$;
- $a \leq b$ implies $b' \leq a'$.

Note that, for any bounded involution lattice \mathbf{L} , the involution $' : L \rightarrow L$ is a dual lattice isomorphism of \mathbf{L}_l .

Definition 2 *A bounded involution lattice $\mathbf{L} = (L, \wedge, \vee, ', 0, 1)$ is a pseudo-Kleene algebra in case it satisfies any of the following two equivalent conditions:*

- (i) *for all $a, b \in L$, if $a \leq a'$ and $b \leq b'$, then $a \leq b'$;*
- (ii) *for all $a, b \in L$, $a \wedge a' \leq b \vee b'$.*

The class of bounded involution lattices is a variety, here denoted by \mathbb{BI} . The involution of a pseudo-Kleene algebra is called *Kleene complement*. The variety of pseudo-Kleene algebras, for which see e.g. [10], is denoted by \mathbb{PKA} . Distributive pseudo-Kleene algebras are variously called *Kleene lattices* or *Kleene algebras* in the literature. Observe that in [14], embracing the terminological usage from [12, p. 12], pseudo-Kleene algebras were referred to as “Kleene lattices”. In [15], however, the authors switched to the less ambiguous “pseudo-Kleene algebras”.

In unsharp quantum logic, there are several competing purely algebraic characterisations of sharp effects [12, Ch. 7]. A quantum effect or property is usually called *sharp* if it satisfies the noncontradiction principle:

Definition 3 *Let \mathbf{L} be a bounded involution lattice.*

- (i) *An element $a \in L$ is said to be Kleene-sharp iff $a \wedge a' = 0$. $S_K(\mathbf{L})$ denotes the class of Kleene-sharp elements of \mathbf{L} .*
- (ii) *\mathbf{L} is an ortholattice iff $S_K(\mathbf{L}) = L$.*
- (iii) *\mathbf{L} is an orthomodular lattice iff \mathbf{L} is an ortholattice and, for all $a, b \in L$, if $a \leq b$, then $b = (b \wedge a') \vee a$.*

The variety of ortholattices is denoted by \mathbb{OL} . Among ortholattices, orthomodular lattices play a crucial role in the standard (sharp) approach to quantum logic. The class of orthomodular lattices is actually a variety, hereafter denoted by \mathbb{OML} .

It is well-known that an ortholattice \mathbf{L} is orthomodular if and only if, for all $a, b \in L$, if $a \leq b$ and $a' \wedge b = 0$, then $a = b$. In the wider setting of bounded involution lattices, the previous condition does not imply the stronger condition of orthomodularity above. We will call this weaker condition *paraorthomodularity*.

Definition 4 *An algebra \mathbf{L} with a bounded involution lattice reduct is said to be paraorthomodular iff, for all $a, b \in L$:*

$$\text{if } a \leq b \text{ and } a' \wedge b = 0, \text{ then } a = b.$$

It turns out that the class of paraorthomodular pseudo-Kleene algebras is a proper quasivariety, whence we cannot help ourselves to the strong universal algebraic properties that characterise varieties. It is then natural to wonder whether there exists an *expansion* of the language of bounded involution lattices where the paraorthomodular condition can be equationally recovered. The appropriate language expansion is provided by including an additional unary operation and moving to the type $(2, 2, 1, 1, 0, 0)$, familiar in unsharp quantum logic from the investigation of *Brouwer-Zadeh lattices* (see [9] or [12, Ch. 4.2]).

Definition 5 *A Brouwer-Zadeh lattice (or BZ-lattice) is an algebra*

$$\mathbf{L} = (L, \wedge, \vee, ', \sim, 0, 1)$$

of type $(2, 2, 1, 1, 0, 0)$, such that:

(i) $(L, \wedge, \vee, ', 0, 1)$ is a pseudo-Kleene algebra;

(ii) for all $a, b \in L$, the following conditions are satisfied:

$$\begin{aligned} (1) \quad & a \wedge a^\sim = 0; & (2) \quad & a \leq a^{\sim\sim}; \\ (3) \quad & a \leq b \text{ implies } b^\sim \leq a^\sim; & (4) \quad & a^{\sim'} = a^{\sim\sim}. \end{aligned}$$

The operation \sim is called the *Brouwer complement* of the BZ-lattice. The class of all BZ-lattices is a variety, denoted by \mathbb{BZL} ; \mathbb{OL} can be identified with the subvariety of \mathbb{BZL} whose relative equational basis w.r.t. \mathbb{BZL} is given by the equation $x^\sim = x'$. In any BZ-lattice, we set $\diamond x = x^{\sim\sim}$ and $\square x = x'^{\sim}$. The following arithmetical lemma, the proof of which is variously scattered in the above-mentioned literature and elsewhere [7, 8], will be used without being referenced in what follows.

Lemma 6 *Let \mathbf{L} be a BZ-lattice. For all $a, b \in L$, the following conditions hold:*

$$\begin{aligned} (i) \quad & a^{\sim\sim\sim} = a^\sim; & (vi) \quad & \square(a \wedge b) = \square a \wedge \square b; \\ (ii) \quad & a^\sim \leq a'; & (vii) \quad & \diamond(a \vee b) = \diamond a \vee \diamond b; \\ (iii) \quad & (a \vee b)^\sim = a^\sim \wedge b^\sim; & (viii) \quad & \diamond(a \wedge b) \leq \diamond a \wedge \diamond b; \\ (iv) \quad & a^\sim \vee b^\sim \leq (a \wedge b)^\sim; & (ix) \quad & \text{if } a' \leq a, \text{ then } a^\sim = 0. \\ (v) \quad & (\square(a'))' = \diamond a; \end{aligned}$$

We remarked above that Kleene-sharpness is not the unique purely algebraic characterisation of a sharp quantum property. Two noteworthy alternatives now become available in our expanded language of BZ-lattices.

Definition 7 *Let \mathbf{L} be a BZ-lattice.*

(i) *An element $a \in L$ is said to be \diamond -sharp iff $a = \diamond a$; the class of all \diamond -sharp elements of \mathbf{L} will be denoted by $S_\diamond(\mathbf{L})$.*

(ii) *An element $a \in L$ is said to be Brouwer-sharp iff $a \vee a^\sim = 1$; the class of all Brouwer-sharp elements of \mathbf{L} will be denoted by $S_B(\mathbf{L})$.*

It is easy to derive from the previous lemma that, in any BZ-lattice \mathbf{L} , $S_\diamond(\mathbf{L}) = \{a^\sim : a \in L\} = \{a \in L : a' = a^\sim\}$. For any BZ-lattice \mathbf{L} , we have that $S_\diamond(\mathbf{L}) \subseteq S_B(\mathbf{L}) \subseteq S_K(\mathbf{L})$. However, in any BZ-lattice of effects of a Hilbert space (under the meet and join operation induced by the spectral ordering) these three classes coincide. Consequently, it makes sense to investigate whether there is a class of BZ-lattices for which this collapse result can be recovered at a purely abstract level. The next definition and theorem answer this question in the affirmative.

Definition 8 *A BZ*-lattice is a BZ-lattice \mathbf{L} that satisfies, for all $a \in L$, the condition*

$$(*) \quad (a \wedge a')^\sim \leq a^\sim \vee \square a.$$

Theorem 9 *Let \mathbf{L} be a paraorthomodular BZ*-lattice. Then,*

$$S_\diamond(\mathbf{L}) = S_B(\mathbf{L}) = S_K(\mathbf{L}).$$

As pleasing as this result may be, the class of paraorthomodular BZ*-lattices still suffers from a major shortcoming: the paraorthomodularity condition is quasiequational. However, the next result shows that it can be replaced by an equation, so that paraorthomodular BZ*-lattices form a variety, which we will denote by \mathbb{PBZL}^* and whose members will be called, in brief, *PBZ*-lattices*.

Theorem 10 *Let \mathbf{L} be a BZ*-lattice. The following conditions are equivalent:*

- (1) \mathbf{L} is paraorthomodular;
- (2) \mathbf{L} satisfies the following \diamond -orthomodularity condition for all $a, b \in L$:

$$(a^\sim \vee (\diamond a \wedge \diamond b)) \wedge \diamond a \leq \diamond b.$$

Every bounded lattice can be embedded as a sublattice into a PBZ*-lattice [14, Lm. 5.3]. Consequently, \mathbb{PBZL}^* satisfies no nontrivial identity in the language of lattices.

The naturalness of the concept of a PBZ*-lattice is further reinforced by the circumstance that BZ-lattices of effects of a Hilbert space, under the spectral ordering, qualify as instances of PBZ*-lattices:

Theorem 11 *Let \mathbf{H} be a complex separable Hilbert space. The algebra*

$$\mathbf{E}(\mathbf{H}) = \langle \mathcal{E}(\mathbf{H}), \wedge_s, \vee_s, ', \sim, \mathbb{O}, \mathbb{I} \rangle,$$

(see the introduction for the notation) is a PBZ-lattice. Moreover,*

$$S_K(\mathbf{E}(\mathbf{H})) = S_\diamond(\mathbf{E}(\mathbf{H})) = S_B(\mathbf{E}(\mathbf{H}))$$

is an orthomodular subuniverse of $\mathbf{E}(\mathbf{H})$ consisting of all the projection operators of \mathbf{H} .

All orthomodular lattices become, of course, PBZ*-lattices when endowed with a Brouwer complement that equals their Kleene complement. In every PBZ*-lattice \mathbf{L} , $S_K(\mathbf{L})$ is always the universe of the largest orthomodular subalgebra $\mathbf{S}_K(\mathbf{L})$ of \mathbf{L} , so that \mathbf{L} is orthomodular iff it satisfies $x^\sim \approx x'$. Further examples of PBZ*-lattices are given by those algebras in this class that are “as far apart as possible” from orthomodular lattices. In any orthomodular lattice \mathbf{L} , $S_K(\mathbf{L}) = L$; on the other hand, by definition, a PBZ*-lattice \mathbf{L} is an *antiortholattice* iff $S_K(\mathbf{L}) = \{0, 1\}$. We denote by \mathbb{AOL} the class of antiortholattices.

Lemma 12

- (i) *A PBZ*-lattice \mathbf{L} belongs to \mathbb{AOL} iff $0^\sim = 1$ and, for all $a \in L \setminus \{0\}$, $a^\sim = 0$.*
- (ii) *Every $\mathbf{L} \in \mathbb{AOL}$ is directly indecomposable.*

(iii) AOL is a proper universal class.

The Brouwer complement of Lemma 12.(i) is called *trivial*.

For all $n \geq 1$, the n -element Kleene chain with universe $D_n = \{0, d_1, d_2, \dots, d_{n-2}, 1\}$, with $0 < d_1 < d_2 < \dots < d_{n-2} < 1$, is an antiortholattice \mathbf{D}_n under the trivial Brouwer complement. To avoid notational overloading, the reduct $(\mathbf{D}_n)_l$ will simply be denoted by \mathbf{D}_n , as well. Note that every finite chain is self-dual both as a bounded lattice and as a Kleene algebra, so the notation \mathbf{D}_n^d is superfluous in these cases; the same can be stated about direct products of finite chains, in particular about Boolean algebras.

The following easy results are observed (sometimes implicitly) in the literature on BZ-lattices, in particular in [14] and in [15]:

Lemma 13 (i) *Any pseudo-Kleene algebra, endowed with the trivial Brouwer complement, becomes a BZ-lattice.*

(ii) *Any paraorthomodular pseudo-Kleene algebra which, endowed with the trivial Brouwer complement, satisfies condition (*), becomes an antiortholattice.*

(iii) *Any pseudo-Kleene algebra in which 0 is meet-irreducible is paraorthomodular and satisfies condition (*) when endowed with the trivial Brouwer complement, whence it becomes an antiortholattice.*

We will repeatedly have the occasion to consider the following identities in the language of BZ-lattices:

SDM (the *Strong de Morgan law*) $(x \wedge y)^\sim \approx x^\sim \vee y^\sim$;

WSDM (*weak SDM*) $(x \wedge y^\sim)^\sim \approx x^\sim \vee \diamond y$;

DIST $x \wedge (y \vee z) \approx (x \wedge y) \vee (x \wedge z)$;

J2 $x \approx (x \wedge (y \wedge y')^\sim) \vee (x \wedge \diamond (y \wedge y'))$;

SK $x \wedge \diamond y \leq \square x \vee y$.

Clearly, SDM implies WSDM. Observe that OML satisfies SDM, J2 and SK. Trivially, AOL satisfies WSDM and J2, whence $\text{OML} \vee V(\text{AOL})$ satisfies these two identities.

We list some useful properties of the variety $V(\text{AOL})$ generated by antiortholattices, including an axiomatisation relative to PBZL^* .

Lemma 14 *Let $\mathbf{A} \in V(\text{AOL})$. Then for all $a, b, c \in A$:*

(i) $a \wedge (b \vee c^\sim) = (a \wedge b) \vee (a \wedge c^\sim)$;

(ii) $a \vee (b \wedge c^\sim) = (a \vee b) \wedge (a \vee c^\sim)$;

(iii) $a^\sim \wedge (b \vee c) = (a^\sim \wedge b) \vee (a^\sim \wedge c)$;

(iv) $a^\sim \vee (b \wedge c) = (a^\sim \vee b) \wedge (a^\sim \vee c)$.

Theorem 15 (i) *An equational basis for $V(\mathbf{AOL})$ relative to \mathbf{PBZL}^* is given by the identities*

$$\begin{aligned} (AOL1) \quad & (x^\sim \vee y^\sim) \wedge (\diamond x \vee z^\sim) \approx ((x^\sim \vee y) \wedge (\diamond x \vee z))^\sim; \\ (AOL2) \quad & x \approx (x \wedge y^\sim) \vee (x \wedge \diamond y); \\ (AOL3) \quad & x \approx (x \vee y^\sim) \wedge (x \vee \diamond y). \end{aligned}$$

(ii) *Every subdirectly irreducible member of $V(\mathbf{AOL})$ is an antiortholattice.*

Clearly $V(\mathbf{AOL}) \cap \mathbf{OML}$ is the variety \mathbf{BA} of Boolean algebras.

The lattice $\mathbf{L}_{\mathbf{PBZL}^*}$ of subvarieties of \mathbf{PBZL}^* has \mathbf{BA} as a unique atom. It is well-known that \mathbf{BA} has a single orthomodular cover [5, Cor. 3.6]: the variety $V(\mathbf{MO}_2)$, generated by the simple modular ortholattice with 4 atoms. Moreover:

Theorem 16 *There is a single non-orthomodular cover of \mathbf{BA} in $\mathbf{L}_{\mathbf{PBZL}^*}$, the variety $V(\mathbf{D}_3)$ generated by the 3-element antiortholattice chain, whose equational basis relative to $V(\mathbf{AOL})$ is given by the identity SK .*

Two other notable subvarieties of $V(\mathbf{AOL})$ are the variety \mathbf{DIST} , whose equational basis relative to $V(\mathbf{AOL})$ (or, equivalently, relative to \mathbf{PBZL}^*) is given by the distribution identity \mathbf{DIST} , and the variety \mathbf{SAOL} , whose equational basis relative to $V(\mathbf{AOL})$ is given by the Strong De Morgan identity \mathbf{SDM} . We have that:

Theorem 17 $V(\mathbf{D}_5) = \mathbf{DIST} \cap \mathbf{SAOL}$.

A more circumscribed study of \mathbf{DIST} and its subvarieties, also focussing on the relationship with known classes of algebras (including Kleene-Stone algebras Łukasiewicz algebras, Heyting-Wajsberg algebras) is currently in preparation [16]

2.3 Subtractive Varieties

Subtractive varieties were introduced by Ursini [23] to enucleate the common features of pointed varieties with a good ideal theory, like groups, rings or Boolean algebras. They were further investigated in [1, 2, 3, 24].

Definition 18 *Let \mathbb{V} a variety of type ν , and let 0 be a nullary term (or equationally definable constant) of type ν . \mathbb{V} is called 0-subtractive if there exists a binary term s , also of type ν , s.t. \mathbb{V} satisfies the identities $s(x, x) \approx 0$ and $s(x, 0) \approx x$. A variety of type ν which is 0-subtractive w.r.t. at least one constant 0 of type ν is called subtractive tout court.*

It is not hard to see that subtractivity is a congruence property: namely, a variety \mathbb{V} is 0-subtractive exactly when in each $\mathbf{A} \in \mathbb{V}$ congruences permute at 0 (meaning that for all θ, φ in $\mathbf{Con}_{\mathbb{V}}(\mathbf{A})$, $0^{\mathbf{A}}/(\theta \circ \varphi) = 0^{\mathbf{A}}/(\varphi \circ \theta)$).

To investigate ideals in this context, first and foremost, we need a workable general notion of ideal encompassing all the intended examples mentioned above (normal subgroups of groups, two-sided ideals of rings, ideals or filters of Boolean algebras). Ursini's candidate for playing this role is defined below.

Definition 19 (i) If \mathbb{K} is a class of similar algebras whose type ν is as in Definition 18, then a term $p(\vec{x}, \vec{y})$ of type ν is a \mathbb{K} -ideal term in \vec{x} iff $\mathbb{K} \models p(0, \dots, 0, \vec{y}) \approx 0$.

(ii) A nonempty subset J of the universe of an $\mathbf{A} \in \mathbb{K}$ is a \mathbb{K} -ideal of \mathbf{A} (w.r.t. 0) iff for any \mathbb{K} -ideal term $p(\vec{x}, \vec{y})$ in \vec{x} we have that $p^{\mathbf{A}}(\vec{a}, \vec{b}) \in J$ whenever $\vec{a} \in J$ and $\vec{b} \in A$.

We will denote by $\mathcal{I}_{\mathbb{K}}(\mathbf{A})$ the set (or the lattice) of all \mathbb{K} -ideals of \mathbf{A} , dropping the subscript whenever \mathbb{K} can be contextually identified; observe that $\{0\}, A \in \mathcal{I}_{\mathbb{K}}(\mathbf{A})$. The main reason that backs our previous claim to the effect that subtractive varieties have a good ideal theory is given by the following result. Let \mathbb{V} be a variety of type ν . Recall that an algebra \mathbf{A} from \mathbb{V} is said to be 0-regular iff the map sending a congruence $\theta \in \text{Con}(\mathbf{A})$ to its 0-class $0^{\mathbf{A}}/\theta$ is injective. The variety \mathbb{V} is said to be 0-regular iff every $\mathbf{A} \in \mathbb{V}$ is 0-regular; this happens exactly when there exists a finite family of binary ν -terms (called *Fichtner terms*) $\{d_i(x, y)\}_{i \leq n}$ such that $\mathbb{V} \models d_1(x, y) \approx 0 \& \dots \& d_n(x, y) \approx 0 \Leftrightarrow x \approx y$.

Theorem 20 (i) Subtractive varieties have normal ideals. That is, if \mathbb{V} is a 0-subtractive variety and $\mathbf{A} \in \mathbb{V}$, then

$$\mathcal{I}_{\mathbb{V}}(\mathbf{A}) = \{I \subseteq A : I = 0^{\mathbf{A}}/\theta \text{ for some } \theta \in \text{Con}_{\mathbb{V}}(\mathbf{A})\}.$$

(ii) If \mathbb{V} is a 0-subtractive and 0-regular variety, then, for every $\mathbf{A} \in \mathbb{V}$, $\text{Con}_{\mathbb{V}}(\mathbf{A})$ is isomorphic to $\mathcal{I}_{\mathbb{V}}(\mathbf{A})$.

Actually, the situation described by the previous theorem can be made more precise as follows. Let \mathbf{A} be an algebra in a 0-subtractive variety \mathbb{V} . Then the following maps are well-defined: $\cdot^{\delta}, \cdot^{\varepsilon} : \mathcal{I}_{\mathbb{V}}(\mathbf{A}) \rightarrow \text{Con}_{\mathbb{V}}(\mathbf{A})$, for all $I \in \mathcal{I}_{\mathbb{V}}(\mathbf{A})$,

$$\begin{aligned} I^{\delta} &= \bigwedge \{\theta \in \text{Con}_{\mathbb{V}}(\mathbf{A}) : 0^{\mathbf{A}}/\theta = I\}, \\ I^{\varepsilon} &= \bigvee \{\theta \in \text{Con}_{\mathbb{V}}(\mathbf{A}) : 0^{\mathbf{A}}/\theta = I\}. \end{aligned}$$

Henceforth, all unnecessary superscripts will be dropped for the sake of conciseness. Note that, for all $I \in \mathcal{I}_{\mathbb{V}}(\mathbf{A})$, we have $0/I^{\delta} = 0/I^{\varepsilon} = I$, so that $I^{\delta} = \min\{\theta \in \text{Con}_{\mathbb{V}}(\mathbf{A}) : 0/\theta = I\}$ and $I^{\varepsilon} = \max\{\theta \in \text{Con}_{\mathbb{V}}(\mathbf{A}) : 0/\theta = I\}$. Moreover, the map $I \mapsto [I^{\delta}, I^{\varepsilon}]$ is a lattice isomorphism from $\mathcal{I}_{\mathbb{V}}(\mathbf{A})$ to the following lattice of intervals of $\text{Con}_{\mathbb{V}}(\mathbf{A})$: $\{[\min(C_{\theta}), \max(C_{\theta})] : \theta \in \text{Con}_{\mathbb{V}}(\mathbf{A})\}$, where, for all $\theta \in \text{Con}_{\mathbb{V}}(\mathbf{A})$, $C_{\theta} = \{\alpha \in \text{Con}_{\mathbb{V}}(\mathbf{A}) : 0/\alpha = 0/\theta\} = \{\alpha \in \text{Con}_{\mathbb{V}}(\mathbf{A}) : 0/\theta \in A/\alpha\}$, which is a complete sublattice of $\text{Con}_{\mathbb{V}}(\mathbf{A})$. Clearly, if \mathbb{V} is in addition 0-regular, then $C_{\theta} = \{\theta\}$ for all $\theta \in \text{Con}_{\mathbb{V}}(\mathbf{A})$, hence all intervals of the form $[I^{\delta}, I^{\varepsilon}]$ for some $I \in \mathcal{I}_{\mathbb{V}}(\mathbf{A})$ are trivial, and Theorem 20.(ii) follows as a special case.

Definition 21 Let \mathbf{A} be an algebra in a 0-subtractive variety \mathbb{V} . \mathbf{A} is said to be reduced iff $\{0\}^\varepsilon = \Delta_A$.

The class of all reduced algebras in \mathbb{V} will be denoted by \mathbb{V}_ε . Clearly, for all $\theta \in \text{Con}_\mathbb{V}(\mathbf{A})$, we have $\mathbf{A}/\theta \in \mathbb{V}_\varepsilon$ iff $\theta = I^\varepsilon$ for some $I \in \mathcal{I}_\mathbb{V}(\mathbf{A})$.

For a 0-subtractive variety, a property that generalises 0-regularity is finite congruentiality. Roughly put, a 0-subtractive variety is finitely congruential if it has a family of terms that do “part of the job” usually dispatched by the Fichtner terms for 0-regularity.

Definition 22 A variety \mathbb{V} , whose type ν is as in Definition 18, is finitely congruential iff there exists a finite set $\{d_i(x, y)\}_{i \leq n}$ of binary ν -terms s.t., whenever $\mathbf{A} \in \mathbb{V}$ and $I \in \mathcal{I}_\mathbb{V}(\mathbf{A})$, we have:

$$I^\varepsilon = \{(a, b) : d_i^{\mathbf{A}}(a, b) \in I \text{ for all } i \leq n\}.$$

The next results establish a significant connection between the theory of subtractive varieties and *abstract algebraic logic* (for information on this research area, the reader is referred to [13]). It turns out that, for a 0-subtractive variety \mathbb{V} , the properties of the 0-assertional logic of \mathbb{V} yield relevant information on the properties of the class of reduced algebras in \mathbb{V} , and conversely.

Theorem 23 If \mathbf{A} is a member of a 0-subtractive variety \mathbb{V} , then $\mathcal{I}_\mathbb{V}(\mathbf{A})$ is the class of all deductive filters on \mathbf{A} of the 0-assertional logic of \mathbb{V} , and for all $I \in \mathcal{I}_\mathbb{V}(\mathbf{A})$, $I^\varepsilon = \Omega^{\mathbf{A}}(I)$.

Theorem 24 [2, Thm. 3.12] For a 0-subtractive variety \mathbb{V} the following are equivalent:

- (i) The 0-assertional logic of \mathbb{V} is equivalential.
- (ii) \mathbb{V}_ε is closed under subalgebras and direct products.

Theorem 25 [2, Thm. 3.16] For a 0-subtractive variety \mathbb{V} the following are equivalent:

- (i) The 0-assertional logic of \mathbb{V} is strongly algebraisable with \mathbb{V}_ε as an equivalent algebraic semantics.
- (ii) \mathbb{V}_ε is a variety.

Important examples of subtractive varieties are *binary discriminator varieties*. Recall that a *discriminator variety* [25] is a variety \mathbb{V} of given type ν for which there exists a ternary ν -term $t(x, y, z)$ that realises the ternary discriminator function

$$t(a, b, c) = \begin{cases} c & \text{if } a = b, \\ a, & \text{otherwise} \end{cases}$$

on any subdirectly irreducible member of \mathbb{V} (equivalently, on any member of some class \mathbb{K} such that $\mathbb{V} = V(\mathbb{K})$). The introduction of *binary discriminator*

varieties by Chajda, Halaš, and Rosenberg [11] was aimed at singling out an appropriate weakening of the ternary discriminator that can vouchsafe some of the strong properties of discriminator varieties (like congruence distributivity or congruence permutability) only “locally”, i.e. at 0. Binary discriminator varieties are paramount among subtractive varieties with *equationally definable principal ideals* [1]; they were thoroughly studied in the unpublished [4].

Definition 26 [11] *Let A be a nonempty set and fix $0 \in A$. The 0-binary discriminator on A is the binary function b_0^A on A defined by:*

$$b_0^A(a, c) = \begin{cases} a & \text{if } c = 0, \\ 0 & \text{otherwise.} \end{cases}$$

An algebra \mathbf{A} with a term definable element 0 is said to be a 0-binary discriminator algebra in case the 0-binary discriminator b_0^A on A is a term operation on \mathbf{A} . A variety $V(\mathbb{K})$ is a 0-binary discriminator variety if it is generated by a class \mathbb{K} of 0-binary discriminator algebras such that the property is witnessed by the same terms for all members of \mathbb{K} .

3 Central Elements

One of the most distinctive and far-reaching chapters in the theory of orthomodular lattices is the study of the commuting relation and of central elements (see e.g. [5, § 2]). Given an orthomodular lattice \mathbf{L} and $a, b \in L$, a is said to *commute* with b in case $(a \wedge b) \vee (a' \wedge b) = b$. Such a relation is reflexive and symmetric. An element $a \in L$ is said to be *central* in \mathbf{L} in case it commutes with all elements of L . The next celebrated result is one of the most useful tools for practitioners of the field:

Theorem 27 (Foulis-Holland) [5, Prop. 2.8] *If \mathbf{L} is an orthomodular lattice and $a, b, c \in L$ are such that a commutes both with b and with c , then the set $\{a, b, c\}$ generates a distributive sublattice of \mathbf{L}_1 .*

Although these investigations were carried out in the special context of orthomodular lattices, the notion of a central element is deeply rooted in universal algebra. Recall that, if \mathbf{A} is an algebra in a double-pointed variety \mathbb{V} with constants $0, 1$, an element $e \in A$ is *central* in \mathbf{A} in case the congruences $Cg(e, 0)$ and $Cg(e, 1)$ are complementary factor congruences of \mathbf{A} [21]. By $C(\mathbf{A})$ we denote the *centre* of A , i.e. the set of central elements of the algebra \mathbf{A} . In particular, if \mathbf{A} is a *Church algebra* [21], namely, if there is an “if-then-else” term operation $q^{\mathbf{A}}$ on \mathbf{A} s.t., for all $a, b \in A$, $q^{\mathbf{A}}(1^{\mathbf{A}}, a, b) = a$ and $q^{\mathbf{A}}(0^{\mathbf{A}}, a, b) = b$, then, by defining

$$x \wedge y = q(x, y, 0), \quad x \vee y = q(x, 1, y) \quad \text{and} \quad x' = q(x, 0, 1),$$

we get:

Theorem 28 [21, Thm. 3.7] *The algebra $c[\mathbf{A}] = (C(\mathbf{A}), \wedge, \vee, ', 0, 1)$ is a Boolean algebra which is isomorphic to the Boolean algebra of factor congruences of \mathbf{A} .*

In Church algebras, central elements can be equationally characterised [21, Prop. 3.6]. When studying a specific variety \mathbb{V} , however, more informative descriptions of central elements in members of \mathbb{V} can sometimes be found either in terms of certain properties of intrinsic interest, or by appropriately streamlining the above-mentioned equational characterisation. As regards members of \mathbb{OML} , for example, it can be shown that the two definitions of central element we have given above are equivalent. For \mathbb{PBZL}^* , on the other hand, a more economical equational description of central elements was provided in [14, Lm. 5.9]. We reproduce this lemma for the reader’s convenience:

Lemma 29 *Let $\mathbf{L} \in \mathbb{PBZL}^*$. Then $e \in L$ is central in \mathbf{L} iff it satisfies the following conditions for all $a, b \in L$:*

- C1** $a = (e \wedge a) \vee (e' \wedge a)$;
- C2** $(a \vee b) \wedge e = (a \wedge e) \vee (b \wedge e)$;
- C3** $(a \vee b) \wedge e' = (a \wedge e') \vee (b \wedge e')$;
- C4** $((e \vee b) \wedge (e' \vee a))^\sim = (e \vee b^\sim) \wedge (e' \vee a^\sim)$.

The aim of the next subsection is to improve on this result, giving a description of the centre in a generic \mathbb{PBZ}^* -lattice that resembles as closely as possible the one we have in orthomodular lattices.

3.1 Central Elements in \mathbb{PBZ}^* -lattices

Throughout the rest of this subsection, unless mentioned otherwise, \mathbf{L} will be an arbitrary \mathbb{PBZ}^* -lattice. Observe that $C(\mathbf{L}) \subseteq S_K(\mathbf{L})$, by C1 for $a = 1$. Since $e' = e^\sim$ for all $e \in S_K(\mathbf{L})$, it follows that, for all $e \in L$: $e \in C(\mathbf{L})$ iff $e^\sim \in C(\mathbf{L})$. In [14, Thm. 5.4] it is also proved that central elements in any member \mathbf{L} of $V(\mathbb{AOL})$ are exactly the members of $S_K(\mathbf{L})$. The problem of finding a manageable description of central elements in \mathbb{PBZ}^* -lattices, that relates in a perspicuous way to the notions of a centre and of a commutator in orthomodular lattice, was however left open. We now set about filling this gap. We show that the property of being central, in a generic \mathbb{PBZ}^* -lattice, is actually two-sided. In order to belong to $C(\mathbf{L})$, an $e \in L$ should not only “commute” with any element of L , in a sense of commuting that appropriately generalises the corresponding (reflexive and symmetric) relation on orthomodular lattices; but it should also be such that, for any $a \in L$, $(e \wedge a)^\sim = e^\sim \vee a^\sim$ and $(e' \wedge a)^\sim = \square e \vee a^\sim$. This latter component is sort of “hidden” in the case of \mathbb{OML} , where the Strong De Morgan identity is satisfied across the board.

We define the following binary relations on L :

$$C_{\mathbf{L}} = \{(a, b) \in L^2 : (a \wedge b) \vee (a' \wedge b) = b\};$$

$$C_{\text{SDM}, \mathbf{L}} = C_{\mathbf{L}} \cap \{(a, b) \in L^2 : (a \wedge b)^\sim = a^\sim \vee b^\sim \text{ and } (a' \wedge b)^\sim = \Box a \vee b^\sim\}.$$

Both relations are clearly reflexive. Also, for all $a \in L$, $(1, a) \in C_{\text{SDM}, \mathbf{L}} \subseteq C_{\mathbf{L}}$, while the following four conditions are mutually equivalent: i) $(a, 1) \in C_{\text{SDM}, \mathbf{L}}$; ii) $(a, 1) \in C_{\mathbf{L}}$; iii) $a \vee a' = 1$; iv) $a \in S_K(\mathbf{L})$. Hence, $C_{\text{SDM}, \mathbf{L}}$ is symmetric exactly when $C_{\mathbf{L}}$ is symmetric, which in turn obtains if and only if \mathbf{L} is orthomodular — in which case, apparently, the two relations coincide.

Definition 30 Let $\mathbf{L} \in \text{PBZL}^*$. An element $a \in L$ is said to be PBZ* -central iff $(a, b) \in C_{\text{SDM}, \mathbf{L}}$ for all $b \in L$.

We will denote by $C_{pbz}(\mathbf{L})$ the set of all PBZ* -central elements of \mathbf{L} :

$$C_{pbz}(\mathbf{L}) = \{a \in L : (\forall b \in L) ((a, b) \in C_{\text{SDM}, \mathbf{L}})\}.$$

We also consider the following subset of L :

$$C_p(\mathbf{L}) = \{a \in L : (\forall b \in L) ((a, b) \in C_{\mathbf{L}})\}.$$

In virtue of the above,

$$C_{pbz}(\mathbf{L}) = C_p(\mathbf{L}) \cap \{a \in L : (\forall b \in L) ((a \wedge b)^\sim = a^\sim \vee b^\sim, (a' \wedge b)^\sim = \Box a \vee b^\sim)\}$$

$$= C_p(\mathbf{L}) \cap \{a \in S_K(\mathbf{L}) : (\forall b \in L) ((a \wedge b)^\sim = a^\sim \vee b^\sim, (a' \wedge b)^\sim = \Box a \vee b^\sim)\}$$

$$\subseteq C_p(\mathbf{L}) \subseteq S_K(\mathbf{L}).$$

Lemma 31 Let $\mathbf{L} \in \text{PBZL}^*$. Then:

- (i) $C_p(\mathbf{L}) = \{a \in S_K(\mathbf{L}) : (\forall b \in L) ((a \wedge b) \vee (a^\sim \wedge b) = b)\} = \{a \in S_K(\mathbf{L}) : (\forall b \in L) ((a \vee b) \wedge (a^\sim \vee b) = b)\};$
- (ii) for all $a \in S_K(\mathbf{L})$, $a \in C_p(\mathbf{L})$ iff $a^\sim \in C_p(\mathbf{L})$; furthermore, $a \in C_{pbz}(\mathbf{L})$ iff $a^\sim \in C_{pbz}(\mathbf{L})$;
- (iii) if \mathbf{L} satisfies WSDM, then $C_{pbz}(\mathbf{L}) = C_p(\mathbf{L})$.

Proof. Let $a \in S_K(\mathbf{L})$, arbitrary.

(i) The fact that $a' = a^\sim$ gives us the first equality. To obtain the second equality, notice that, for all $b \in L$, we have: $(a, b) \in C_{\mathbf{L}}$ iff $(a \wedge b) \vee (a' \wedge b) = b$ iff $(a' \vee b') \wedge (a \vee b') = b'$ iff $(a \vee b') \wedge (a^\sim \vee b') = b'$, hence: $(a, b) \in C_{\mathbf{L}}$ for all $b \in L$ iff $(a \vee b) \wedge (a^\sim \vee b) = b$ for all $b \in L$.

(ii) From (i), along with the fact that $a = \diamond a$.

(iii) From the fact that $C_p(\mathbf{L}) \subseteq S_K(\mathbf{L})$. ■

Note from this proof that $C_{\mathbf{L}}$ and $C_{\text{SDM}, \mathbf{L}}$ preserve the Kleene complement and that, under WSDM, the relation $C_{\text{SDM}, \mathbf{L}} \cap (S_K(\mathbf{L}) \times L) = C_{\mathbf{L}} \cap (S_K(\mathbf{L}) \times L)$ preserves the Brouwer complement. Some useful properties of members of $C_p(\mathbf{L})$ follow in the next two lemmas.

Lemma 32 Let $\mathbf{L} \in \mathbb{PBZL}^*$ and let $a \in L$ be such that $a^\sim \in C_p(\mathbf{L})$. Then, for all $b, c \in L$:

- (i) $a^\sim \vee b = a^\sim \vee (\diamond a \wedge b)$ and $a^\sim \wedge b = a^\sim \wedge (\diamond a \vee b)$;
- (ii) $b \vee (c \wedge \diamond a) = (b \vee c \vee a^\sim) \wedge (b \vee \diamond a)$ and $b \wedge (c \vee \diamond a) = (b \wedge c \wedge a^\sim) \vee (b \wedge \diamond a)$;
- (iii) $a^\sim \wedge (b \vee c) = a^\sim \wedge (b \vee (a^\sim \wedge c))$ and $a^\sim \vee (b \wedge c) = a^\sim \vee (b \wedge (a^\sim \vee c))$.

Proof. By adapting the corresponding proofs in [14, Lm. 5.10]. ■

Lemma 33 Let $\mathbf{L} \in \mathbb{PBZL}^*$, and let $a \in L$ be such that $a^\sim \in C_p(\mathbf{L})$. Then the following hold for all $b, c \in L$:

- (i) $b \vee (c \wedge a^\sim) = (b \vee c) \wedge (b \vee a^\sim)$ and $b \wedge (c \vee a^\sim) = (b \wedge c) \vee (b \wedge a^\sim)$;
- (ii) $a^\sim \wedge (b \vee c) = (a^\sim \wedge b) \vee (a^\sim \wedge c)$ and $a^\sim \vee (b \wedge c) = (a^\sim \vee b) \wedge (a^\sim \vee c)$.

Proof. We prove the first equalities in each pair. The lattice duals are derived similarly.

(i) Let $x = b \vee (c \wedge a^\sim)$. Then:

$$\begin{aligned}
x &= x \wedge (x \vee (c \wedge (x \vee \diamond a))) && \text{Absorption} \\
&= (x \vee a^\sim) \wedge (x \vee \diamond a) \wedge (x \vee (c \wedge (x \vee \diamond a))) && \text{Lemma 31} \\
&= (x \vee a^\sim) \wedge (x \vee (c \wedge (x \vee \diamond a))) && \text{Lattice prop.} \\
&= (b \vee a^\sim) \wedge (x \vee (c \wedge (x \vee \diamond a))) && \text{Absorption} \\
&= (b \vee a^\sim) \wedge (x \vee c) \\
&= (b \vee c) \wedge (b \vee a^\sim) && \text{Absorption}
\end{aligned}$$

The penultimate equality is obtained by observing that, according to Lemma 32.(i), since $\diamond(a^\sim) = a^\sim$, we have: $c \leq b \vee c \vee \diamond a = b \vee \diamond a \vee (a^\sim \wedge c) = x \vee \diamond a$.

(ii)

$$\begin{aligned}
a^\sim \wedge (b \vee c) &= a^\sim \wedge (b \vee (a^\sim \wedge c)) && \text{Lemma 32.(iii)} \\
&= ((a^\sim \wedge c) \vee a^\sim) \wedge (b \vee (a^\sim \wedge c)) && \text{Absorption} \\
&= (a^\sim \wedge b) \vee (a^\sim \wedge c) && \text{(i)}
\end{aligned}$$

■

Theorem 34 Let $\mathbf{L} \in \mathbb{PBZL}^*$. Then $C_{pbz}(\mathbf{L}) = C(\mathbf{L})$.

Proof. Let $e \in L$.

Assume that $e^\sim \in C_{pbz}(\mathbf{L}) \subseteq C_p(\mathbf{L})$. We verify all the conditions C1 through C4 in Lemma 29. C1 holds because $(e^\sim, a) \in C_{SDM, \mathbf{L}} \subseteq C_{\mathbf{L}}$ for all $a \in L$. C2 holds by Lemma 33.(ii). C3 holds by Lemma 33.(ii) and Lemma 31, which ensures us that $e^{\sim'} = \diamond e \in C_p(\mathbf{L})$. From the latter fact, the equality $e^\sim \vee \diamond e = 1$ and Lemma 33.(i), we obtain:

$$\begin{aligned}
a \wedge b &\leq (a \vee b) \wedge (a \vee \diamond e) \wedge (b \vee e^\sim) \\
&= (a \vee b) \wedge (e^\sim \vee b) \wedge (\diamond e \vee a) \wedge (e^\sim \vee \diamond e) \\
&= (b \vee (a \wedge e^\sim)) \wedge (\diamond e \vee (a \wedge e^\sim)) \\
&= (a \wedge e^\sim) \vee (b \wedge \diamond e),
\end{aligned}$$

whence, also using the equality $e^\sim \wedge \diamond e = 0$ and Lemma 31.(ii),

$$\begin{aligned}
((e^\sim \vee b) \wedge (\diamond e \vee a))^\sim &= ((e^\sim \wedge a) \vee (\diamond e \wedge b) \vee (b \wedge a))^\sim && \text{Lemma 33} \\
&= ((e^\sim \wedge a) \vee (\diamond e \wedge b))^\sim \\
&= (e^\sim \wedge a)^\sim \wedge (\diamond e \wedge b)^\sim \\
&= (e^\sim \vee b^\sim) \wedge (\diamond e \vee a^\sim) && e^\sim, \diamond e \in C_{pbz}(\mathbf{L})
\end{aligned}$$

Conversely, if e^\sim is central in \mathbf{L} , then by C1 we have that for all $a \in L$, $a = (a \wedge e^\sim) \vee (a \wedge \diamond e)$, whereby $e^\sim \in C_p(\mathbf{L})$. Now, recall that if e^\sim is central, then so is $\diamond e$, whereby C4 holds for both elements — i.e., for all $a, b \in L$:

$$\begin{aligned}
\text{(i)} \quad &((e^\sim \vee b) \wedge (\diamond e \vee a))^\sim = (e^\sim \vee b^\sim) \wedge (\diamond e \vee a^\sim); \\
\text{(ii)} \quad &((\diamond e \vee b) \wedge (e^\sim \vee a))^\sim = (\diamond e \vee b^\sim) \wedge (e^\sim \vee a^\sim).
\end{aligned}$$

Letting $b = e^\sim$ in (i) and using Lemma 33.(ii), we have that $(e^\sim \wedge a)^\sim = ((e^\sim \vee e^\sim) \wedge (\diamond e \vee a))^\sim = \diamond e \vee a^\sim$. Similarly, letting $b = \diamond e$ in (ii), we obtain $(\diamond e \wedge a)^\sim = e^\sim \vee a^\sim$. ■

3.2 Central Elements in the Variety Generated by Antiortholattices

In any member \mathbf{L} of $V(\mathbb{AOL})$ central elements are exactly the sharp elements: $C(\mathbf{L}) = S_K(\mathbf{L})$. Therefore, for any $a \in \mathbf{L}$, \mathbf{L} is decomposable as $\mathbf{L}/Cg(a^\sim, 0) \times \mathbf{L}/Cg(a^\sim, 1)$, or equivalently as $\mathbf{L}/Cg(a^\sim, 0) \times \mathbf{L}/Cg(\diamond a, 0)$. However, the mentioned results in [21] do not provide us with a uniform recipe to obtain an explicit description of the factors in this decomposition. This situation marks a sharp contrast with the case of orthomodular lattices, where the following result is available:

Theorem 35 [5, Lm. 2.7] *Let $\mathbf{L} \in \mathbb{OML}$ and let $e \in C(\mathbf{L})$. Then $\mathbf{L} \simeq \mathbf{L}_1 \times \mathbf{L}_2$, where $\mathbf{L}_1, \mathbf{L}_2$ are algebras whose universes are the intervals $[0, e]$ and $[0, e']$, respectively.*

In this subsection, we similarly characterise the factors in these decompositions in terms of algebras on intervals in \mathbf{L} .

Lemma 36 *Let $\mathbf{L} \in V(\mathbb{AOL})$ and let $a \in L$. Define:*

$$\begin{aligned}
\mathbf{L}_1 &= ([0, a^\sim], \wedge, \vee, {}^{/1}, {}^{\sim 1}, 0, a^\sim); \\
\mathbf{L}_2 &= ([0, \diamond a], \wedge, \vee, {}^{/2}, {}^{\sim 2}, 0, \diamond a),
\end{aligned}$$

where for all b in $[0, a^\sim]$, $b^{/1} = b' \wedge a^\sim$, $b^{\sim 1} = b^\sim \wedge a^\sim$, while for all c in $[0, \diamond a]$, $c^{/2} = c' \wedge \diamond a$, $c^{\sim 2} = c^\sim \wedge \diamond a$. Then the algebras \mathbf{L}_1 and \mathbf{L}_2 are in $V(\mathbb{AOL})$.

Proof. Clearly, \mathbf{L}_1 and \mathbf{L}_2 are bounded lattices. We now verify the remaining properties for \mathbf{L}_1 ; by replacing a by a^\sim , we obtain our claim for \mathbf{L}_2 .

(\mathbf{L}_1 is a pseudo-Kleene algebra). Let $b, c \in [0, a^\sim]$. Then, using Lemma 14.(iii), $b^{/1/1} = (b' \wedge a^\sim)' \wedge a^\sim = (b \vee \diamond a) \wedge a^\sim = b \wedge a^\sim = b$. Moreover,

$(b \wedge c)^{\prime 1} = (b \wedge c)' \wedge a^\sim = (b' \vee c') \wedge a^\sim = (b' \wedge a^\sim) \vee (c' \wedge a^\sim) = b^{\prime 1} \vee c^{\prime 1}$.
 Finally, since $b \wedge b' \leq c \vee c'$ in \mathbf{L} , we use Lemma 14.(ii) and obtain

$$b \wedge b^{\prime 1} = b \wedge b' \wedge a^\sim \leq (c \vee c') \wedge (c \vee a^\sim) = c \vee (c' \wedge a^\sim) = c \vee c^{\prime 1}.$$

(\mathbf{L}_1 is paraorthomodular). Let $b, c \in [0, a^\sim]$. Suppose that $b \leq c$ and that $b^{\prime 1} \wedge c = b' \wedge a^\sim \wedge c = 0$. Since $c \leq a^\sim$, $b' \wedge a^\sim \wedge c = b' \wedge c$, whence paraorthomodularity of \mathbf{L} yields the desired result.

(\mathbf{L}_1 is a BZ-lattice). Let $b, c \in [0, a^\sim]$. Clearly, $b \wedge b^{\sim 1} = 0$. Further, $b^{\sim 1 \sim 1} = (b^\sim \wedge a^\sim)^\sim \wedge a^\sim$. Since b^\sim and a^\sim are sharp elements, $(b^\sim \wedge a^\sim)^\sim = (b^\sim \wedge a^\sim)' = \diamond b \vee \diamond a$, hence $b^{\sim 1 \sim 1} = (b^\sim \wedge a^\sim)^\sim \wedge a^\sim = (b^\sim \wedge a^\sim)' \wedge a^\sim = b^{\sim 1 \prime 1}$. By Lemma 14.(iii) $b^{\sim 1 \sim 1} = (\diamond b \vee \diamond a) \wedge a^\sim = \diamond b \wedge a^\sim \geq b$. Finally, it is easily seen that if $b \leq c$, then $c^{\sim 1} \leq b^{\sim 1}$.

(\mathbf{L}_1 is a BZ*-lattice). Let $b \in [0, a^\sim]$. Then, by Lemma 14.(iii) and WSDM, we have that:

$$\begin{aligned} (b \wedge b^{\prime 1})^{\sim 1} &= (b \wedge b' \wedge a^\sim)^\sim \wedge a^\sim \\ &= (b \wedge b')^\sim \wedge a^\sim \\ &= (b^\sim \vee \square b) \wedge a^\sim \\ &= (b^\sim \wedge a^\sim) \vee (\square b \wedge a^\sim) \\ &= (b^\sim \wedge a^\sim) \vee ((b' \wedge a^\sim)^\sim \wedge a^\sim) \\ &= b^{\sim 1} \vee b^{\prime 1 \sim 1}. \end{aligned}$$

($\mathbf{L}_1 \in V(\mathbb{A}\mathbb{O}\mathbb{L})$). By way of example, we check that the reformulation of AOL2 in terms of the new operations \sim^1 and \prime^1 is satisfied in any antiortholattice \mathbf{M} . Thus, for $a, b, c \in \mathbf{M}$, consider the element

$$t^{\mathbf{M}}(a, b, c) = (b \wedge c^\sim \wedge a^\sim) \vee (b \wedge (c^\sim \wedge a^\sim)^\sim \wedge a^\sim).$$

If $c > 0$, then $t^{\mathbf{M}}(a, b, c) = 0 \vee (b \wedge 1 \wedge a^\sim) = b \wedge a^\sim$. If $c = 0$, then

$$t^{\mathbf{M}}(a, b, c) = (b \wedge a^\sim) \vee (b \wedge \diamond a \wedge a^\sim) = (b \wedge a^\sim) \vee 0 = b \wedge a^\sim.$$

Since $\mathbf{L} \in V(\mathbb{A}\mathbb{O}\mathbb{L})$, it follows that for all $a, b, c \in L$, $t^{\mathbf{L}}(a, b, c) = b \wedge a^\sim$, which equals b whenever $b \leq a^\sim$, hence \mathbf{L}_1 satisfies AOL2. ■

Theorem 37 *Let $\mathbf{L} \in V(\mathbb{A}\mathbb{O}\mathbb{L})$ and let $a \in L$. Then $\mathbf{L} \simeq \mathbf{L}_1 \times \mathbf{L}_2$, where $\mathbf{L}_1, \mathbf{L}_2$ are defined as in Lemma 36.*

Proof. Let $\varphi : L \rightarrow L_1 \times L_2$ be defined, for any $b \in L$, by $\varphi(b) = (b \wedge a^\sim, b \wedge \diamond a)$. We first show that φ is a bijection. If $\varphi(b) = \varphi(c)$ for some $b, c \in L$, then $b \wedge a^\sim = c \wedge a^\sim$ and $b \wedge \diamond a = c \wedge \diamond a$. Thus, by AOL2,

$$b = (b \wedge a^\sim) \vee (b \wedge \diamond a) = (c \wedge a^\sim) \vee (c \wedge \diamond a) = c.$$

Now, let $(x, y) \in L_1 \times L_2$. Then $x \leq a^\sim$ and $y \leq \diamond a$, whence $y \wedge a^\sim \leq \diamond a \wedge a^\sim = 0$. Let us compute $\varphi(x \vee y)$. Using Lemma 14.(iii), we obtain

$$\begin{aligned} (x \vee y) \wedge a^\sim &= (x \wedge a^\sim) \vee (y \wedge a^\sim) \\ &= x \vee 0 = x. \end{aligned}$$

Similarly, $(x \vee y) \wedge \diamond a = y$ and thus φ is onto.

Next, we show that φ preserves meets and the unary operations (observe that this is sufficient in virtue of Lemma 36). For meets,

$$\begin{aligned}\varphi(b \wedge^{\mathbf{L}} c) &= (b \wedge c \wedge a^{\sim}, b \wedge c \wedge \diamond a) \\ &= (b \wedge a^{\sim}, b \wedge \diamond a) \wedge^{\mathbf{L}_1 \times \mathbf{L}_2} (c \wedge a^{\sim}, c \wedge \diamond a) \\ &= \varphi(b) \wedge^{\mathbf{L}_1 \times \mathbf{L}_2} \varphi(c).\end{aligned}$$

With regards to Kleene complements, resorting again to Lemma 14.(iii),

$$\begin{aligned}\varphi(b^{\mathbf{L}}) &= (b' \wedge a^{\sim}, b' \wedge \diamond a) \\ &= ((b' \vee \diamond a) \wedge a^{\sim}, (b' \vee a^{\sim}) \wedge \diamond a) \\ &= ((b \wedge a^{\sim})' \wedge a^{\sim}, (b \wedge \diamond a)' \wedge \diamond a) \\ &= \left((b \wedge a^{\sim})'^1, (b \wedge \diamond a)'^2 \right) \\ &= \varphi(b)^{\mathbf{L}_1 \times \mathbf{L}_2}.\end{aligned}$$

For Brouwer complements the computation is similar. It is safely left to the reader, who is warned that the WSDM identity $(x \wedge y^{\sim})^{\sim} \approx x^{\sim} \vee \diamond y$ will be needed somewhere down the line. ■

4 Ideal Theory

It is well-known from the theory of orthomodular lattices that $\mathbb{O}ML$ -ideals admit a manageable characterization in terms of lattice ideals *closed under perspectivity*, for short *p-ideals*: in other words, in terms of lattice ideals I of an orthomodular lattice \mathbf{L} such that, whenever $a \in I$, then also $b \pitchfork a = b \wedge (b' \vee a) \in I$ for all $b \in L$ [5, Prop. 4.7]. The aim of this section is to generalise this idea within the expanded language of $\mathbb{P}BZL^*$. Unfortunately, in the general case these p-ideals do not even coincide with 0-classes of congruences — and, a fortiori, no isomorphism result between the lattices of p-ideals and of congruences can be attained. The situation improves if we restrict ourselves to the subvariety $\mathbb{S}DM$ of $\mathbb{P}BZL^*$, axiomatised relative to $\mathbb{P}BZL^*$ by the Strong De Morgan identity. In fact, in any $\mathbf{L} \in \mathbb{S}DM$ p-ideals coincide with Ursini $\mathbb{S}DM$ -ideals, hence with 0-classes of congruences, given the fact that $\mathbb{P}BZL^*$ (and thus, all the more so, $\mathbb{S}DM$) is a 0-subtractive variety.

4.1 Ideals in $\mathbb{P}BZ^*$ -lattices

We start by defining the notion of a p-ideal for generic $\mathbb{P}BZ^*$ -lattices.

Definition 38 *Let \mathbf{L} be a $\mathbb{P}BZ^*$ -lattice. $I \subseteq L$ is a p-ideal iff it is a lattice ideal of \mathbf{L} s.t. if $a \in I$, then $\diamond b \pitchfork \diamond a = \diamond b \wedge (b^{\sim} \vee \diamond a) \in I$ for all $b \in L$.*

Lemma 39 *Let \mathbf{L} be a $\mathbb{P}BZ^*$ -lattice, let I be a p-ideal of \mathbf{L} , and let $a, b \in L$. Then: (i) if $a \in I$, then $\diamond a \in I$; (ii) $\diamond a \pitchfork \diamond b \in I$ iff $\diamond b \pitchfork \diamond a \in I$.*

Proof. (i) Let $b = 1$ in Definition 38.

(ii) Suppose that $\diamond b \wedge (b^\sim \vee \diamond a) \in I$. Then, since I is a p-ideal,

$$\diamond a \wedge (a^\sim \vee \diamond (\diamond b \wedge (b^\sim \vee \diamond a))) = \diamond a \wedge (a^\sim \vee (\diamond b \wedge (b^\sim \vee \diamond a))) \in I.$$

By Theorem 27 applied to $\mathbf{S}_K(\mathbf{L})$, however,

$$\diamond a \wedge (a^\sim \vee (\diamond b \wedge (b^\sim \vee \diamond a))) = \diamond a \wedge (a^\sim \vee \diamond b),$$

whence our conclusion. ■

We now aim at defining a customary notion of equivalence between elements in a PBZ* -lattice modulo a given p-ideal. Note that $\{0\}$ is a p-ideal; thus,

Definition 40 Let \mathbf{L} be a PBZ* -lattice, and let I be a p-ideal of \mathbf{L} . The elements $a, b \in L$ are said to be I -modally equivalent iff

$$(\diamond a)^\sim \pitchfork \diamond b, (\diamond b)^\sim \pitchfork \diamond a, (\square a)^\sim \pitchfork \square b, (\square b)^\sim \pitchfork \square a \in I.$$

The elements $a, b \in L$ are said to be modally equivalent iff they are $\{0\}$ -modally equivalent, namely iff $\diamond a = \diamond b$ and $\square a = \square b$.

Definition 41 Let \mathbf{L} be a PBZ* -lattice, and let I be a p-ideal of \mathbf{L} . We define the following binary relation on L :

$$\rho(I) = \{(a, b) \in L : (\diamond a)^\sim \pitchfork \diamond b, (\diamond b)^\sim \pitchfork \diamond a, (\square a)^\sim \pitchfork \square b, (\square b)^\sim \pitchfork \square a \in I\}.$$

Thus, $(a, b) \in \rho(I)$ iff a and b are I -modally equivalent, and, in particular, $(a, b) \in \rho(\{0\})$ iff a and b are modally equivalent. The relation $\rho(I)$ also admits a less cumbersome description:

Theorem 42 Let \mathbf{L} be a PBZ* -lattice, and let I be a p-ideal of \mathbf{L} . For $a, b \in L$ the following conditions are equivalent:

- (i) $(\diamond a \vee \diamond b) \wedge (a^\sim \vee b^\sim), ((\square a)^\sim \vee (\square b)^\sim) \wedge (\square a \vee \square b) \in I$;
- (ii) there exist $s, t \in I$ such that $\diamond a \vee s = \diamond b \vee s$ and $\square a \vee t = \square b \vee t$;
- (iii) $(a, b) \in \rho(I)$.

Proof. (i) implies (ii). Let $s = (\diamond a \vee \diamond b) \wedge (a^\sim \vee b^\sim)$ and $t = ((\square a)^\sim \vee (\square b)^\sim) \wedge (\square a \vee \square b)$. Then:

$$\begin{aligned} \diamond a \vee s &= \diamond a \vee ((\diamond a \vee \diamond b) \wedge (a^\sim \vee b^\sim)) \\ &= \diamond a \vee \diamond b && \text{(Thm. 27 in } \mathbf{S}_K(\mathbf{L}) \text{)} \\ &= \diamond b \vee s. \end{aligned}$$

Similarly, $\square a \vee t = \square b \vee t$.

(ii) implies (iii). Suppose that there exist $s, t \in I$ such that $\diamond a \vee s = \diamond b \vee s$ and $\square a \vee t = \square b \vee t$. First, we prove $(\diamond b)^\sim \pitchfork \diamond a \in I$.

$$\begin{aligned}
(\diamond b)^\sim \pitchfork \diamond a &= b^\sim \wedge (\diamond b \vee \diamond a) \\
&\leq b^\sim \wedge (\diamond s \vee \diamond b \vee \diamond a) \\
&= b^\sim \wedge (\diamond (s \vee \diamond a) \vee \diamond b) \\
&= b^\sim \wedge (\diamond (s \vee \diamond b) \vee \diamond b) \\
&= b^\sim \wedge (\diamond b \vee \diamond s) \\
&= (\diamond b)^\sim \pitchfork \diamond s.
\end{aligned}$$

Since I is a p-ideal of \mathbf{L} and $s \in I$, we have that $(\diamond b)^\sim \pitchfork \diamond s \in I$ and therefore $(\diamond b)^\sim \pitchfork \diamond a \in I$. The remaining conditions are proved similarly and thus $(a, b) \in \rho(I)$.

(iii) implies (i). Suppose $(a, b) \in \rho(I)$; we prove that $(\diamond a \vee \diamond b) \wedge (a^\sim \vee b^\sim) \in I$. By assumption,

$$(a^\sim \wedge (\diamond a \vee \diamond b)) \vee (b^\sim \wedge (\diamond b \vee \diamond a)) = ((\diamond a)^\sim \pitchfork \diamond b) \vee ((\diamond b)^\sim \pitchfork \diamond a) \in I.$$

By Theorem 27 applied to $\mathbf{S}_K(\mathbf{L})$, however,

$$(a^\sim \wedge (\diamond a \vee \diamond b)) \vee (b^\sim \wedge (\diamond b \vee \diamond a)) = (\diamond a \vee \diamond b) \wedge (a^\sim \vee b^\sim),$$

whence our claim follows. A similar proof establishes the other claim. ■

Theorem 43 *Let \mathbf{L} be a PBZ* -lattice, and let I be a p-ideal of \mathbf{L} . Then $\rho(I)$ is an equivalence relation on L that preserves the operations $'$ and \sim .*

Proof. Since, for all $a \in L$, $(\diamond a)^\sim \pitchfork \diamond a$, $(\square a)^\sim \pitchfork \square a = 0 \in I$, $\rho(I)$ is reflexive. Symmetry is trivial. For transitivity, suppose $(a, b), (b, c) \in \rho(I)$. By Theorem 42, there exist:

- $s_1, t_1 \in I$ such that $\diamond a \vee s_1 = \diamond b \vee s_1$ and $\square a \vee t_1 = \square b \vee t_1$;
- $s_2, t_2 \in I$ such that $\diamond b \vee s_2 = \diamond c \vee s_2$ and $\square b \vee t_2 = \square c \vee t_2$.

Thus $s_1 \vee s_2 \in I$ and $\diamond a \vee s_1 \vee s_2 = \diamond b \vee s_1 \vee s_2 = \diamond c \vee s_1 \vee s_2$, and similarly for the other condition, whence by Theorem 42 again, $(a, c) \in \rho(I)$. The unary operations are clearly preserved. ■

Although $\rho(I)$ is always an equivalence relation, it need not always be a congruence, as the next example shows.

Example 44 *Consider the distributive antiortholattice whose lattice reduct is the ordinal sum of \mathbf{D}_2^2 with itself, with atoms a, b and the fixpoint $c = c'$. Observe that $(a, c) \in \rho(\{0\})$, because $\diamond a = \diamond c = 1$ and $\square a = \square c = 0$. However, $\diamond(a \wedge b) = \diamond 0 = 0$ and $\diamond(c \wedge b) = \diamond b = 1$, whence $\rho(\{0\})$ does not preserve meets.*

Our next goal is to tweak the notion of p-ideal in such a way that its associated equivalence is necessarily a congruence.

Definition 45 Let \mathbf{L} be a PBZ^* -lattice, and let I be a p -ideal of \mathbf{L} . I is a weak De Morgan ideal iff for all $a, b \in L$, whenever $(a, b) \in \rho(I)$, then for all $c \in L$ it is the case that $\diamond(a \wedge c) \sim \mathfrak{m} \diamond(b \wedge c) \in I$.

Lemma 46 Let \mathbf{L} be a PBZ^* -lattice, and let I be a p -ideal of \mathbf{L} . The following conditions are equivalent:

- (i) $\rho(I)$ is a congruence;
- (ii) I is a weak De Morgan ideal.

Proof. (i) implies (ii). Suppose that $\rho(I)$ is a congruence and let $(a, b) \in \rho(I)$, $c \in L$. Then $(\diamond(a \wedge c), \diamond(b \wedge c)) \in \rho(I)$. It follows that $(\diamond(a \wedge c) \sim \mathfrak{m} \diamond(b \wedge c), 0) \in \rho(I)$, which implies $\diamond(a \wedge c) \sim \mathfrak{m} \diamond(b \wedge c) \in I$.

(ii) implies (i). Let I be a weak De Morgan ideal. By Theorem 43, to attain our conclusion it will suffice to show that $\rho(I)$ preserves meets. Thus, let $(a, b) \in \rho(I)$ and $c \in L$. In virtue of our assumption, $\diamond(a \wedge c) \sim \mathfrak{m} \diamond(b \wedge c) \in I$ and, taking into account the symmetry of $\rho(I)$, $\diamond(b \wedge c) \sim \mathfrak{m} \diamond(a \wedge c) \in I$. It remains to show that

$$(\Box(a \wedge c)) \sim \mathfrak{m} \Box(b \wedge c), (\Box(b \wedge c)) \sim \mathfrak{m} \Box(a \wedge c) \in I.$$

However, since $(a, b) \in \rho(I)$ and $\rho(I)$ preserves the unary operations, $(\Box a, \Box b) \in \rho(I)$. Given that I is a weak De Morgan ideal, thus,

$$\begin{aligned} (\Box(a \wedge c)) \sim \mathfrak{m} \Box(b \wedge c) &= \diamond(a' \vee c') \mathfrak{m} (b' \vee c') \sim \\ &= (\Box a \wedge \Box c) \sim \mathfrak{m} \diamond(\Box b \wedge \Box c) \\ &= \diamond(\Box a \wedge \Box c) \sim \mathfrak{m} \diamond(\Box b \wedge \Box c) \in I. \end{aligned}$$

Similarly, $(\Box(b \wedge c)) \sim \mathfrak{m} \Box(a \wedge c) \in I$. ■

4.2 Ideals in the Strong De Morgan Subvariety

The subvariety of $PBZL^*$ that is axiomatised relative to $PBZL^*$ by the Strong De Morgan law SDM , here labelled SDM , includes OML and stands out for its smooth theory of ideals. In fact, we have that:

Lemma 47 Let $\mathbf{L} \in \mathcal{SDM}$, and let I be a p -ideal of \mathbf{L} . Then I is a weak De Morgan ideal and therefore $\rho(I)$ is a congruence.

Proof. If $(a, b) \in \rho(I)$, then $(a \sim, b \sim) \in \rho(I)$ by Theorem 43. Thus, Theorem 42 guarantees that there is $s \in I$ such that $a \sim \vee s = \diamond a \sim \vee s = \diamond b \sim \vee s = b \sim \vee s$. Then, for an arbitrary $c \in L$, $a \sim \vee c \sim \vee s = b \sim \vee c \sim \vee s$. Applying SDM , we have that $\Box((a \wedge c) \sim) \vee s = (a \wedge c) \sim \vee s = (b \wedge c) \sim \vee s = \Box((b \wedge c) \sim) \vee s$. A further recourse to Theorem 42 yields $((a \wedge c) \sim, (b \wedge c) \sim) \in \rho(I)$, whence $(\diamond(a \wedge c), \diamond(b \wedge c)) \in \rho(I)$, which implies, in particular, that I is weak De Morgan. Lemma 46 takes care of the remaining claim. ■

We are now in a position to prove that within the boundaries of this subvariety, p -ideals coincide with ideals in the sense of Ursini.

Theorem 48 *If $\mathbf{L} \in \text{SDM}$, then the class of p -ideals of \mathbf{L} coincides with $\mathcal{I}_{\text{SDM}}(\mathbf{L})$.*

Proof. Let $I \in \mathcal{I}_{\text{SDM}}(\mathbf{L})$, whence by Theorem 20.(i) $I = 0/\theta$ for some $\theta \in \text{Con}_{\text{BZL}}(\mathbf{L})$. Clearly, I is a lattice ideal of \mathbf{L} . Furthermore, if $a \in I$, then $(\diamond a, 0) \in \theta$ and then $\diamond a \in I$. What remains to show is that, for an arbitrary $b \in L$, $\diamond b \mathbin{\frown} \diamond a \in I$. Since $\diamond a \in I = 0/\theta$, $(\diamond b \mathbin{\frown} \diamond a, 0) = (\diamond b \mathbin{\frown} \diamond a, \diamond b \mathbin{\frown} 0) \in \theta$, which means $\diamond b \mathbin{\frown} \diamond a \in I$. Conversely, it will be enough to prove that if I is a p -ideal of \mathbf{L} , then $I = 0/\rho(I)$. However, by Theorem 42,

$$\begin{aligned} 0/\rho(I) &= \{a \in L : (a, 0) \in \rho(I)\} \\ &= \{a \in L : \diamond a \leq s, \square a \leq t \text{ for some } s, t \in I\}. \end{aligned}$$

If $a \in I$, then choose $s = t = \diamond a \in I$ to obtain $a \in 0/\rho(I)$. If $a \in 0/\rho(I)$, then there is $s \in I$ such that $a \leq \diamond a \leq s$, whence $a \in I$. ■

Observe that, by Lemma 47 and Theorem 48, whenever $\mathbf{L} \in \text{SDM}$, all members of $\mathcal{I}_{\text{SDM}}(\mathbf{L})$ are weak De Morgan ideals.

Theorem 49 *Let $\mathbf{L} \in \text{SDM}$, and let $I \in \mathcal{I}_{\text{SDM}}(\mathbf{L})$. Then $\rho(I) = I^\varepsilon$.*

Proof. By the proof of Theorem 48 $0/\rho(I) = I$, whence $\rho(I) \subseteq I^\varepsilon$. For the converse inequality, suppose $(a, b) \in I^\varepsilon$. Since I^ε is a congruence,

$$((\diamond a \vee \diamond b) \wedge (a^\sim \vee b^\sim), 0), ((\square a \vee \square b) \wedge ((\square a)^\sim \vee (\square b)^\sim), 0) \in I^\varepsilon.$$

So $(\diamond a \vee \diamond b) \wedge (a^\sim \vee b^\sim) \in 0/I^\varepsilon = I$ and $(\square a \vee \square b) \wedge ((\square a)^\sim \vee (\square b)^\sim) \in 0/I^\varepsilon = I$. By Theorem 42, this means that $(a, b) \in \rho(I)$. ■

Corollary 50 *SDM is finitely congruential.*

Proof. We have to find a finite set of terms $\{d_i(x, y)\}_{i \leq n}$ that witnesses finite congruentiality according to Definition 22. Thus, let

$$\begin{aligned} d_1(x, y) &= (\diamond x)^\sim \mathbin{\frown} \diamond y, & d_2(x, y) &= (\diamond y)^\sim \mathbin{\frown} \diamond x, \\ d_3(x, y) &= (\square x)^\sim \mathbin{\frown} \square y, & d_4(x, y) &= (\square y)^\sim \mathbin{\frown} \square x. \end{aligned}$$

If $\mathbf{L} \in \text{SDM}$ and $I \in \mathcal{I}_{\text{SDM}}(\mathbf{L})$, then by Theorems 49 and 42 $\rho(I) = I^\varepsilon$. As a result, $(a, b) \in I^\varepsilon = \rho(I)$ iff $d_i^{\mathbf{A}}(a, b) \in I$, for all $i \leq 4$. ■

Theorem 51 *The 0-assertional logic of PBZL^* is not equivalential.*

Proof. Consider again the antiortholattice of Example 44. Being simple, this antiortholattice belongs to $\text{PBZL}_\varepsilon^*$. Moreover, the set $\{0, a, a', 1\}$ is a subuniverse of such, isomorphic to \mathbf{D}_4 , and its middle congruence, that collapses only a and a' , is a nonzero pseudo-identical congruence. Our claim follows then from Theorem 24. ■

Lemma 52 *Let $\mathbf{L} \in \text{SDM}$. The following are equivalent:*

$$(i) \rho(\{0\}) = \Delta_{\mathbf{L}}.$$

(ii) \mathbf{L} satisfies the quasi-identity $\Box x \leq \Box y \ \& \ \Diamond x \leq \Diamond y \Rightarrow x \leq y$.

(iii) \mathbf{L} satisfies the identity SK.

Proof. (i) implies (ii). $\rho(\{0\}) = \Delta_{\mathbf{L}}$ means that modally equivalent elements of L are identical. Now, let $\Box a \leq \Box b$ and $\Diamond a \leq \Diamond b$. By SDM, this implies that $a \wedge b$ and a are modally equivalent, whence $a \leq b$.

(ii) implies (iii). Using SDM, we have that for all $a, b \in L$,

$$\begin{aligned}\Box(a \wedge \Diamond b) &= \Box a \wedge \Diamond b \leq \Box a \vee \Box b = \Box(\Box a \vee b); \\ \Diamond(a \wedge \Diamond b) &= \Diamond a \wedge \Diamond b \leq \Box a \vee \Diamond b = \Diamond(\Box a \vee b).\end{aligned}$$

By our assumption, then, $a \wedge \Diamond b \leq \Box a \vee b$.

(iii) implies (i). Suppose $\Box a = \Box b$ and $\Diamond a = \Diamond b$. Then

$$a = a \wedge \Diamond a = a \wedge \Diamond b \leq \Box a \vee b = \Box b \vee b = b.$$

Similarly, $b \leq a$, whence our conclusion. ■

Let us call \mathbf{SK} the subvariety of \mathbf{SDM} that is axiomatised relative to \mathbf{SDM} by the identity SK.

Theorem 53 *The 0-assertional logic of \mathbf{SDM} is strongly algebraisable with equivalent variety semantics $\mathbf{SDM}_\varepsilon = \mathbf{SK}$.*

Proof. By Theorem 25, it suffices to establish that \mathbf{SDM}_ε is a variety, which would follow if we were to show that $\mathbf{SDM}_\varepsilon = \mathbf{SK}$. By Theorem 49, whenever \mathbf{L} belongs to \mathbf{SDM} , $\{0\}^\varepsilon = \rho(\{0\})$. Thus $\mathbf{L} \in \mathbf{SDM}_\varepsilon$ iff $\rho(\{0\}) = \Delta_{\mathbf{L}}$, and by Lemma 52, this happens exactly when $\mathbf{L} \in \mathbf{SK}$. ■

By Theorems 23 and 48, in any member \mathbf{L} of \mathbf{SDM} the Ursini ideals of \mathbf{L} coincide with its p -ideals and with the deductive filters on \mathbf{L} of the 0-assertional logic of \mathbf{SDM} . By [13, Thm. 3.58], therefore, we obtain:

Corollary 54 *Let $\mathbf{L} \in \mathbf{SDM}$. Then the lattice of p -ideals of \mathbf{L} is isomorphic to the lattice of all congruences θ on \mathbf{L} such that $\mathbf{L}/\theta \in \mathbf{SK}$.*

Observe that, although SK implies SDM in the context of $V(\mathbf{AOL})$ [15, Lm. 3.8] this is not the case in the more general context of \mathbf{PBZL}^* . In fact, consider the \mathbf{PBZ}^* -lattice \mathbf{L} whose lattice reduct is the 5-element modular and non-distributive lattice \mathbf{M}_3 with atoms a, a', b , where $b = b'$ and $b^\sim = 0$. Then \mathbf{L} satisfies SK but fails SDM - actually, it fails even WSDM because $a \in S_K(\mathbf{L})$ and $(a \wedge b)^\sim = 1$ but $a^\sim \vee b^\sim = a'$. For future reference, we make a note of the fact that \mathbf{L} satisfies J2.

4.3 Ideals in $V(\mathbf{AOL})$

Another subvariety of \mathbf{PBZL}^* where our description of ideals can be considerably simplified is the variety $V(\mathbf{AOL})$ generated by all antiortholattices. Bignall and Spinks first observed that the variety of distributive BZ-lattices is a binary discriminator variety [4]. We extend their observation by noticing that $V(\mathbf{AOL})$ is itself a binary discriminator variety.

Proposition 55 $V(\mathbb{AOL})$ is a 0-binary discriminator variety.

Proof. Referring to Definition 26 for notation and terminology, let $b_0^L(x, y) = x \wedge y^\sim$. Then for any antiortholattice \mathbf{L} and, for any $a, c \in L$, $b_0^L(a, 0) = a \wedge 0^\sim = a \wedge 1 = a$, while, if $c > 0$, then $b_0^L(a, c) = a \wedge c^\sim = a \wedge 0 = 0$. ■

Among the consequences of this remark we have a very slender description of $V(\mathbb{AOL})$ -ideals. In fact, recall from [4] that if \mathbf{A} is an algebra in a 0-binary discriminator variety \mathbb{V} and $b_0(x, y)$ is the term witnessing this property for \mathbb{V} , $I \subseteq A$ is a \mathbb{V} -ideal of \mathbf{A} exactly when, for any $a, c \in A$, if $c \in I$ and $b_0^{\mathbf{A}}(a, c) \in I$, then $a \in I$. Therefore:

Proposition 56 Let $\mathbf{L} \in V(\mathbb{AOL})$. For a lattice ideal $I \subseteq L$ the following are equivalent:

- (i) I is a $V(\mathbb{AOL})$ -ideal.
- (ii) For any $a, b \in L$, if $b \in I$ and $a \wedge b^\sim \in I$, then $a \in I$.
- (iii) I is closed w.r.t. all interpretations in \mathbf{L} of the $V(\mathbb{AOL})$ -ideal term (in y, z):

$$u(x, y, z) = x \wedge \diamond(y \vee z).$$

Proof. The equivalence of (i) and (ii) follows from the remarks immediately preceding this lemma. Suppose now that (ii) holds, and that $a, b \in I$. Then $a \vee b \in I$. On the other hand, for any c in L ,

$$0 = c \wedge \diamond(a \vee b) \wedge (a \vee b)^\sim \in I,$$

whence by our hypothesis $c \wedge \diamond(a \vee b) \in I$. Conversely, under the assumption (iii), let $b \in I$ and $a \in L$ be such that $a \wedge b^\sim \in I$. Then, using WSDM and Lemma 14 several times,

$$\begin{aligned} u^{\mathbf{A}}(a, b, a \wedge b^\sim) &= a \wedge \diamond(b \vee (a \wedge b^\sim)) \\ &= a \wedge (\diamond b \vee \diamond(a \wedge b^\sim)) \\ &= a \wedge (\diamond b \vee (\diamond a \wedge b^\sim)) \\ &= a \wedge (\diamond b \vee \diamond a) \\ &= (a \wedge \diamond b) \vee a = a, \end{aligned}$$

hence $a \in I$. ■

5 Axiomatic Bases for Some Subvarieties

The goal of this final section is to simplify the axiomatisation of $V(\mathbb{AOL})$ given in [14] and to solve a problem (here called the Join Problem) posed in [15], where it was observed that the varietal join $\mathbb{OML} \vee V(\mathbb{AOL})$ in the lattice of subvarieties of \mathbb{PBZL}^* was strictly included in \mathbb{PBZL}^* , but no axiomatic basis for such a join was given.

5.1 A Streamlined Axiomatisation for $V(\mathbb{AOL})$

In Theorem 15.(i) we recalled that an equational basis for $V(\mathbb{AOL})$ relative to \mathbb{PBZL}^* is given by the identities AOL1-AOL3, here reproduced for the reader's convenience:

$$\begin{aligned} \text{(AOL1)} \quad & (x^\sim \vee y^\sim) \wedge (\diamond x \vee z^\sim) \approx ((x^\sim \vee y) \wedge (\diamond x \vee z))^\sim; \\ \text{(AOL2)} \quad & x \approx (x \wedge y^\sim) \vee (x \wedge \diamond y); \\ \text{(AOL3)} \quad & x \approx (x \vee y^\sim) \wedge (x \vee \diamond y). \end{aligned}$$

The aim of this subsection is showing that AOL2 suffices to derive the remaining two axioms. For a start, we notice that Lemma 14 does not depend on AOL1, whence it holds for any subvariety of \mathbb{PBZL}^* that satisfies AOL2 and AOL3.

Lemma 57 *Let \mathbf{L} be a member of \mathbb{PBZL}^* that satisfies AOL2 and AOL3. Then, for any $a, b, c \in L$: (i) $a \wedge b \leq (a \wedge c^\sim) \vee (b \wedge \diamond c)$; (ii) $(a \wedge \diamond b)^\sim \vee \diamond b = 1$; (iii) $\diamond a \leq (b \wedge a^\sim)^\sim$.*

Proof. (i) In fact, using Lemma 14,

$$a \wedge b \leq (a \vee c^\sim) \wedge (b \vee \diamond c) \wedge (a \vee b) = (a \wedge c^\sim) \vee (b \wedge \diamond c).$$

(ii) Since $a \wedge \diamond b \leq \diamond b$, it follows that $b^\sim \leq (a \wedge \diamond b)^\sim$, whence $1 = b^\sim \vee \diamond b \leq (a \wedge \diamond b)^\sim \vee \diamond b$.

(iii) Since $b \wedge a^\sim \leq a^\sim$, our conclusion follows. ■

Lemma 58 *Let \mathbf{L} be a member of \mathbb{PBZL}^* that satisfies AOL2 and AOL3. Then \mathbf{L} satisfies AOL1 iff it satisfies WSDM.*

Proof. From left to right, WSDM can be obtained by taking $y = 0$ and applying Lemma 14.(iii). Conversely, let \mathbf{L} satisfy WSDM, and let $a, b, c \in L$. Then:

$$\begin{aligned} ((a^\sim \vee b) \wedge (\diamond a \vee c))^\sim &= ((a^\sim \wedge c) \vee (\diamond a \wedge b) \vee (b \wedge c))^\sim && \text{Lm. 14} \\ &= ((a^\sim \wedge c) \vee (\diamond a \wedge b))^\sim && \text{Lm. 57.(i)} \\ &= (a^\sim \wedge c)^\sim \wedge (\diamond a \wedge b)^\sim && \text{Lm. 6.(iii)} \\ &= (a^\sim \vee b^\sim) \wedge (\diamond a \vee c^\sim) && \text{WSDM} \end{aligned}$$

■

Theorem 59 *An equational basis for $V(\mathbb{AOL})$ relative to \mathbb{PBZL}^* is given by the single identity AOL2.*

Proof. By Theorem 15.(i), an equational basis for $V(\mathbb{AOL})$ relative to \mathbb{PBZL}^* is given by the identities AOL1-AOL3. To attain our conclusion, taking into account Lemma 58, it will suffice to show that: i) any subvariety of \mathbb{PBZL}^* that satisfies AOL2 and AOL3 also satisfies WSDM; ii) any subvariety of \mathbb{PBZL}^* that satisfies AOL2 also satisfies AOL3. We establish these claims in reverse order.

i) Let \mathbf{L} belong to any subvariety of \mathbb{PBZL}^* that satisfies AOL2, and let $a, b \in L$. Then $a' = (a' \wedge b^\sim) \vee (a' \wedge \diamond b)$, whence

$$a = ((a' \wedge b^\sim) \vee (a' \wedge \diamond b))' = (a' \wedge b^\sim)' \wedge (a' \wedge \diamond b)' = (a \vee \diamond b) \wedge (a \vee b^\sim).$$

ii) Let \mathbf{L} belong to any subvariety of \mathbb{PBZL}^* that satisfies AOL2 (thus also AOL3, by the previous item), and let $a, b \in L$. Then:

$$\begin{aligned} a^\sim \vee \diamond b &= ((a \wedge b^\sim) \vee (a \wedge \diamond b))^\sim \vee \diamond b && \text{AOL2} \\ &= ((a \wedge b^\sim)^\sim \wedge (a \wedge \diamond b)^\sim) \vee \diamond b && \text{Lm. 6.(iii)} \\ &= ((a \wedge b^\sim)^\sim \vee \diamond b) \wedge ((a \wedge \diamond b)^\sim \vee \diamond b) && \text{Lm. 14.(iv)} \\ &= ((a \wedge b^\sim)^\sim \vee \diamond b) && \text{Lm. 57.(ii)} \\ &= (a \wedge b^\sim)^\sim && \text{Lm. 57.(iii)}. \end{aligned}$$

■

Taking into account Lemma 31 and Theorem 34, we have that:

Corollary 60 (i) $V(\mathbf{AOL})$ is the class of all \mathbb{PBZ}^* -lattices \mathbf{L} such that $C_p(\mathbf{L}) = S_K(\mathbf{L})$.

(ii) $V(\mathbf{AOL})$ is the class of all \mathbb{PBZ}^* -lattices \mathbf{L} that satisfy WSDM and are such that $C_{pbz}(\mathbf{L}) = S_K(\mathbf{L})$.


(iii) The class of the directly indecomposable members of $V(\mathbf{AOL})$ is \mathbf{AOL} .

5.2 The Join Problem

We round off this paper by axiomatising the variety $\mathbf{OML} \vee V(\mathbf{AOL})$, as well as some of its notable subvarieties. Let:

- \mathbb{V}_1 be the variety of \mathbb{PBZ}^* -lattices that is axiomatised relative to \mathbb{PBZL}^* by the identities J2 and WSDM;
- \mathbb{V}_2 be the variety of \mathbb{PBZ}^* -lattices that is axiomatised relative to \mathbb{PBZL}^* by the identities J2 and SDM;
- \mathbb{V}_3 be the variety of \mathbb{PBZ}^* -lattices that is axiomatised relative to \mathbb{PBZL}^* by the identities J2, WSDM, and SK.

Taking into account the results in [15] and [14], as well as Theorem 59, in $V(\mathbf{AOL})$, SK implies SDM, that is AOL2 and SK imply SDM, while AOL2 and SDM do not imply SK. We observed in Section 4 that J2 and SK do not imply WSDM (all the more so, thus, SDM). The following \mathbb{PBZ}^* -lattice:



satisfies SK and SDM and fails J2. The 4-element antiortholattice chain \mathbf{D}_4 fails SK but satisfies SDM and AOL2, thus also J2. Therefore, in the sets of axioms $\{\text{J2}, \text{SK}, \text{SDM}\}$ and $\{\text{J2}, \text{SK}, \text{WSDM}\}$, each axiom is independent from the other two.

Given any $\mathbf{L} \in \text{PBZL}^*$, it will be expedient to denote by $T(\mathbf{L})$ the set $\{x \in L : x^\sim = 0\} \cup \{0\}$.

Lemma 61 *Let \mathbf{L} be a PBZ*-lattice that satisfies WSDM and such that $S_K(\mathbf{L}) \cup T(\mathbf{L}) = L$. Then if $b \in S_K(\mathbf{L})$ and $c \notin S_K(\mathbf{L})$, it follows that either $b = 1$ or $b \leq c$.*

Proof. If $b \wedge c \in S_K(\mathbf{L})$, then

$$b \wedge c = \diamond(b \wedge c) = \diamond b \wedge \diamond c = b \wedge 1 = b,$$

where WSDM can be applied to obtain the second equality because $b \in S_K(\mathbf{L})$, while the third equality follows from the fact that $c \notin S_K(\mathbf{L})$, whence $c^\sim = 0$. On the other hand, if $b \wedge c \notin S_K(\mathbf{L})$, then we apply again WSDM (as $b \in S_K(\mathbf{L})$) and the assumption that $S_K(\mathbf{L}) \cup T(\mathbf{L}) = L$, obtaining

$$0 = (b \wedge c)^\sim = b^\sim \vee c^\sim = b^\sim \vee 0 = b^\sim,$$

whereby $b = 1$ since $b \in S_K(\mathbf{L})$. ■

Proposition 62 *Any directly indecomposable $\mathbf{L} \in \mathbb{V}_1$ is either orthomodular or an antiortholattice.*

Proof. Let \mathbf{L} be as in the statement of the proposition, and suppose that \mathbf{L} is directly indecomposable, but is neither orthomodular nor an antiortholattice. By Theorem 34, the only PBZ*-central elements of \mathbf{L} are 0 and 1. By WSDM and Lemma 31.(iii) we conclude that 0 and 1 are the only sharp elements a such that $b = (b \wedge a^\sim) \vee (b \wedge \diamond a)$ for all $b \in L$.

Now, we want to show that $S_K(\mathbf{L}) \cup T(\mathbf{L}) = L$. Let $x \in L$. The element $\diamond(x \wedge x')$ is sharp and, by J2, we have that

$$b = (b \wedge (x \wedge x')^\sim) \vee (b \wedge \diamond(x \wedge x')) = (b \wedge \diamond(x \wedge x')^\sim) \vee (b \wedge \diamond \diamond(x \wedge x'))$$

for all $b \in L$. So, $\diamond(x \wedge x') \in \{0, 1\}$. If $\diamond(x \wedge x') = 0$, then $x \wedge x' \leq \diamond(x \wedge x') = 0$, whence $x \in S_K(\mathbf{L})$. If $\diamond(x \wedge x') = 1$, then

$$x^\sim \leq x^\sim \vee \square x = (x \wedge x')^\sim = 0,$$

and $x \in T(\mathbf{L})$. Our claim is therefore settled.

Recall that \mathbf{L} is directly indecomposable but fails to be an antiortholattice — whence by Corollary 60.(iii) there exist $a, b \in L$ such that $a > (a \wedge b^\sim) \vee (a \wedge \diamond b)$. Also, recall throughout the remainder of this proof that $S_K(\mathbf{L}) \cup T(\mathbf{L}) = L$. If $b \notin S_K(\mathbf{L})$, then

$$a > (a \wedge b^\sim) \vee (a \wedge \diamond b) = a,$$

a contradiction. Therefore $b \in S_K(\mathbf{L})$ and we can apply Lemma 61: either $b = 1$, or $b \leq x$ for every $x \notin S_K(\mathbf{L})$. If $b = 1$, then $a > a$, a contradiction again. If there is some $c \notin S_K(\mathbf{L})$, then $b \leq c$, whence $b = \Box b \leq \Box c = 0$, which yields again the contradiction $a > a$. Therefore $L = S_K(\mathbf{L})$ and \mathbf{L} is orthomodular, against our assumption. ■

Theorem 63 (i) $\mathbb{V}_1 = \text{OML} \vee V(\text{AOL})$.

(ii) $\mathbb{V}_2 = \text{OML} \vee \text{SAOL}$.

(iii) $\mathbb{V}_3 = \text{OML} \vee V(\mathbf{D}_3)$.

Proof. (i) It will suffice to show that any subdirectly irreducible $\mathbf{L} \in \mathbb{V}_1$ is either an orthomodular lattice or an antiortholattice. However, since \mathbf{L} is directly indecomposable, Proposition 62 applies and we obtain our conclusion.

(ii) Any subdirectly irreducible, and thus directly indecomposable, member of \mathbb{V}_2 is either orthomodular, or an antiortholattice satisfying SDM; since SAOL is generated by such antiortholattices, our claim follows.

(iii) This follows, as above, from the fact that $V(\mathbf{D}_3)$ is axiomatised by SK relative to $V(\text{AOL})$ [15, Cor. 3.3]. ■

An upshot of this theorem is that $\mathbb{V}_3 \subset \mathbb{V}_2 \subset \mathbb{V}_1 \subset \text{SDM} \vee V(\text{AOL})$, where the last strict inclusion is witnessed by the PBZ*-lattice \mathbf{H} above which satisfies SDM, thus also WSDM, but fails J2, thus showing in passing that $\mathbb{V}_2 \subset \text{SDM}$. Note, also, that \mathbb{V}_3 is the unique cover of OML in the lattice of subvarieties of PBZL^* , because any member of such which is not included in OML contains \mathbf{D}_3 [14, Thm. 5.5].

Acknowledgement 64 *All authors gratefully acknowledge the following funding sources: the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 689176 (project "Syntax Meets Semantics: Methods, Interactions, and Connections in Substructural logics"); the project "Proprietà d'ordine nella semantica algebrica delle logiche non classiche", Regione Autonoma della Sardegna, L. R. 7/2007, n. 7, 2015, CUP: F72F16002920002. Discussions with Antonio Ledda and Matthew Spinks were extremely helpful in shaping up this paper.*

References

- [1] Aglianò P., Ursini A., "On subtractive varieties II: General properties", *Algebra Universalis*, 36, 1996, pp. 222-259.
- [2] Aglianò P., Ursini A., "On subtractive varieties III: From ideals to congruences", *Algebra Universalis*, 37, 1997, pp. 296-333.
- [3] Aglianò P., Ursini A., "On subtractive varieties IV: Definability of principal ideals", *Algebra Universalis*, 38, 1997, pp. 355-389.

- [4] Bignall R., Spinks M., "On binary discriminator varieties, I, II, and III", typescript.
- [5] Bruns G., Harding J., "Algebraic aspects of orthomodular lattices", in B. Coecke et al. (Eds.), *Current Research in Operational Quantum Logic*, Springer, Berlin, 2000, pp. 37-65.
- [6] Burris S., Sankappanavar H. P., *A Course in Universal Algebra*, Graduate Texts in Mathematics, 78, Springer-Verlag, New York-Berlin, 1981.
- [7] Cattaneo G., Dalla Chiara M. L., Giuntini R., "Some algebraic structures for many-valued logics", *Tatra Mountains Mathematical Publications*, 15, 1998, pp. 173-196.
- [8] Cattaneo G., Giuntini R., Pilla R., "BZMV and Stonian MV algebras (applications to fuzzy sets and rough approximations)", *Fuzzy Sets and Systems*, 108, 1999, pp. 201-222.
- [9] Cattaneo G., Nisticò G., "Brouwer-Zadeh posets and three-valued Lukasiewicz posets", *Fuzzy Sets and Systems*, 33, 2, 1989, pp. 165-190.
- [10] Chajda I., "A note on pseudo-Kleene algebras", *Acta Univ. Palackay Olomouc*, 55, 1, 2016, pp. 39-45.
- [11] Chajda I., Halaš R., Rosenberg I.G., "Ideals and the binary discriminator in universal algebra", *Algebra Universalis* 42, 1999, pp. 239-251.
- [12] Dalla Chiara M.L., Giuntini R., Greechie R.J., *Reasoning in Quantum Theory*, Kluwer, Dordrecht, 2004.
- [13] Font J.M., *Abstract Algebraic Logic: An Introductory Textbook*, College Publications, London, 2016.
- [14] Giuntini R., Ledda A., Paoli F., "A new view of effects in a Hilbert space", *Studia Logica*, 104, 2016, pp. 1145-1177.
- [15] Giuntini R., Ledda A., Paoli F., "On some properties of PBZ* -lattices", *International Journal of Theoretical Physics*, 56, 12, 2017, pp 3895-3911.
- [16] Giuntini R., Mureşan C., Paoli F., "Distributive PBZ* -lattices", in preparation.
- [17] Grätzer G., *Universal Algebra*, Second Edition, Springer Science+Business Media, LLC, New York, 2008.
- [18] de Groote H.F., "On a canonical lattice structure on the effect algebra of a von Neumann algebra", arXiv:math-ph/0410018v2, 2005.
- [19] Kreyszig E., *Introductory Functional Analysis with Applications*, Wiley, New York, 1978.

- [20] Olson M.P., "The self-adjoint operators of a von Neumann algebra form a conditionally complete lattice", *Proceedings of the American Mathematical Society*, 28, 1971, pp. 537-544.
- [21] Salibra A., Ledda A., Paoli F., Kowalski T., "Boolean-like algebras", *Algebra Universalis*, 69, 2, 2013, pp. 113–138.
- [22] Stroock D.W., *A Concise Introduction to the Theory of Integration* (3rd ed.), Birkhäuser, Basel, 1998.
- [23] Ursini A., "On subtractive varieties, I", *Algebra Universalis*, 31, 1994, pp. 204-222.
- [24] Ursini A., "On subtractive varieties, V: congruence modularity and the commutators", *Algebra Universalis*, 43, 2000, pp. 51-78.
- [25] Werner H., *Discriminator Algebras*, Studien zur Algebra und ihre Anwendungen, Band 6, Akademie-Verlag, Berlin, 1978.