OLIVER KNILL

ABSTRACT. Defined by a single axiom, finite abstract simplicial complexes belong to the simplest constructs of mathematics. We look at a few theorems.

Theorems

1. Simplicial complexes

1.1. A finite abstract simplicial complex G is a finite set of non-empty sets which is closed under the process of taking finite non-empty subsets. The **Barycentric** refinement G_1 of G is the set of finite subsets of the power set of G which are pairwise contained into each other. The new complex G_1 defines a finite simple graph $\Gamma = (V, E)$, where V = G and E are the pairs where one is contained in the other. G_1 agrees with the Whitney complex of Γ , the collection of vertex sets of complete sub graphs of Γ .

Theorem: Barycentric refinements are Whitney complexes.

- 1.2. Examples of complexes not coming directly from graphs are buildings or matroids. Oriented matroids are examples of elements of the ring \mathcal{R} generated by simplicial complexes. Still, the Barycentric refinement G_1 of G always allows to study G with the help of graph theory.
- 1.3. A subset H of G is called a **sub-complex**, if it is itself a simplicial complex. Any subset H **generates** a sub-complex, the smallest simplicial complex in G containing H. The set G of sub-complexes is a Boolean lattice because it is closed under intersection and union. The **f-vector** of G is $f = (v_0, v_1, \ldots, v_r)$, where v_k is the number of elements in G with cardinality k + 1. The integer r is the **maximal dimension** of G.

2. Poincaré-Hopf

2.1. A real-valued function $f: G \to \mathcal{R}$ is **locally injective** if $f(x) \neq f(y)$ for any $x \subset y$ or $y \subset x$. In other words, it is a **coloring** in the graph Γ representing G_1 . The **unit sphere** S(x) of $x \in G$ is the set $\{y \in G | (x,y) \in E(\Gamma)\}$. It is the unit sphere in the metric space G, where the distance is the geodesic distance in the graph representing G_1 . Define the stable unit sphere $S_f^-(x) = \{y \in S(x) \mid f(y) < f(x)\}$ and the **index** $i_f(x) = \chi(S_f^-(x))$. The **Poincaré-Hopf theorem** is

Theorem: $\sum_{x} i_f(x) = \chi(G)$.

- 2.2. Classically, for a smooth function with isolated critical points on a Riemannian manifold M, the same definitions and result apply for $i_f(x) = \lim_{r\to 0} \chi(S_{r,f}^-(x))$, where S_r is the geodesic sphere of radius r in M centered at x.
- 2.3. If $f(x) = \dim(x)$, then $i_f(x) = \omega(x)$. Poincaré-Hopf tells then that $\chi(G) = \chi(G_1)$. If $f(x) = -\dim(x)$, then $i_f(x) = \omega(x)(1 \chi(S(x)))$. For complexes for which every unit sphere is a 2d-sphere, we have $i_{\dim} = -i_{-\dim}$ implying $\chi(G) = 0$.

3. Gauss-Bonnet

3.1. Any probability space Ω of locally injective functions defines a **curvature** $\kappa(x) = \mathrm{E}[i_f(x)]$. As we have integrated over f, the curvature value $\kappa(x)$ only depends on x.

Theorem:
$$\sum_{x} \kappa(x) = \chi(G)$$
.

- 3.2. If Ω is the product space $\prod_{x \in G} [-1,1]$ with product measure so that $f \to f(x)$ are independent identically distributed random variables, then $\kappa(x) = K(x)$ is the **Levitt curvature** $1 + \sum_{k=0} (-1)^k v_k(S(x))/(k+1)$. The same applies if the probability space consists of all colorings. If $f = 1 + v_0 t + v_1 t^2 + \ldots$ is the generating function of the f-vector of the unit sphere, with anti-derivative $F = t + v_0 t^2/2 + v_1 t^3/3...$, then $\kappa = F(0) F(-1)$. Compare $\chi(G) = f(0) f(-1)$ and $\sum_x \chi(S(x)) = f'(0) f'(-1)$.
- 3.3. If P is the Dirac measure on $f(x) = \dim(x)$, then the curvature is $\omega(x)$. If P is the Dirac measure on $f(x) = -\dim(x)$, then the curvature is $\omega(x)(1 \chi(S(x)))$.

4. Valuations

4.1. A real-valued function X on \mathcal{G} is called a **valuation** if $X(A \cap B) + \chi(A \cup B) = \chi(A) + \chi(B)$ for all $A, B \in \mathcal{G}$. It is called an **invariant valuation** if X(A) = X(B) if A and B are isomorphic. Let \mathcal{G}_r denote the set of complexes of dimension r. The **discrete Hadwiger theorem** assures:

Theorem: Invariant valuations on \mathcal{G}_r have dimension r+1.

4.2. A basis of the space of invariant valuations is given by $v_k : \mathcal{G} \to \mathbb{R}$. Every vector $X = (x_0, \dots, x_r)$ defines a valuation $X(G) = X \cdot f(G)$ on \mathcal{G}_r .

5. The Stirling formula

5.1. The f-vectors transform linearly under Barycentric refinements. Let Stirling(x, y) denote the **Stirling numbers** of the second kind. It is the number of times one can partition a set of x elements into y non-empty subsets. The map $f \to Sf$ is the **Barycentric refinement operator**

Theorem: $f(G_1) = Sf$, where S(x, y) = Stirling(y, x)x!.

5.2. The matrix is upper triangular with diagonal entries k! the factorial. If $X(G) = \langle X, f(G) \rangle = X(G_1) = \langle X, Sf(G) \rangle = \langle S^T X f(G), \text{ then } X = S^T X \text{ so that } X \text{ is an eigenvector to the eigenvalue 1 of } S^T$. The valuation with X = (1, -1, 1, -1, ...) is the **Euler characteristic** $\chi(G)$. This shows that Euler characteristic is unique:

Theorem: If X(1) = 1 and $X(G) = X(G_1)$ for all G, then $X = \chi$.

6. The unimodularity theorem

6.1. A finite abstract simplicial complex G of n sets defines the $n \times n$ connection matrix L(x,y) = 1 if $x \cap y \neq \emptyset$ and L(x,y) = 0 if $x \cap y = \emptyset$. The unimodularity theorem is:

Theorem: For all $G \in \mathcal{G}$, the matrix L is unimodular.

7. Wu characteristic

7.1. Using the notation $x \sim y$ if $x \cap y \neq \emptyset$, define the **Wu characteristic**

$$\omega(G) = \sum_{x \sim y} \omega(x)\omega(y) .$$

For a complete complex K_{d+1} we have $\omega(K^{d+1}) = (-1)^d$. As every $x \in G$ defines a simplicial complex generated by $\{x\}$, the notation $\omega(x)$ is justified.

7.2. A complex G is a d-complex if every unit sphere is a (d-1)-sphere. A complex G is a d-complex with boundary if every unit sphere S(x) is either (d-1) sphere or a d-1-ball. The sets for which S(x) is a ball form the boundary of G. A complex without boundary is **closed**.

Theorem: For a *d*-complex *G* with boundary, $\omega(G) = \chi(G) - \chi(\delta G)$.

7.3. For any d one can define higher **Wu** characteristic

$$\omega_k(G) = \sum_{x_1 \sim \dots x_k} \omega(x_1) \cdots \omega(x_d)$$

summing over all simultaneously intersecting sets in G.

8. The energy theorem

8.1. As L has determinant 1 or -1, the inverse $g = L^{-1}$ is a matrix with integer entries. The entries g(x,y) are the **potential energy values** between the simplices x,y.

Theorem: For any complex G, we have $\sum_{x} \sum_{y} g(x, y) = \chi(G)$.

8.2. This **energy theorem** assures that the total potential energy of a complex is the Euler characteristic.

9. Номотору

9.1. The graph $1 = K_1$ is **contractible**. Inductively, a graph is **contractible** if there exists a vertex x such that both S(x) and G - x are contractible. The step $G \to G - x$ is a homotopy step. Two graphs are **homotopic** if there exists a sequence of homotopy steps or inverse steps which brings one into the other. Contractible is not the same than homotopic to 1. A graph G is a **unit ball** if there exists a vertex such that B(x) = G.

Theorem: If G is a unit ball then it is contractible.

9.2. It is proved by induction. It is not totally obvious. A **cone extension** G = D + x for the **dunce hat** D obtained by attaching a vertex x to D is a ball but we can not take x away. Any other point y can however be taken away by induction as G - y is a ball with less elements.

Theorem: Contractible graphs have Euler characteristic 1.

9.3. The proof is done by induction starting with G = 1. It is not true that the Wu characteristic $\sum_{x \sim y} \omega(x)\omega(y)$ is a homotopy invariant as $\omega(K_{n+1}) = (-1)^n$.

10. Spheres

10.1. The empty graph 0 is the (-1) sphere. A d-sphere G is a d-graph for which all S(x) are (d-1) spheres and for which there exists a vertex x such that both G-x is contractible. The **1-skeleton graphs** of the octahedron and the icosahedron are examples of 2-spheres. Circular graphs with more than 3 vertices are 1-spheres. A simplicial complex G is a d-sphere, if the graph G_1 is a d-sphere. Here is the **polished Euler Gem**

Theorem: $\chi(G) = 1 + (-1)^d$ for a *d*-sphere G.

Theorem: The join of a *p*-sphere with a *q*-sphere is a p + q + 1-sphere.

10.2. The **generating function** of G is $f_G(t) = 1 + \sum_{k=0}^{\infty} v_k(G) t^{k+1}$ with $v_k(G)$ being the number of k-dimensional sets in G. It satisfies

Theorem: $f_{G+H}(t) = f_G(t) + f_H(t) - 1$ and $f_{G \oplus H}(t) = f_G(t) f_H(t)$.

For example, for $P_2 \oplus P_2 = S_4$ we have $(1 + 2t)(1 + 2t) = 1 + 4t + 4t^2$.

10.3. Given a d-graph. The function $\dim(x)$ has every point a critical point and $S^-(x) = \{y \in S(x) \mid f(y) < f(x)\}$ and $S^+(x) = \{y \in S(x) \mid f(y) > f(x)\}$ then $S(x) = S^-(x) + S^+(x)$.

10.4. Since by definition, a sphere becomes contractible after removing one of its points:

Theorem: d-spheres admit functions with exactly two critical points.

Spheres are the d-graphs for which the minimal number of critical points is 2. There are no d-graphs for which the minimal number of critical points is 1.

11. PLATONIC COMPLEXES

- 11.1. A **combinatorial CW complex** is an empty or finite ordered sequence of spheres $G = \{c_1, \ldots, c_n\}$ such that $G_n = \{c_1, \ldots, c_n\}$ is obtained from $G_{n-1} = \{c_1, \ldots, c_{n-1}\}$ by selecting a sphere c_n in G_{n-1} such that c_n is either empty or different from any c_j . We identify c_j with the cell filling out the sphere. Its dimension is 1 plus the dimension of the sphere. The Barycentric refinement G_1 of G is the Whitney complex of the graph with vertex set G and where two vertices a, b are connected if one is a sub sphere of the other.
- 11.2. G is a d-sphere if G_1 is a d-sphere as a simplicial complex. A subset H of G is a **sub-complex** of G if $H_1 \subset G_1$ for the refinements.
- 11.3. The **Levitt curvature** of a cell c_j is F(0)-F(-1), where F is the anti-derivative of the f-generating function $f = 1 + tv_0 + t^2v_1 + \ldots$ of the sphere $S(c_j)$. The curvature of a cell x in a 2-sphere is $1 v_0(S(x))/2 + v_1(S(x))/3 = 1 v_0(S(x))/6$. The curvature of a cell in a 3 sphere is 0. Gauss-Bonnet assures that the sum of the curvatures is the Euler characteristic.
- 11.4. A d-sphere G is called a **Platonic** d-**polytope** if for every $0 \le k \le d$ and any cell c_j of dimension k, there exists a Platonic (d-1)-sphere P_k such that S(x) is isomorphic to P_k . The -1-dimensional sphere 0 is assumed to be Platonic. The 0-dimensional sphere consisting of two isolated points is Platonic too. The 1-dimensional complexes C_k with $k \ge 3$ are the Platonic 1-spheres. With C_3 one denotes the 1-skeleton complex of K_3 . Let $P = (p(-1), p(0), p(1), p(2), \ldots$ denote the number of Platonic d-polytopes. In the CW case, we have the familiar classification:

Theorem: Platonic_{CW} =
$$(1, 1, \infty, 5, 6, 3, 3, 3, ...)$$
.

11.5. The classification of Platonic polytopes of dimension d which are simplicial complexes is easier. There is a unique platonic solid in each dimension except in dimensions 1, 2, 3. In the 1-dimensional case there are infinitely many. In the two-dimensional case, only the Octahedron and Icosahedron are Platonic. In the three dimensional case, there is only the 600 cell and the 16 cells. After that the curvature condition brings it down to the cross polytopes.

Theorem: Platonic_{SC} =
$$(1, 1, \infty, 2, 2, 1, 1, 1, 1, \dots)$$
.

12. Dehn-Sommerville relations

- 12.1. Given a d-dimensional complex G, any integer vector (X_0, \ldots, X_d) in Z^{d+1} defines a **valuation** $X(G) = X_0v_0 + \ldots X_dv_d$. By distributing the values X_k attached to each k-simplex in G equally to its k+1 vertices, we get the **curvature** $K(x) = \sum_{k=0}^d X_k v_{k-1}(S(x))/(k+1)$ for the valuation X and graph G at the vertex x. The formula $\sum_{x \in V} K(x) = X(G)$ is the **Gauss-Bonnet theorem** for X.
- 12.2. In the case $X(G) = v_1$, the curvature is the vertex degree divided by 2 and the formula reduces to the "Euler handshake". If $X = v_d$ is the volume of G, then K(x) is the number of d-simplices attached to x divided by d+1. In the case $X = (1, -1, 1, -1, \ldots)$, X is the **Euler characteristic** and K is the discrete analogue of the Euler form in differential geometry entering the Gauss-Bonnet-Chern theorem. For d-graphs, there are some valuations which are zero. Define the **Dehn-Sommerville** valuations

$$X_{k,d} = \sum_{j=k}^{d-1} (-1)^{j+d} {j+1 \choose k+1} v_j(G) + v_k(G) .$$

Theorem: For d-graphs, the Dehn-Sommerville curvatures are zero.

12.3. The proof is by noticing that the curvature of $X_{k,d}$ is $K(x) = X_{k-1,d-1}(S(x))$. This follows from the relation

$$X_{k+1,d+1}(l+1)/(l+1) = X(k,d)(l)/(k+2)$$
.

Use Gauss-Bonnet and induction using the fact that the unit sphere of a geometric graph is geometric and that for d=1, a geometric graph is a cyclic graph C_n with $n \geq 4$. For such a graph, the Dehn-Sommerville valuations are zero.

13. Dual Connection Matrix

13.1. Define the dual connection matrix $\overline{L}(x,y) = 1 - L(x,y)$. It is the adjacency graph of a dual connection graph, where two simplices are connected, if they do not intersect.

Theorem:
$$1 - \chi(G) = \det(-L\overline{L})$$
.

13.2. Let E be the constant matrix E(x,y) = 1. The result follows from unimodularity $\det(L) = \det(g)$ and the energy theorem telling that $\overline{L}g = (E-L)g = Eg-1$ has the eigenvalues of Eg minus 1 which are $\chi(G)$ and 0. Assume G has n sets:

Theorem: $-L\overline{L}$ has n-1 eigenvalues 1 and one eigenvalue $1-\chi(G)$.

13.3. The above formula is not the first one giving the Euler characteristic as a determinant of a Laplacian. [10] show, using a formula of Stanley, that if A(x, y) = 1 if x is not a subset of y and A(x, y) = 0 else, then $1 - \chi(G) = \det(A)$.

14. Alexander Duality

14.1. The **Alexander dual** of G is the simplicial complex $G^* = \{x \in V \mid x^c \notin G\}$. It is the complex generated by the complements x^c of the sets x in G. For the complete complex K_d , the dual is the empty complex. In full generality one has for the Betti numbers $b_k(G)$

Theorem:
$$b_k(G) = b_{n-3-k}(G), k = 1, ..., n-1$$

14.2. In order to have content, this needs $n \geq 5$. It works for $G = C_5$ already, where G^* is the complement of a circle in a 3-sphere. The combinatorial Alexander duality is due to Kalai and Stanley.

15.1. Given a locally injective function f on a graph G = (V, E), define for $c \notin f(V)$ the **level surface** $\{f = c\}$ as the subgraph of the Barycentric refinement of G generated by simplices x on which f changes sign. Remember that G is a d-graph if every unit sphere S(x) is a (d-1)-sphere. A discrete Sard theorem is:

Theorem: For a d-graph, every level surface is a (d-1)-graph.

If G is a finite abstract simplicial complex, then $f: G \to R$ defines a function on the Barycentric refinement G_1 and the level surface is defined like that. This result has practical value as we can define discrete versions of classical surfaces.

15.2. Given a finite set of functions f_1, \ldots, f_k on the vertex sets of Barycentric refinements G_1, \ldots, G_k of a simplicial complex, we can now look at the (d-k)-graph $\{f=c\}=\{f_1=c_1,\ldots,f_k=c_k\}$. Unlike in the continuum case, where the result only holds for almost all c, this holds for all c disjoint from the range.

Theorem: Given
$$f_j: G_j \to \mathcal{R}, j \leq k$$
, then $\{f = c\}$ is a $(d-k)$ -complex.

16. Bonnet and Synge

- 16.1. The topic of positive curvature complexes is analog to the continuum. Still, it would be nice to have entirely combinatorial proofs of the results in the continuum.
- 16.2. Let G be a d-complex so that every unit sphere is a (d-1) sphere. A **geodesic** 2-surface is a subcomplex if the embedded graph does not contain a 3-simplex. G has positive sectional curvature if every geodesic embedded wheel graph W(x) has interior curvature $\geq 5/6$. The **geomag lemma** is that any wheel graph in a positive curvature G can be extended to an embedded 2-sphere.
- 16.3. An elementary analog of the **Bonnet theorem**

Theorem: A positive curvature complex has diameter ≤ 4 .

16.4. The simplest analog of **Synge theorem** is

Theorem: A positive curvature complex is simply connected.

- 16.5. The reason for both statements is the **geomag lemma** stating that any closed geodesic curve can be extended to a 2-complex which is a sphere and so simply connected. The strict curvature assumption as we can not realize a projective plane yet with so few cells. With weaker assumptions getting closer to the continuum, we also have to work harder:
- 16.6. Define more generally the **sectional curvature to be** $\geq \kappa$ if there exists M such that the total interior curvature of any geodesic embedded 2-disk with M interior points is $\geq \delta \cdot M$ and such that every geodesic embedded wheel graph W(x) has nonnegative interior curvature. A complex has **positive curvature** if there exists $\kappa > 0$ such that G has sectional curvature $\geq \kappa$. The maximal κ which is possible is then the "sectional curvature bound".
- 16.7. An embedded 2-surface of positive sectional curvature κ must then have surface area $\leq 2/\kappa$. The classical theorem of Bonnet assures that a Riemannian manifold of positive sectional curvature is compact and satisfies an upper diameter bound π/\sqrt{k} . An analog bound C/\sqrt{k} should work in the discrete.
- 16.8. Having a notion of sectional curvature allows to define **Ricci curvature** of an edge e as the average over all sectional curvatures over all wheel graphs passing through e. The **scalar curvature** at a vertex x is the average Ricci curvatures over all edges e containing x. The **Hilbert functional** is then the total scalar curvature. Unlike in Regge calculus, all these notions are combinatorial and do not depend on an embedding.

17. An inverse spectral result

17.1. Let p(G) denote the number of positive eigenvalues of the connection Laplacian L and let n(G) the number of negative eigenvalues of L.

Theorem: For all
$$G \in \mathcal{G}$$
 we have $\chi(G) = p(G) - n(G)$.

17.2. The proof checks this by deforming L when adding a new cell. This result implies that Euler characteristic is a logarithmic potential energy of the origin with respect to the spectrum of iL.

Theorem:
$$\chi(G) = \operatorname{tr}(\log(iL))(2\pi/i)$$
.

17.3. The proof shows also that after a CW ordering of the sets in a finite abstract simplicial complex, one can assign to every simplex a specific eigenvalue and so eigenvector of L.

18. The Green star formula

18.1. Given a simplex $x \in G$, the **stable manifold** of the dimension functional $\dim(x)$ is $W^-(x) = \{y \in G \mid y \in x\}$. The **unstable manifold** $W^+(x) = \{y \in G \mid x \in y\}$ is known as the **star** of x. Unlike $W^-(x)$ which is always a simplicial complex, the star $W^+(x)$ is in general not a sub complex of G.

Theorem: $g(x,y) = \omega(x)\omega(x)\chi(W^+(x)\cap W^+(y)).$

18.2. In comparison, we have $W^-(x) \cap W^+(x) = \omega(x)$ and $L(x,y) = \chi(W^-(x) \cap W^-(y))$. The to L similar matrix $M(x,y) = \omega(x)\omega(x)\chi(W^-(x) \cap W^-(y))$ satisfies $\sum_x \sum_y M(x,y) = \omega(G)$, the Wu characteristic.

19. Wu characteristic

- 19.1. The **Euler characteristic** $\chi(G) = \omega_1(G) = \sum_{x \in G} \omega(x)$ of G is the simplest of a sequence of combinatorial invariants $\omega_k(G)$. The second one, $\omega(G) = \sum_{x,y,L(x,y)=1} \omega(x)\omega(y)$, is the **Wu characteristic** of G. The valuation χ is an example of a linear valuation, while ω is a **quadratic valuation**. The Wu characteristic also defines an **intersection number** $\omega(A,B)$ between sub-complexes.
- 19.2. All multi-linear valuations feature Gauss-Bonnet and Poincaré-Hopf theorems, where the curvature of Gauss-Bonnet is an index averaging. For example, with $K(v) = \sum_{v \in x, x \sim y} \omega(x) \omega(y) / (|x| + 1)$ The Gauss-Bonnet theorem for Wu characteristic is

Theorem: $\omega(G) = \sum_{v} K(v)$.

20. The boundary formula

- 20.1. We think of the **internal energy** $E(G) = \chi(G) \omega(G)$ as a sum of **potential energy** and **kinetic energy**. A **d-complex** is a simplicial complex G for which every S(x) is a (d-1)-sphere. A **d-complex with boundary** is a complex S(x) is either a (d-1)-sphere or a (d-1)-ball for every $x \in G$.
- 20.2. The *d*-complexes are **discrete** *d*-manifolds and *d*-complexes with boundary is a discrete version of a *d*-manifold with boundary. We denote by δG the boundary of G. It is the d-1 complex consisting of boundary points. By definition, $\delta \delta G = 0$, the empty complex. The reason is that the boundary of a complex is closed, has no boundary. We can reformulate the formula given below as

Theorem: If G is a d-complex with boundary then $E(G) = \chi(\delta(G))$.

20.3. If G is a d-ball, then δG is a (d-1)-sphere and $E(G) = 1 + (-1)^{d-1}$, by the **polished Euler gem formula**.

21. Zeta function

21.1. For a one-dimensional complex G, there is a **spectral symmetry** which will lead to a **functional equation**:

Theorem: If dim(G) = 1, then $\sigma(L^2) = \sigma(L^{-2})$.

21.2. If H is a Laplacian operator with non-negative spectrum like the **Hodge operator** H or **connection operator** L, one can look at its **zeta function**

$$\zeta_H(s) = \sum_{\lambda \neq 0} \lambda^{-s} ,$$

where the sum is over all non-zero eigenvalues of H or L^2 . In the connection case, we take L^2 to have all eigenvalues positive.

21.3. The case of the connection Laplacian is especially interesting because one does not have to exclude any zero eigenvalue. The **connection zeta function** of G is defined as $\zeta(s) = \sum_{\lambda} \lambda^{-s}$, where the sum is over all eigenvalues λ of L^2 . It is an entire function in s.

Theorem: If $\dim(G) = 1$, then $\zeta(s) = \zeta(-s)$.

21.4. When doing Barycentric refinement steps, the zeta function converges to an explicit function.

$$\zeta(it) = \int_0^1 \frac{2\cos\left(2t\log\left(\sqrt{4v^2 + 1} + 2v\right)\right)}{\pi\sqrt{1 - v}\sqrt{v}} dv.$$

It is a hypergeometric series $\zeta(2s) = \pi_4 F_3\left(\frac{1}{4}, \frac{3}{4}, -s, s; \frac{1}{2}, \frac{1}{2}, 1; -4\right)$.

22. The Hydrogen formula

22.1. Given a simplicial complex G, let $\Lambda_k(G)$ denote the set of real valued functions on k-dimensional simplices. It is a v_k -dimensional vector space. Define the $v_k \times v_{k+1}$ matrices $d_k(x,y) = 1$ if $x \subset y$ and $d_k(x,y) = 0$ else. It is the **sign-less incidence matrix**. It can be extended to a $n \times n$ matrix d so that $d = d_0 + d_1 + \cdots + d_r$ and $D = d + d^*$ and $H = (d + d^*)^2$, the **sign-less Dirac** and **sign-less Hodge operator**. In the one-dimensional case, we have $H = d^*d + dd^*$. The **Hydrogen relations** are

Theorem: If
$$\dim(G) = 1$$
, then $L - L^{-1} = H$.

- 22.2. The relation allows to relate the spectra of L and H. It allows to estimate the spectral radius or give explicit formulas for the spectrum of the connection Laplacian in the circular case. This is needed to get the explicit **dyadic zeta function**
- 22.3. Let S(x) denote the **unit sphere** of a simplex $x \in G$. While S(x) is at first a subset of G, it generates a sub-complex in G_1 . As $g(x,x) = 1 \chi(S(x)) = \chi(W^+(x))$, we have a functional $\sum_x \chi(S(x))$ of Dehn-Sommerville type. With $f(t) = 1 + \sum_{k=1}^{\infty} v_{k-1}t^k = 1 + v_0t + v_1t^2 + v_2t^3 + \cdots$, the Euler characteristic of G_1 can be written as $\chi(G) = f(0) f(-1)$. The following result holds for any simplicial complex:

Theorem:
$$tr(L - L^{-1}) = \sum_{x} \chi(S(x)) = f'(0) - f'(-1).$$

22.4. Compare that the Levitt curvature at a point x was F(0) - F(-1), where F is the anti-derivative of the generating function of S(x).

23. Brouwer-Lefschetz

- 23.1. The **exterior derivative** d for G defines the **Dirac operator** $D = d + d^*$ of d. The Hodge Laplacian $H = D^2$ splits into a direct sum $H_0 \oplus H_1 \cdots H_d$. The null space of H_k is isomorphic to the k'th cohomology group $H^k(G) = \ker(d_k)/\operatorname{im}(d_{k-1})$. Its dimension b_k is the k'th Betti number. The **Euler-Poincaré** relation assures that the cohomological Euler characteristic $\sum_k (-1)^k b_k$ is equal to the Euler characteristic.
- 23.2. An **endomorphism** T of G is a map from G to G which preserves the order structure. It is an automorphism if it is bijective. An endomorphism T induces a linear map on cohomology $H^k(G)$. The super trace of this map is the **Lefschetz number** $\chi(T,G)$ of T. Given a fixed point $x \in G$ of T, its **Brouwer index** is defined as $i_T(x) = \omega(x) \operatorname{sign}(T|x)$. Now

Theorem:
$$\chi(T,G) = \sum_{x=T(x)} i_T(x)$$
.

23.3. A special case is T = 1, where $\chi(1, G) = \chi(G)$ and $i_T(x) = \omega(x)$. The Brouwer-Lefschetz fixed point theorem is then the Euler-Poincaré theorem.

24. McKean-Singer

24.1. The super trace $\operatorname{str}(A)$ of a $n \times n$ matrix defined for a complex with n sets is defined as $\sum_{x \in G} \omega(x) L(x,x)$. By definition, we have $\operatorname{str}(1) = \operatorname{str}(L)$. For the Hodge operator $H = D^2 = (d + d^*)^2$ we have the McKean-Singer formula:

Theorem:
$$str(exp(-tH)) = \chi(G)$$
 for all t .

- 24.2. The reason is that $str(H^k) = 0$ for k > 0, implying $str(exp(tH)) = str(1) = \chi(G)$. The McKean-Singer identity is very important as it allows to give almost immediate **proofs** of the Lefschetz formulas in any framework in which the identity holds. We proposed in [108] to define a discrete version of a **differential complex** as McKean-Singer enables Atiyah-Singer or Atiyah-Bott like extensions of Gauss-Bonnet or Lefschetz. They are caricatures of the heavy theorems in the continuum.
- 24.3. The Hodge operator $H=(d+d^*)^2$ and the connection operator L live on the same finite dimensional Hilbert space. There is no cohomology associated to L. But for the connection operator L, there is still a localized version of McKean-Singer:

Theorem:
$$str(L^k) = \chi(G)$$
 for $k = -1, 0, 1$.

25. Barycentric limit

25.1. A matrix L with eigenvalues $\lambda_0 \leq \lambda_2 \leq \cdots \leq \lambda_{n-1}$ defines a **spectral function** $F(x) = \lambda_{[nx]}$ on [0,1), where [t] is the floor function giving the largest integer smaller or equal than t. The inverse function $k(x) = F^{-1}(x)$ is called the **integrated density** of states of L and $\mu = k'$ is the **density of states**. The sequence G_k of Barycentric refinements of G defines a sequence of operators L_k and so a sequence of spectral

functions $F_n(x)$. Let \mathcal{G}_r denote the set of complexes of dimension r. The following spectral universality is a central limit theorem:

Theorem: $\exists F = F(r) \text{ such that } F_n(G) \to_{L^1} F \text{ for all } G \in \mathcal{G}_r.$

- 25.2. For r=1, we know $F(x)=4\sin^2(\pi x/2)$. The function is important as it conjugates the **Ulam map** $z\to 4x(1-x)$ to a linear function T(F(x))=F(2x). The measure μ maximizes **metric entropy** of the Ulam map and is equal to the **topological entropy** which is $\log(2)$ for T.
- 25.3. We think of $G_n \to G_{n+1}$ as a renormalization step like adding and normalizing two independent random variables. The result can be seen as a **central limit** theorem.

26. The join monoid

- 26.1. The **join** G+H of two complexes G, H is the complex $G \cup H \cup \{x \cup y, x \in H, y \in G\}$. For graphs it is known as the **Zykov sum**. Given graphs G = (V, E), H = (W, F) then the sum is $(V \cup W, E \cup F \cup \{(a, b) \mid a \in V, b \in W\})$. If \overline{G} denotes the **complement graph** and + the disjoint union, then $\overline{G \oplus H} = \overline{G} + \overline{H}$.
- 26.2. The **join** of two simplicial complexes G, H is defined as the complex generated by $G + H = G \cup H \cup \{x \cup y \mid x \in G, y \in H\}$. Let $f_G(t) = 1 + v_0 t + v_1 t^2 + \ldots$ denote the **generating function** of G: then we have the multiplication formula:

Theorem: $f_{G+H}(t) = f_G(t) f_H(t)$.

26.3. This gives $1 - \chi(G) = f_G(-1)$. The dimension function on G not only defines a coloring on G_1 , it also defines a hyperbolic splitting of the unit spheres. Let $S^-(x) = \{y \in S(x), \dim(y) < \dim(x)\}$ and $S^+(x) = \{y \in S(x), \dim(y) > \dim(x)\}$. We call them the **stable sphere** and **unstable sphere**.

Theorem: $S(x) = S^{-}(x) + S^{+}(x)$.

- 26.4. It follows that $g(x,x) = 1 \chi(S(x)) = (1 \chi(S^-(x)))(1 \chi(S^+(x))) = \omega(x)(1 \chi(S^+(x)))$. This implies that $\operatorname{str}(L^{-1}) = \sum_x (1 \chi(S^+(x))) = \chi(G)$ because this is the sum over the Poincaré-Hopf indices of the function —dim.
- 26.5. The join monoid is isomorphic to the additive monoid of disjoint union. The **zero element** is 0, the -1 sphere. One can show by induction that if H is contractible and K arbitrary then H + K is contractible. This implies:

Theorem: The join G of two spheres H + K is a sphere.

26.6. For example, the join of two zero dimensional spheres P_2 is the circle C_4 . The join of two circles a three sphere. It is not the dimension but the **clique number** $\dim(G) + 1$ which is additive. The clique number of the -1 sphere 0 is 0.

27. The strong ring

27.1. The addition A+B of two complexes is the disjoint union. The empty complex 0 is the **zero element**. The **Cartesian product** $G \times H$ is not a simplicial complex any more. We can look at the ring \mathcal{R} generated by simplicial complexes. It has the one point complex $1 = K_1$ as **one element**. Connected elements are the additive **primes**, simplicial complexes are **multiplicative primes**. The **Hodge operator** H and the **connection operator** L can both be extended to the ring \mathcal{R} .

Theorem:
$$\sigma(H(A \times B)) = \sigma(H(A)) + \sigma(H(B)),$$

27.2. Furthermore:

Theorem:
$$\sigma(L(A \times B)) = \sigma(L(A)) \cdot \sigma(L(B))$$

28. Kuenneth formula

28.1. The **Betti numbers** of a signed complex $b_k(G)$ are now signed with $b_k(-G) = -b_k(G)$. The maps assigning to G its Poincaré polynomial $p_G(t) = \sum_{k=0} b_k(G)t^k$ or **Euler polynomial** $e_G(t) = \sum_{k=0} v_k(G)t^k$ are ring homomorphisms from R to $\mathbb{Z}[t]$. Also $G \to \chi(G) = p(-1) = e(-1) \in \mathbb{Z}$ is a ring homomorphism.

Theorem: e_G and p_G are ring homomorphisms $\mathcal{R} \to \mathbb{Z}[t]$.

- 28.2. The **Kuenneth formula** for cohomology groups $H^k(G)$ is explicit via Hodge: a basis for $H^k(A \times B)$ is obtained from a basis of the factors. The product in R produces the strong product for the connection graphs. These relations generalize to Wu characteristic. R is a subring of the full **Stanley-Reisner ring** S, a subring of a quotient ring of the polynomial ring $Z[x_1, x_2, \ldots]$. An object $G \in R$ can be visualized by ts Barycentric refinement G_1 and its connection graph G'.
- 28.3. Theorems like Gauss-Bonnet, Poincaré-Hopf or Brouwer-Lefschetz for Euler and Wu characteristic extend to the strong ring. The isomorphism $G \to G'$ to a subring of the strong Sabidussi ring shows that the multiplicative primes in R are the simplicial complexes and that connected elements in R have a unique prime factorization.
- 28.4. The **Sabidussi ring** is dual to the Zykov ring. The Zykov join was the addition which is a sphere-preserving operation. The Barycentric limit theorem implies that the connection Laplacian remains invertible in the limit.

29. Dimension

29.1. The **inductive dimension** of a graph is defined inductively as $\dim(G) = 1 + \sum_{v \in V} \dim(S(x))/|V|$. For a general complex G we can define $\dim(G) = \dim(G_1)$, as G_1 is now the Whitney complex of a graph. We have $\dim(G) \leq \max_{x \in G} (|x| - 1)$, where the right hand side is the **maximal dimension**.

Theorem:
$$\dim(A \times B) = \dim(A) + \dim(B)$$
.

29.2. Under Barycentric refinements, the inductive dimension can only increase.

Theorem: $\dim(G_1) \ge \dim(G)$

- 29.3. The reason is that higher dimensional complexes have more off-springs than smaller dimensional ones.
- 29.4. This implies a inequality which resembles the corresponding inequality for **Hausdorff dimension** in the continuum:

Theorem: $\dim((A \times B)_1) \ge \dim(A) + \dim(B)$.

30. Random complexes

30.1. Given a probability space of complexes, one can study the expectations of random variables. The simplest probability space is the **Erdös-Rényi space** E(n,p) of random graphs equipped with the Whitney complex. Define the polynomials $d_n(p)$ of degree $\binom{n}{2}$ as

$$d_{n+1}(p) = 1 + \sum_{k=0}^{n} {n \choose k} p^{k} (1-p)^{n-k} d_{k}(p) ,$$

where $d_0 = -1$. We can now estimate the inductive dimension.

Theorem: $E_{G(n,p)}[\dim] = d_n(p).$

30.2. As the Euler characteristic is one of the most important functionals, we want to estimate its expectation:

Theorem:

$$E_{G(n,p)}[\chi] = \sum_{k=1}^{n} (-1)^{k+1} \binom{n}{k} p^{\binom{k}{2}}.$$

30.3. We don't yet know the expectation value of the Wu characteristic on E(n, p).

31. Lusternik-Schnirelmann

- 31.1. A complex G is **contractible** if there exists $x \in G$ such that both the unit sphere S(x) as well as the complex $G \setminus x$ are contractible. A complex is **homotopic** to K=1 if there there exists a complex H such that H is contractible to both G and K. The **dunce hat** is an example of a complex homotopic to 1 which is not contractible. The minimal number of contractible subcomplexes of G covering G is called the Lusternik-Schnirelman category of G.
- 31.2. A $x \in G$ is called a **critical point** of a function f if $S_f^-(x)$ is not contractible. The minimal number of critical points which a function f on G can have is denoted by cri(G).

- 31.3. There is a graded multiplication $H^k(G) \times H^l(G) \to H^{k+l}(G)$ called the **cup product**. If m-1 is the maximal number of p > 0-forms f_1, \ldots, f_{m-1} for which $f_1 \cup \cdots \cup f_{m-1}$ is not zero, then m is called the **cup length** of G.
- 31.4. The following result, established with Josellis in 2012 is completely analog to the continuum.

Theorem: $cup(G) \le cat(G) \le cri(G)$.

31.5. For any critical point x_i , we can form the maximal complex G_i which does not contain an other critical point. Each U_i is contractible and cover G. This proves $\operatorname{cat}(G) \leq \operatorname{cri}(G)$. If $\operatorname{cat}(G) = n$, let $\{U_k\}_{k=1}^n$ be a **Lusternik-Schnirelmann cover**. Given a collection of $k_j \geq 1$ -forms f_j with $f_1 \wedge f_2 \cdots \wedge \ldots f_n \neq 0$. Using coboundaries we can achieve that for any simplex $y_k \in U_k$, we can change f in the same cohomology class f so that $f(y_k) = 0$. Because U_k are contractible in G, we can render f zero in U_k . This shows that we can choose f_k in the relative cohomology groups $H^k(G, U_k)$ meaning that we can find representatives k_j forms f_j which are zero on each p_{k_j} simplices in the in G contractible sets U_k . But now, taking these representatives, we see $f_1 \wedge \cdots \wedge f_n = 0$. This shows $\operatorname{cup}(G) \leq n$.

32. Morse inequality

32.1. A locally injective scalar function f on the vertex set of a d-graph is called a **Morse function**, if $S_f^-(x)$ is a sphere for every x. The **Morse index** is $m(x) = 1 + \dim(S_f^-(x))$. The **Poincaré-Hopf index** is $(-1)^{m(x)}$. For example, if d = 2, and $S_f^-(x)$ is 0-dimensional, then m(x) = 1 and $i_f(x) = -1$. A function f on an abstract simplicial d-complex G is a Morse function if it is a Morse function on the graph G_1 .

Theorem: Every d-complex admits a Morse function.

32.2. We can build up G as a **discrete** CW-**complex**. The number at which a simplex x has been added is a Morse function as S(x) and $S^{-}(x)$ are both spheres. Also the function $\dim(x)$ is a Morse function. For d-complexes, the stars of two simplices intersect in a simplex so that:

Theorem: For a *d*-complex, the Green function takes values 1, -1, 0.

We have $g(x,y) = \omega(x)\omega(y)\chi(W^+(x)\cap W^+(y))$. We have $W^+(x)\cap W^+(y) = (1-S^+(x))(1-\chi(S^+(x)))$ which is in $\{-1,1\}$ if there is an intersection and 0 if not. Let $b_k(G)$ denote the k'th Betti number. Let $c_k(G)$ denote the number of critical points of index k. Here are the **weak Morse inequalities**:

Theorem: $b_k(G) \leq c_k(G)$.

We even have the strong Morse inequalities

Theorem: $(-1)^p \sum_{k=0}^p (-1)^k (c_k - b_k) \ge 0$

By Euler-Poincaré, this is zero for the entire sum. It appears as if the Witten deformatin proof (see e.g. [32]) works in the discrete too.

33. ISOSPECTRAL DEFORMATION

33.1. If d is the exterior derivative, the operator $D = d + d^*$ is the **Dirac operator** of G. The Dirac operator D admits an **isospectral Lax deformations** D' = [B, D] = BD - DB, where $B = d - d^* + \gamma ib$, if $D = d + d^* + b$. The parameter γ is a tuning parameter. For $\gamma = 0$ the deformation stays real. For $\gamma \neq 0$, it is allowed to become complex. The Dirac operator D(t) defines for every t an **elliptic complex** $D: E \to F$ meaning that we have a splitting $D(t): E \to F$ such that McKean-Singer relation holds.

Theorem: The Lax system for the Dirac operator is integrable.

- 33.2. The spectrum of D(t) stays constant. Actually, $L = D(t)^2$ stays constant.
- 33.3. We have a deformation of the complex for which all classical geometry like the wave equation stays the same because L does not change. It is only the underlying d which changes. The **Connes formula** $\sup_{|Df|_{\infty}=1} |f(x) f(y)|$ allows to re-interpret the isospectral deformation as a deformation of the metric.

34. Trees and Forests

- 34.1. Given a finite simple graph G, a **rooted spanning tree** is a subgraph H of G which is a tree with the same vertex set together with a base point x. A **rooted spanning forest** is a subgraph H of G which is a forest with the same vertex set together with a base point x. Let K be the **Kirchhoff Laplacian** of the graph and Det(K) the **pseudo determinant**, the product of the non-zero eigenvalues of K. It is $\exp(-\zeta'(0))$ for the zeta function of K.
- 34.2. The **tree number** of a graph G is the number of rooted spanning tree in G. The **forest number** of a graph is the number of rooted spanning forests. The first part of the following theorem is the **Kirchhoff matrix tree theorem**. The second part of the theorem is the **Chebotarev-Shamis forest theorem**.

Theorem: Det(K) is the tree number. det(K+1) is the forest number.

- 34.3. By Baker-Norine theory, the tree number is also the order of the **Picard group** which appears in the context of discrete Riemann-Roch.
- 34.4. If F, G are arbitrary $n \times m$ matrices. Assume $p(x) = p_0(-x)^m + p_1(-x)^{m-1} + \cdots + p_k(-x)^{m-k} + \cdots + p_m$ is the **characteristic polynomial** of the $m \times m$ matrix F^TG with $p_0 = 1$. The **generalized Cauchy-Binet theorem** is

Theorem:
$$p_k = \sum_{|P|=k} \det(F_P) \det(G_P)$$

where the sum is over k-minors and where p_k are the coefficients of the characteristic polynomial of F^TG . It implies the polynomial identity $\det(1 + zF^TG) = \sum_P z^{|P|} \det(F_P) \det(G_P)$ in which the sum is over all minors A_P including the empty one |P| = 0 for which $\det(F_P) \det(G_P) = 1$.

35. Wave equation

35.1. Because the Hodge Laplacian is a square $L = D^2 = (d + d^*)^2$, the wave equation $u_{tt} = Lu$ has an explicit d'Alembert solution. Let D^{-1} be the pseudo inverse of D. It is defined as $\sum_{k,\lambda_k\neq 0} u_k u_k^T/\lambda_k$, where $Du_k = \lambda_k u_k$ with an orthonormal eigenbasis $\{u_k\}$ of D.

Theorem:
$$u(t) = \cos(Dt)u(0) + i\sin(Dt)D^{-1}u'(0)$$

35.2. With the **complex wave** $\psi(t) = u(t) - iDu'(0)$, we can write the solution of the **real wave** equation of u as a solution of the **Schrödinger equation**.

Theorem:
$$\psi(t) = e^{iDt}\psi(0)$$
.

35.3. Just use the Euler identity $e^{iDt} = \cos(Dt) + i\sin(Dt)$ and plug in $\psi(t) = u(t) - iDu'(0)$ to see that the relation holds.

36.1. Let $\Lambda^p(G)$ be the functions from $G_p = \{x \in G \mid \dim(x) = k \}$ to R which are anti-symmetric. The **exterior derivatives**

$$d_p f(x_0, x_1, \dots, x_p) = \sum_j (-1)^j f(x_0, \dots, \hat{x_j}, \dots, x_p)$$

define linear map $d: \Lambda(G) \to \Lambda(G)$, where $\Lambda(G)$ is the Hilbert space of dimension n = |G|. Since $d^2 = 0$, the **cohomology groups** $H^p(G) = \ker(d_p)/\operatorname{im}(d_{p-1})$ are defined. Their dimensions are the Betti numbers $b_p(G)$. The matrix $H = (d + d^*)^2$ decomposes into blocks $H_k(G)$. We have the **Hodge relations**:

Theorem:
$$\dim(\ker(H_k)) = \dim(H^k)$$
.

36.2. Define the **Poincaré polynomial** $p_G(t) = \sum_{k=0} b_k(G)t^k$. The **cohomological Euler characteristic** is $p_G(-1) = b_0(G) - b_1(G) + b_2(G) - \cdots$. If the f-vector of G is (v_0, v_1, v_2, \ldots) , then the **Euler polynomial** is $e_G(t) = \sum_{k=0} v_k(G)t^k$. By definition, we have $d_G(-1) = \chi(G)$. The **Euler-Poincaré theorem** tells that the combinatorial and cohomological Euler characteristic agree.

Theorem:
$$\chi(G) = e_G(-1) = p_G(-1)$$
.

37. Interaction cohomology

37.1. Let $\Lambda_2^p(G)$ be the functions from $G_p = \{(x,y) \mid x \cap y \neq \emptyset, \dim(x) + \dim(y) = p\}$ which are anti-symmetric. Like **Stokes theorem** $df(x) = f(\delta x)$ for simplicial cohomology, we define the **exterior derivative** $df((x,y)) = f(\delta x,y) + (-1)^{\dim}(x)f(x,\delta y)$ with the understanding that $f(\delta x,y) = 0$ if $\delta x \cap y = \emptyset$ or $f(x,\delta y) = 0$ if $x \cap \delta y = \emptyset$. It defines a linear map $d: \Lambda_2(G) \to \Lambda_2(G)$, where $\Lambda_2(G)$ has as dimension the number of intersecting simplices (x,y) in G. Again, we can define the **Dirac operator** $D = d + d^*$ and the **Hodge operator** $H = D^2$ and decompose the later into blocks H_k . As before:

Theorem: $\dim(\ker(H_k)) = \dim(H^k)$.

37.2. The quadratic Poincaré polynomial $p_G(t) = \sum_{k=0} b_k(G)t^k$ and quadratic Euler polynomial $e_G(t) = \sum_{k=0} v_k(G)t^k$ are defined in the same way. By definition, we have $d_G(-1) = \chi(G)$. The Euler-Poincaré theorem tells that the combinatorial and cohomological Wu characteristic agree.

Theorem: $\omega(G) = e_G(-1) = p_G(-1)$.

38. Stokes theorem

- 38.1. Examples of **orientation oblivious** measurements are valuations F like $F(A) = v_k(A)$ measuring the k dimensional volume of a subcomplex A of G or $\chi(A)$ giving the Euler characteristic of a subcomplex. The **length** of a subcomplex A for example is $v_1(A)$. In the continuum, such quantities are accessible via **integral geometry**, like **Crofton type formulas**. In the discrete one refers to it also as]bf geometric probability theory.
- 38.2. If valuations are done after an orientation has been chosen on the elements of G, we get a **calculus** which features a **fundamental theorem**. Given an arbitrary choice of orientation of the sets in G, the boundary δA of a subcomplex is in general no more a subcomplex, it becomes a **chain**. Given a form $F \in \Lambda$, we can still compute $F(\delta A)$. If G is **orientable** d-complex and A is a k-subcomplex with **boundary** δA , then δA is a complex. **Stokes theorem** tells that for any k-subcomplex A with boundary δA , and any k-form F

Theorem: $dF(A) = F(\delta A)$.

- 38.3. For k=1, we talk about the fundamental theorem of line integrals, for k=2 we have Stokes theorem and k=3 goes under the name divergence theorem. The derivative $d_0: \Lambda^0 \to \Lambda^1$ is the gradient, the derivative $d_1: \Lambda^1 \to \Lambda^2$ is the curl and $d_2: \Lambda^2 \to \Lambda^3$ is the divergence (often just identified with the dual $d_0^*: \Lambda^1 \to \Lambda^0$, as 2-forms and 1-forms in three dimensions are dual to each other). This Stokes theorem holds both for the familiar simplicial calculus related to Euler characteristic $\chi(G)$ as well as the connection calculus related to the Wu characteristics $\omega_k(G)$.
 - 39. Quadratic Lefschetz fixed point
- 39.1. Given an automorphism T, define the quadratic Lefschetz number $\chi_T(G)$, the super trace of the induced map on cohomology.

Theorem: $\chi_T(G) = \sum_{x \sim y, (x,y) = (T(x), T(y))} i_T(x,y)$

39.2. We can especially look at the case when G is a ball. This is cohomologically non-trivial.

Theorem: An endomormorphism of a ball G has a fixed $(x, y), x \cap y \neq \emptyset$.

40. Eulerian spheres

40.1. Let \mathcal{G}_d be the class of d-graphs, \mathcal{S}_d the class of d-spheres, \mathcal{B}_d the class of d-balls, and \mathcal{C}_k the class of graphs with **chromatic number** k. Note that all Barycentric refinements of a complex are Eulerian. We call the class $\mathcal{S}_d \cap \mathcal{C}_{d+1}$ the class of **Eulerian** spheres and $\mathcal{B}_d \cap \mathcal{C}_{d+1}$ the class of Eulerian disks. The 0-sphere 2 is Eulerian. Eulerian 1-spheres are cyclic graphs with an even number of vertices.

Theorem: Every unit sphere of an Eulerian sphere is Eulerian.

40.2. The **dual graph** \hat{G} of a d-sphere G is the graph in which the d-simplices are the vertices and where two simplices are connected, if one is contained in the other. A graph (V, E) is **bipartite** if $V = (A \cup B \text{ with disjoint } A, B \text{ such } E = \{(a, b) \mid a \in A, b \in B\}$. Every Barycentric refinement of a complex is a bipartite graph as we can take $A = \{x \in G \dim(x) \text{ even}\}$ and $B = \{x \in G \dim(x) \text{ odd}\}$.

Theorem: For $G \in \mathcal{S}_d$, then \hat{G} is bipartite if and only if G is Eulerian.

41. RIEMANN-HURWITZ

41.1. The **automorphism group** $\operatorname{Aut}(G)$ of a simplicial complex is the group of all **automorphisms** of G. An **endomorphism** T is a simplicial map $G \to G$. If an endomorphism T is restricted to the **attractor** $\bigcap_k T^k(G)$ is an automorphism. An automorphism T of G induces automorphisms on Barycentric refinements and so **graph automorphisms**. The equivalence classes G_1/A are graphs.

Theorem: If $A \subset Aut(G)$, then G_1/A is a simplicial complex.

- 41.2. We can see G_1 as a **branched cover** G_1/A , **ramified** over some points. If G was a d-graph, then G_1/A is a discrete **orbifold**. If there are no ramification points, then the cover $G \to G/A$ is a fibre bundle with structure group A.
- 41.3. Given an automorphism T, define the **ramification index** $e(x) = 1 \sum_{T \neq 1, T(x) = x} \omega(x)$ of X. The following remark was obtained with Tom Tucker. It is a discrete **Riemann-Hurwitz** result:

Theorem:
$$\chi(G) = |A|\chi(G/A) - \sum_{x \in G} (e(x) - 1)$$

- 41.4. For every subset \mathcal{G}_k of indices of fixed dimension k, we have by the **Burnside lemma** $\sum_{T \in A} \sum_{x \in \mathcal{G}_k, T(x) = x} 1 = |A| |\mathcal{G}_k|$. The super sum gives $\sum_{T \in A} \sum_{x, T(x) = x} \omega(x) = |A| \chi(H)$. This gives $\sum_{T \neq 1} \sum_{x \in G} \omega(x) + \sum_{x \in G} \omega(x) = |A| \chi(H)$.
- 41.5. Let $\chi(G,T)$ denote the **Lefschetz number** of T. From the Lefschetz fixed point formula we get

Theorem:
$$\chi(G/A) = \frac{1}{|A|} \sum_{T \in A} L(G,T)$$

42. RIEMANN-ROCH

42.1. A divisor X is an integer-valued function on G. The simplex Laplacian L is defined as $L(x,y) = \omega(x)\omega(y)H_0(x,y)$, where H_0 is the Kirchhoff Laplacian of the simplex graph in which G is the vertex set and two x,y are connected if one is contained in the other and the dimensions differ by 1. The simplex graph is one-dimensional as it has no triangles. A divisor X is called **principal** if X = Lf for some integer valued function f. We think of a divisor as a geometric object and define the Euler characteristic $\chi(G) = \sum_x \omega(x)X(x)$. A divisor is **essential** if $\omega(x)X(x) \geq 0$ for all x. The **linear system** |X| of X is the set of f for which X + (f) is essential. Its **dimension** l(X) is the maximal $k \geq 0$ such that for every m < k and every Y of $\chi(Y) = m$, the divisor X - Y is essential. Define the **canonical divisor** K(x) = 0. The simplest Riemann-Roch theorem is

Theorem:
$$l(X) - l(K - X) = \chi(X)$$
.

- 42.2. This is **Baker-Norine theory**, slightly adapted to change the perspective: classically a divisors appear one a one dimensional connected curve (Riemann surface or 1-dimensional graph) G and $\deg(X) + \chi(G) = \chi(G)$. Centering at the geometric underlying object gives the canonical divisor K = -2 which is in the case when G is one-dimensional is linearly equivalent to the negated curvature function $K(v) = -2 + \deg(v)$ on the **vertices** of G. Riemann-Roch tells that the signed distance to the surface $\chi(G) = 0$ is $\chi(G)$.
- 42.3. Reflecting at 0 rather than at usual canonical divisor representing the curve G allows to have a Riemann-Roch for arbitrary dimensions. Generalizing Baker-Norine naively to higher dimensional simplicial complexes does not work, as the curvature κ of $\chi(G)$ has only in the one-dimensional case the property that $K = -2\kappa$ is a divisor. Classically l(X), L(K-X) have cohomological interpretations. Also here, Riemann-Roch appears like a fancy Euler-Poincaré formula, but it is deeper than the later, as surface $\ker(\chi)$ is **bumpy**: it contains both **generic divisors** as well as **special divisors**.
- 42.4. The image of L is a linear subspace of the set $\ker(G) = \chi(G) = 0$. The quotient $\ker(\chi)/\operatorname{im}(L)$ is the **Picard group** or **divisor class group**. The equivalence classes of divisors can be represented by rooted spanning trees in the simplex graph. This defines a group structure on **rooted spanning trees**. That there is a bijective identification between divisor classes and spanning trees is the subject of:

Theorem: The Picard group is isomorphic to the tree group.

References

42.5. For the history of topology[34, 65] and graph theory [135, 65, 48] and discrete geometry [19]. See [55, 154, 147] for notations in algebraic topology, [54, 14, 20] for graph theory.

- 42.6. Abstract simplicial complexes appeared in 1907 by Dehn and Heegaard [22, 132]. In [2] they appeared under the name **unrestricted skeleton complex**. In [161], J.H.C. Whitehead calls them **symbolic complexes**.
- 42.7. Some of the results generalize to Δ sets or simplicial sets. Some connection calculus however does not. Some connection calculus does not go over yet. The unimodularity theorem does not hold for simplicial sets, at least for the approaches we tried so far.
- 42.8. Homotopy theory as developed by [161] uses elementary expansions and contractions. Homotoptic complexes are said to have the same "nucleus". [161] uses "collapsible" for "homotopic to a point". See also [160]. The notions appearing for simplices described by graph theory, see [64, 63, 24].
- 42.9. Dimension theory has a long history [31]. The inductive definition of graphs appeared first in [76]. We studied the average in [73].
- 42.10. **Random graphs** were first studied in [37]. The average Euler characteristic appears in [73].
- 42.11. The idea of seeing geometric quantities as expectations is central in **integral geometry**. The first time, that curvature was seen as an expectation of indices is Banchoff [7, 8]. Random methods in geometry is part of integral geometry as pioneered by Crofton and Blaschke [18, 134]. We have used in in [96, 80] and [79]. Having curvature given as an expectation allows to deform it. Given a unitary flow U_t on functions for example produces a deformation of the curvature.
- 42.12. Discrete curvature traces back to a combinatorial curvature considered by Heesch [13] in the context of graph coloring and extended in [47]. The formula $K(p) = 1 V_1(p)/6$ and for graphs on the sphere appears also in [141, 142], where it is also pointed out that $\sum_p K(p) = 2$ is Gauss-Bonnet formula. Discrete curvature was used in [58] and unpublished work of Ishida from 1990. Higushi use $K(p) = 1 \sum_{y \in S(p)} (1/2 1/d(y))$, where d(y) are the cardinalities of the neighboring face degrees in the sphere S(p). For two dimensional graphs, where all faces are triangles, this simplifies to $d_j = 3$ so that K = 1 |S|/6, where |S| is the cardinality of the sphere S(p). In [76] second order curvatures were used. The **Levitt curvature** in arbitrary dimension appears in [127]. We rediscovered it in [74] after tackling dimension by dimension separately, not aware of Levitt. We got into the topic while working on [76]. Chern's proof is [25] followed [3, 38]. See [145, 32] for modern proofs. Historical remarks are in [26].
- 42.13. The Erdös Rényi probability space were introduced in [37]. The formulas for the average dimension and Euler characteristic has been found in [73]. The recursive dimension was first used in [76]. We looked at more functionals in [93].
- 42.14. The **discrete Hadwiger Theorem** appears in [69]. The continuous version is [53]. For integral geometry and geometric probability, see [148]. The theory of valuations on distributive lattices has been pioneered by Klee [70] and Rota [146] who proved that there is a unique valuation such that X(x) = 1 for any join-irreducible element. See also [45].

- 42.15. Wu characteristic appeared in [158] and was discussed in [49]. We worked on it in [104] and announced cohomology in [119] and [120]. For the **connection cohomology** belonging to Wu characteristic, see [114].
- 42.16. For discrete Poincaré-Hopf see [78] and an attempt to popularize it in [82] or Mathematica demonstrations [75, 77]. It got pushed a bit more in [79]. For the classical Poincaré-Hopf, see [156]. For the classical case, Poincaré covered the 2-dimensional case in chapter VIII of [139] It got extended by Hopf in arbitrary dimensions [60]. It is pivotal in the proof of Gauss-Bonnet theorems for smooth Riemannian manifolds (i.e. [52, 155, 59, 56, 35, 12]).
- 42.17. Discrete McKean-Singer was covered in [81]. The best proof in the continuum is [32]. The classical result is [129]. In [108], the suggestion appeared to define elliptic discrete complexes using McKean-Singer.
- 42.18. The Zykov sum (join) was introduced in [163] to graph theory. The strong ring was covered in [110, 113].
- 42.19. The Brouwer-Lefschetz theorem is [83]. It generalizes the 1-dimensional case [137]. The classical result is [126]. See also [61].
- 42.20. The classical **Kuenneth formula** is [123]. The graph version [102], uses the Barycentric refinement $(A \times B)_1$ of the Cartesian product $A \times B$.
- 42.21. About the history of discrete notions of manifolds, see [151]. The Evako definition of a sphere as a cell complex for which every unit sphere is a n-1 sphere and such that removing one point makes it contractible was predated by approaches of Vietoris or van Kampen. The later would have accepted homology spheres as unit spheres.
- 42.22. The classical Sard theorem is [149]. The discrete version was remarked in [103].
- 42.23. For the spectral universality, see [100] and [100]. It uses a result of Lidskii-Last [153] which assures if $||\mu \lambda||_1 \le \sum_{i,j=1}^n |A B|_{ij}$ for any two symmetric $n \times n$ matrices A, B with eigenvalues $\alpha_1 \le \alpha_2 \le \cdots \le \alpha_n$.
- 42.24. The discrete exterior derivative goes back to Betti and Poincaré and was already anticipated by Kirchhoff. As pointed out in [83], the discrete Hodge point is [36]. It appeared also in [62]. The **discrete Dirac operator** was stressed in [85].
- 42.25. The unimodularity theorem $|\det(L)| = 1$ was discovered in February 2016, announced in [121] and proven in [105]. An other proof was given in [131].
- 42.26. We have looked at the arithmetic of unit spheres in [112], especially in the context of the diagonal Green function entries. The other Green function entries are covered in [117].
- 42.27. The result $\chi(G) = p(G) n(G)$ was proven in [111, 117]. The functional equation for the spectral zeta function of the connection Laplacian was proven in [115]. Earlier work in the Hodge Zeta case is [118]. The zeta function is called Dyadic because the Barycentric limit is in an ergodic setup a **von Neumann-Kakutani system** [72], which has the Prüfer group as the spectrum. The system is a group translation on the dyadic group of integers and also known as the **adding machine**.

- 42.28. The **Hydrogen relation** $H = L L^{-1}$ for one-dimensional complexes was studied in [107, 109] and [116].
- 42.29. An earlier talk [94] summarizes things also. [86] is an earlier snapshot about the linear algebra part. [97, 82] summarize the calculus.
- 42.30. The matrix tree theorem is [67]. It is based on the **Cauchy-Binet theorem** [23, 15]. A generalization [92] gives the coefficients of the characteristic polynomial. The **Chebotarev-Shamis theorem** is [138, 143]. See also [84], where we initially were not aware of the work of Chebotarev and Shamis.
- 42.31. The **Lax deformation** of exterior derivatives was introduced in [89, 88] and was motivated by **Witten deformation** [162, 32]. Lax systems were introduced first to [125]. Commutation relations of that form have appeared earlier when describing **free tops** L' = [B, L], where $B = I^{-1}L$ is the angular velocity and L the angular velocity in so(n), which are geodesics in SO(n) [4].
- 42.32. The Connes formula [28] is elementary but crucial in the process of generalizing Riemannian geometry to **non-commutative geometry**.
- 42.33. After finding a multiplication completing the Zykov addition to a ring in [110], we realized it is the dual to the Sabidussi ring. In [113], we looked at the ring generated by the Cartesian product. It is a subring and consists of discrete CW complexes. Unlike for simplicial sets, the classical theorems like Gauss-Bonnet and energy theorem go over.
- 42.34. Riemann-Roch for graphs is [5]. See also [6]. We worked on Riemann-Hurwitz in [122]. The usual approach for Riemann-Hurwitz in graph theory is to see them as discrete analogues of algebraic curves or Riemann surfaces see [130].
- 42.35. [159] first looked for a combinatorial definition of spheres. Forman [42] defined spheres through the Reeb as objects admitting 2 critical points. See also [43]. More on discrete Morse theory in [44, 46].
- 42.36. We used data fitting to get first heuristically the Stirling formula then proved it. It is however considered "well known" [21]. It appears also in [157, 128, 57].
- 42.37. The history of polytopes is a "delicate task" [33]. The Euler polyhedron formula (Euler's gem) was discussed in [144]. The early proofs of Schläfli and Staudt had still gaps according to [22]. The difficulty is also explained in [124, 50].
- 42.38. The story of polyhedra is told in [144, 30]. Historically, it was developed in [150], [152], [140]. Coxeter [30] defines a polytop as a convex body with polygonal faces. [51] also works with convex polytopes in \mathbb{R}^n where the dimension is the dimension of the affine span.
- 42.39. The perils of a general definition of a polytop were known since Poincaré (see [1, 144, 27, 124]). Polytop definitions are given in [150, 30, 51, 66]. Topologists started with new definitions [2, 41, 29, 154], and define first a simplicial complex and then polyhedra as topological spaces which admit a **triangularization** by a simplicial complex.

- 42.40. Dehn-Sommerville relations have traditionally been formulated for convex polytopes and then been generalized to situations where unit spheres can be realized as convex polytopes. See [71, 136, 133, 128, 21, 57, 68] or [11].
- 42.41. We started to think about graph coloring during the project [99]. The reports [95] and [101] explored this a bit more. It is related to Fisk theory [40, 39].
- 42.42. Some special graphs appearing when counting was considered in [106]. When writing this, we were not aware that the cell complex introduced already in [16] which goes much further than what we did. Other classes of complexes called **orbital networks** [87, 90, 91] were studied first with Montasser Ghachem.
- 42.43. For the Alexander duality, see [17]. Originally established by Alexander in 1922, it was formulated by Kalai and Stanley in combinatorial topology. We formulated it with cohomology rather than homology and cohomology. As such it is an identity where we have numbers on both sides.

Questions

43. Inverse spectral questions

43.1. We have seen that the spectrum of L does not determine the Betti numbers in general but that for a Barycentric refinement of G, the Betti numbers b_0, b_1 can be read of from the spectrum as the number of eigenvalues 1 and -1.

Question: Does the spectrum of L determine b_k for $k \geq 2$.

Question: Does the spectrum of L determine the Wu characteristic $\omega(G)$?

44. Barycentric Limit

We have seen that the limiting spectral measure can be computed in the case d = 1. It is a smooth measure. In higher dimensions, we see spectral gaps. These gaps have first been seen in the **BeKeNePaPeTe paper** [9].

Question: Prove spectral gaps in limiting spectral measure for $d \geq 2$.

45. Coloring

45.1. The **four color theorem** is equivalent to the statement that all 2-spheres are 4-colorable.

Question: Are all d-spheres (d+2)-colorable?

Question: Are all 2-graphs 5 colorable?

46. Connection Cohomology

46.1. While we know that connection cohomology is not a homotopy invariant, we have not yet proven that it is a topological invariant. We have introduced a notion of homeomorphism in [98]. One can also use the notion whether geometric realizations are homeomorphic to ask:

Question: Is connection cohomology a topological invariant?

46.2. We would like to find more examples of triangulations of non-homeomorphic d-manifolds with different connection cohomology which can not be distinguished by other means:

Question: Can one distinguish homology spheres with Wu cohomology?

46.3. Something we have only started to look at"

Question: Is there a duality for connection cohomology?

46.4. As connection cohomology is not a homotopy invariant, the naive generalization does not work.

47. Random complexes

47.1. The probability spaces E(n, p) of graphs define natural random spaces of simplicial complexes as we can take the Whitney complex of a graph. While we have a formula for the expectation of Euler characteristic, this is not yet available for Wu characteristic numbers ω_k .

Question: What is the expected value of ω_k on E(n, p)?

47.2. We would also like to know the expectations of the Betti numbers:

Question: What is the expectation of $b_k(G)$ on E(n,p)?

48. Zeta function

48.1. While various equivalent expressions exist for the **connection zeta function** in the Barycentric limit of a one-dimensional complex, we don't yet have found a reference about where the roots of ζ are:

Question: The limiting zeta function ζ has roots on the imaginary axes.

Bibliography

References

- [1] A. Aczel. Descartes's secret notebook, a true tale of Mathematics, Mysticism and the Quest to Understand the Universe. Broadway Books, 2005.
- [2] P. Alexandroff. *Combinatorial topology*. Dover books on Mathematics. Dover Publications, Inc, 1960. Three volumes bound as one.
- [3] C. Allendoerfer and A. Weil. The gauss-bonnet theorem for riemannian polyhedra. *Transactions of the American Mathematical Society*, 53:101–129, 1943.
- [4] V.I. Arnold. Mathematical Methods of classical mechanics. Springer Verlag, New York, second edition, 1980.
- [5] M. Baker and S. Norine. Riemann-Roch and Abel-Jacobi theory on a finite graph. Advances in Mathematics, 215:766-788, 2007.
- [6] M. Baker and S. Norine. Harmonic morphisms and hyperelliptic graphs. *International Mathematics Research Notices*, pages 2914–2955, 2009.
- [7] T. Banchoff. Critical points and curvature for embedded polyhedra. J. Differential Geometry, 1:245–256, 1967.
- [8] T. F. Banchoff. Critical points and curvature for embedded polyhedral surfaces. *Amer. Math. Monthly*, 77:475–485, 1970.
- [9] M. Begue, D.J. Kelleher, A. Nelson, H. Panzo, R. Pellico, and A. Teplyaev. Random walks on barycentric subdivisions and the Strichartz hexacarpet. *Exp. Math.*, 21(4):402–417, 2012.
- [10] S. Bera and S.K. Mukherjee. Combinatorial proofs of some determinantal identities. *Linear and Multilinear algebra*, 0(0):1–9, 2017.
- [11] M. Berger. Jacob's Ladder of Differential Geometry. Springer Verlag, Berlin, 2009.
- [12] M. Berger and B. Gostiaux. Differential geometry: manifolds, curves, and surfaces, volume 115 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1988.
- [13] H-G. Bigalke. Heinrich Heesch, Kristallgeometrie, Parkettierungen, Vierfarbenforschung. Birkhäuser, 1988.
- [14] N. Biggs. Algebraic Graph Theory. Cambridge University Press, 1974.
- [15] J.P.M. Binet. Mémoire sur un systeme de formules analytiques, et leur application à des considerations géométriques. J. de l'Ecole Polytechnique IX, Cahier 16:280–302, page 287, 1813.
- [16] A. Björner. A cell complex in number theory. Advances in Appl. Math., (46):71-85, 2011.
- [17] A. Björner and M. Tancer. Combinatorial alexander duality a short and elementary proof. Discrete Comput. Geom., 42:586–593, 2009.
- [18] W. Blaschke. Vorlesungen über Integralgeometrie. Chelsea Publishing Company, New York, 1949.
- [19] A. Bobenko and Y. Suris. Discrete Differential Geometry, Integrable Structure, volume 98 of Graduate Studies in Mathematics. AMS, 2008.
- [20] J. Bondy and U. Murty. Graph theory, volume 244 of Graduate Texts in Mathematics. Springer, New York, 2008.
- [21] F. Brenti and V. Welker. f-vectors of barycentric subdivisions. Math. Z., 259(4):849–865, 2008.
- [22] G. Burde and H. Zieschang. Development of the concept of a complex. In *History of Topology*. Elsevier, 1999.
- [23] A. Cauchy. Memoire sur le nombre de valeurs qu'une fonction peut obtenir. J. de l'Ecole Polytechnique X, pages 51–112, 1815.
- [24] B. Chen, S-T. Yau, and Y-N. Yeh. Graph homotopy and Graham homotopy. *Discrete Math.*, 241(1-3):153–170, 2001. Selected papers in honor of Helge Tverberg.
- [25] S.-S. Chern. A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds. *Annals of Mathematics*, 45, 1944.
- [26] S-S. Chern. Historical remarks on Gauss-Bonnet. In Analysis, et cetera, pages 209–217. Academic Press, Boston, MA, 1990.
- [27] P.R. Comwell. *Polyhedra*. Cambridge University Press, 1997.
- [28] A. Connes. Noncommutative geometry. Academic Press, 1994.
- [29] J.B. Conway. Mathematical Connections: A Capstone Course. American Mathematical Society, 2010.
- [30] H.S.M. Coxeter. Regular Polytopes. Dover Publications, New York, 1973.

- [31] T. Crilly. The emergence of topological dimension theory. In J. James, editor, *History of Topology*, 1999.
- [32] H.L. Cycon, R.G.Froese, W.Kirsch, and B.Simon. Schrödinger Operators—with Application to Quantum Mechanics and Global Geometry. Springer-Verlag, 1987.
- [33] S. Devadoss and J. O'Rourke. Discrete and Computational Geometry. Princeton University Press, 2011.
- [34] J. Dieudonne. A History of Algebraic and Differential Topology, 1900-1960. Birkhäuser, 1989.
- [35] M.P. do Carmo. *Differential forms and applications*. Universitext. Springer-Verlag, Berlin, 1994. Translated from the 1971 Portuguese original.
- [36] B. Eckmann. The Euler characteristic a few highlights in its long history. In *Mathematical Survey Lectures:* 1943-2004, 1999.
- [37] P. Erdö and A. Rényi. On random graphs. I. Publ. Math. Debrecen, 6:290-297, 1959.
- [38] W. Fenchel. On total curvatures for riemannian manifolds (i). J. London Math. Soc, 15:15, 1940.
- [39] S. Fisk. Geometric coloring theory. Advances in Math., 24(3):298–340, 1977.
- [40] S. Fisk. Variations on coloring, surfaces and higher-dimensional manifolds. *Advances in Mathematics*, pages 226–266, 1977.
- [41] A. Fomenko. Visual Geometry and Topology. Springer-Verlag, Berlin, 1994. From the Russian by Marianna V. Tsaplina.
- [42] R. Forman. A discrete Morse theory for cell complexes. In Geometry, topology, and physics, Conf. Proc. Lecture Notes Geom. Topology, IV, pages 112–125. Int. Press, Cambridge, MA, 1995.
- [43] R. Forman. Morse theory for cell complexes. Adv. Math., page 90, 1998.
- [44] R. Forman. Combinatorial differential topology and geometry. New Perspectives in Geometric Combinatorics, 38, 1999.
- [45] R. Forman. The Euler characteristic is the unique locally determined numerical invariant of finite simplicial complexes which assigns the same number to every cone. *Discrete Comput. Geom*, 23:485–488, 2000.
- [46] R. Forman. Bochner's method for cell complexes and combinatorial ricci curvature. *Discrete Comput. Geometry*, pages 323–374, 2003.
- [47] M. Gromov. Hyperbolic groups. In Essays in group theory, volume 8 of Math. Sci. Res. Inst. Publ., pages 75–263. Springer, 1987.
- [48] J. Gross and J. Yellen, editors. *Handbook of graph theory*. Discrete Mathematics and its Applications (Boca Raton). CRC Press, Boca Raton, FL, 2004.
- [49] B. Grünbaum. Polytopes, graphs, and complexes. Bull. Amer. Math. Soc., 76:1131–1201, 1970.
- [50] B. Grünbaum. Are your polyhedra the same as my polyhedra? In *Discrete and computational geometry*, volume 25 of *Algorithms Combin.*, pages 461–488. Springer, Berlin, 2003.
- [51] B. Grünbaum. Convex Polytopes. Springer, 2003.
- [52] V. Guillemin and A. Pollack. Differential topology. Prentice-Hall, Inc., New Jersey, 1974.
- [53] H. Hadwiger. Vorlesungen über Inhalt, Oberfläche und Isoperimetrie. Springer Verlag, Berlin, 1957, 1957.
- [54] F. Harary. Graph Theory. Addison-Wesley Publishing Company, 1969.
- [55] A. Hatcher. Algebraic Topology. Cambridge University Press, 2002.
- [56] M. Henle. A combinatorial Introduction to Topology. Dover Publications, 1994.
- [57] G. Hetyei. The Stirling polynomial of a simplicial complex. Discrete and Computational Geometry, 35:437–455, 2006.
- [58] Y. Higuchi. Combinatorial curvature for planar graphs. J. Graph Theory, 38:220–229, 2001.
- [59] M.W. Hirsch. Differential topology. Graduate texts in mathematics. Springer-Verlag, Berlin, 1976.
- [60] H. Hopf. Uber die Curvatura integra geschlossener Hyperflaechen. Mathematische Annalen, 95:340–367, 1926.
- [61] H. Hopf. A new proof of the Lefschetz formula on invariant points. Proc. Nat. Acad. Sci., 14:149– 153, 1928.

- [62] D. Horak and J. Jost. Spectra of combinatorial Laplace operators on simplicial complexes. Adv. Math., 244:303–336, 2013.
- [63] A. Ivashchenko. Contractible transformations do not change the homology groups of graphs. Discrete Math., 126(1-3):159–170, 1994.
- [64] A.V. Ivashchenko. Graphs of spheres and tori. Discrete Math., 128(1-3):247-255, 1994.
- [65] J. James. History of topology. In *History of Topology*, 1999.
- [66] C. Goodman-Strauss J.H. Conway, H.Burgiel. The Symmetries of Things. A.K. Peterse, Ltd., 2008.
- [67] G. Kirchhoff. Über die Auflösung der Gleichungen auf welche man bei der Untersuchung der linearen Verteilung galvanischer Ströme geführt wird. Ann. Phys. Chem., 72:497–508, 1847.
- [68] D. Klain. Dehn-Sommerville relations for triangulated manifolds. http://faculty.uml.edu/dklain/ds.pdf, 2002.
- [69] D.A. Klain and G-C. Rota. *Introduction to geometric probability*. Lezioni Lincee. Accademia nazionale dei lincei, 1997.
- [70] V. Klee. The Euler characteristic in combinatorial geometry. The American Mathematical Monthly, 70(2):pp. 119–127, 1963.
- [71] V. Klee. A combinatorial analogue of Poincaré's duality theorem. Canadian J. Math., 16:517–531, 1964.
- [72] O. Knill. Renormalization of of random Jacobi operators. Communications in Mathematical Physics, 164:195–215, 1995.
- [73] O. Knill. The dimension and Euler characteristic of random graphs. http://arxiv.org/abs/1112.5749, 2011.
- [74] O. Knill. A graph theoretical Gauss-Bonnet-Chern theorem. http://arxiv.org/abs/1111.5395, 2011.
- [75] O. Knill. Dimension and Euler characteristics of graphs. demonstrations.wolfram.com/DimensionAndEulerCharacteristicsOfGraphs, 2012.
- [76] O. Knill. A discrete Gauss-Bonnet type theorem. Elemente der Mathematik, 67:1–17, 2012.
- [77] O. Knill. Gauss-Bonnet and Poincare-Hopf for graphs. demonstrations.wolfram.com/GaussBonnetAndPoincareHopfForGraphs, 2012.
- [78] O. Knill. A graph theoretical Poincaré-Hopf theorem. http://arxiv.org/abs/1201.1162, 2012.
- [79] O. Knill. An index formula for simple graphs http://arxiv.org/abs/1205.0306, 2012.
- [80] O. Knill. On index expectation and curvature for networks. http://arxiv.org/abs/1202.4514, 2012.
- [81] O. Knill. The McKean-Singer Formula in Graph Theory. http://arxiv.org/abs/1301.1408, 2012.
- [82] O. Knill. The theorems of Green-Stokes, Gauss-Bonnet and Poincare-Hopf in Graph Theory. http://arxiv.org/abs/1201.6049, 2012.
- [83] O. Knill. A Brouwer fixed point theorem for graph endomorphisms. Fixed Point Theory and Appl., 85, 2013.
- [84] O. Knill. Counting rooted forests in a network. http://arxiv.org/abs/1307.3810, 2013.
- [85] O. Knill. The Dirac operator of a graph. http://arxiv.org/abs/1306.2166, 2013.
- [86] O. Knill. The Dirac operator of a graph. http://http://arxiv.org/abs/1306.2166, 2013.
- [87] O. Knill. Dynamically generated networks. http://arxiv.org/abs/1311.4261, 2013.
- [88] O. Knill. An integrable evolution equation in geometry. http://arxiv.org/abs/1306.0060, 2013.
- [89] O. Knill. Isospectral deformations of the Dirac operator. http://arxiv.org/abs/1306.5597, 2013.

- [90] O. Knill. Natural orbital networks. http://arxiv.org/abs/1311.6554, 2013.
- [91] O. Knill. On quadratic orbital networks. http://arxiv.org/abs/1312.0298, 2013.
- [92] O. Knill. A Cauchy-Binet theorem for Pseudo determinants. Linear Algebra and its Applications, 459:522-547, 2014.
- [93] O. Knill. Characteristic length and clustering. http://arxiv.org/abs/1410.3173, 2014.
- [94] O. Knill. Classical mathematical structures within topological graph theory. http://arxiv.org/abs/1402.2029, 2014.
- [95] O. Knill. Coloring graphs using topology. http://arxiv.org/abs/1410.3173, 2014.
- [96] O. Knill. Curvature from graph colorings. http://arxiv.org/abs/1410.1217, 2014.
- [97] O. Knill. If Archimedes would have known functions http://arxiv.org/abs/1403.5821, 2014.
- [98] O. Knill. A notion of graph homeomorphism. http://arxiv.org/abs/1401.2819, 2014.
- [99] O. Knill. On the chromatic number of geometric graphs. Report on HCRP work with Jenny Nitishinskaya, www.math.harvard.edu/~knill/4color, 2014.
- [100] O. Knill. The graph spectrum of barycentric refinements. http://arxiv.org/abs/1508.02027, 2015.
- [101] O. Knill. Graphs with Eulerian unit spheres. http://arxiv.org/abs/1501.03116, 2015.
- [102] O. Knill. The Künneth formula for graphs. http://arxiv.org/abs/1505.07518, 2015.
- [103] O. Knill. A Sard theorem for graph theory. http://arxiv.org/abs/1508.05657, 2015.
- [104] O. Knill. Gauss-Bonnet for multi-linear valuations. http://arxiv.org/abs/1601.04533, 2016.
- [105] O. Knill. On Fredholm determinants in topology. https://arxiv.org/abs/1612.08229, 2016.
- [106] O. Knill. On primes, graphs and cohomology. https://arxiv.org/abs/1608.06877, 2016.
- [107] O. Knill. On a Dehn-Sommerville functional for simplicial complexes. https://arxiv.org/abs/1705.10439, 2017.
- [108] O. Knill. On Atiyah-Singer and Atiyah-Bott for finite abstract simplicial complexes. https://arxiv.org/abs/1708.06070, 2017.
- [109] O. Knill. On Helmholtz free energy for finite abstract simplicial complexes. https://arxiv.org/abs/1703.06549, 2017.
- [110] O. Knill. On the arithmetic of graphs. https://arxiv.org/abs/1706.05767, 2017.
- [111] O. Knill. One can hear the Euler characteristic of a simplicial complex. https://arxiv.org/abs/1711.09527, 2017.
- [112] O. Knill. Sphere geometry and invariants. https://arxiv.org/abs/1702.03606, 2017.
- [113] O. Knill. The strong ring of simplicial complexes. https://arxiv.org/abs/1708.01778, 2017.
- [114] O. Knill. The cohomology for Wu characteristics. https://arxiv.org/abs/1803.1803.067884, 2018.
- [115] O. Knill. An elementary Dyadic Riemann hypothesis. https://arxiv.org/abs/1801.04639, 2018.
- [116] O. Knill. The hydrogen identity for laplacians. https://arxiv.org/abs/1803.01464, 2018.

- [117] O. Knill. Listening to the cohomology of graphs. https://arxiv.org/abs/1802.01238, 2018.
- [118] O. Knill. The zeta function for circular graphs. http://arxiv.org/abs/1312.4239, December 2013.
- [119] O. Knill. A case study in interaction cohomology. http://www.math.harvard.edu/knill/graphgeometry/papers/interactioncohomology.pdf, March, 18, 2016.
- [120] O. Knill. Wu characteristic. http://www.math.harvard.edu/knill/graphgeometry/papers/mathtable.pdf, March, 8, 2016.
- [121] O. Knill. Bowen-Lanford Zeta functions. http://www.math.harvard.edu/knill/graphgeometry/papers/mathtable_fredholm.pdf, October, 18, 2016.
- [122] O. Knill and T. Tucker. A Riemann-Hurwitz theorem in graph theory. Draft Notes, December 2012.
- [123] H. Künneth. Über die Bettischen Zahlen einer Produktmannigfaltigkeit. Math. Ann., 90(1-2):65–85, 1923.
- [124] I. Lakatos. Proofs and Refutations. Cambridge University Press, 1976.
- [125] P.D. Lax. Integrals of nonlinear equations of evolution and solitary waves. Courant Institute of Mathematical Sciences AEC Report, January 1968.
- [126] S. Lefschetz. Intersections and transformations of complexes and manifolds. *Trans. Am. Math. Soc*, 28:1–49, 1926.
- [127] N. Levitt. The Euler characteristic is the unique locally determined numerical homotopy invariant of finite complexes. *Discrete Comput. Geom.*, 7:59–67, 1992.
- [128] A. Luzon and M.A. Moron. Pascal triangle, Stirling numbers and the unique invariance of the euler characteristic. arxiv.1202.0663, 2012.
- [129] H.P. McKean and I.M. Singer. Curvature and the eigenvalues of the Laplacian. *J. Differential Geometry*, 1(1):43–69, 1967.
- [130] A.D. Mednykh and L.A. Mednykh. On 7-hyperellipticity of graphs. To appear, 2013.
- [131] S.K. Mukherjee and S. Bera. A simple elementary proof of The Unimodularity Theorem of oliver knill. *Linear Algebra and Its applications*, pages 124–127, 2018.
- [132] E.S. Munkholm and H.J. Munkholm. Poul heegaard, the Dehn-Heegaard Enzyklopädie article (1907). http://www.imada.sdu.dk/hjm/heegaard3.stor/heegaard3.stor.html, 1998, Accessed, October 4, 2017.
- [133] S. Murai and I. Novik. Face numbers of manifolds with boundary. http://arxiv.org/abs/1509.05115, 2015.
- [134] L. Nicolaescu. Lectures on the Geometry of Manifolds. World Scientific, second edition, 2009.
- [135] R.J. Wilson N.L. Biggs, E.K. Lloyd. Graph Theory, 1736-1936. Clarendon Press, Oxford, second edition, 1998.
- [136] I. Novik and E. Swartz. Applications of Klee's Dehn-Sommerville relations. *Discrete Comput. Geom.*, 42(2):261–276, 2009.
- [137] R. Nowakowski and I. Rival. Fixed-edge theorem for graphs with loops. J. Graph Theory, 3:339–350, 1979.
- [138] P.Chebotarev and E. Shamis. Matrix forest theorems. arXiv:0602575, 2006.
- [139] H. Poincaré. Sur les courbes definies par les equation differentielle III. *Journal de Mathematique pures et appliquées*, pages 167–244, 1885.
- [140] I. Polo-Blanco. Alicia Boole Stott, a geometer in higher dimension. *Historia Mathematica*, 35(2):123 139, 2008.
- [141] E. Presnov and V. Isaeva. Positional information as symmetry of morphogenetic fields. *Forma*, 5:59–61, 1990.
- [142] E. Presnov and V. Isaeva. Local and global aspects of biological morphogenesis. *Speculations in Science and Technology*, 14:68, 1991.
- [143] E.V. Shamis P.Yu, Chebotarev. A matrix forest theorem and the measurement of relations in small social groups. *Avtomat. i Telemekh.*, 9:125–137, 1997.

- [144] D.S. Richeson. *Euler's Gem.* Princeton University Press, Princeton, NJ, 2008. The polyhedron formula and the birth of topology.
- [145] S. Rosenberg. The Laplacian on a Riemannian Manifold, volume 31 of London Mathematical Society, Student Texts. Cambridge University Press, 1997.
- [146] G-C. Rota. On the combinatorics of the Euler characteristic. In *Studies in Pure Mathematics* (Presented to Richard Rado), pages 221–233. Academic Press, London, 1971.
- [147] J.J. Rotman. An introduction to Algebraic Topology. Graduate Texts in Mathematics. Springer.
- [148] L.A. Santalo. Introduction to integral geometry. Hermann and Editeurs, Paris, 1953.
- [149] A. Sard. The measure of the critical values of differentiable maps. *Bull. Amer. Math. Soc.*, 48:883–890, 1942.
- [150] L. Schläffi. Theorie der Vielfachen Kontinuität. Cornell University Library Digital Collections, 1901.
- [151] E. Scholz. The concept of manifold, 1850-1950. In History of Topology. Elsevier, 1999.
- [152] P.H. Schoute. Analytical treatment of the polytopes regularly derived from the regular polytopes. Johannes Mueller, 1911.
- [153] B. Simon. Trace Ideals and their Applications. AMS, 2. edition, 2010.
- [154] E.H. Spanier. Algebraic Topology. Springer Verlag, 1966.
- [155] M. Spivak. A comprehensive Introduction to Differential Geometry I-V. Publish or Perish, Inc, Berkeley, third edition, 1999.
- [156] M. Spivak. A comprehensive Introduction to Differential Geometry V. Publish or Perish, Inc, Berkeley, third edition, 1999.
- [157] R. Stanley. Enumerative Combinatorics, Vol. I. Wadworth and Brooks/Cole, 1986.
- [158] Wu W-T. Topological invariants of new type of finite polyhedrons. *Acta Math. Sinica*, 3:261–290, 1953.
- [159] H. Weyl. Riemanns geometrische Ideen, ihre Auswirkung und ihre Verknüpfung mit der Gruppentheorie. Springer Verlag, 1925, republished 1988.
- [160] J. H. C. Whitehead. Combinatorial homotopy. I. Bull. Amer. Math. Soc., 55:213-245, 1949.
- [161] J.H.C. Whitehead. Simplicial spaces, nuclei and m-groups. Proc. London Math. Soc., 45(1):243–327, 1939.
- [162] E. Witten. Supersymmetry and Morse theory. J. of Diff. Geometry, 17:661–692, 1982.
- [163] A.A. Zykov. On some properties of linear complexes. (russian). Mat. Sbornik N.S., 24(66):163–188, 1949.

Department of Mathematics, Harvard University, Cambridge, MA, 02138