

ORBIFOLD KÄHLER GROUPS RELATED TO ARITHMETIC COMPLEX HYPERBOLIC LATTICES

PHILIPPE EYSSIDIEUX

ABSTRACT. We study fundamental groups of toroidal compactifications of non compact ball quotients and show that the Shafarevich conjecture on holomorphic convexity for these complex projective manifolds is satisfied in dimension 2 provided the corresponding lattice is arithmetic and small enough. The method is to show that the Albanese mapping on an étale covering space generates jets on the interior, if the lattice is small enough. We also explore some specific examples of Picard-Eisenstein type.

1. INTRODUCTION

1.1. Given X a compact Kähler manifold, the question raised by Shafarevich whether the universal covering space of X is holomorphically convex, also known as the Shafarevich Conjecture on holomorphic convexity $SC(X)$, and the study of the Serre problem of characterizing the finitely presented groups arising as fundamental groups of complex algebraic manifolds lead to consider several properties the fundamental group of X may or may not satisfy:

- (1) $Inf(X)$: $\pi_1(X)$ is infinite.
- (2) $Inf_{et}(X)$: Assuming $Inf(X)$, the profinite completion $\hat{\pi}_1(X)$ is infinite.
- (3) $RF(X)$: $\pi_1(X)$ is residually finite.
- (4) $Q(X)$: Assuming $Inf(X)$, $\pi_1(X)$ has a finite rank representation in a complex vector space whose image is infinite.
- (5) $SSC(X)$: For every $f : Z \rightarrow X$ where Z is a compact connected complex analytic space and f is holomorphic $\#\text{Im}(\pi_1(Z) \rightarrow \pi_1(X)) = +\infty \Leftrightarrow \exists N \in \mathbb{N}^* \exists \rho : \pi_1(X) \rightarrow GL_N(\mathbb{C}) \#\rho(\text{Im}(\pi_1(Z) \rightarrow \pi_1(X))) = +\infty$.
- (6) $L(X)$: $\pi_1(X)$ is a linear group.

One has the following implications, the first one being a classical result of Malčev, the last one was proved in increasing generality in [EKPR12, CCE15, Eys17]:

$$L(X) \Rightarrow RF(X), SSC(X), SSC(X) \Rightarrow Q(X) \Rightarrow Inf_{et}(X), SSC(X) \Rightarrow SC(X).$$

The counterexamples to $RF(_)$ -hence $L(_)$ - [Tol93, CK92] do not give rise to counterexamples of $SSC(_)$, actually all the other statements hold trivially true for the complex projective manifolds considered there.

All these properties make sense if X is replaced by a compact Kähler smooth Deligne-Mumford stack, see e.g. [Eys17], a complex algebraic manifold (say quasi projective) or a smooth separated Deligne-Mumford stack with quasi projective or quasi Kähler moduli space. Using orbifold compactifications of a given quasi-Kähler manifold U , one can produce compact Kähler orbifolds with **potentially** interesting fundamental groups if $\pi_1(U)$ has a sufficiently rich normal subgroup lattice. These orbifold Kähler groups can sometimes be proven to be fundamental groups of related compact Kähler manifolds. This happens if the inertia morphisms are injective after

Date: April, 30 2018.

This research was partially supported by the ANR project Hodgefun ANR-16-CE40-0011-01.

passing to the profinite completion. Checking this property requires at least *some* understanding of the structure of the orbifold fundamental group.

A first non trivial class of U , complements of line arrangements in projective space, was analysed in [Eys17] where $SSC(_)$ was settled affirmatively in the equal weight case and it seems much more difficult to settle $L(_)$. The resulting orbifolds are abelian quotients of the Hirzebruch algebraic surfaces ramified over the arrangement [Hir83, BHH87]. A general theory of the fundamental group of these surfaces in the unequal weight case seems to be out of reach except in specific cases: in the case of $CEVA(2)$, which was investigated in depth for the construction of complex hyperbolic lattices, the theory is fascinating [DeMo86, DeMo93, Mos86].

1.2. The present article investigates a second non-trivial class, where a lot of the most beautiful examples come precisely from the aforementioned work of Mostow and Deligne-Mostow: finite covolume non compact quotients of a complex hyperbolic space. Then $\pi_1(U)$ is a non-uniform complex hyperbolic lattice, and the manifolds we investigate are the toroidal compactifications of non compact ball quotients [AMRT75, Mok11]. These objects have attracted a lot of attention in recent years from several perspectives, construction of interesting lattices [DPP16, Sto15, DiCSto18], their position in the classification of algebraic varieties [DD15, BT15], Kobayashi hyperbolicity [Cad16, Cad17].

In contrast to the case of rank ≥ 2 where the fundamental group of a toroidal compactification of an irreducible hermitian locally symmetric space is finite, a toroidal compactification of a ball quotient can have a large fundamental group. For instance, [HS96] constructs a Riemannian metric of non positive curvature on the toroidal compactification of a small enough lattice. These toroidal compactifications are thus $K(\pi, 1)$, the universal covering space being diffeomorphic to a real affine space and the fundamental group has exponential growth. But it does not seem possible to use the methods in [HS96] to prove that the universal covering space is Stein (which what SC predicts) or to construct linear representations of the fundamental group.

1.3. Let us describe the content of this article.

Given $\Gamma < PU(n, 1)$ a non uniform lattice, $n \geq 2$, decorating the construction of [Hol87, p. 29-30] with its natural stack structure, or interpreting [Hol98, Ch. 4] and [Ulu07] in a more flexible language, we construct an orbifold compactifications $[\Gamma \backslash \mathbb{H}_{\mathbb{C}}^n] \subset \mathcal{X}_{\Gamma}^{tor}$ with no codimension 1 ramification at infinity and a singular DM-stack compactification $[\Gamma \backslash \mathbb{H}_{\mathbb{C}}^n] \subset \mathcal{X}_{\Gamma}^{BBS}$. When Γ is neat this is the usual construction from [AMRT75].

Let $C \subset \partial \mathbb{H}_{\mathbb{C}}^n$ be the set of the preimages of the cusps in $[\Gamma \backslash \mathbb{H}_{\mathbb{C}}^n]^{BBS}$. The finite set $\Gamma \backslash C$ is the set of cusps of Γ .

Theorem 1.1. *For each $c \in C$ of Γ denote by H_c (resp. Z_c) the intersection of Γ with the unipotent radical of the parabolic subgroup attached to c (resp. its center). Then:*

- (1) $\pi_1(\mathcal{X}_{\Gamma}^{BBS}) = \Gamma^{BBS} = \Gamma / \langle H_c, c \in C \rangle$.
- (2) $\pi_1(\mathcal{X}_{\Gamma}^{tor}) = \Gamma^{tor} = \Gamma / \langle Z_c, c \in C \rangle$.

Here $\langle _ \rangle$ stands for the subgroup generated by $_$ which is normal in the two above cases. We have not been able so far to find any piece of information on these rather natural quotients of Γ in the literature. In spite of the fact that the credit for this result should be given to [LKMSS15] and [KS15], we nevertheless display it in the introduction, in order to translate the problem we study here in terms familiar to complex hyperbolic geometers.

Then, we focus on constructing virtually abelian linear representations of the fundamental group Γ^{tor} , hence on studying the virtual first Betti number. We will give evidence for the following conjecture:

Conjecture 1. *If \bar{X}_Γ is the toroidal compactification of a non uniform arithmetic lattice $\Gamma < PU(n, 1)$, which is torsion free and torsion free at infinity¹, $SSC(\bar{X}_\Gamma)$ holds and the representations can be taken to be virtually abelian. Furthermore, there is a finite étale Galois covering $\bar{X}_{\Gamma'} \rightarrow \bar{X}_\Gamma$ with Galois group G such that the Stein factorization of the quotient Albanese morphism $\bar{X}_\Gamma \rightarrow G \backslash Alb(\bar{X}_{\Gamma'})$ is the Shafarevich morphism of \bar{X}_Γ .*

Theorem 1.2. *If $\Gamma'' < PU(2, 1)$ is arithmetic, there is a finite index subgroup $\Gamma' <_{fi} \Gamma''$ such that if $\Gamma <_{fi} \Gamma'$, \bar{X}_Γ has Albanese dimension 2 and its image contains no translate of an elliptic curve at its generic point. It satisfies SC, Q .*

With the notations of Theorem 1.2, $SSC(\bar{X}_\Gamma)$ would follow from the following statement:

(†) $\exists \Gamma_* <_{fi} \Gamma'' \quad \forall c \in C \quad H_c \cap \Gamma_* / Z_c \cap \Gamma_* \rightarrow H_1(\Gamma_*^{tor}, \mathbb{Q})$ is injective,

and the universal covering space of \bar{X}_Γ would be a Stein manifold. It is enough to look at all c in a finite representative set for $\Gamma'' \backslash C$. We write $<_{fi}$ to when we want to emphasize a subgroup inclusion has finite index.

Theorem 1.3. *If $n \geq 3$ and $\Gamma'' < PU(n, 1)$ is arithmetic, there is a finite index subgroup $\Gamma' < \Gamma''$ such that if $\Gamma <_{fi} \Gamma'$, \bar{X}_Γ has Albanese dimension n and its image contains no translate of an abelian variety at its generic point. It satisfies Q .*

As a corollary of our approach, we get a new proof of the following known facts:

Theorem 1.4. *Under the assumptions of Theorems 1.2 1.3, \bar{X}_Γ has ample cotangent bundle modulo its boundary, is Kobayashi hyperbolic modulo its boundary, and the rational points over a number field over which it is defined are finite modulo its boundary.*

One would like effective versions of these results and quantify them in terms of the ramification indices at infinity in the spirit of [BT15, Cad16] which but our method is inherently non-effective. The examples we have studied so far, most notably the 2-dimensional Picard-Eisenstein case where the hard work has been done by Feustel and Holzapfel [Hol87], suggest much better statements. The article finishes with a detailed discussion of the Picard-Eisenstein commensurability class from the present perspective. A very strong version of (†) holds in this class.

In order to streamline the discussion, we introduce in Definition 4.12 an equivalence relation, *refined commensurability*, on the set of all lattices in a commensurability class, namely that their (orbifold) toroidal compactification are related by an étale correspondance. There is a natural partial ordering on its the quotient set, which is induced by reverse inclusion of lattices. The questions we study here depend only of the refined commensurability class and as in [Eys17] are more delicate for small classes.

1.4. When writing this article we were not sure whether the statement about rational points was new, it is not a corollary of [Ull04]. In the final stage of the redaction, Y. Brunebarbe informed us that, if $n = 2$, it follows from [DimRam15, Theorem 0.3] a paper we were not aware of. The Kobayashi hyperbolicity statement is not new since it follows from [Nad89] and an effective, hence better, version was proved in [BT15] - and even more precise results follow from [DimRam15] if $n = 2$. On the other hand [DimRam15] does not imply SC .

¹Actually, torsion can and will be dealt with orbifold methods.

This article follows the same basic idea as [DimRam15] where N. Fakhruddin is credited for it. [DimRam15, Proposition 3.8] gives an explicit Γ in each commensurability class such that $q(X_\Gamma) > 2$. Here, with Lemma 4.8, we go a little further in the study of the differential geometry of the Albanese mapping, using as the only automorphic input the classical fact [Wal84] that one may achieve $q(X_\Gamma) > 0$. We did not find a reference for the analogous property in the non arithmetic case and will refrain from making any conjecture in that case. Our method however seems to be hopelessly non-effective and relies on the commensurator property of arithmetic lattices.

1.5. These results say that small covolume arithmetic lattices and, unsurprisingly, non-arithmetic lattices are the most interesting ones from the present perspective and we hope to come back to their study in future work.

The author would like to thank Y. Brunenbarbe, B. Cadorel, B. Claudon, M. Deraux, B. Klingler, F. Sala and G. Wüstholtz for useful conversations related to this article and the Freiburg Institute for Advanced Studies for hospitality during its preparation.

2. ORBIFOLD PARTIAL COMPACTIFICATIONS

2.1. As advocated by [Noo05, Ler10], we define an orbifold to be a smooth Deligne-Mumford stack with trivial generic isotropy groups relative to the category of complex analytic spaces² with the classical topology: we assume that the moduli space is Hausdorff and that the inertia groups are finite. There is an analytification 2-functor from DM-stacks over \mathbb{C} to complex analytic DM-stacks and an underlying topological stack functor from complex analytic DM-stacks to topological DM-stacks [Noo05].

2.2. An orbifold \mathcal{X} is said to be *developable* if its universal covering stack [Noo05]³ is an ordinary manifold, which is equivalent to the injectivity of every local inertia morphism $I_x = \pi_1^{loc}(\mathcal{X}, x) \rightarrow \pi_1(\mathcal{X}, x)$, x being an orbifold point of \mathcal{X} . If x_0 is a base point of \mathcal{X} , we have a conjugacy class of morphisms $I_x \rightarrow \pi_1^{et}(\mathcal{X}, x_0)$, also called the local inertia morphisms. We will abuse notation and drop the base point dependency of π_1 when harmless. The orbifold \mathcal{X} is said to be *uniformizable* whenever the profinite completion $I_x \rightarrow \pi_1^{et}(\mathcal{X})$ of every local inertia morphism is injective⁴. When $\pi_1(\mathcal{X})$ is residually finite, uniformizability and developability are equivalent properties. The condition ‘residually finite’ cannot be dropped, a counterexample is given in [Eys17]. The fundamental group of a compact Kähler uniformizable orbifold occurs as the fundamental group of a compact Kähler manifold [Eys13].

2.3. **The fundamental group of a weighted DCN.** Let us recall the simplest examples of orbifold compactification of quasi-Kähler manifolds described in [Eys17].

2.3.1. *Root stacks.* Let M be a (Hausdorff second countable) complex analytic space and D be an effective Cartier divisor. Let $r \in \mathbb{N}^*$. Then, one can construct $P \rightarrow M$ the principal \mathbb{C}^* -bundle attached to $\mathcal{O}_M(-D)$ and the tautological section $s_D \in H^0(M, \mathcal{O}_M(D))$ can be lifted to a holomorphic function $f_D : P \rightarrow \mathbb{C}$ satisfying $f_D(p.\lambda) = \lambda f_D(p)$ for every $\lambda \in \mathbb{C}^*$, $p \in P$. Define a complex analytic

²Except in specific cases, one should not work relatively to the category of complex manifolds, since the Yoneda functor should distinguish a complex space and the normalisation of underlying reduced complex space.

³The covering theory of [Noo05] has nothing to do with the complex structure.

⁴The literature also uses *good* orbifold for developable and *very good* orbifold for uniformizable.

space $Y := Y_D \subset P \times \mathbb{C}_z$ by the equation $z^r = f_D(p)$. One can define a \mathbb{C}^* -action on Y by $(p, z) \cdot \lambda = (p\lambda^r, \lambda \cdot z)$. The complex analytic stack

$$M(\sqrt[r]{D}) := [Y_D/\mathbb{C}^*]$$

(see e.g. [EySa16, BN06]) is then a Deligne-Mumford separated complex analytic stack with trivial generic isotropy groups whose moduli space is M and an orbifold if M and D are smooth. The non-trivial isotropy groups live over the points of D and are isomorphic to μ_r , the group of r -roots of unity. In the smooth case, the corresponding differentiable stack can be expressed as the quotient by the natural infinitesimally free $U(1)$ action on the restriction UY of Y to the unit subbundle for (any hermitian metric) of P which is a manifold indeed. It is straightforward to see that this is an analytic version of Vistoli's root stack construction:

Lemma 2.1. *If (M, D) is the analytification of $(\mathcal{M}, \mathcal{D})$ a pair consisting of a \mathbb{C} - (separated) scheme and a Cartier divisor, then $M(\sqrt[r]{D})$ is the analytification of $\mathcal{M}_{O(\mathcal{D}), s_D, r}$ in the notation of [Cad07, Section 2].*

A Cartier divisor D on a scheme M should be thought of as a pair of an invertible sheaf \mathcal{L} and a section $s_D : \mathcal{O} \rightarrow \mathcal{L}$ such that $[s_D] = D$. In other words a map of algebraic stacks $\mu : M \rightarrow [\mathbb{C}/\mathbb{C}^*]$. We have the natural r -th power map $\cdot^r : [\mathbb{C}/\mathbb{C}^*] \rightarrow [\mathbb{C}/\mathbb{C}^*]$ and an equivalence $M(\sqrt[r]{D}) \rightarrow M \times_{[\mathbb{C}/\mathbb{C}^*]_{\mu, \cdot^r}} [\mathbb{C}/\mathbb{C}^*]$. Using this we may even promote M to be a stack and allow for $s_D = 0$. In the latter case we get a $\mathbb{Z}/r\mathbb{Z}$ -gerbe on M whose class is the reduction mod r of $c_1(\mathcal{L})$. The main property is treated in the scheme-theoretic setting by [Cad07]:

Lemma 2.2. *If S is a complex analytic stack*

$$\text{Hom}(S, M(\sqrt[r]{D})) = \{f : S \rightarrow M \mid \mathbb{C} - \text{analytic}, \exists D_S \text{ Cartier on } S \text{ s.t. } D_S = r \cdot f^* D\}.$$

2.3.2. Let \bar{X} be a compact Kähler manifold and $x_0 \in X$ a base point, $n = \dim_{\mathbb{C}}(X)$, and let $D := D_1 + \dots + D_l$ be a simple normal crossing divisor whose smooth irreducible components are denoted by D_i . We assume for simplicity $x_0 \notin D$. For each choice of weights $d := (d_1, \dots, d_l)$, $d_i \in \mathbb{N}^*$, one may construct as [Cad07, Definition 2.2.4] does in the setting of scheme theory the compact Kähler orbifold (Compact Kähler DM stack with trivial generic isotropy)

$$\mathcal{X}(\bar{X}, D, d) := \bar{X}(\sqrt[r]{D_1}) \times_{\bar{X}} \dots \times_{\bar{X}} \bar{X}(\sqrt[r]{D_l}).$$

In other words, $\mathcal{X}(\bar{X}, D, d) = [Y_{D_1} \times_{\bar{X}} \dots \times_{\bar{X}} Y_{D_l}/\mathbb{C}^{*l}]$. Denote by X the quasi-Kähler manifold $X := \bar{X} \setminus D$. View $\mathcal{X}(\bar{X}, D, d)$ as an orbifold compactification of X and denote by $j_d : X \hookrightarrow \mathcal{X}(\bar{X}, D, d)$ the natural open immersion.

By Zariski-Van Kampen, the fundamental group $\pi_1(\bar{X}, x_0)$ is the quotient of $\pi_1(X, x_0)$ by the normal subgroup generated by the γ_i , where γ_i is a meridian loop for D_i . Zariski-Van Kampen generalizes to orbifolds, see e.g. [Noo04, Zoo02], and $\pi_1(\mathcal{X}(\bar{X}, D, d), x_0)$ is the quotient of $\pi_1(X, x_0)$ by the normal subgroup generated by the $\gamma_i^{d_i}$.

Remark 2.3. *If (X, Δ) is an orbifold in the sense of Campana, the Campana orbifold fundamental group is the fundamental group of the root stack on $X \setminus D^1$ where D^1 is the log-singular set of (X, Δ) . It may happen that the fundamental group of the trace of the root stack on $X \setminus D^1$ of a small ball centered on D^1 is finite and in this case we get a DM-stack compactification which may be singular. In dimension 2, the list of local configurations giving rise to an orbifold compactification with a smooth moduli space is given in [Ulu07].*

Remark 2.4. *The orbifolds constructed here are specified up to equivalence by their moduli space and ramification indices in codimension 1 by [GeSa17]. Hence, in*

dimension 2, these orbifolds carry the same information as Holzappel's orbital surfaces or Uludag's orbifaces. One could also use orbifolds in the sense of Thurston in higher dimension. However, it is convenient to have at our disposal maps of stacks (defined as functors of fibered categories), moduli spaces, substacks, (2-)fiber products hence fibers, basic homotopy theory [Noo05] and many differential geometric constructions [EySa16].

3. TOROIDAL COMPACTIFICATIONS OF COMPLEX HYPERBOLIC ORBISURFACES

3.1. Orbifold Toroidal compactifications.

3.1.1. The following constructions are well known to the experts [BT15] but we recall them in order to fix the notations. Let $\mathbb{H}_{\mathbb{C}}^2 := PU(1, 2)/P(U(1) \times U(2))$ be the complex hyperbolic plane. Let $\Gamma < PU(1, 2)$ be non uniform lattice. Then $X_{\Gamma} := \Gamma \backslash \mathbb{H}_{\mathbb{C}}^2$ is an algebraic quasi-projective variety by Baily-Borel's theorem in the arithmetic case, and [Mok11] for the non arithmetic case. It has a complex projective compactification $X_{\Gamma} \subset X_{\Gamma}^{BBS}$ obtained by adding a finite number of points we shall refer to as cusps.

Each cusp c has some neighborhood V_c such that the preimage of $U_c := V_c \setminus \{c\}$ is a disjoint union of horoballs $W_{\tilde{c}}$ of $\mathbb{H}_{\mathbb{C}}^2$ labelled by $\partial\mathbb{H}_{\mathbb{C}}^2 \ni \tilde{c} := \partial\mathbb{H}_{\mathbb{C}}^2 \cap \overline{W_{\tilde{c}}}$ where the closure and boundary are taken in the euclidean topology of the projective plane \mathbb{P} dual to $\mathbb{H}_{\mathbb{C}}^2$. The set C of all such \tilde{c} is acted upon with finitely many orbits by Γ . For such a \tilde{c} , let $\Gamma_{\tilde{c}}$ be the stabilizer of \tilde{c} in Γ .

Since the stabilizer $S_{\tilde{c}}$ of \tilde{c} in $PU(1, 2)$ has a 3 dimensional Heisenberg group of unipotent 3×3 upper triangular matrices with real coefficients as its unipotent radical $H_{\tilde{c}}$ and \mathbb{C}^* has its Levi component, see [Par09], we have an exact sequence:

$$1 \rightarrow H_{\tilde{c}} \cap \Gamma_{\tilde{c}} \rightarrow \Gamma_{\tilde{c}} \rightarrow \mu_{k_c} \rightarrow 1$$

where $\mu_k < \mathbb{C}^*$ is group of k_c -roots of unity. Whenever this causes no confusion, we use a shorthand notation $k = k_c$. Then, $H_{\tilde{c}} \cap \Gamma_{\tilde{c}} < H_{\tilde{c}}$ is a lattice in $H_{\tilde{c}}$. One has $\gamma W_{\tilde{c}} \cap W_{\tilde{c}} \neq \emptyset \Leftrightarrow \gamma \in \Gamma_{\tilde{c}}$. The center $Z_{\tilde{c}}$ of $H_{\tilde{c}} \cap \Gamma_{\tilde{c}}$ is a cyclic infinite subgroup and we have $Z_{\tilde{c}} = Z(H_{\tilde{c}}) \cap \Gamma_{\tilde{c}}$.

The group $\Lambda_{\tilde{c}} := Z_{\tilde{c}} \backslash H_{\tilde{c}} \cap \Gamma_{\tilde{c}}$ is a lattice in the real 2-dimensional additive group $A_{\tilde{c}}^{\tau} := Z(H_{\tilde{c}}) \backslash H_{\tilde{c}}$ which has a natural structure of an affine complex line $A_{\tilde{c}}$. Hence $A_{\tilde{c}}^{\tau}$ is naturally a one dimensionnal complex additive group acting as the translation group on $A_{\tilde{c}}$. This complex structure comes from the fact that $Z(H_{\tilde{c}})$ stabilizes a unique complex geodesic having \tilde{c} as a boundary point. All such complex geodesics are the trace of a complex projective line in \mathbb{P} through \tilde{c} and form a single orbit of $H_{\tilde{c}}$. In particular we have a bijection $A_{\tilde{c}}^{\tau} \rightarrow \mathbb{P}(T_{\tilde{c}}\mathbb{P}) \setminus \{l_{\tilde{c}}\}$ where $l_{\tilde{c}}$ is the complex line tangent to $\partial\mathbb{H}_{\mathbb{C}}^2$ at \tilde{c} . The linear projection from \tilde{c} give an equivariant holomorphic map $W_{\tilde{c}} \rightarrow A_{\tilde{c}}$ where $S_{\tilde{c}}$ acts through its quotient group $Z_{\tilde{c}} \backslash S_{\tilde{c}} \simeq A_{\tilde{c}}^{\tau} \rtimes \mathbb{C}^*$. The latter group acts as the complex affine group of $A_{\tilde{c}}$.

The natural map $\psi : \tilde{V}_{\tilde{c}}^1 := Z_{\tilde{c}} \backslash W_{\tilde{c}} \rightarrow A_{\tilde{c}}$ is a holomorphic fiber bundle whose fiber at $\lambda \in A_{\tilde{c}}$ is $Z_{\tilde{c}} \backslash \lambda \cap W_{\tilde{c}}$ a pointed disk. There is an effective action of $Z_{\tilde{c}} \backslash S_{\tilde{c}}$ on $\tilde{V}_{\tilde{c}}^1$ hence an action of $A_{\tilde{c}}^{\tau}$ such that ψ is equivariant. Now, consider the genus one Riemann Surface $E_{\tilde{c}} := \Lambda_{\tilde{c}} \backslash A_{\tilde{c}}$. The commutator gives a natural symplectic form on $\Lambda_{\tilde{c}}$ with values in $Z_{\tilde{c}}$ hence, with an adequate choice of a generator of $Z_{\tilde{c}}$, a polarization $\Theta_{\tilde{c}}$ of the weight -1 Hodge structure on $\Lambda_{\tilde{c}}$ induced by the complex structure on $\Lambda_{\tilde{c}} \otimes_{\mathbb{Z}} \mathbb{R}$. There is also a map $\psi' : V_{\tilde{c}}^1 := \Lambda_{\tilde{c}} \backslash \tilde{V}_{\tilde{c}}^1 \rightarrow E_{\tilde{c}}$ which is a holomorphic fiber bundle in pointed disks. A coordinate calculation enables to see that it is biholomorphic to the complement of the zero section of a unit disk bundle of a hermitian line bundle $(L_{\tilde{c}}, h_{\tilde{c}})$ on $E_{\tilde{c}}$ with constant curvature $c_1(L_{\tilde{c}}) = -\Theta_{\tilde{c}}$. The degree of $\Theta_{\tilde{c}}$ is the index in $Z_{\tilde{c}}$ of the image of the symplectic form on $\Lambda_{\tilde{c}}$.

Hence there is a partial compactification $V_{\tilde{c}}^1 \simeq U_{\tilde{c}}^{1,tor} \setminus D_{\tilde{c}}^{1,tor}$ where $U_{\tilde{c}}^{1,tor}$ is the full unit bundle of $(L_{\tilde{c}}, h_{\tilde{c}})$ and $D_{\tilde{c}}^{1,tor}$ is the zero section, a smooth divisor isomorphic to $E_{\tilde{c}}$.

The quotient action of the quotient a group μ_k on the hermitian line bundle $(L_{\tilde{c}}, h_{\tilde{c}})$ gives an action on $U_{\tilde{c}}^{1,tor}$ in such a way that the natural retraction $\pi_{\tilde{c}}^1 : (U_{\tilde{c}}^{1,tor}, D_{\tilde{c}}^{1,tor}) \rightarrow E_{\tilde{c}}$ is equivariant. In particular the group μ_k acts on $E_{\tilde{c}}$ in such a way that the action on $H^0(E_{\tilde{c}}, \Omega^1)$ is given by complex multiplication. Since it is an action by automorphisms and since the Lefschetz number of an automorphism is an integer it follows that $k \in \{1, 2, 3, 4, 6\}$ ⁵.

Dividing out $\pi_{\tilde{c}}^1$ by μ_k we get a retraction $\pi_{\tilde{c}}^1 : U_{\tilde{c}}^{tor} \rightarrow D_{\tilde{c}}^{tor}$ in such a way that $U_c \simeq U_{\tilde{c}}^{tor} \setminus D_{\tilde{c}}^{tor}$ and a map $U_{\tilde{c}}^{tor} \rightarrow U_c$ contracting $D_{\tilde{c}}^{tor}$ to \tilde{c} .

Since the construction is independent of \tilde{c} we may drop this dependency in our notation and redefine $(U_c^{1,tor}, D_c^{1,tor}, E_c) := (U_{\tilde{c}}^{1,tor}, D_{\tilde{c}}^{1,tor}, E_{\tilde{c}})$ for some \tilde{c} .

Gluing the U_c^{tor} with X_{Γ} along U_c gives a normal surface X_{Γ}^{tor} with a family $(D_c)_{c \in C}$ of disjoint curves with an isomorphism $X_{\Gamma} \rightarrow X_{\Gamma}^{tor} \setminus \bigcup_{c \in C} D_c$ and a map $X_{\Gamma}^{tor} \rightarrow X_{\Gamma}^{BBS}$ contracting D_c to c . Then, X_{Γ}^{tor} is a normal surface with quotient singularities and is projective algebraic too [AMRT75, Mok11].

It is actually better to glue the orbifold $\mathcal{X}_{\Gamma} := [\Gamma \backslash \mathbb{H}_{\mathbb{C}}^2]$ (stack theoretic quotient) with $[\mu_k \backslash U_{\tilde{c}}^{1,tor}]$ along $[\Gamma_{\tilde{c}} \backslash W_{\tilde{c}}]$ to get a compact orbifold $\mathcal{X}_{\Gamma}^{tor}$ containing \mathcal{X}_{Γ} as the complement of a smooth divisor \mathcal{D}_{Γ} consisting of a finite number of disjoint smooth substacks $(\mathcal{D}_c)_{c \in C}$ whose generic points have no inertia. One also has a stack theoretic compactification $\mathcal{X}_{\Gamma}^{BBS}$ obtained by adding the disjoint union of the $(B\mu_{k_c})_{c \in C}$ and a contraction map $\mathcal{X}_{\Gamma}^{tor} \rightarrow \mathcal{X}_{\Gamma}^{BBS}$.

When the lattice is neat the stack we constructed is equivalent to the usual smooth toroidal compactification of [AMRT75, Mok11]. On the other other hand, $\mathcal{X}_{\Gamma}^{tor}$ is NOT the quotient stack of the toroidal compactification attached to a neat normal sublattice: the generic point of the boundary has trivial isotropy.

3.1.2. Some comments have to be made regarding the gluing construction we are performing. First of all, gluing DM topological stacks along open substacks is always possible thanks to [Noo05, Cor. 16.11, p. 57] - we are using the class of local homeomorphisms as **LF**. In order to have a better picture of the toroidal compactifications, they are smooth complex DM-stacks, we can present the open substacks as the quotient of their frame bundles by the general linear group. The frame bundle is indeed representable, e.g. an ordinary complex manifold, and carries an *infinitesimally* free proper action of the general linear group. If this action is free the stack is equivalent to an ordinary manifold. An equivalence of smooth DM stacks then gives rise to an isomorphism of the frame bundles intertwining the infinitesimally free actions and these glue perfectly well along invariant open subsets.

Also, the construction can be performed with a non-effective finite kernel action whose image is a lattice the price being that one has to consider general smooth Deligne-Mumford stacks. This seems to be inevitable if one wants to work with lattices in $U(2, 1)$ as in [Hol87].

3.2. Fundamental Groups of orbifold toroidal compactifications. The moduli space of \mathcal{X}_{Γ}^* is X_{Γ}^* for $* = \emptyset, BBS, tor$. We will denote by $m : \mathcal{X}_{\Gamma}^* \rightarrow X_{\Gamma}^*$ the moduli map. Using Van Kampen [Zoo02], we get:

Lemma 3.1. *Let x_0 be a base point of \mathcal{X}_{Γ} . Then $\pi_1(\mathcal{X}_{\Gamma}, x_0) = \Gamma$,*

- $\pi_1(\mathcal{X}_{\Gamma}^{tor}, x_0)$ is the quotient Γ^{tor} of Γ by the subgroup normally generated by the Z_c

⁵A torsion free lattice will be called neat if $k = 1$ for every cusp.

- $\pi_1(\mathcal{X}_\Gamma^{BBS}, x_0)$ is the quotient Γ^{BBS} of Γ by the subgroup normally generated by the $H_c \cap \Gamma_c$.

Corollary 3.2 ([KMR92]). *The natural map $H^1(\Gamma^{tor}, \mathbb{Q}) \rightarrow H^1(\Gamma, \mathbb{Q})$ is an isomorphism, consequently the Deligne MHS on $H^1(\Gamma \backslash \mathbb{H}_\mathbb{C}^2, \mathbb{Z})$ is pure of weight one.*

Proof: Dually, it is enough to show that $H_1(\Gamma^{tor}, \mathbb{Q}) \leftarrow H_1(\Gamma, \mathbb{Q})$ is an isomorphism. This amounts to proving that the image of $H_1(Z_c) \rightarrow H_1(\Gamma)$ is torsion. The group Z_c contains the commutator subgroup of $H_c \cap \Gamma_c$ as a finite index subgroup. In particular it maps to the torsion subgroup of $H_1(\Gamma) = \Gamma/[\Gamma, \Gamma]$. \square

The moduli map m presents the fundamental group of X_Γ^* is the quotient of the fundamental group of \mathcal{X}_Γ^* by the normal subgroup generated by the images of the inertia morphisms, see [Noo04], and gives an isomorphism on cohomology with rational coefficients.

Remark 3.3. *Lemma 3.1 is not new, the first point is the easiest special case of [KS15], the second point results from [LKMS15]. Note that in the rank ≥ 2 case, Margulis' normal subgroup theorem implies that the fundamental group of a toroidal compactification of a \mathbb{R} -rank ≥ 2 irreducible locally hermitian symmetric space is finite.*

Question 3.4. *Can Γ^{tor} be finite?*

3.3. Ramification of the natural Orbifold Toroidal compactifications maps.

Lemma 3.5. *Let $\Gamma' < \Gamma$ be a finite index subgroup. Then, the finite covering map $X_{\Gamma'} \rightarrow X_\Gamma$ lifts to an orbifold map $\mathcal{X}_{\Gamma'}^{tor} \rightarrow \mathcal{X}_\Gamma^{tor}$ which restricts over X_Γ to an étale map $\mathcal{X}_{\Gamma'} \rightarrow \mathcal{X}_\Gamma$.*

- Let c', c be cusps such that the mapping $\eta : X_{\Gamma'}^{BBS} \rightarrow X_\Gamma^{BBS}$ maps c' to c . The ramification index of $\mathcal{D}_{c'}$ over \mathcal{D}_c is $d_{c',c} := [Z_{c'} : Z_c]$.
- If Γ' is normal in Γ and $G = \Gamma' \backslash \Gamma$ is the quotient subgroup, G acts ⁶ on $\mathcal{X}_{\Gamma'}^{tor}$, $d_{c',c} := d_c$ depends only on c and $[G \backslash \mathcal{X}_{\Gamma'}^{tor}] = \mathcal{X}_\Gamma^{tor}(\mathcal{D}_\Gamma, (d_c)_{c \in C})$.

Due to ramification, the Γ^{tor} , $\Gamma \in \mathcal{C}_d$, split into infinitely many commensurability classes.

3.4. Hermitian forms over $\mathbb{Q}(\sqrt{-d})$.

Let $d \in \mathbb{N}^*$ be a squarefree positive integer. The imaginary quadratic field $\mathbb{Q}(\sqrt{-d})$ has the complex conjugation as its Galois isomorphism. A non degenerate hermitian form H over $\mathbb{Q}(\sqrt{-d})$ defines a \mathbb{Q} -algebraic group $U(H)$ which is a form of $GL(\dim H)$. The $\mathbb{Q}(\sqrt{-d})$ vector space V_H underlying H will be denoted by V_H . The signature of H is the signature of the corresponding hermitian form also denoted by H on $V := V_H \otimes_{\mathbb{Q}(\sqrt{-d})} \mathbb{C}$ where $\mathbb{Q}(\sqrt{-d}) \rightarrow \mathbb{C}$ is a given embedding.

Let $\mathcal{O}_d \subset \mathbb{Q}(\sqrt{-d})$ be the subring of its quadratic integers. Let L be a free \mathcal{O}_d module which is a lattice in the $\mathbb{Q}(\sqrt{-d})$ vector space V_H underlying H . Then $\Gamma_{H,L} = PU(H) \cap PAut(L)$ is an arithmetic subgroup and the $\Gamma_{H,L}$ all belong to a commensurability class \mathcal{C}_d of lattices in $PU(H)$. Denote by $\Gamma_{\mathbb{Q}}$ be the group of \mathbb{Q} -points of $PU(H)$ which is dense in $PU(H_{\mathbb{R}}) \simeq PU(2, 1)$ with respect to the classical topology.

The study of the groups $\pi_1(\mathcal{X}_{\Gamma'}^{tor})$ and $\pi_1(\mathcal{X}_{BBS}^{tor})$ for $\Gamma \in \mathcal{C}_d$ does not seem to have been carried out systematically in the litterature even in that simple case.

It is known that all non uniform commensurability classes of arithmetic lattices in $PU(2, 1)$ are of the form \mathcal{C}_d .

⁶in the sense of [Rom95]. Actually G acts on the frame bundle of $\mathcal{X}_{\Gamma'}^{tor}$, an ordinary complex manifold, and the action commutes with the right action of the general linear group, from this it is easy to find an étale chart with a strict action of G on the corresponding étale groupoid.

4. PROOF OF THE MAIN THEOREMS

4.1. Commensurability classes with non vanishing virtual b_1 . By commensurability, we mean commensurability in the wide sense:

Definition 4.1. *Two lattices $\Gamma_1, \Gamma_2 < PU(2, 1)$ are commensurable if there exists a finite index torsion free lattice $\Gamma'_i < \Gamma_i$ and a holomorphic isometry $\Gamma'_1 \backslash \mathbb{H}_{\mathbb{C}}^2 \rightarrow \Gamma'_2 \backslash \mathbb{H}_{\mathbb{C}}^2$.*

Definition 4.2. *Say a commensurability class \mathcal{C} of non uniform lattices in $PU(2, 1)$ has non vanishing virtual b_1 if some member Γ_0 satisfies $b_1(\Gamma_0, \mathbb{Q}) > 0$.*

Each commensurability class of non uniform arithmetic lattices in $PU(2, 1)$ has non vanishing virtual b_1 [Wal84], generalizing [Kaz75]. Since these lattices are finitely generated and passing to a sublattice increases $b_1(_, \mathbb{Q})$:

Lemma 4.3. *If \mathcal{C} has non vanishing virtual b_1 every member Γ has a finite index normal subgroup Γ_1 such that $b_1(\Gamma_1, \mathbb{Q}) > 0$.*

4.2. Freeness of virtual Albanese Mappings. We will now fix an **arithmetic** lattice which is torsion free and has unipotent monodromies so that the toroidal compactification \bar{X} is a complex projective manifold. Using $b_1(\Gamma, \mathbb{Q}) = b_1(\Gamma^{tor}, \mathbb{Q})$ we conclude that there is a non zero closed holomorphic one-form α on \bar{X} . We restrict α to X lift it to $\mathbb{H}_{\mathbb{C}}^2$ to construct a closed one form $\tilde{\omega} \in \Omega^1(\mathbb{H}_{\mathbb{C}}^2)$.

Lemma 4.4. *Every element of the vector space V spanned by the $\Gamma_{\mathbb{Q}} \cdot \tilde{\omega}$ is the lift of a holomorphic closed one form on $X_{\Gamma'}^{tor}$ for some normal finite index subgroup $\Gamma' \leq \Gamma$.*

Proof: Indeed $\Gamma_{\mathbb{Q}}$ is the commensurator of Γ . Here we use arithmeticity in a crucial way. \square

Lemma 4.5. *The closure \bar{V} in the Fréchet space $\Omega^1(\mathbb{H}_{\mathbb{C}}^2)$ is a non zero vector space of closed holomorphic 1-forms which is preserved by the action of $PU(H_{\mathbb{R}})$.*

Proof: Indeed $\Gamma_{\mathbb{Q}}$ is dense in $PU(H_{\mathbb{R}})$. \square

Corollary 4.6. *For every point o of $\mathbb{H}_{\mathbb{C}}^2$ there are two elements of V which gives a coframe of the tangent space at o .*

Proof: The restriction $r : \bar{V} \rightarrow \Omega_{\mathbb{H}_{\mathbb{C}}^2, o}^1$ is equivariant under the stabilizer K of o . Hence the image of this linear map is K -equivariant and no zero, the surjectivity of r follows from the irreducibility of the isotropy action on the cotangent space. \square

Corollary 4.7. *There is a finite index normal subgroup Γ' such that every $\Gamma'' < \Gamma'$, has an Albanese map which is unramified on $X_{\Gamma''}$. In particular $\bar{X}_{\Gamma''}^{tor}$ satisfies the Shafarevich conjecture.*

Proof: The immersivity outside the boundary is an immediate consequence of Corollary 4.6 using noetherian induction. I learned from [Sto15] that the fact that the Albanese map does not factor through a curve follows from [Clo93]. The application to Shafarevich conjecture is then a consequence of [Nap90] since an **irreducible** connected component of a fibre of the Albanese mapping cannot give rise to a Nori chain. Note that *SSC* need not be satisfied. \square

Lemma 4.8. *The space $W = \int \bar{V} \subset \mathcal{O}(\mathbb{H}_{\mathbb{C}}^2)$ of primitives of the elements of \bar{V} is infinite dimensional. Actually the map $W \rightarrow \mathcal{O}_{\mathbb{H}_{\mathbb{C}}^2, o} / \mathfrak{m}^N$ is surjective for all $N > 0$.*

Proof: W is invariant under the whole group $PU(H_{\mathbb{R}})$. The evaluation map $ev : W \rightarrow \mathcal{O}_{\mathbb{H}_{\mathbb{C}}^2, o}$ is K -equivariant. So is the completed evaluation map: $W \rightarrow \hat{\mathcal{O}}_{\mathbb{H}_{\mathbb{C}}^2, o}$.

But this is K equivariantly isomorphic to $\mathbb{C}[[\mathfrak{m}/\mathfrak{m}^2]] = \sum_{n \in \mathbb{N}} \text{Sym}^n \mathfrak{m}/\mathfrak{m}^2$. To construct this isomorphism we have chosen the polynomials as a K -invariant subspace of holomorphic functions and the linear functions as generators of the maximal ideal \mathfrak{m} . The K equivariance implies $ev(W) \supset ev(W) \cap \text{Sym}^n \mathfrak{m}/\mathfrak{m}^2$ for all n . If W were finite there would be a finite number of nontrivial $ev(W) \cap \text{Sym}^n \mathfrak{m}/\mathfrak{m}^2$ which would give a direct sum decomposition of $ev(W)$. Hence W would consist of polynomials. Restricting to a generic line through the origin (viewed as a complex geodesic curve isomorphic to a unit disk) there would be a non zero space of polynomials in one variable, generated by monomials, fixed by the homographies which are the automorphisms of the unit disk, which is obviously absurd.

We have already proved surjectivity of $\bar{V} \rightarrow \mathcal{O}_{\mathbb{H}_{\mathbb{C}}^2, o}/\mathfrak{m}^2$. We now use induction on N and assume the theorem is proved for some N . Consider the smallest integer $M > N$ such that $ev(W) \cap \text{Sym}^M \mathfrak{m}/\mathfrak{m}^2 \neq 0$. M exists since W is infinite dimensional. Using the irreducibility of the isotropy representation on $\text{Sym}^M \mathfrak{m}/\mathfrak{m}^2$ we see that $ev(W) \cap \text{Sym}^M \mathfrak{m}/\mathfrak{m}^2 = \text{Sym}^M \mathfrak{m}/\mathfrak{m}^2$. Hence there is a monomial say z_1^M in W . Now one of the infinitesimal generators of the Lie algebra of $PU(H_{\mathbb{R}})$ takes the form $\xi = \frac{\partial}{\partial z_1} + \xi'$ where $\xi' \in \mathfrak{m}\mathcal{T}_{\mathbb{H}_{\mathbb{C}}^2}$. If $M > N + 1$, $\xi.z_1^M \in W$ would contradict the minimality of M . Hence $M = N + 1$ which is the desired conclusion. \square

Corollary 4.9. *Fix $M \in \mathbb{N}$. There is a finite index normal subgroup Γ' such that for all $\Gamma'' < \Gamma'$, $\bar{X}_{\Gamma''}^{\text{tor}}$ has an Albanese map which generates M jets at all points of $X_{\Gamma''}$.*

In particular, if $M = 2$ the genus zero curves in the Albanese image lie on the image of the boundary and we recover special cases of classical results:

Corollary 4.10 ([Nad89, BT15]). *$\bar{X}_{\Gamma''}^{\text{tor}}$ satisfies the Green-Griffiths-Lang conjecture*

Proof: Immediate corollary of [Och77, Theorem D]. \square

Corollary 4.11. *$X_{\Gamma''}$ has finitely many rational points on any number field of definition.*

Proof: Immediate corollary of Faltings' solution of Lang's conjecture [Fal94], see also [HinSil00, Theorem F.1.1, p. 436]. \square

4.3. Refined commensurability classes. The following definition is a slight generalization of the notion of completely étale map between negatively elliptic bounded surfaces [Hol86, p.256]:

Definition 4.12. *Two (commensurable) lattices $\Gamma_1, \Gamma_2 < PU(2, 1)$ are refined commensurable if there exists a finite index lattice $\Gamma'_i < \Gamma_i$ and a holomorphic isometry $\Gamma'_1 \backslash \mathbb{H}_{\mathbb{C}}^2 \rightarrow \Gamma'_2 \backslash \mathbb{H}_{\mathbb{C}}^2$ such that $\mathcal{X}_{\Gamma'_1}^{\text{tor}} \rightarrow \mathcal{X}_{\Gamma'_2}^{\text{tor}}$ is étale.*

If $\mathcal{X}_{\Gamma'_i}^{\text{tor}}$ is uniformizable we can assume that Γ'_i is torsion free and neat. The questions (1)-(6) (and the orbifold Kodaira dimension) in the introduction depend only on the refined commensurability class.

Lemma 4.13. *Two commensurable arithmetic lattices Γ_1, Γ_2 having a common finite index lattice are refined commensurable iff for every $\tilde{c} \in C \subset \partial \mathbb{H}_{\mathbb{C}}^2$ there is a common finite index subgroup Γ_3 such that $Z(H_{\tilde{c}}) \cap \Gamma_1 = Z(H_{\tilde{c}}) \cap \Gamma_2 = Z(H_{\tilde{c}}) \cap \Gamma_3$.*

There is an order on the set \mathcal{C}_r of refined commensurability classes of a given commensurability class. We say $\mathcal{C}_1 \prec \mathcal{C}_2$ if there is some member of \mathcal{C}_2 is a finite index subgroup of a member of \mathcal{C}_1 . It is clear that (\mathcal{C}_r, \prec) is a filtering order and that the questions investigated in the introduction are more difficult for small elements (with the exception of $L(_)$).

Question 4.14. *Is (C_r, \prec) Artinian? Is every initial segment finite?*

Definition 4.15. *A refined commensurability class is said to be small if the common universal covering stack of the corresponding $\mathcal{X}_\Gamma^{\text{tor}}$ is not a Stein manifold. Its Γ -dimension is the dimension of the Campana quotient of the moduli space of the universal covering stack by its compact complex subvarieties.*

They are the most interesting classes from the present perspective.

4.4. Higher dimensions. The only missing ingredient being [Nap90], we get if $n \geq 3$:

Theorem 4.16. *If $\Gamma < PU(n, 1)$ is arithmetic and small enough in its commensurability class, \bar{X}_Γ has Albanese dimension n and its image contains no translate of an abelian subvariety at its generic point. It satisfies Q , has big cotangent bundle, is Kobayashi hyperbolic modulo its boundary, and the rational points over a number field over which it is defined are finite modulo its boundary.*

5. THE PICARD-EISENSTEIN COMMENSURABILITY CLASS

5.1. The Picard-Eisenstein Lattice. Let us look at the Picard-Eisenstein lattices using its beautiful two-generator presentation [FP06]. This group, denoted by $PU(H_0, \mathbb{Z}[\omega])$ is in \mathcal{C}_3 and is actually of the form $\Gamma_{H,L}$ for the hermitian form whose matrix is $H_0 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$ and $L = \mathcal{O}_3^{\oplus 3} = \mathbb{Z}[\frac{-1+i\sqrt{3}}{2}]^{\oplus 3}$ is the standard lattice.

Proposition 5.1. *$PU(H_0, \mathbb{Z}[\omega])^{\text{tor}}$ has a surjective morphism to a $(2, 3, 6)$ orbifold group.*

Proof: The presentation in [FP06] is

$$\langle P, Q, R \mid R^2 = [PQ^{-1}, R] = (RP)^3 = P^3Q^{-2} = (QP^{-1})^6 \rangle.$$

We are imposing $P^3 = Q^2 = 1$ in this presentation by [FP06, p. 258]. Further imposing $R = 1$ gives $\langle P, Q \mid P^3 = Q^2 = (PQ^{-1})^6 = 1 \rangle$ as a quotient group. \square

It may be reassuring to get a confirmation of the:

Corollary 5.2. *The Picard-Eisenstein commensurability class \mathcal{C}_3 has non vanishing virtual b_1 .*

5.2. Hirzebruch's example of a configuration of elliptic curves in an abelian surface whose complement is complex hyperbolic. It is a well known theorem of Holzapfel [Hol86] that the following example is in \mathcal{C}_3 .

5.2.1. Let $E \simeq \mathbb{C}/\mathbb{Z}[j]$ be the elliptic curve isomorphic to Fermat's cubic curve and let \bar{X} be the blow up of $(0_E \times_E)$. We denote by D_i the strict transforms in \bar{X} of the elliptic curves E_i where:

$$E_1 = E \times \{0_E\}, E_2 = \{0_E\} \times E, E_3 = \Delta = Gr(id_E), E_4 = Gr(-j)$$

and we obtain a SNC divisor (\bar{X}, D) where $D = D_1 + D_2 + D_3 + D_4$. The D_i are pairwise disjoint.

Theorem 5.3. *(Hirzebruch, [BHH87]) $X \simeq \Gamma_{Hirz} \backslash \mathbb{H}_{\mathbb{C}}^2$ where $\Gamma_{Hirz} \subset PU(2, 1)$ is a non uniform lattice. \bar{X} is its toroidal compactification.*

The notation (\bar{X}, D) will be used throughout this section to denote this particular configuration.

Corollary 5.4. *There is a neat lattice $\Gamma' \in \mathcal{C}_3$ such that for every $\Gamma'' < \Gamma'$, $\bar{X}_{\Gamma''}^{tor}$ has a finite Albanese map. In particular the universal covering space of $\bar{X}_{\Gamma''}^{tor}$ is Stein.*

Proof: Corollary 4.7 implies that the only curves contracted by the Albanese map lie on the boundary. It is thus enough to replace the Γ' in Corollary 4.7 with $\Gamma' \cap \Gamma_{Hirz}$ for which the Albanese map does not contract any boundary curve. \square

Remark 5.5. [DeMo86, Prop 15.17, P. 155] *This configuration of elliptic curves has an order 72 complex reflection group H of automorphisms which is a μ_6 central extension of A_4 acting as the symmetry group of the configuration of 4 points in \mathbb{P}^1 given by the points $0, 1, \infty, e^{\frac{2\pi i}{6}}$. A_4 acts as an alternating group on the set whose elements are the 4 elliptic curves.*

5.2.2. *The Orbifold attached to a Normal subgroup of Γ_{Hirz} .* Let us make Lemma 3.5 more explicit in the present case.

Let $\Gamma' \triangleleft_{fi} \Gamma_{Hirz}$ be a finite index normal subgroup. Then Γ' is torsion-free and torsion-free at infinity so that the toroidal compactification $\bar{X}' \supset X' = \Gamma' \backslash \mathbb{H}_{\mathbb{C}}^2$ is a complex projective manifold with an effective $G = \Gamma_{Hirz}/\Gamma'$ -action which is fixed point free on X' . It is clear that $G \backslash \bar{X}' = \bar{X}$ and we denote the corresponding quotient orbifold by $\mathcal{X}' = [G \backslash \bar{X}']$ and the corresponding quotient map by $\pi : \bar{X}' \rightarrow \bar{X}$.

Lemma 5.6. *Let $p'_i \subset E'_i$ be a flag consisting of a point $p_i \in \pi^{-1}(E_i)$ and E'_i the irreducible component of $\pi^{-1}(E_i) = \sum_j F_{ij}$ via p_i . Consider*

$$S'_i = \text{Stab}_G(p'_i) < H_i = \text{Stab}_G(E'_i) < G.$$

*Then S'_i is a cyclic central subgroup of H_i of order $d_i = d_i(\Gamma')$, $G_i = H_i/S'_i$ acts effectively on E'_i without fixed points, $E_i = G_i \backslash E'_i$ and $p^*E_i = \sum_{j \in H_i \backslash G} d_i F_{ij}$.*

Furthermore \mathcal{X}' is equivalent to $\mathcal{X}(\bar{X}, D, d)$ with $d = (d_1, \dots, d_4)$.

We shall adopt the notations of Lemma 5.6 in the rest of section 5.2.

Lemma 5.7. *Conversely every proper finite étale mapping $p : \bar{Y} \rightarrow \mathcal{X}(\bar{X}, D, d)$ comes from a finite index subgroup $\Gamma' = \text{Im}(\pi_1(\bar{Y} \setminus p^{-1}(D)) \rightarrow \pi_1(\bar{X} \setminus D))$ such that $p^*D = \sum_i d_i \text{Supp}(p^{-1}(E_i))$.*

The support $\text{Supp}(D)$ of an effective Cartier divisor D is the sum of its irreducible components with multiplicity one.

The existence of \bar{Y} is equivalent to $\mathcal{X}(\bar{X}, D, d)$ being uniformizable.

Question 5.8. *For which $d \in \mathbb{N}_{\geq 1}^4$ is $\mathcal{X}(\bar{X}, D, d)$ uniformizable?*

Proposition 5.9. *$\mathcal{X}(\bar{X}, D, d)$ is not developpable if $\{d_1, \dots, d_4\} = \{1, 1, 1, m\}$ or $\{1, 1, m, m'\}$ with $1 < m < m'$.*

Proof: In the listed case the exceptional curve in \bar{X} gives a sub-orbifold equivalent to $\mathbb{P}(m, [0])$ or to $\mathbb{P}(m, [0] + m'[1])$ which are non developpable, which obstructs the developpability of the ambient orbifold. \square

Lemma 5.10. *Assume $d \in \mathbb{N}_{\geq 1}^4$ is such that $\mathcal{X}(\bar{X}, D, d)$ uniformizable. Let $Z := \text{Exc}$ be the exceptional curve and $\mathcal{Z} = (\mathbb{P}^1, d_1, d_2, d_3, d_4)$ be the natural suborbifold. $SS(\mathcal{X}(\bar{X}, D, d))$ holds iff it holds for \mathcal{Z} .*

5.2.3. $d_3 = d_4 = 1$. This corresponds to studying the pair $(\bar{X}, D'' := D_1 + D_2)$. In order to fix notations, we denote by U'' the complement of $E_1 + E_2$ in $E \times E$ and by X'' the complement of $D_1 + D_2$ in \bar{X} . Plainly $U'' = X'' \setminus \text{Exc}$.

Lemma 5.11. *The fundamental group of U'' is a product of two finite groups on 2 generators $\pi_1(U'') = \mathbb{F}_2(a, b) \times \mathbb{F}_2(c, d)$. The fundamental group of X'' is the quotient of $\pi_1(U'')$ by the normal subgroup generated by $[a, b].[c, d]$.*

Proof: Elementary calculation. \square

Corollary 5.12. *The natural morphism $F_2(a, b) \rightarrow \pi_1(X'')$ factors through $H_{\mathbb{Z}}$ the Heisenberg group of unipotent 3×3 upper triangular matrices with integer coefficients .*

Proof: Since a and b commute with c and d the relation $[a, b] = [c, d]^{-1}$ implies that a and b commute with $[a, b]$. But $F_2(a, b)/\langle\langle [a, [a, b]], [b, [a, b]] \rangle\rangle \simeq H_{\mathbb{Z}}$. \square

We have a non trivial central extension:

$$1 \rightarrow \mathbb{Z}_{[a, b]} = Z(H_{\mathbb{Z}}) \rightarrow H_{\mathbb{Z}} \rightarrow \mathbb{Z}_{a, b}^2 \rightarrow 1.$$

The geometric interpretation is clear. Consider the boundary B of a regular neighborhood of D_1 in \bar{X} . If the base point is in B a and b can be homotoped in $B \cap U''$. However $B \subset X''$ and $\pi_1(B) = H_{\mathbb{Z}}$ since $D_1^2 = -1$.

Corollary 5.13. *The fundamental group $\pi_1(X'')$ is the quotient group of $H_{\mathbb{Z}}^2$ by the diagonal subgroup of its center $\Delta_{\mathbb{Z}} : \mathbb{Z} \rightarrow \mathbb{Z}^2$. Thus, we have the central extension:*

$$1 \rightarrow \mathbb{Z} \rightarrow \pi_1(X'') \rightarrow \mathbb{Z}^4 = H_1(X'') \rightarrow 1,$$

$[a, b]$ mapping to a generator of the center and $[c, d]$ to its opposite.

Proof: The map $H_{\mathbb{Z}}^2 \rightarrow \pi_1(X'')$ comes from the previous lemma which can also be applied with c, d and all these groups have naturally isomorphic abelianization. \square

Corollary 5.14. *$\mathcal{X}(\bar{X}, D, n, n, 1, 1)$ is uniformizable.*

Proof: The new relations to get $\pi_1(\mathcal{X}(\bar{X}, D, n, n, 1, 1))$ are $[a, b]^n = [c, d]^n = 1$. It thus suffices to consider the quotient map

$$\pi_1(\mathcal{X}(\bar{X}, D, n, n, 1, 1)) \rightarrow H_{\mathbb{Z}/n\mathbb{Z}}^2 / \Delta_{\mathbb{Z}}(\mathbb{Z}/n\mathbb{Z}).$$

\square

Thanks to lemma 5.10 $SSC(\mathcal{X}(\bar{X}, D, n, n, 1, 1))$ is trivially true since the fundamental group of $\mathcal{Z} \simeq \mathbb{P}^1(\sqrt[n]{0 + \infty})$ is finite.

5.2.4. $d_4 = 1$. This corresponds to studying the pair $(\bar{X}, D' := D_1 + D_2 + D_3)$. In order to fix notations, we denote by U' the complement of $E_1 + E_2 + E_3$ in $E \times E$ and by X' the complement of $D_1 + D_2 + D_3$ in \bar{X} . Plainly $U' = X' \setminus Exc$.

Lemma 5.15. *The fundamental group $G' = \pi_1(U')$ has 5 generators a, b, c, d, e and is presented by the relations:*

$$\begin{aligned} a^{-1}ca &= c, a^{-1}ea = e, a^{-1}da = c^{-1}dce, \\ b^{-1}db &= d, b^{-1}eb = e, b^{-1}cb = d^{-1}cde^{-1}. \end{aligned}$$

Proof: Omitted. Easy calculation. \square

It is a semi-direct product of $F_3(c, d, e)$ by $F_2(a, b)$.

One has to set $\alpha = c^{-1}a$ $\beta = d^{-1}b$ $e = 1$ to recover the previous case.

Lemma 5.16. *Let $V' \subset U'$ the trace on U' of a regular neighborhood of Exc . Then $\pi_1(V') \hookrightarrow \pi_1(U')$ is generated by a_1, a_2, a_3 subject to the relations:*

$$a_3a_2a_1 = a_2a_1a_3 = a_1a_3a_1$$

where

$$a_3 = e, a_1 = c^{-1}d^{-1}cde^{-1}, a_2 = \alpha^{-1}\beta^{-1}e^{-1}\alpha\beta.$$

The center $Z(\pi_1(V'))$ is infinite cyclic generated by the element $\gamma_Z = a_3a_2a_1 = eaba^{-1}b^{-1}$.

Proof: Omitted. In principle the calculation is easy, but it turned out to be rather messy. \square

Lemma 5.17. *The fundamental group $G := \pi_1(X')$ has 5 generators a, b, c, d, e and is presented by the relations 5.15 plus:*

$$eaba^{-1}b^{-1} = 1.$$

The fundamental group $\pi_1(\mathcal{X}(\bar{X}, D, d_1, d_2, d_3, 1))$ has 5 generators a, b, c, d, e and is presented by the relations 5.15 plus:

$$a_3a_2a_1 = a_1^{d_1} = a_2^{d_2} = a_3^{d_3} = 1.$$

The map $\pi_1(\mathcal{Z}) = F_2(a_1, a_2) / \langle\langle a_1^{d_1}, a_2^{d_2}, (a_2a_1)^{d_3} \rangle\rangle \rightarrow \pi_1(\mathcal{X}(\bar{X}, D, d_1, d_2, d_3, 1))$ maps a_1, a_2 to their above expressions.

Hence by Lemma 5.10 SSC holds true in that case if and only if we can find a finite index normal subgroup H of $G(d_1, d_2, d_3) := \pi_1(\mathcal{X}(\bar{X}, D, d_1, d_2, d_3, 1))$ and $\eta \in H_1(H, \mathbb{Q})$ which does not vanish on $\langle a_1, a_2 \rangle \cap H$. The worst possible choice is when $G(d_1, d_2, d_3)/H$ is abelian. Unfortunately since $H_1(G(d_1, d_2, d_3))$ has rank 4 a lot of the H one gets with `LowIndexSubgroups` in GAP or MAGMA have that property.

Lemma 5.18. *$\mathcal{X}(\bar{X}, D, (n, n, n, 1))$ is uniformizable.*

Proof: Immediate consequence of Corollary 5.14. Indeed, thanks to Lemma 2.2, we have a map

$$\mathcal{X}(\bar{X}, D, (n, n, n, 1)) \rightarrow \mathcal{X}(\bar{X}, D, (n, n, 1, 1))$$

and a map $\mathcal{X}(\bar{X}, D, (n, n, n, 1)) \rightarrow \mathcal{X}(\bar{X}, D, (1, n, n, 1))$ thanks to Remark 5.5 which gives a group morphism in a finite group

$$\pi_1(\bar{X}, D, (n, n, n, 1)) \rightarrow (H_{\mathbb{Z}/n\mathbb{Z}}^2 / \Delta_{\mathbb{Z}}(\mathbb{Z}/n\mathbb{Z}))^2$$

which is injective on the isotropy groups. \square

Hence $G_n = \pi_1(\mathcal{X}(\bar{X}, D, n, n, n, 1))$ is the fundamental group of a complex projective surface. Let us introduce the following quotient of G_n :

$$G_n^- := G_n / \langle\langle a^n, b^n, c^n, d^n \rangle\rangle.$$

where a, b, c, d denote the natural images of the generators of group G .

Lemma 5.19. *The nilpotent group⁷ of class 2 $G_n^- / \gamma_3(G_n^-)$ is isomorphic to $H_{\mathbb{Z}/n\mathbb{Z}}^2$.*

Proof: Using MAGMA [BCP06], we get an isomorphism:

$$G / \gamma_3(G) \simeq H_{\mathbb{Z}}^2,$$

whence the result since we are killing the n -th power of the lifted generators of $H_1(G) = G / \gamma_2(G)$. A file containing the MAGMA code is available on my webpage. \square

Proposition 5.20. *SSC($\mathcal{X}(\bar{X}, D, (3, 3, 3, 1))$) holds and the universal covering space is Stein.*

Proof:

The abelianization A_3 of $K_3 := \ker(\phi)$, where we denote by ϕ the resulting epimorphism $\phi : G_3 \rightarrow H_{\mathbb{Z}/3\mathbb{Z}}^2$ is free of rank 10, thanks to MAGMA. MAGMA computes the image of $a_1a_3a_1^{-1}a_3^{-1}$ to be the row vector $(0, -1, -1, 1, -1, 0, 0, 0, -1, 2)$ in MAGMA's basis of A_3 .

\square

⁷We use the following notation for the central series of a group G : $\gamma_1(G) = G$, $\gamma_2(G) = [G, G]$, $\gamma_{k+1}(G) = [\gamma_k(G), G]$

Corollary 5.21. *For all $k, l, m \in \mathbb{N}^*$, $SSC(\mathcal{X}(\bar{X}, D, (3k, 3l, 3m, 1)))$ holds and the moduli space of the universal covering stack is Stein, hence the universal covering space is Stein provided the orbifold is indeed developable.*

Proof: Use the natural map $\mathcal{X}(\bar{X}, D, (3k, 3l, 3m, 1)) \rightarrow \mathcal{X}(\bar{X}, D, (3, 3, 3, 1))$ given by Lemma 2.2 to deduce that the Albanese map is virtually finite. \square

5.2.5. *General case.* We have no general results on uniformizability. It follows from the above that $\mathcal{X}(\bar{X}, D, (n, n, n, n))$ is uniformizable for all $n \in \mathbb{N}^*$, and $SSC(\mathcal{X}(\bar{X}, D, (3k, 3l, 3m, 3p)))$ holds.

5.3. Some small refined commensurability classes in \mathcal{C}_3 .

5.3.1. Quite confusingly, the Picard-Eisenstein lattice studied in [Hol86, Hol87] is not the same as the one studied in [FP06]. Also the relationship with Γ_{Hirz} and other lattices in \mathcal{C}_3 is slightly involved. Holzapfel's Picard modular group is (conjugate by a transposition matrix to) $\Gamma_{H_1, L_{st}}$ for the hermitian form whose matrix is H_1 and $L_{st} = \mathcal{O}_3^{\oplus 3} = \mathbb{Z}[\frac{1+i\sqrt{3}}{2}]^{\oplus 3}$ is the standard lattice. By definition $H_0 = {}^t \bar{g}H_1g$ where we have used the notations

$$H_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad g = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1/2 \\ 1 & 0 & -1/2 \end{pmatrix}.$$

As \mathbb{Q} -algebraic groups $PU(H_1) \simeq PU(H_0)$ - more precisely $gPU(H_0)g^{-1} = PU(H_1)$ but the $PU(H_i, \mathcal{O}_3)$ $i = 0, 1$ are different lattices. As we will see, the refined commensurability classes of these two lattices have a very similar behaviour but we did not complete the elementary but lengthy calculations to check that they are equal.

5.3.2. The article [Hol86] uses another special elliptic configuration in $E \times E$. Let $0_E, Q_1, Q_2$ be the fixed points of the automorphism $j : E \rightarrow E$ where \mathcal{O}_3^\times acts by multiplication on $E = \mathbb{C}/\mathcal{O}_3$ and $\mathcal{O}_3 = \mathbb{Z}[j] \subset \mathbb{C}$ is the inclusion given by our choice of $i = \sqrt{-1}$ and the formula $j = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$. Then $\{0_E, Q_1, Q_2\}$ defines a group of translations $T := \mathbb{Z}/3\mathbb{Z}$ of E . Let T acts diagonally on $E \times E$. It turns out that there is an isomorphism $T \backslash E \times E \simeq E \times E$ such that the inverse image of the Hirzebruch configuration is the union of 6 elliptic curves. This elliptic curve configuration has $\{0_E \times 0_E, Q_1 \times Q_1, Q_2 \times Q_2\}$ as its multiple (quadruple) points. These 6 elliptic curves $\{S_k\}_{k=1}^6$ are the graph of the automorphisms $1, j, j^2$ and the horizontal factors $\{E \times 0_E, E \times Q_1, E \times Q_2\}$. Thus there is an index 3 subgroup $\Gamma_{Holz} \subset \Gamma_{Hirz}$ which can be defined by the Galois correspondance

$$\Gamma_{Holz} = Im(\pi_1(Bl_{T, (0_E \times 0_E)}(E \times E) \setminus \cup_{k=1}^6 S'_k) \rightarrow \pi_1(\bar{X} \setminus D) = \Gamma_{Hirz}) \triangleleft \Gamma_{Hirz}$$

where S'_k is the strict transform of S_k and the notation (\bar{X}, D) of subsection 5.2 still applies.

One of the facts [Hol86] uses is that we have a composition sequence:

$$\Gamma_{Holz} \triangleleft P\tilde{\Gamma}'_{Holz} := P(SU(H_1, \mathcal{O}_3)(1-j)) \triangleleft PU(H_1, \mathcal{O}_3)$$

with graded quotients

$$P\tilde{\Gamma}'_{Holz}/\Gamma_{Holz} = \mu_3 \times \mu_3, \quad PU(H_1, \mathcal{O}_3)/P(SU(H_1, \mathcal{O}_3)(1-j)) = S_4.$$

Furthermore, thanks to [GeSa17], we can interpret a crucial ingredient in [Hol87] as an equivalence

$$\begin{aligned} [P\tilde{\Gamma}'_{Holz} \backslash \mathbb{H}_{\mathbb{C}}^2] &\simeq \mathcal{X}(\mathbb{P}^2 \setminus \{4pts\}, CEVA(2), (3, 3, 3, 3, 3, 3)) \\ [S_4 \backslash [P\tilde{\Gamma}'_{Holz} \backslash \mathbb{H}_{\mathbb{C}}^2]] &\simeq [PU(H_1, \mathcal{O}_3) \backslash \mathbb{H}_{\mathbb{C}}^2], \end{aligned}$$

where $CEVA(2)$ is the complete quadrangle built on the 4 marked points (in general linear position) in \mathbb{P}^2 and $S_4 < PGL(3, \mathbb{C})$ permutes these 4 points.

5.3.3. Using the equivalence $[\mu_3 \setminus E] \simeq \mathbb{P}^1(\sqrt[3]{0+1+\infty})$ we see easily that the quotient orbifold $[\mu_3 \times \mu_3 \setminus Bl_{T,(0_E \times 0_E)}(E \times E)]$ is the following orbisurface: the moduli space is $Bl_{(0,0),(1,1),(\infty,\infty)}\mathbb{P}^1 \times \mathbb{P}^1$ and the ramification has order 3 on the 9 curves given by the 3 exceptional curves and the strict transforms of the vertical and horizontal factors through the 3 blown up points. There is no ramification over the strict transform of the diagonal.

In particular $\mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor}$ is the orbifold whose moduli space is $Bl_{(0,0),(1,1),(\infty,\infty)}\mathbb{P}^1 \times \mathbb{P}^1$ and which ramifies at order 3 on the 6 (-1)-rational curves given by the 3 exceptional curves and the 3 strict transforms of the vertical factors. The 4 boundary components are the strict transforms of the diagonal and the 3 strict transforms of the horizontal factors carry multiplicity 1 since our construction of the orbifold toroidal compactifications precisely excludes orbifold behaviour at the general point of the boundary.

If $CEVA'(2)$ denotes the strict transform of $CEVA(2)$ in the blow up surface $Bl_{4pts}(\mathbb{P}^2)$ we see using the familiar contraction of these 4 disjoint rational curves and [GeSa17] an equivalence:

$$\mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor} \simeq \mathcal{X}(Bl_{4pts}(\mathbb{P}^2), CEVA'(2), (3, 3, 3, 3, 3, 3))$$

Using the language of [BHH87] we assign weight 3 to the strict transforms of the lines in $CEVA(2)$ and the weight 1 to the exceptional curves. Since the map $\mathcal{X}_{\Gamma_{Holz}}^{tor} \rightarrow [\mu_3 \times \mu_3 \setminus Bl_{3pts}(E \times E)]$ is étale, $\mathcal{X}_{\Gamma_{Holz}} \rightarrow \mathcal{X}_{P\tilde{\Gamma}'_{Holz}}$ is the only non-étale map in the orbifold version of the main diagram in [Hol86]:

$$\begin{array}{ccc} & \mathcal{X}_{\Gamma_{Holz}}^{tor} = Bl_{3pts}(E \times E) & \\ & \swarrow \text{ram} & \searrow \text{et} \\ \mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor} & & \mathcal{X}_{\Gamma_{Hirz}}^{tor} = Bl_0(E \times E) \\ \swarrow \text{et} & & \\ \mathcal{X}_{PU(H_1, \mathcal{O}_3)}^{tor} & & \end{array}$$

In other words, we have in \mathcal{C}_{3r} :

$$[PU(H_1, \mathcal{O}_3)] = [P\tilde{\Gamma}'_{Holz}] \prec [\Gamma_{Holz}] = [\Gamma_{Hirz}].$$

Both classes are small of Γ -dimensions 1 and 2.

Proposition 5.22. $(P\tilde{\Gamma}'_{Holz})^{tor} \cong \pi_1(\mathbb{P}^1(3, 3, 3)) \cong \mathbb{Z}[j] \rtimes \mu_3$. $PU(H_1, \mathcal{O}_3)^{tor}$ is virtually abelian of rank 2 sitting in an exact sequence:

$$\{1\} \rightarrow \mathbb{Z}[j] \rtimes \mu_3 \rightarrow PU(H_1, \mathcal{O}_3)^{tor} \rightarrow S_4 \rightarrow \{1\}.$$

Proof: Let us consider the linear system $|I_{4pts}(2)|$ of conics through the four points. It defines a rational map $\phi : \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ which becomes regular on $Bl_{4pts}(\mathbb{P}^2)$, the exceptional curves are then isomorphically mapped ϕ and $CEVA'(2)$ has three connected components which have 2 irreducible as components and coincide to the three singular fibers of ϕ . The generic fibre is a smooth conic with no deleted points. In other words,

$$\phi : \mathbb{P}^2 \setminus CEVA'(2) \rightarrow \mathbb{P}^1 \setminus \{3pts\}$$

is a projective smooth conic bundle. Lemma 2.2 gives a map:

$$\bar{\phi} : \mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor} \rightarrow \mathbb{P}^1(3, 3, 3)$$

whose general fiber is a smooth rational curve. This gives an isomorphism

$$(P\tilde{\Gamma}'_{Holz})^{tor} \xrightarrow{\cong} \pi_1(\mathbb{P}^1(3, 3, 3)).$$

The other statements are immediate consequences granted the geometric description above. \square

Corollary 5.23. *When Γ lies in $[PU(H_1, \mathcal{O}_3)]$ or $[\Gamma_{Hirz}]$, Γ^{tor} is infinite virtually abelian and linear.*

5.3.4. In the notations of [Mos86], the group $PU(H_0, \mathcal{O}_3)$ can be described as Γ_{μ, S_3} with $\mu = \frac{1}{6}(2, 2, 2, 1, 5)$ [Der]. Since this ball 5-uple satisfies INT, the orbifold $[PU(H_0, \mathcal{O}_3) \setminus \mathbb{H}_{\mathbb{C}}^2]$ can be easily described.

The moduli space of $[\Gamma_{\mu} \setminus \mathbb{H}_{\mathbb{C}}^2]$ is $\mathbb{P}^2 \setminus \{P\}$ this point P being a triple point of $CEVA(2)$. In $[\Gamma_{\mu} \setminus \mathbb{H}_{\mathbb{C}}^2]$ the 3 lines through P have orbifold weight 3, the three remaining lines have orbifold weight 2 and the 3 triple points have a non abelian order 36 inertia group [Ulu07, pp 392-393].

Then $\mathcal{X}_{\Gamma_{\mu}}^{tor}$ has $Bl_P(\mathbb{P}^2)$ as its moduli space and the only modification is that we should affect the weight 1 to the exceptional curve.

When we mod out by the action of S_3 fixing P and permuting the other triple points we observe that there is no ramification on the exceptional curve. In particular $[\mathcal{X}_{PU(H_0, \mathcal{O}_3)}^{tor}] \simeq S_3 \setminus \mathcal{X}_{\Gamma_{\mu}}^{tor}$. In terms of refined cohomology classes $[PU(H_0, \mathcal{O}_3)] = [\Gamma_{\mu}]$.

The central projection to P defines a map $\mathcal{X}_{\Gamma_{\mu}}^{tor} \rightarrow \mathbb{P}^1(3, 3, 3)$ whose general fiber F is a $\mathbb{P}^1(2, 2, 2)$ an elliptic orbifold whose fundamental group is the Vierergruppe. This implies that $PU(H_0, \mathcal{O}_3)^{tor}$ is virtually abelian of rank 2 and that the morphism of Proposition 5.1 has a finite kernel. Actually, $\pi_1(F)$ injects thanks to:

Proposition 5.24. *There is a (split) exact sequence*

$$1 \rightarrow (P\tilde{\Gamma}'_{Holz})^{tor} \rightarrow \Gamma_{\mu}^{tor} \rightarrow K_4 \rightarrow 1$$

and the refined commensurability classes of $PU(H_0, \mathcal{O}_3)$ and $PU(H_1, \mathcal{O}_3)$ are the same.

Proof: The group S_4 acts on the linear system $|I_{4pts}(2)|$ by the projectivities preserving the 4 points. On the 3 singular members it acts as S_3 where $S_3 = S_4/K_4$ where K_4 is the Vierergruppe or Klein group isomorphic to $(\mathbb{Z}/2)^2$. In particular K_4 acts as automorphisms of the map ϕ (see also [Hol87, I.6.2]) and its orbifold compactification $\bar{\phi}$ in Proposition 5.22. It is then easy to interpret [Hol87, I.3.6.3] and see that $[K_4 \setminus \mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor}] = [\mathcal{X}_{\Gamma_{\mu}}^{tor}]$. \square

5.3.5. *The universal covering stack attached to $[PU(H_0, \mathcal{O}_3)]$.*

Proposition 5.25. *$[\mathcal{X}_{PU(H_0, \mathcal{O}_3)}^{tor}]$ is not developable.*

Proof: We consider the étale map $e : [\mu_3 \setminus E] \xrightarrow{\cong} \mathbb{P}^1(3, 3, 3)$. Then

$$E \times_{e, \bar{\phi}} \mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor} \simeq \mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor}$$

where $P\tilde{\Gamma}'_{Holz}$ is the lattice in $PU(2, 1)$ corresponding to the group Γ' in the notations of [Hol87, p. 27]. The moduli space of that stack is a surface birational to $E \times \mathbb{P}^1$ with 3 A_2 singular points which is actually isomorphic to $\mu_3 \times \{1\} \setminus Bl_{T, 0_E \times 0_E} E \times E$. The orbifold structure of $\mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor}$ a μ_3 inertia group at the singular points. So there are orbifold points in the fiber of the map $\mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor} \rightarrow E$ which is an isomorphism on π_1 . In particular the universal covering stack is equivalent to $\mathbb{C} \times_E \mathcal{X}_{P\tilde{\Gamma}'_{Holz}}^{tor}$ and has infinitely many μ_3 orbifold points. \square

5.3.6. $SSC(\mathcal{X}(\bar{X}, D, (n, n, n, n)))$. The results in subsection 5.2.5 were slightly unsatisfying but one can settle SSC in the case where the d_i have a common factor:

Proposition 5.26. *If $n \in \mathbb{N}_{\geq 2}$, $SSC(\mathcal{X}(\bar{X}, D, (n, n, n, n)))$ holds and the universal covering space is a Stein manifold.*

Proof: It is enough to prove this for the 3-1 étale cover \mathcal{X}' given by the blow up at 3 points of $E \times E$ with weights n on the strict transform of the Holzapfel configuration of 6 elliptic curves. Let us take the (étale) quotient stack by the action of $\mu_3 \times \mu_3$ we have already encountered. The resulting orbifold \mathcal{Y} can be described in the language of [BHH87] as follows: the moduli space is \mathbb{P}^2 blown up at 4 points, the strict transforms of the lines in CEVA(2) have weight 3, one exceptional curve has weight n , the 3 other exceptional curves have weight $3n$.

Let us now consider the linear projection from this blown up \mathbb{P}^2 to \mathbb{P}^1 with center the point whose exceptional curve has weight n . It is a regular map which has 3 special fibers which are isomorphic to a nodal conic, the irreducible constituents carrying weights 3 and $3n$. Hence there is a map $\mathcal{Y} \rightarrow \mathbb{P}^1(3, 3, 3)$. Composing with the natural étale map $\mathcal{X}' \rightarrow \mathcal{Y}$, we get a map $\mathcal{X}' \rightarrow \mathbb{P}^1(3, 3, 3)$ which is not constant on the 3 preimages of \mathcal{Z} . Since $\mathbb{P}^1(3, 3, 3)$ is elliptic and has virtually abelian rank 2 fundamental group the proposition follows.

□

6. CONCLUDING REMARKS

We conclude by a discussion of some interesting examples from the literature.

In [Sto15, DicSto17], bielliptic smooth toroidal compactifications of ball quotients are constructed. They satisfy $L(_)$ since the fundamental group is virtually abelian hence linear. The Shafarevich conjecture is established for surfaces of Kodaira dimension ≤ 1 by [GurSha85] and their argument gives that the fundamental group is linear.

More to the point, [DiCSto18, Theorem 1.3] asserts that a smooth toroidal compactification of a ball quotient which is birational to an abelian or a bielliptic surface is the blow up in finitely many distinct points of the minimal surface. In particular, it has a finite (abelian) cover such that the connected components of the Albanese fibres are smooth hence irreducible. Hence, one cannot construct a counterexample to $SC(_)$ by ramifying along these connected components as in subsection 5.2. It is not clear whether $SSC(_)$ is satisfied.

REFERENCES

- [AMRT75] A. Ash, D. Mumford, M. Rapoport, Y.S. Tai *Smooth Compactifications of Locally Symmetric Varieties* Lie Groups: History Frontiers and Applications **4**, Math. Sci. Press, Brookline, MA (1975).
- [BT15] B. Bakker, J. Tsimerman *The Kodaira dimension of complex hyperbolic manifolds with cusps* arXiv:1503.05654v1.
- [BHH87] G. Barthel, F. Hirzebruch, T. Höfer *Geradenkonfigurationen und Algebraische Flächen* Aspekte der Mathematik, Band **D4** (1987), Vieweg.
- [BN06] K. Behrend and B. Noohi, *Uniformization of Deligne-Mumford curves*, J. Reine Angew. Math., 599 (2006), pp. 111–153.
- [BCP06] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput., 24 (1997), 235–265.
- [Cad07] C. Cadman *Using stacks to impose tangency conditions on curves*, Amer. J. Math., 129 (2007), pp. 405–427.
- [Cad16] B. Cadorel *Symmetric differentials on complex hyperbolic manifolds with cusps* arXiv:1606.05470.
- [Cad17] B. Cadorel *Jet differentials on toroidal compactifications of ball quotients* arXiv:1707.07875.

- [CCE15] F. Campana, B. Claudon, P. Eyssidieux *Représentations linéaires des groupes kähleriens : Factorisations et conjecture de Shafarevich linéaire*, doi:10.1112/S0010437X14007751, *Compositio Mathematica*, **151** (2015) 351-376.
- [CK92] F. Catanese, J. Kollár *Trento examples in Classification of irregular varieties, Trento 1990*, Lecture Notes in Mathematics **1515**, Springer (1992), 134–139.
- [Clo93] L. Clozel *Produits dans la cohomologie holomorphe des variétés de Shimura.II. Calculs et applications*. *Journal für die reine und angewandte Mathematik* **444** (1993), 1–15.
- [DeMo86] P. Deligne, G.D. Mostow *Monodromy of hypergeometric functions and non-lattice integral monodromy*, *Pub. Math. I.H.E.S.* **63** (1986), 5–90.
- [DeMo93] P. Deligne, G.D. Mostow *Commensurabilities among Lattices in $PU(1, n)$* , *Annals of Math. Stud.* **132**, Princeton Univ. Press, 1993.
- [Der] M. Deraux, Personal communication (2017).
- [DPP16] M. Deraux, J.R. Parker, J. Paupert *On commensurability classes of non-arithmetic complex hyperbolic lattices* arXiv:1611.00330.
- [DD15] L.F. Di Cerbo, G. Di Cerbo *Effective results for complex hyperbolic manifolds* *J. of the London Math. Soc.* **91** (2015), 89–104.
- [DicSto17] L.F. Di Cerbo, M. Stover *Bielliptic ball quotients compactifications and lattices in $PU(2,1)$ with finitely generated commutator subgroup* *Ann. Inst. Fourier* **67** (2017) 315–328.
- [DiCSto18] L.F. Di Cerbo, M. Stover *Punctured Spheres in complex hyperbolic surfaces and bielliptic ball quotient compactifications* arxiv: 1801.1575v1.
- [DimRam15] M. Dimitrov, D. Ramakrishnan *Arithmetic quotients of the complex ball and a conjecture of Lang* *Documenta Mathematica* **20** (2015) 1185–1205.
- [Eys13] P. Eyssidieux *On the Uniformisation of Compact Kähler orbifolds* *Vietnam J. Math.* (2013), **41**. (2013) 399-407.
- [Eys17] P. Eyssidieux *Orbifold Kähler Groups and the Shafarevich Conjecture for Hirzebruch’s covering surfaces with equal weights*, *Asian J. Math.*, to appear, special volume in the honor of N. Mok, arxiv: 1611.09178.
- [EKPR12] P. Eyssidieux, L. Katzarkov, T. Pantev, and M. Ramachandran, *Linear Shafarevich Conjecture*, *Annals of Math.* **176** (2012), 1545–1581.
- [EySa16] P. Eyssidieux, F. Sala, *Instantons and framed sheaves on Kähler Deligne-Mumford stacks*, arxiv 1404.3504 , *Annales de la Faculté des Sciences de Toulouse (Mathématiques)* to appear.
- [FP06] E. Falbel, R. Parker *The geometry of the Eisenstein-Picard modular group*, *Duke J. Math.* **131** (2006), 249–289.
- [Fal94] G. Faltings *The general case of Lang’s conjecture* in *Symposium in Algebraic Geometry*, Barsotti ed., Academic Press (1994), 175–192.
- [GeSa17] A. Geraschenko, M. Satriano *A “bottom-up” characterization of smooth Deligne-Mumford stacks* *Int. Math. Res. Not. Volume 2017 Issue 21* (2017), 6469–6483.
- [GurSha85] R. V. Gurjar, A. R. Shastri, *Covering spaces of an elliptic surface*, *Compositio Math.* **54** (1985), no. 1, 95–104.
- [HinSil00] M. Hindry, J. Silvermann *Diophantine Geometry, An introduction* *Grad. Texts in Math.* **201** Springer (2000).
- [Hir83] F. Hirzebruch *Arrangements of lines and algebraic surfaces* in *Arithmetic and Geometry-Papers dedicated to I. R. Shafarevich (II)*, *Progress in Mathematics* **36** (1983), 113-140, Birkhäuser.
- [Hol86] R. P. Holzapfel *Chern numbers of algebraic surfaces- Hirzebruch’s examples are Picard Modular Surfaces* *Math. Nachr.* **196** (1996) 255–273.
- [Hol87] R. P. Holzapfel *Geometry and Arithmetic around Euler Partial Differential Equations*, *Mathematics and its applications: East European Series* **11** Reidel Publ. Co. (1987).
- [Hol98] R. P. Holzapfel *Ball and Surface Arithmetics* *Aspects of Mathematic* **E29**, Vieweg (1998).
- [HS96] C. Hummel, V. Schröder *Cusp closing in rank one symmetric spaces* *Inventiones Math.* **123** (1996), 283–307.
- [LKMSS15] L. Ji, V. Kumar Murty, L. Saper, J. Scherk *The fundamental group of Reductive Borel-Serre and Satake compactifications*, *Asian J. Math.* **19** (2015), 465–486.
- [KS15] A. Kasparian, G. Sankaran *Fundamental groups of toroidal compactifications* arxiv 1501.00053v2.
- [Kaz75] D. Kazhdan *Some applications of the Weil representation* *Journal d’Analyse Mathématique* **32** (1975), 235–248.
- [KMR92] V. Kumar Murty, D. Ramakrishnan *The Albanese of unitary Shimura varieties*, in *The zeta functions of Picard modular surfaces*, , R.P. Langlands, D. Ramakrishnan ed., CRM Montreal, (1992) 445–464.
- [Ler10] E. Lerman, *Orbifolds as stacks?*, *Enseign. Math.* (2), **56** (2010), pp. 315–363.

- [Mok11] N. Mok *Projective algebraicity of minimal compactifications of complex hyperbolic space forms of finite volume* in *Perspectives in Analysis, Geometry, and Topology: On the Occasion of the 60th Birthday of Oleg Viro*, Ilia Itenberg, Burglind Jöricke, Mikael Passare eds., Birkhäuser (2011).
- [Mos86] G. D Mostow *Generalized Picard lattices arising from half-integral conditions* Publications Mathématiques de l’IHÉS, **63** (1986), 91–106
- [Nad89] A. Nadel *The Nonexistence of Certain Level Structures on Abelian Varieties over Complex Function Fields* Annals of Mathematics **129**(1) (1989), 161–178.
- [Nap90] T. Napier, *Convexity properties of coverings of smooth projective varieties*, Math. Ann. 286 (1990), no. 1-3, 433–479.
- [Noo04] B. Noohi *Fundamental groups of algebraic stacks* Journal of the Institute of Mathematics of Jussieu **3** (2004), 69-103.
- [Noo05] B. Noohi *Foundations of topological stacks* Arxiv math/0503247 (2005).
- [Och77] T. Ochiai *On holomorphic curves in Algebraic varieties with ample irregularity* Inv. Math **43** (1977), 83–96.
- [Par09] J. Parker *Complex Hyperbolic Lattices in Discrete Groups and Geometric Structures*, Amer. Math. Soc. (2009), 1–42.
- [Sto15] M. Stover *Cusp and b_1 growth for ball quotients and maps onto \mathbb{Z} with finitely generated kernel*, arxiv 1506.0612v2.
- [Rom95] M. Romagny *Group actions on stacks and applications* Michigan Math. J. **53** (2005), 209-236.
- [Tol93] D. Toledo *Projective varieties with non-residually finite fundamental group*, Pub. Math. IHES **77** (1993), 103-119.
- [Ull04] E. Ullmo *Rational points of Shimura varieties* Vol.2004, Issue **76**, (2004) 4109–4125, <https://doi.org/10.1155/S1073792804140415>.
- [Ulu07] A.M. Uludag *Orbifolds and their uniformization in Arithmetic and Geometry around Hypergeometric functions*, R.P. Holzapfel, A.M. Uludag, M. Yoshida eds, Progress in Mathematics **260**, Birkhäuser (2007).
- [Wal84] N. Wallach *L^2 - Automorphic forms and cohomology classes on arithmetic quotients of $SU(p, q)$* Math. Annalen, **266** (1984), 261-278.
- [Zoo02] V. Zoonekynd *Théorème de Van Kampen pour les champs algébriques*, Ann. Math. Blaise Pascal **9** n. 1 (2002), 101–145.

Philippe Eyssidieux

Université de Grenoble-Alpes. Institut Fourier. 100 rue des Maths, BP 74, 38402 Saint Martin d’Hères Cedex, France

philippe.eyssidieux@univ-grenoble-alpes.fr <http://www-fourier.ujf-grenoble.fr/~eyssi/>