

# Resistors in dual networks

Martina Furrer, Norbert Hungerbühler, and Simon Jantschi

Department of Mathematics, ETH Zürich, 8092 Zürich,  
Switzerland

## Abstract

Let  $G$  be a finite plane multigraph and  $G'$  its dual. Each edge  $e$  of  $G$  is interpreted as a resistor of resistance  $R_e$ , and the dual edge  $e'$  is assigned the dual resistance  $R_{e'} := 1/R_e$ . Then the equivalent resistance  $r_e$  over  $e$  and the equivalent resistance  $r_{e'}$  over  $e'$  satisfy  $r_e/R_e + r_{e'}/R_{e'} = 1$ . We provide a graph theoretic proof of this relation by expressing the resistances in terms of sums of weights of spanning trees in  $G$  and  $G'$  respectively.

**Keywords:** dual graphs, electrical networks, equivalent resistance

**AMS classification:** 05C50 (05C05 05C12 15A15 15A18)

## 1 Introduction

The systematic study of electrical resistor networks goes back to the German physicist Gustav Robert Kirchhoff in the middle of the 19th century. In particular, Kirchhoff's two circuit laws and Ohm's law allow to fully describe the electric current and potential in a given static network of resistors and voltage sources. In the course of his investigations, Kirchhoff discovered the Matrix Tree Theorem, which states that the number of spanning trees in a graph  $G$  is equal to any cofactor of the Laplacian matrix of  $G$ . Surprisingly, this purely graph theoretic fact has a deep connection to the physical question of the equivalent resistance between two vertices of an electric network.

In the simplest situation, a finite simple graph  $G$  can be interpreted as an electrical network by considering each edge as a resistor of 1 Ohm. Then, one of Kirchhoff's results states that the equivalent resistance over an edge  $e$  in this graph is given by the quotient of the number of spanning trees containing the edge  $e$  divided by the total number of spanning trees in  $G$ .

More generally, we may consider a finite multigraph  $G$  and assign to each edge  $e$  of  $G$  a weight  $R_e > 0$  interpreted as resistance of  $e$ . Also in this case, the equivalent resistance between two vertices can be expressed in terms of sums of weights of spanning trees (see Section 2).

Consider a cube as a graph with unit resistance on each edge and the dual polyhedron, the octahedron, in the same way. The equivalent resistance over an

edge of the cube turns out to be  $7/12$  Ohm, the equivalent resistance over an edge of the octahedron is  $5/12$  Ohm. Observe, that these values add up to 1! The same phenomenon occurs for the dodecahedron with equivalent resistance of  $19/30$  Ohm over each edge and the dual graph, the icosahedron, with  $11/30$  Ohm, or the rhombic dodecahedron with equivalent resistance of  $13/24$  Ohm over each edge and the dual graph, the cuboctahedron, with  $11/24$  Ohm. This is not just a coincidence: Suppose that a planar graph and its dual are both interpreted as electrical networks with unit resistance for all edges. Now, if  $r_e$  is the equivalent resistance over an edge  $e$  and  $r'_e$  is the equivalent resistance over the dual edge  $e'$ , then  $r_e + r'_e = 1$  (see [1, Exercise 7, Section 9.5], [6, Theorem 2.3]). The aim of this article is to generalize this formula to plane networks with arbitrary resistors (see Theorem 5) and to give a graph theoretic proof.

## 2 Preliminaries

Let  $G$  be a finite connected graph with  $n \geq 3$  vertices and without loops. Multiple edges are allowed. Each edge  $e$  is considered as a resistor of resistance  $R_e > 0$ . Then, we consider the weighted Laplace matrix  $L = (\ell_{ij})$  of  $G$  defined as

$$\ell_{ij} := \begin{cases} -\sum \frac{1}{R_e} & \text{where the sum runs over all edges } e \\ & \text{between the vertices } i \text{ and } j^1, \\ a_{ii} & \text{if } i = j \end{cases}$$

where the diagonal values  $a_{ii}$  are chosen such that the sum of all rows of  $L$  vanishes. The weight of a subgraph  $H$  of  $G$  is defined as

$$\Pi(H) := \prod_{e \text{ an edge of } H} \frac{1}{R_e}.$$

Then we recall the following:

**Proposition 1.** (i) *The value of each cofactor  $L_{ij}$  of  $L$  is the sum of the weights of all spanning trees of  $G$ .*

(ii) *If  $L_{ij,ij}$  denotes the determinant of the matrix  $L$  with rows  $i, j$  and columns  $i, j$  deleted, then the quotient  $L_{ij,ij}/R_e$  equals the sum of the weights of all spanning trees of  $G$  which contain the edge  $e$  between the vertices  $i$  and  $j$ .*

*Proof.* The first part of the proposition follows directly from a general version of the Matrix Tree Theorem (see, e.g., [5, Theorem VI.27]).

For the second part we proceed as follows: Let  $S(G)$  denote the set of all spanning trees of  $G$ . Observe, that by using the edge  $e$  between the vertices  $i$  and  $j$  we can split up the sum of the weights of all spanning trees of  $G$  as follows:

$$\sum_{T \in S(G)} \Pi(T) = \sum_{\substack{T \in S(G) \\ e \in T}} \Pi(T) + \sum_{\substack{T \in S(G) \\ e \notin T}} \Pi(T). \quad (1)$$

---

<sup>1</sup>By convention, an empty sum is 0.

Furthermore, the second sum on the right-hand side of (1) corresponds to the sum of the weights of all spanning trees of  $G - e$ , i.e.  $G$  with edge  $e$  removed. Using the first part of the proposition, we get from (1) that

$$\sum_{\substack{T \in \mathcal{S}(G) \\ e \in T}} \Pi(T) = L_{ii} - L_{ii}^e,$$

where  $L^e = (\ell_{hk}^e)$  is the weighted Laplace matrix of  $G - e$ . We have

$$\ell_{ij}^e = \ell_{ji}^e = \ell_{ij} + \frac{1}{R_e}, \quad \ell_{ii}^e = \ell_{ii} - \frac{1}{R_e}, \quad \ell_{jj}^e = \ell_{jj} - \frac{1}{R_e}$$

and  $\ell_{hk}^e = \ell_{hk}$  for all other  $h, k$ . The term  $L_{ii} - L_{ii}^e$  can be computed using Laplace's cofactor expansion. Expanding both  $L_{ii}$  and  $L_{ii}^e$  along the  $j$ -th row yields

$$\sum_{\substack{T \in \mathcal{S}(G) \\ e \in T}} \Pi(T) = \sum_{k \neq i} \ell_{jk} (L_{ii})_{jk} - \sum_{k \neq i} \ell_{jk}^e (L_{ii}^e)_{jk}.$$

Since the cofactors  $(L_{ii})_{jk}$  and  $(L_{ii}^e)_{jk}$  are equal for all  $k$  we are left with

$$\sum_{\substack{T \in \mathcal{S}(G) \\ e \in T}} \Pi(T) = (\ell_{jj} - \ell_{jj}^e) (L_{ii})_{jj} = \frac{L_{ij,ij}}{R_e}. \quad \square$$

**Remark.** The second part of Proposition 1 follows also quite easily from the All Minors Matrix Tree Theorem (see [3]).

The connection to the equivalent resistance is given by the following

**Proposition 2.** *The equivalent resistance  $r_e$  over the edge  $e$  connecting the vertices  $i$  and  $j$  is given by*

$$r_e = \frac{L_{ij,ij}}{L_{11}}. \quad (2)$$

*Remark.* Recall, that by Proposition 1(i) the denominator in (2) can be replaced by any other cofactor  $L_{hk}$ .

*Proof.* Observe, that the Laplace matrix of the weighted multigraph  $G$  corresponds to the Laplace matrix of a weighted simple graph  $H$  where the multiple edges  $e_1, \dots, e_k$  between each two vertices  $i$  and  $j$  of  $G$  are collapsed to a single edge  $e$  with weight

$$R_e = \frac{1}{\frac{1}{R_{e_1}} + \dots + \frac{1}{R_{e_k}}}.$$

However this value corresponds exactly to the equivalent resistance of the parallel resistors  $R_{e_1}, \dots, R_{e_k}$ . Thus, the equivalent resistance over the vertices  $i$  and  $j$  in  $G$  equals the equivalent resistance over the vertices  $i$  and  $j$  in  $H$  and the claim follows from [2, Lemma 2] and Proposition 1.  $\square$

From now on we assume that is  $G$  a finite planar multigraph with dual graph  $G'$ . Recall that in general  $G'$  depends on the embedding of  $G$  in the plane.

**Definition 3.** Let  $G'$  be the dual of a planar embedded multigraph  $G$ , and let  $G$  be interpreted as an electrical network by associating to each edge  $e$  a resistance  $R_e > 0$ . For each edge  $e$  of  $G$ , we define the electrical resistance  $R_{e'}$  of the dual edge  $e'$  to be the conductance of  $e$ , i.e.  $R_{e'} := 1/R_e$ . Then,  $G'$  equipped with these resistances is called the *dual electrical network* of  $G$ .

Observe, that the Laplace matrix  $L' = (\ell'_{ij})$  of the dual electrical network  $G'$  is given by

$$\ell'_{ij} := \begin{cases} -\sum \frac{1}{R_e} = -\sum R_e & \text{where the sum runs over all edges } e' \\ & \text{between the vertices } i \text{ and } j \text{ of } G', \\ a_{ii} & \text{if } i = j \end{cases}$$

where the diagonal values  $a_{ii}$  are chosen such that the sum of all rows of  $L'$  vanishes.

Similarly the weight of a subgraph  $H'$  of  $G'$  is

$$\Pi(H') := \prod_{e' \text{ an edge of } H'} \frac{1}{R_{e'}} = \prod_{e' \text{ an edge of } H'} R_e.$$

Then we have:

**Proposition 4.** (i) *The value of an arbitrary cofactor  $L'_{ij}$  of  $L'$  is equal to the sum of the weights of all spanning trees in  $G'$  and there holds:*

$$L'_{ij} = L_{ij}\Pi(G') \quad (3)$$

where  $\Pi(G') = \prod R_k$  is the total weight of  $G'$ .

(ii) *The product  $R_e L'_{ij,ij}$  equals the sum of the weights of the spanning trees in  $G'$  which contain the dual edge  $e'$  of edge  $e$ .*

*Proof.* We only have to show equation (3), since all other statements follow from Proposition 1. Let  $S(G)$  and  $S(G')$  denote the set of all spanning trees of  $G$  and its dual  $G'$  respectively. Furthermore, let  $\Psi$  denote the canonical bijection from  $S(G')$  to  $S(G)$  given by  $\Psi(T') = \{e \in G \mid e' \in G' - T'\}$ . Observe, that the weight of a spanning tree  $T'$  of  $G'$  can be expressed as

$$\Pi(T') = \frac{\Pi(G')}{\Pi(G' - T')}. \quad (4)$$

Using the bijection  $\Psi$  and Definition 3 in (4), namely the fact, that the electrical resistance of an edge in  $G'$  is equal to the conductance of the dual edge in  $G$ , we get

$$\Pi(T') = \Pi(G')\Pi(\Psi(T')). \quad (5)$$

The characterization of  $L_{ij}$  and  $L'_{ij}$  as the sum of the weights of all spanning trees of  $G$  and  $G'$  respectively leads, together with (5), to

$$\begin{aligned} L'_{ij} &= \sum_{T' \in S(G')} \Pi(T') = \sum_{T' \in S(G')} \Pi(G')\Pi(\Psi(T')) = \\ &= \Pi(G') \sum_{T \in S(G)} \Pi(T) = \Pi(G')L_{ij}, \end{aligned}$$

where we have used the bijectivity of  $\Psi$  in the penultimate equality. This completes the proof.  $\square$

### 3 The sum formula in dual networks

The main result is now the following:

**Theorem 5.** *Let  $R_e$  be the resistance of an edge  $e$  and  $R_{e'} = 1/R_e$  the resistance of the dual edge  $e'$  in the dual electrical network. Let  $r_e$  denote the equivalent resistance over edge  $e$  and  $r_{e'}$  denote the equivalent resistance over edge  $e'$ . Then*

$$\frac{r_e}{R_e} + \frac{r_{e'}}{R_{e'}} = 1.$$

For a proof of this formula based upon physical arguments see [4]. Here, we provide a purely graph theoretic proof.

*Proof.* Let  $e$  be an edge between the vertices  $i$  and  $j$  in  $G$ , and we may assume that the vertices are numbered such that  $e'$  also runs between the vertices  $i$  and  $j$  in  $G'$ . In a first step, we are going to derive a new expression for  $L'_{ij,ij}$ . Let  $S(G)$  and  $S(G')$  denote the set of all spanning trees of  $G$  and its dual  $G'$  respectively, and let  $\Psi$  be the canonical bijection from  $S(G')$  to  $S(G)$  as defined in the proof of Proposition 4. By part ii of Proposition 4 we have

$$L'_{ij,ij} = \frac{1}{R_e} \sum_{\substack{T' \in S(G') \\ e' \in T'}} \Pi(T'). \quad (6)$$

The identity (5) and the fact, that  $\Psi(T')$  does not contain the edge  $e$  if  $T'$  contains the dual edge  $e'$  allow to rewrite the right-hand side of (6) as follows

$$\frac{1}{R_e} \sum_{\substack{T' \in S(G') \\ e' \in T'}} \Pi(T') = \frac{\Pi(G')}{R_e} \sum_{\substack{T \in S(G) \\ e \notin T}} \Pi(T). \quad (7)$$

Furthermore, the sum on the right-hand side of (7) can be expressed as the difference between the sum of the weights of all spanning trees of  $G$  and the sum of the weights of the spanning trees, that contain the edge  $e$ . Therefore, it holds that

$$L'_{ij,ij} = \frac{\Pi(G')}{R_e} \left( \sum_{T \in S(G)} \Pi(T) - \sum_{\substack{T \in S(G) \\ e \in T}} \Pi(T) \right). \quad (8)$$

Using Proposition 1 in (8) yields the following identity:

$$L'_{ij,ij} = \frac{\Pi(G')}{R_e} \left( L_{ii} - \frac{L_{ij,ij}}{R_e} \right). \quad (9)$$

Now, in a second step, it follows from Proposition 2 and Definition 3, that


$$\frac{r_e}{R_e} + \frac{r_{e'}}{R_{e'}} = \frac{L_{ij,ij}}{L_{ii}R_e} + R_e \frac{L'_{ij,ij}}{L'_{ii}}. \quad (10)$$

Using the first part of Proposition 4 and (9), we can rewrite the right-hand side of (10) and simplify the resulting expression to arrive at

$$\frac{r_e}{R_e} + \frac{r_{e'}}{R_{e'}} = \frac{L_{ij,ij}}{L_{ii}R_e} + R_e \frac{\frac{\Pi(G')}{R_e} \left( L_{ii} - \frac{L_{ij,ij}}{R_e} \right)}{L_{ii}\Pi(G')} = 1,$$

as claimed.  $\square$

**Example 6.** Let us consider the following electrical network:



The corresponding Laplace matrix  $L$  is

$$L = \begin{pmatrix} \frac{1}{R_1} + \frac{1}{R_2} & -\frac{1}{R_2} & 0 & -\frac{1}{R_1} \\ -\frac{1}{R_2} & \frac{1}{R_2} + \frac{1}{R_3} & -\frac{1}{R_3} & 0 \\ 0 & -\frac{1}{R_3} & \frac{1}{R_3} + \frac{1}{R_4} + \frac{1}{R_5} & -\frac{1}{R_4} - \frac{1}{R_5} \\ -\frac{1}{R_1} & 0 & -\frac{1}{R_4} - \frac{1}{R_5} & \frac{1}{R_1} + \frac{1}{R_4} + \frac{1}{R_5} \end{pmatrix}$$

The cofactor


$$L_{11} = \frac{1}{R_1 R_2 R_3 R_4 R_5} (R_4(R_1 + R_2 + R_3) + R_5(R_1 + R_2 + R_3 + R_4))$$

corresponds indeed to the total weight of all spanning trees of  $G$  as one easily checks directly. For the edge  $e$  between the vertices 3 and 4 with resistance  $R_4$ , we get

$$L_{34,34} = \frac{1}{R_1 R_2} + \frac{1}{R_1 R_3} + \frac{1}{R_2 R_3}$$

which, divided by  $R_4$ , gives the sum of the weights of the trees which contain  $e$ , as stated in Proposition 1.

The dual network looks as follows:



and the corresponding Laplace matrix is

$$L' = \begin{pmatrix} R_1 + R_2 + R_3 + R_4 & -R_4 & -(R_1 + R_2 + R_3) \\ -R_4 & R_4 + R_5 & -R_5 \\ -(R_1 + R_2 + R_3) & -R_5 & R_1 + R_2 + R_3 + R_5 \end{pmatrix}.$$

The cofactor

$$L'_{11} = R_4(R_1 + R_2 + R_3) + R_5(R_1 + R_2 + R_3 + R_4)$$

is the total weight of the spanning trees of  $G'$ . And indeed, we have

$$L'_{11} = L_{11}R_1R_2R_3R_4R_5$$

as predicted by Proposition 4. Furthermore, we get

$$L'_{12,12} = R_1 + R_2 + R_3 + R_5$$

which gives according to Proposition 1, after multiplication by  $R_4$ , the total weight of the trees in  $G'$  which contain the dual edge  $e'$ .

Now, the equivalent resistances over edge  $e$  and  $e'$  respectively are, according to Proposition 2,

$$r_4 = \frac{L_{34,34}}{L_{11}} = \frac{R_4R_5(R_1 + R_2 + R_3)}{R_4(R_1 + R_2 + R_3) + R_5(R_1 + R_2 + R_3 + R_4)}$$

$$r'_4 = \frac{L'_{12,12}}{L'_{11}} = \frac{R_1 + R_2 + R_3 + R_5}{R_4(R_1 + R_2 + R_3) + R_5(R_1 + R_2 + R_3 + R_4)}$$

and finally indeed, with  $R'_4 = 1/R_4$ ,

$$\frac{r_4}{R_4} + \frac{r'_4}{R'_4} = 1.$$

## References

- [1] Ravindra B. Bapat. *Graphs and matrices*. Universitext. Springer, London; Hindustan Book Agency, New Delhi, second edition, 2014.
- [2] Ravindra B. Bapat and Somit Gupta. Resistance distance in wheels and fans. *Indian J. Pure Appl. Math.*, 41(1):1–13, 2010.
- [3] Seth Chaiken. A combinatorial proof of the all minors matrix tree theorem. *SIAM J. Algebraic Discrete Methods*, 3(3):319–329, 1982.
- [4] Martina Furrer. Widerstandssumme in dualen Netzwerken. Mentoriere Arbeit, 2017. ETH Zürich.
- [5] William T. Tutte. *Graph theory*, volume 21 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2001. With a foreword by Crispin St. J. A. Nash-Williams, Reprint of the 1984 original.
- [6] Yu Jun Yang. An identity on resistance distances. In *Material and Manufacturing Technology IV*, volume 748 of *Advanced Materials Research*, pages 1024–1027. Trans Tech Publications, 10, 2013.