POLES OF THE COMPLEX ZETA FUNCTION OF A PLANE CURVE

GUILLEM BLANCO

ABSTRACT. We study the poles and residues of the complex zeta function f^s of a plane curve. We prove that most non-rupture divisors do not contribute to poles of f^s or roots of the Bernstein-Sato polynomial $b_f(s)$ of f. For plane branches we give an optimal set of candidates for the poles of f^s from the rupture divisors and the characteristic sequence of f. We prove that for generic plane branches f_{gen} all the candidates are poles of f_{gen}^s . As a consequence, we prove Yano's conjecture for any number of characteristic exponents if the eigenvalues of the monodromy of f are different.

1. INTRODUCTION

Let k be either $\mathbb R$ or $\mathbb C$ and let $\varphi(x) \in C_c^{\infty}(k^n)$ be an infinitely many times differentiable function with compact support. Define the archimedean zeta function f^s of a non-constant polynomial $f(x) \in k[x_1, \ldots, x_n]$ as the distribution

(1.1)
$$
\langle f^s, \varphi \rangle = \int_{k^n} |f(x)|^{\delta s} \varphi(x) dx,
$$

for $s \in \mathbb{C}$, Re $(s) > 0$, where $\delta = 1$ if $k = \mathbb{R}$ and $\delta = 2$ if $k = \mathbb{C}$. In the 1954 edition of the International Congress of Mathematicians, I. M. Gel'fand [\[31\]](#page-33-0) posed the following problem: first, determine whether f^s is a meromorphic function of s with poles forming several arithmetic progressions; second, study the residues at those poles. The problem is solved for some specific polynomials having simple singularities in the book of Gel'fand and Shilov $[32]$, by regularizing the integral in Equation (1.1) . It is not after Hironaka's resolution of singularities [\[36,](#page-33-2) [37\]](#page-33-3), that Bernstein and S. I. Gel'fand [\[11\]](#page-32-0), and independently Atiyah [\[3\]](#page-32-1), give a positive answer to Gel'fand's first question. Both results use resolution of singularities to reduce the problem to the monomial case, already settled in [\[32\]](#page-33-1), and give a sequence of candidates poles for f^s from the resolution data.

A different approach to the same problem is considered by Bernstein [\[9,](#page-32-2) [10\]](#page-32-3), who develops the theory of D-modules and proves the existence of the Bernstein-Sato polynomial $b_f(s)$ of f and its functional equation,

(1.2)
$$
P(s) \cdot f^{s+1} = b_f(s) f^s,
$$

with $P(s) \in D[s]$ being a differential operator. The existence of the Bernstein-Sato polynomial in the local case is due to Björk [\[12\]](#page-32-4). The global $b_f(s)$ is equal to the least common multiple of all the local Bernstein-Sato polynomials $b_{f,p}(s), p \in kⁿ$, see [\[54\]](#page-34-0). The Bernstein-Sato polynomial coincides with the b-function in the theory of prehomogeneous vectors spaces developed by Sato [\[64,](#page-34-1) [63\]](#page-34-2) in the 1960s, hence the name. The rationality of the roots of the Bernstein-Sato polynomial is established by Malgrange [\[52\]](#page-34-3) for isolated singularities and by Kashiwara [\[42\]](#page-33-4) in general, using resolution of singularities. One

The author is supported by Spanish Ministerio de Economía y Competitividad MTM2015-69135-P and Generalitat de Catalunya 2017SGR-932 projects.

verifies, using the functional equation in Equation [\(1.2\)](#page-0-1) and integration by parts, that the poles of f^s are among the rationals $s = \alpha - k$, with $b_f(\alpha) = 0$ and $k \in \mathbb{Z}_{\geq 0}$. Loeser [\[49\]](#page-34-4) shows the equality between both sets for reduced plane curves and isolated quasihomogeneous singularities.

In this paper we examine the original questions of Gel'fand and we use resolution of singularities to study the possible poles and the residues of the complex zeta function of general plane curves, generalizing the ideas and results of Lichtin in $[47, 48]$ $[47, 48]$ $[47, 48]$. The main results of this work are the following:

- For any candidate pole σ of f^s , we give a formula for its residue expressed as an integral along the exceptional divisor associated to σ , see Proposition [5.4.](#page-18-0)
- In Theorem [5.10,](#page-23-0) we prove that most non-rupture divisors do not contribute to the poles of f^s , and consequently to the roots of $b_f(s)$.

This result answers, for reduced plane curves, a question raised by Kollár $[45]$ on which exceptional divisors contribute to roots of the Bernstein-Sato polynomial. It is already well-known that, for plane curves, non-rupture divisors do not contribute to topological invariants such as the eigenvalues of the monodromy $[1, 56]$ $[1, 56]$, the jumping numbers $[65]$, or the poles of Igusa's local zeta functions [\[50\]](#page-34-10). For irreducible plane curves, we use Teissier's monomial curve $[68]$ associated to the semigroup of f to refine our previous results:

- In Theorem [6.7,](#page-28-0) we obtain an optimal set of candidates for the poles of f^s in terms of the rupture divisors, the characteristic sequence, and the semigroup of f.
- From this, in Theorem [6.8,](#page-30-0) we prove that if f_{gen} is generic among all plane branches with fixed characteristic sequence (in the sense that the coefficients of a μ -constant deformation are generic), all the candidates are indeed poles of f_{gen}^s .
- As a consequence, in Corollary [6.9,](#page-32-6) we prove a conjecture of Yano [\[73\]](#page-35-1) about the b-exponents of a plane branch for any number of characteristic exponents under the assumption that the eigenvalues of the monodromy of f are pairwise different.

Both the roots of $b_f(s)$ and the poles of f^s are related to the local geometry of f. By a series of results of Malgrange [\[52,](#page-34-3) [53\]](#page-34-11), first in the isolated singularity case and later in general, for every root α of $b_f(s)$, the value $\exp(2\pi i\alpha)$ is an eigenvalue of the local monodromy at some point of $f^{-1}(0)$ and every eigenvalue is obtained in this way. For an isolated singularity these results imply that the degree of $b_f(s)$ is at most the Milnor number. Therefore, in general, every pole σ of the archimedean zeta function f^s has that $\exp(2\pi i\sigma)$ is an eigenvalue of the monodromy at some point of $f^{-1}(0)$. Barlet [\[4\]](#page-32-7) proves that all eigenvalues are obtained in this way if $k = \mathbb{C}$. Moreover, Barlet [\[6\]](#page-32-8) shows that if the monodromy action in the q -th cohomology group of the Milnor fiber at some point of $f^{-1}(0)$ has an eigenvalue $\exp(-2\pi i\alpha)$ with a $k \times k$ Jordan block, then f^s has a pole at $-q-\alpha, \alpha \in [0,1)$, of order at least k.

A non-archimedean version of the zeta function of a polynomial f can be defined by a p-adic version of the integral in Equation (1.1) . These zeta functions over the p-adic fields are usually called Igusa's local zeta functions. They were first studied by Igusa in [\[38,](#page-33-5) [39\]](#page-33-6), where he proves that they are rational functions. The theory of Igusa's zeta functions is vast and has many connection with Singularity and Number Theory. For instance, Igusa's local zeta functions are related to the number of solutions of f modulo p^m . In [\[40\]](#page-33-7), Igusa conjectures that if s is a pole of a p-adic zeta function of f, Re(s) is a root of $b_f(s)$. This conjecture is proved by Loeser for plane curves [\[50\]](#page-34-10) and for many non-degenerate singularities [\[51\]](#page-34-12). There is a version of this conjecture for the monodromy instead of the Bernstein-Sato polynomial. By the results of Malgrange, the former implies the later. Little more is known in this case, see the work of Artal Bartolo, Cassou-Noguès, Luengo and Melle Hernández $[7]$ on quasi-ordinary singularities, Bories and Veys $[14]$ on non-degenerate surface singularities, and Budur, Mustată and Teitler $[20]$ on hyperplane arrangements. These two conjectures imply the same sort of conjectures for the topological zeta function introduced by Denef and Loeser [\[29\]](#page-33-9), for which more cases are known. The reader is referred to the classical reports of Denef [\[28\]](#page-33-10) and Igusa [\[41\]](#page-33-11), and the references therein for the concrete definitions, results and conjectures in the theory of Igusa's zeta

There are many singularity invariants related to the poles of f^s and the roots of $b_f(s)$. The log-canonical threshold of f [\[45\]](#page-34-7) is minus the largest pole of f^s and the largest root of $b_f(s)$ and it sets the maximal region of holomorphy of the integral in Equation [\(1.1\)](#page-0-0). For isolated singularities, it coincides with the complex singularity index, a concept that dates back to Arnold [\[2\]](#page-32-11). Using resolution of singularities, one defines the multiplier ideals and the associated jumping numbers [\[30\]](#page-33-12). The log-canonical threshold appears as the smallest jumping number. The opposites in sign to the jumping numbers in $(0, 1]$ are always roots of the Bernstein-Sato polynomial $[30]$, see also $[21]$. The spectral numbers $[66, 67]$ $[66, 67]$ are a set of logarithms of the eigenvalues of the monodromy constructed using the mixed Hodge structure of the cohomology of the Milnor fiber. In the isolated singularity case the spectral and jumping numbers in $(0, 1]$ coincide, for non-isolated singularities see [\[19\]](#page-33-14). Furthermore, for isolated singularities the opposites in sign to the spectral numbers are poles of f^s , [\[49\]](#page-34-4).

functions. A survey of Meuser [\[55\]](#page-34-13) includes the more recent developments.

There exist algorithms to compute the Bernstein-Sato polynomial $b_f(s)$ for arbitrary polynomials f , see [\[58,](#page-34-16) [57,](#page-34-17) [46\]](#page-34-18). However, even computationally, determining the roots of $b_f(s)$ is a hard problem. For bounds on the roots and their multiplicities see the results of Saito $[61]$. For candidates for the roots, we refer to the result of Kashiwara $[42]$ and a refinement given by Lichtin [\[48\]](#page-34-6). Very few formulas for the Bernstein-Sato polynomial are known. If f is smooth at $p \in k^n$, then $b_{f,p}(s) = s + 1$. Hence, if f is everywhere smooth, then $b_f(s) = s + 1$ and the converse is also true, [\[16\]](#page-33-15). The monomial case is obtained by a straightforward computation. Yano worked out many interesting examples in [\[72\]](#page-35-2). For isolated quasi-homogeneous singularities, see [\[52,](#page-34-3) [72\]](#page-35-2). For isolated semi-quasihomogeneous singularities, see $[60, 15]$ $[60, 15]$. The case of hyperplane arrangements has been studied by Walther [\[71\]](#page-35-3) and Saito [\[62\]](#page-34-21).

The case of plane curves has attracted a lot of attention, see [\[73,](#page-35-1) [43,](#page-34-22) [44,](#page-34-23) [23,](#page-33-16) [24,](#page-33-17) [25,](#page-33-18) [35,](#page-33-19) [17\]](#page-33-20). It is well-known that the roots of the Bernstein-Sato polynomial, hence the poles of f^s , can change within a deformation with constant Milnor number of a plane curve, see, for instance, the examples in [\[43,](#page-34-22) [44\]](#page-34-23). This contrasts with the fact that the spectral numbers and, therefore, the monodromy eigenvalues and the jumping numbers, remain constant in a deformation with constant Milnor number, [\[69\]](#page-35-4). For irreducible plane curve singularities, there is a conjecture of Yano [\[73\]](#page-35-1) asserting that if the plane branch is generic among those with fixed characteristic sequence, the so-called b-exponents, see definition in Section [2.3,](#page-7-0) are constant and depend only on the characteristic sequence. In addition, Yano conjectures a closed formula for the b-exponents. This conjecture has been verified by

Cassou-Noguès $[25]$ for a single characteristic exponent. Recently, Artal Bartolo, Cassou-Noguès, Luengo and Melle Hernández $[8]$ proved Yano's conjecture for plane branches with two characteristic exponents and monodromy with different eigenvalues.

This paper is organized as follows. In Section [2,](#page-3-0) we first review the classical results on the regularization and analytic continuation of f^s for a general complex polynomial f. We then focus on the connection with the Bernstein-Sato polynomial and its basic properties, specially for isolated singularities. Yano's conjecture is presented at the end. In Section [3,](#page-7-1) we present all the relevant results about the characteristic sequence and the semigroup of a plane branch. Then, we introduce Teissier's monomial curve and its deformations. Throughout Section [4](#page-11-0) we review the results on resolution of singularities of plane curves that will be needed in the following sections. Section [5](#page-15-0) deals with the poles and the formulas for the residues along an exceptional divisor of the complex zeta function of a general plane curve, specially the case of non-rupture exceptional divisors. In the last section, we focus on irreducible plane curves and the optimal sequence of candidates coming from their rupture divisors. Finally, we work on the generic case and we prove Yano's conjecture.

Acknowledgments. The author would like to thank his advisors, Maria Alberich-Carramiñana and Josep Alvarez Montaner, for the fruitful discussions, the helpful comments and suggestions, and the constant support during the development of this work.

2. Analytic continuation of complex powers

In this section, we will review the basic results on regularization of complex powers appearing in the book of Gel'fand and Shilov [\[32\]](#page-33-1). We will see how resolution of singularities is used to construct the analytic continuation of the complex zeta function of an arbitrary polynomial $f \in \mathbb{C}[z_1,\ldots,z_n]$. The Bernstein-Sato polynomial [\[10\]](#page-32-3) is presented next and it is used to construct the analytic continuation of f^s in a different way. In order to state Yano's conjecture [\[73\]](#page-35-1), we introduce the equivalent definition of the Bernstein-Sato polynomial for isolated singularities by Malgrange [\[52\]](#page-34-3) in terms of the Brieskorn lattice.

2.1. Regularization of complex powers. We will take the set of test functions of complex variable as the set of smooth, compactly supported functions $\varphi : \mathbb{C}^n \longrightarrow \mathbb{C}$. The space of such functions is denoted by $C_c^{\infty}(\mathbb{C}^n)$. Alternatively, we can consider the larger space of test functions consisting of Schwartz functions. From the analytic continuation principle one deduces that there are no holomorphic compactly supported functions. Therefore, any $\varphi \in C_c^{\infty}(\mathbb{C}^n)$ has an holomorphic an antiholomorphic part, i.e. $\varphi = \varphi(z, \bar{z})$.

Let $f(z) \in \mathbb{C}[z_1,\ldots,z_n]$ be a non-constant polynomial. We define a parametric family of distributions of complex variable $f^s: C_c^{\infty}(\mathbb{C}^n) \longrightarrow \mathbb{C}$ given by

(2.1)
$$
\langle f^s, \varphi \rangle := \int_{\mathbb{C}^n} \varphi(z, \bar{z}) |f(z)|^{2s} dz d\bar{z},
$$

which is well-defined for any $s \in \mathbb{C}$ with $\text{Re}(s) > 0$. The dependence of f^s on the parameter s is holomorphic as we can differentiate under the integral symbol to obtain another well-defined distribution, namely

$$
\frac{d}{ds}\langle f^s, \varphi \rangle = \int_{\mathbb{C}^n} \varphi(z, \bar{z}) |f(z)|^{2s} \log |f(z)|^2 dz d\bar{z} = \left\langle \frac{df^s}{ds}, \varphi \right\rangle, \quad \text{Re}(s) > 0.
$$

The distribution f^s or the function $\langle f^s, \varphi \rangle$ are usually called the *complex zeta function* of f. This name goes back to I. M. Gel'fand $[31]$. In $[32]$, it is shown how one can obtain the analytic continuation of $\langle f^s, \varphi \rangle$ by *regularizing* the integral in Equation [\(2.1\)](#page-3-1), for some classes of polynomials. The concept of regularization is better understood after the following example.

If one takes the function $z^{-3/2}$, in general, its integral $\langle z^{-3/2}, \varphi \rangle$ against a test function $\varphi(z,\bar{z})$ will diverge. However, if $\varphi(z,\bar{z})$ vanishes at zero, the integral converges. Any distribution whose action on the elements of $C_c^{\infty}(\mathbb{C})$ vanishing at zero coincides with the action of $z^{-3/2}$ is a regularization of $z^{-3/2}$. The regularization of a function with algebraic singularities is unique up to functionals concentrated in the zero locus, see [\[32,](#page-33-1) I.1.7]. For a fixed $s \in \mathbb{C}$, the *canonical regularization* of the function z^s , in the sense that it is the most natural, is presented in the following proposition.

Proposition 2.1 ([\[32,](#page-33-1) B1.2], Gel'fand-Shilov regularization). For any $m \in \mathbb{Z}_{\geq 0}$, the regularization of the distribution $z^s: C_c^{\infty}(\mathbb{C}) \longrightarrow \mathbb{C}$ is given by

(2.2)

$$
\langle z^s, \varphi \rangle = \int_{|z| \le 1} \left[\varphi(z, \bar{z}) - \sum_{k+l=0}^{m-1} \varphi^{(k,l)}(0, 0) \frac{z^k \bar{z}^l}{k!l!} \right] |z|^{2s} dz d\bar{z} + \int_{|z| > 1} \varphi(z, \bar{z}) |z|^{2s} dz d\bar{z} - 2\pi i \sum_{k=0}^{m-1} \frac{\varphi^{(k,k)}(0, 0)}{(k!)^2 (s + k + 1)}, \quad \text{Re}(s) > -m - 1,
$$

where $\varphi^{(i,j)} := \partial^{i+j} \varphi / \partial z^i \partial \bar{z}^j$. Hence, z^s has poles at $s = -k-1$ for $k \in \mathbb{Z}_{\geq 0}$ with residues

$$
\operatorname{Res}_{s=-k-1} z^s = -\frac{2\pi i}{(k!)^2} \delta_0^{(k,k)},
$$

where $\delta_0^{(i,j)}$ $\int_0^{(i,j)}$ are the distributional derivatives of the Dirac's delta function defined by $\langle \delta_0^{(i,j)} \rangle$ $\langle u^{(i,j)}_0, \varphi \rangle := (-1)^{i+j} \varphi^{(i,j)}(0,0)$. Furthermore, in the strip $-m-1 < \text{Re}(s) < -m$, Equation [\(2.2\)](#page-4-0) reduces to

$$
\langle z^s, \varphi \rangle = \int_{\mathbb{C}} \left[\varphi(z, \bar{z}) - \sum_{k+l=0}^{m-1} \varphi^{(k,l)}(0, 0) \frac{z^k \bar{z}^l}{k!l!} \right] |z|^{2s} dz d\bar{z}.
$$

For a fixed $\varphi \in C_c^{\infty}(\mathbb{C})$, Proposition [2.1](#page-4-1) gives the meromorphic continuation to the whole complex plane of the holomorphic function of s defined by the integral $\langle z^s, \varphi \rangle$. For any polynomial f , we will talk indistinguishably about the meromorphic continuation or the (canonical) regularization of its complex zeta function f^s .

Remark 2.2. Although in Proposition [2.1](#page-4-1) the test function φ is assumed to be in $C_c^{\infty}(\mathbb{C})$, the proof of the result only uses the fact that φ is infinitely differentiable near 0 and compactly supported. This means that the same result works for a meromorphic φ with poles away from 0 and compact support. In particular, if $\varphi(z, \bar{z}; s) \in C^{\infty}(U)$, where U is a neighborhood of 0, and compactly supported, the poles of $\langle z^s, \varphi(z, \bar{z}; s) \rangle$ will be the negative integers $\mathbb{Z}_{<0}$ together with the poles of $\varphi(z,\bar{z};s)$ in s away from U.

Resolution of the singularities in $f \in \mathbb{C}[z_1,\ldots,z_n]$ is used in [\[11\]](#page-32-0) and [\[3\]](#page-32-1) to reduce the problem of finding the analytic continuation of f^s to the monomial case considered in Proposition [2.1.](#page-4-1) Let $\pi : X' \longrightarrow \mathbb{C}^n$ be a resolution of f with $F_{\pi} := \sum N_i D_i$ the total transform divisor and $K_\pi := \sum k_i E_i$ the relative canonical divisor, and suppose that

 $E := \text{Exc}(\pi) = \sum E_i$ is the exceptional locus. Let $\{U_\alpha\}_{\alpha \in \Lambda}$ be an affine open cover of X'. Take $\{\eta_{\alpha}\}\$ a partition of unity subordinated to the cover $\{U_{\alpha}\}_{{\alpha}\in{\Lambda}}\$. That is, $\eta_{\alpha}\in C^{\infty}(\mathbb{C}^n)$ (not necessarily with compact support), $\sum \eta_{\alpha} \equiv 1$, with only finitely many η_{α} being non-zero at a point of X' and $\text{Supp}(\eta_{\alpha}) \subseteq U_{\alpha}$. Then, with a small abuse of notation,

$$
\langle f^s, \varphi \rangle = \int_{X'} |\pi^* f|^{2s} (\pi^* \varphi) |d\pi|^2
$$

(2.3)

$$
= \sum_{\alpha \in \Lambda} \int_{U_{\alpha}} |z_1|^{2(N_{1,\alpha}s + k_{1,\alpha})} \cdots |z_n|^{2(N_{n,\alpha}s + k_{n,\alpha})} |u_\alpha(z)|^{2s} |v_\alpha(z)|^2 \varphi_\alpha(z, \bar{z}) dz d\bar{z},
$$

where $\varphi_{\alpha} := \eta_{\alpha} \pi^* \varphi$ for each $\alpha \in \Lambda$ and $u_{\alpha}(0), v_{\alpha}(0) \neq 0$. The resolution morphism π being proper implies that both E and $\pi^{-1}(\text{Supp}(\varphi))$ are compact sets. Since the singularities of the integral $\langle f^s, \varphi \rangle$ are produced by the zero set of f, in order to study the poles of f^s , it is enough to consider a finite affine open cover $\{U_{\alpha}\}_{{\alpha}\in {\Lambda}}$ of E consisting of neighborhoods of points $p_{\alpha} \in E$ and such that $\text{Supp}(\varphi) \subseteq \pi(\cup_{\alpha} U_{\alpha}).$

From Equation [\(2.3\)](#page-5-0) and Proposition [2.1,](#page-4-1) we see that each divisor D_i in the support of F_{π} generates a set of candidate poles of f^s , namely

$$
-\frac{k_i+1+\nu}{N_i}, \quad \nu \in \mathbb{Z}_{\geq 0}.
$$

The opposite in sign to the largest pole is the *log-canonical threshold* $lct(f)$ of f and sets the maximal region of holomorphy of $\langle f^s, \varphi \rangle$ for a general $\varphi \in C_c^{\infty}(\mathbb{C}^n)$. This solves Gel'fand's first question in [\[31\]](#page-33-0). However, nothing is said about the residues of f^s at those poles. Moreover, the set of candidates is usually large compared with the set of roots of the Bernstein-Sato polynomial or the actual poles of f^s .

2.2. The Bernstein-Sato polynomial. Let X be a complex manifold of dimension n with \mathcal{O}_X the sheaf of regular functions and \mathcal{D}_X the sheaf of differentials operators. We set $\mathcal{D}_X[s] = \mathcal{D}_X \otimes_{\mathbb{C}} \mathbb{C}[s]$, where s is an indeterminate commuting with all differential operators. Fix $f \in \mathcal{O}_X$ a non-zero regular function on X. The Bernstein-Sato functional equation [\[10\]](#page-32-3) asserts the existence of a differential operator $P(s) \in \mathcal{D}_X[s]$ and a non-zero polynomial $b(s) \in \mathbb{C}[s]$ such that

$$
(2.4) \t\t P(s) \cdot f^{s+1} = b(s)f^s.
$$

Although neither $P(s)$ or $b(s)$ are necessarily unique, all the polynomials $b(s) \in \mathbb{C}[s]$ satisfying Equation [\(2.4\)](#page-5-1) form an ideal. The unique monic generator of this ideal is called the Bernstein-Sato polynomial $b_f(s)$ of f. Everything remains true in the local case, see [\[13\]](#page-32-14). The only two general results about the structure of the roots of the Bernstein-Sato polynomial are the following.

Theorem 2.3 ([\[42\]](#page-33-4), [\[48\]](#page-34-6), Rationality of the roots). Let $f \in \mathcal{O}_X$ be non-constant and let $\pi: X' \longrightarrow X$ be a resolution of f with $F_{\pi} = \sum N_i E_i$ and $K_{\pi} = \sum k_i E_i$ the resolution and relative canonical divisors. Then, the roots of $b_f(s)$ are among the numbers

$$
-\frac{k_i+1+\nu}{N_i}-k,
$$

for $\nu \in \{0, \ldots, N_i-1\}$ and $k \in \mathbb{Z}_{\geq 0}$. Therefore, all the roots are negative rational numbers.

As in the case of the candidate poles of the complex zeta function f^s , the set of can-didates provided by Theorem [2.3](#page-5-2) is usually larger than the set of roots of $b_f(s)$. Indeed, there are many exceptional divisors of the resolution not contributing to actual poles of f^s or roots of $b_f(s)$. Since $s = -1$ is always a root of $b_f(s)$, it is sometimes useful to work with the *reduced* Bernstein-Sato polynomial, $\widetilde{b}_f(s) := b_f(s)/(s+1)$.

Theorem 2.4 ([\[61\]](#page-34-19)). Let $f \in \mathcal{O}_X$ be non-constant. Then, for any root α of $\widetilde{b}_f(s)$, $\alpha \in [-n + \mathrm{lct}(f), -\mathrm{lct}(f)].$ If m_{α} denotes the multiplicity of α , $m_{\alpha} \leq n - \mathrm{lct}(f) - \alpha + 1$.

The bounds in Theorem [2.4](#page-6-0) are optimal for isolated quasi-homogeneous singularities.

The Bernstein-Sato functional equation together with integration by parts can be used to obtain the analytic continuation of f^s in a different way.

Proposition 2.5. The complex zeta function f^s admits a meromorphic continuation to $\mathbb C$ with poles among the rational numbers $\alpha - k$ with $b_f(\alpha) = 0$ and $k \in \mathbb{Z}_{\geq 0}$.

Proof. We can use the functional Equation (2.4) and integration by parts to analytically continue Equation (2.1) in the following way

(2.5)
\n
$$
\langle f^s, \varphi \rangle = \int_{\mathbb{C}^n} \varphi(z, \bar{z}) |f(z)|^{2s} dz d\bar{z} = \frac{1}{b_f^2(s)} \int_{\mathbb{C}^n} \varphi(z, \bar{z}) [P(s) \cdot f^{s+1}(z)] [\overline{P}(s) \cdot f^{s+1}(\bar{z})] dz d\bar{z}
$$
\n
$$
= \frac{1}{b_f^2(s)} \int_{\mathbb{C}^n} \overline{P}^* P^*(s) (\varphi(z, \bar{z})) |f(z)|^{2(s+1)} dz d\bar{z}.
$$

The last term of Equation [\(2.5\)](#page-6-1) defines an analytic function whenever $\text{Re}(s) > -1$, except for possible poles at b_f^{-1} $f_f^{-1}(0)$, and it is equal to $\langle f^s, \varphi \rangle$ in $\text{Re}(s) > 0$. If a differential operator has the form $P(s) = \sum_{\beta} a_{\beta}(s, z) \left(\frac{\partial}{\partial z}\right)^{\beta}$, we have considered

$$
\overline{P}(s) := \sum_{\beta} a_{\beta}(\overline{s}, \overline{z}) \left(\frac{\partial}{\partial \overline{z}}\right)^{\beta}, \quad P^*(s) := \sum_{\beta} (-1)^{|\beta|} \left(\frac{\partial}{\partial z}\right)^{\beta} a_{\beta}(z, s),
$$

the *conjugate* and *adjoint operator* of $P(s)$, respectively. Iterating the process we get

(2.6)
$$
\langle f^s, \varphi \rangle = \frac{\langle f^{s+k+1}, \overline{P}^* P^*(s+k) \cdots \overline{P}^* P^*(s)(\varphi) \rangle}{b_f^2(s) \cdots b_f^2(s+k)}, \quad \text{Re}(s) > -k-1,
$$

and the result follows.

The set of poles of the complex zeta function f^s is known to be exactly the set $\alpha - k$ with $b_f(\alpha) = 0$ and $k \in \mathbb{Z}_{\geq 0}$ for reduced plane curve singularities and isolated quasi-homogeneous singularities, see [\[49,](#page-34-4) Th. 1.9]. Therefore, at least in these cases, the divisors contributing to poles of the complex zeta function f^s are the same divisors that contribute to roots of the Bernstein-Sato polynomial $b_f(s)$. However, even in these cases, it is not straightforward to relate poles of f^s with roots of $b_f(s)$. In general, from Theorem [2.4](#page-6-0) and Proposition [2.5,](#page-6-2) one has that,

Corollary 2.6. Every pole $\sigma \in [-n + \text{lct}(f), -\text{lct}(f)]$ of f^s such that $\sigma + k$ is not a root of $b_f(s)$ for all $k \in \mathbb{Z}_{>0}$ is a root of $b_f(s)$.

2.3. Yano's conjecture. We assume now that f has an isolated singularity at 0 and we work locally around this point. Let $\mu := \dim_{\mathbb{C}} \mathcal{O}_{X,0}/\langle \partial f/\partial z_1,\ldots,\partial f/\partial z_n \rangle$ be the Milnor number of f at 0. The Brieskorn lattice, [\[18\]](#page-33-21) $H''_{f,0} := \Omega_{X,0}^n/df \wedge d\Omega_{X,0}^{n-1}$, of f at 0 has a structure of $\mathbb{C}\lbrace t \rbrace$ -module given by the multiplication by f, moreover, it is a free module of dimension μ and carries a connection ∂_t . Malgrange shows in [\[52\]](#page-34-3) that if $\widetilde{H}''_{f,0} := \sum_{k\geq 0} (\partial_t t)^k H''_{f,0}$ is the saturation of the Brieskorn lattice, then $\widetilde{b}_{f,0}(s)$ is the minimal polynomial of the endomorphism

$$
-\partial_t t : \widetilde{H}_{f,\mathbf{0}}^{\prime\prime}/t\widetilde{H}_{f,\mathbf{0}}^{\prime\prime} \longrightarrow \widetilde{H}_{f,\mathbf{0}}^{\prime\prime}/t\widetilde{H}_{f,\mathbf{0}}^{\prime\prime}.
$$

Therefore, the degree of the Bernstein-Sato polynomial of an isolated singularity is at most $\prod_j (s - \exp(2\pi i \widetilde{\alpha}_j))$ is a divisor of the characteristic polynomial of the monodromy and μ . If $\widetilde{\alpha}_1, \ldots, \widetilde{\alpha}_r, r \leq \mu$ are the roots of $\widetilde{b}_{f,0}$, Malgrange also proves that the polynomial a multiple of the minimal polynomial of the monodromy. Following the terminology of Yano, the *b-exponents* of an isolated singularity f are the μ roots $\alpha_1, \ldots, \alpha_{\mu}$ of the characteristic polynomial of the action of $\partial_t t$ on $\overline{H}''_{f,0}/t\overline{H}''_{f,0}$.

Yano's conjecture [\[73\]](#page-35-1) on the b-exponents of generic irreducible plane curves reads as follows. Let $(n, \beta_1, \ldots, \beta_q)$ be the characteristic sequence of a plane branch with $g \geq 1$ being the number of characteristic exponents. With the same notation as in [\[73\]](#page-35-1) define,

$$
e_0 := n, \quad e_i := \gcd(n, \beta_1, \dots, \beta_i), \quad i = 1, \dots, g,
$$

\n
$$
r_i := \frac{\beta_i + n}{e_i}, \quad R_i := \frac{\beta_i e_{i-1} + \beta_{i-1} (e_{i-2} - e_{i-1}) + \dots + \beta_1 (e_0 - e_1)}{e_i},
$$

\n(2.7)
\n
$$
r'_0 := 2, \quad r'_i := r_{i-1} + \left[\frac{\beta_i - \beta_{i-1}}{e_{i-1}} \right] + 1 = \left[\frac{r_i e_i}{e_{i-1}} \right] + 1,
$$

\n
$$
R'_0 := n, \quad R'_i := R_{i-1} + \beta_i - \beta_{i-1} = \frac{R_i e_i}{e_{i-1}}.
$$

Inspired by A'Campo's formula [\[1\]](#page-32-5) for the eigenvalues of the monodromy, Yano defines the following polynomial with fractional powers in t

(2.8)
$$
R((n, \beta_1, ..., \beta_g), t) := \sum_{i=1}^g t^{\frac{r_i}{R_i}} \frac{1-t}{1-t^{\frac{1}{R_i}}} - \sum_{i=0}^g t^{\frac{r'_i}{R'_i}} \frac{1-t}{1-t^{\frac{1}{R'_i}}} + t,
$$

and proves that $R((n, \beta_1, ..., \beta_n), t)$ has non-negative coefficients. Finally,

Conjecture (Yano, [\[73\]](#page-35-1)). For generic curves among all irreducible plane curves with characteristic sequence $(n, \beta_1, \ldots, \beta_g)$ the b-exponents $\alpha_1, \ldots, \alpha_\mu$ are given by the generating function $R((n, \beta_1, \ldots, \beta_n), t)$. That is,

$$
\sum_{i=1}^{\mu} t^{\alpha_i} = R((n, \beta_1, \ldots, \beta_g), t).
$$

3. The semigroup of a plane branch and its monomial curve

The characteristic sequence of an irreducible plane curve is a complete topological invariant. Two germs of a curve are topologically equivalent, if and only if, they have the same characteristic sequence. Alternatively, one defines the semigroup of a plane branch from its associated valuation. The aim of this section is to first describe all the 3.1. The semigroup of a plane branch. We begin by fixing a germ of a plane branch $f: (\mathbb{C}^2, \mathbf{0}) \longrightarrow (\mathbb{C}, 0)$ with *characteristic sequence* $(n, \beta_1, \ldots, \beta_g), n, \beta_i \in \mathbb{Z}_{>0}$. The characteristic sequence can be obtained from the Puiseux parameterization of f . After an analytic change of variables we can always assume that $n < \beta_1 < \cdots < \beta_q$. Define the integers $e_i := \gcd(e_{i-1}, \beta_i), e_0 := n$, with n being the multiplicity of f. Notice that they satisfy $e_0 > e_1 > \cdots > e_g = 1$ and $e_{i-1} \nmid \beta_i$.

are Zariski's book [\[74\]](#page-35-5) and its appendix [\[68\]](#page-35-0) by Teissier.

We set $n_i := e_{i-1}/e_i$ for $i = 1, \ldots, g$ and, by convention, $\beta_0 := 0$ and $n_0 := 0$. The integers n_1, \ldots, n_q are strictly larger than 1 and we have that $e_{i-1} = n_i n_{i+1} \cdots n_q$ for $i = 1, \ldots, g$. In particular, $n = n_1 \cdots n_g$. The fractions $m_i/n_1 \cdots n_i$, with m_i defined as $m_i := \beta_i/e_i$, are the reduced *characteristic exponents* appearing in the Puiseux series of f. The tuples (m_i, n_i) satisfy $gcd(m_i, n_i) = 1$ and are usually called the *Puiseux pairs*.

Let \mathcal{O}_f be the local ring of f. The Puiseux parameterization of f gives an injection $\mathcal{O}_f \hookrightarrow \mathbb{C}{t}$. We denote the t-adic valuation of \mathcal{O}_f by ν and $\Gamma \subseteq \mathbb{Z}_{\geq 0}$ denotes the associated semigroup

$$
\Gamma := \big\{ \nu(g) \in \mathbb{Z}_{\geq 0} \mid g \in \mathcal{O}_f \setminus \{0\} \big\}.
$$

Since f is irreducible there exists a minimum integer $c \in \mathbb{Z}_{>0}$, the *conductor* of Γ, such that $(t^c) \cdot \mathbb{C}{t} \subseteq \mathcal{O}_f$. As a result, any integer in $[c, \infty)$ must belong to Γ , which implies that $\mathbb{Z}_{\geq 0} \setminus \Gamma$ is finite. Since $\mathbb{Z}_{\geq 0} \setminus \Gamma$ is finite, we can find a minimal generating set $\langle \beta_0, \beta_1, \ldots, \beta_g \rangle$ of Γ , i.e. β_i are the minimal integers such that $\beta_i \notin \langle \beta_0, \beta_1, \ldots, \beta_{i-1} \rangle$, with $\beta_0 < \beta_1 < \cdots < \beta_g$ and $gcd(\beta_0, \beta_1, \ldots, \beta_g) = 1$.

The semigroup generators can be computed from the characteristic sequence in the following way, see $[74, \text{II}.3]$,

$$
(3.1) \qquad \overline{\beta}_i = (n_1 - 1)n_2 \cdots n_{i-1}\beta_1 + (n_2 - 1)n_3 \cdots n_{i-1}\beta_2 + \cdots + (n_{i-1} - 1)\beta_{i-1} + \beta_i,
$$

for $i = 2, \ldots, g$ and with $\overline{\beta}_0 = n, \overline{\beta}_1 = \beta_1$. Recursively this can be expressed as

(3.2)
$$
\overline{\beta}_i = n_{i-1}\overline{\beta}_{i-1} - \beta_{i-1} + \beta_i, \qquad i = 2, \ldots, g.
$$

By Equation [\(3.1\)](#page-8-0), $gcd(e_{i-1}, \beta_i) = e_i$ with $e_0 = \beta_0 = n$ and $e_{i-1} \not\mid \beta_i$. In the same way as before, we define the sequence of integers $\overline{m}_i := \beta_i/e_i$ which will be useful in the sequel. The integers \overline{m}_i , $i = 1, \ldots, g$ can be obtained recursively using Equation [\(3.2\)](#page-8-1), namely

(3.3)
$$
\overline{m}_i = n_i n_{i-1} \overline{m}_{i-1} - n_i m_{i-1} + m_i, \quad i = 2, ..., g,
$$

with $\overline{m}_0 = 1, \overline{m}_1 = m_1$. Note that Equation [\(3.3\)](#page-8-2) implies $gcd(\overline{m}_i, n_i) = 1$ for $i = 1, \ldots, g$. Finally, we define $q_i := (\beta_i - \beta_{i-1})/e_i = m_i - n_i m_{i-1}$ for $i = 1, \ldots, g$. Alternatively, by Equation [\(3.2\)](#page-8-1), these quantities are $q_i = (\beta_i - n_{i-1}\beta_{i-1})/e_i = \overline{m}_i - n_i n_{i-1} \overline{m}_{i-1}$ for $i=2,\ldots,g$ and $q_1=\overline{m}_1=m_1$.

The following lemma is a fundamental property of the semigroups coming from plane branches.

Lemma 3.1 ([\[68,](#page-35-0) I.2]). If $\Gamma = \langle \beta_0, \beta_1, \dots, \beta_g \rangle$ is the semigroup of a plane branch, then $n_i\beta_i \in \langle \beta_0, \beta_1, \ldots, \beta_{i-1} \rangle.$

The property in Lemma [3.1](#page-8-3) together with the fact that $n_i\overline{\beta}_i < \overline{\beta}_{i+1}$, which follows directly from $\beta_i < \beta_{i+1}$ and Equation [\(3.2\)](#page-8-1), characterize the semigroups of plane branches.

Proposition 3.2 ([\[68,](#page-35-0) I.3.2]). $\Gamma = \langle \overline{\beta}_0, \overline{\beta}_1, \ldots, \overline{\beta}_g \rangle \subseteq \mathbb{Z}_{\geq 0}$ is the semigroup of a plane branch, if and only if, $gcd(\beta_0, \beta_1, \ldots, \beta_g) = 1, n_i \beta_i \in \langle \beta_0, \beta_1, \ldots, \beta_{i-1} \rangle$ for $i = 1, \ldots, g$, and $n_i \overline{\beta}_i < \overline{\beta}_{i+1}$, for $1 \leq i < g$.

The conductor c of Γ can be computed as $c = n_g \overline{\beta}_g - \beta_g - (n-1)$, see [\[74,](#page-35-5) II.3]. Combining this formula with Equation [\(3.2\)](#page-8-1), we get the following formula for the conductor

(3.4)
$$
c = \sum_{i=1}^{g} (n_i - 1)\overline{\beta}_i - n + 1.
$$

Finally, we would like to notice that the Milnor number, $\mu = \dim_{\mathbb{C}} \mathbb{C}\{x,y\}/\langle \partial f/\partial x, \partial f/\partial y \rangle$, of the branch f coincides with its conductor c, see [\[22,](#page-33-22) 6.4]. Therefore, μ can be computed from a semigroup $\Gamma = \langle \beta_0, \beta_1, \dots, \beta_g \rangle$ using Equation [\(3.4\)](#page-9-0).

3.2. The monomial curve and its deformations. Let $\Gamma = \langle \beta_0, \beta_1, \ldots, \beta_g \rangle \subseteq \mathbb{Z}_{\geq 0}$ be a semigroup such that $\mathbb{Z}_{\geq 0} \setminus \Gamma$ is finite, that is $gcd(\overline{\beta}_0, \ldots, \overline{\beta}_g) = 1$, not necessarily the semigroup of a plane branch. We use the notations and definitions from Section [3.1.](#page-8-4) Following [\[68\]](#page-35-0), let $(C^{\Gamma}, 0) \subset (X, 0)$ be the curve defined via the parameterization

$$
C^{\Gamma}: u_i = t^{\overline{\beta}_i}, \quad i \le 0 \le g,
$$

where $X := \mathbb{C}^{g+1}$. The germ $(C^{\Gamma}, \mathbf{0})$ is irreducible since $gcd(\overline{\beta}_0, \ldots, \overline{\beta}_g) = 1$ and its local ring $\mathcal{O}_{C^{\Gamma},0}$ equals

$$
\mathbb{C}\big\{C^{\Gamma}\big\}=\mathbb{C}\big\{t^{\overline{\beta}_0},\ldots,t^{\overline{\beta}_g}\big\} \hookrightarrow \mathbb{C}\{t\},\,
$$

which has a natural structure of graded subalgebra of $\mathbb{C}{t}$.

Theorem 3.3 ([\[68,](#page-35-0) I.1]). Every branch $(C, 0)$ with semigroup Γ is isomorphic to the generic fiber of a one parameter complex analytic deformation of $(C^{\Gamma}, 0)$.

With extra structure on the semigroup Γ it is possible to obtain equations for $(C^{\Gamma}, 0)$.

Proposition 3.4 ([\[68,](#page-35-0) I.2]). If Γ satisfies Lemma [3.1,](#page-8-3) the branch $(C, 0) \subset (X, 0)$ is a quasi-homogeneous complete intersection with equations

(3.5)
$$
h_i := u_i^{n_i} - u_0^{l_0^{(i)}} u_1^{l_1^{(i)}} \cdots u_{i-1}^{l_{i-1}^{(i)}} = 0, \quad 1 \le i \le g,
$$

and weights $\beta_0, \beta_1, \dots, \beta_g$, where

$$
n_i\overline{\beta}_i = \overline{\beta}_0 l_0^{(i)} + \cdots + \overline{\beta}_{i-1} l_{i-1}^{(i)} \in \langle \overline{\beta}_0, \ldots, \overline{\beta}_{i-1} \rangle.
$$

Applying the theory of miniversal deformations to the previous results Teissier proves the following result.

Theorem 3.5 ([\[68,](#page-35-0) I.2]). There exists a germ of a flat morphism

 $p:(X_{\Gamma},\mathbf{0})\subset(X\times\mathbb{C}^{\tau_{-}},\mathbf{0})\longrightarrow(\mathbb{C}^{\tau_{-}},\mathbf{0})$

consisting on the second projection from X_{Γ} , such that it is a miniversal semigroup constant deformation of $(C^{\Gamma}, 0)$ with the property that, for any branch $(C, 0)$ with semigroup Γ , there exists $\boldsymbol{v}_C \in \mathbb{C}^{\tau_-}$ such that $(p^{-1}(\boldsymbol{v}_C), \boldsymbol{0})$ is analytically isomorphic to $(C, 0)$.

The miniversal deformation in Theorem [3.5](#page-9-1) can be made explicit, see [\[68,](#page-35-0) I.2]. Consider the Tjurina module of the complete intersection $(C^{\Gamma}, 0)$,

$$
T_{C^{\Gamma},\mathbf{0}}^{1} = \mathcal{O}_{X,\mathbf{0}}^{g} / \left(J\mathbf{h} \cdot \mathcal{O}_{X,\mathbf{0}}^{g+1} + \langle h_1,\ldots,h_g \rangle \cdot \mathcal{O}_{X,\mathbf{0}}^{g} \right),
$$

where $Jh \cdot \mathcal{O}_{X,\mathbf{0}}^{g+1}$ is the submodule of $\mathcal{O}_{X,\mathbf{0}}^g$ generated by the columns of the Jacobian matrix of the morphism $\mathbf{h} = (h_1, \ldots, h_g)$. Since $(C^{\Gamma}, \mathbf{0})$ is an isolated singularity, $T^1_{C^{\Gamma}, \mathbf{0}}$ is a finite dimensional C-vector space of dimension τ . Moreover, since $(C^{\Gamma}, \mathbf{0})$ is Gorenstein, $\tau = 2 \cdot \#(\mathbb{Z}_{>0} \setminus \Gamma)$, see [\[68,](#page-35-0) Prop. 2.7].

Let ϕ_1, \ldots, ϕ_μ be a basis of $T_{X,0}^1$. It is easy to see that we can take representatives for the vectors ϕ_r in \mathcal{O}_X^g having only one non-zero monomial entry $\phi_{r,i}$. Since $(C^{\Gamma}, 0)$ is quasi-homogeneous, we can endow $T_{X,\mathbf{0}}^1$ with a structure of graded module in such a way that using only the elements $\phi_1, \ldots, \phi_{\tau_-}$ with negative weight, X_{Γ} is defined from Equation (3.5) by

$$
H_i := h_i + \sum_{r=1}^{\tau_-} v_r \phi_{r,i}(u_0, \dots, u_g) = 0, \quad 1 \le i \le g,
$$

with the weight of $\phi_{r,i}$ strictly bigger than $n_i\beta_i$, see [\[68,](#page-35-0) Th. 2.10]. One can check that the classes of the vectors $(u_2, 0, \ldots, 0), (0, u_3, 0, \ldots, 0), \ldots, (0, 0, \ldots, u_g, 0)$ are C-independent in $T_{C_{\Gamma,\mathbf{0}}}^1$. Thus, if Γ is a plane branch semigroup, then $n_i\overline{\beta}_i < \overline{\beta}_{i+1}, 1 \leq i < g$ and these vectors are part of the miniversal semigroup constant deformation of $(C^{\Gamma}, 0)$.

For $\Gamma = \langle \beta_0, \beta_1, \dots, \beta_g \rangle$ the semigroup of a plane branch, consider the following semigroup constant deformation of $(C^{\Gamma}, 0)$

$$
C: H'_i := h_i - u_{i+1} = 0, \quad 1 \le i \le g.
$$

Define $f_0 := x, f_1 := y$ and set recursively,

$$
(3.6) \t f_{i+1} := f_i^{n_i} - f_0^{l_0^{(i)}} f_1^{l_1^{(i)}} \cdots f_{i-1}^{l_{i-1}^{(i)}} \t with \t n_i \overline{\beta}_i = \overline{\beta}_0 l_0^{(i)} + \cdots + \overline{\beta}_{i-1} l_{i-1}^{(i)},
$$

for $1 \leq i \leq g$. Then, we can embedded $(C, 0)$ in the plane in such a way that it has equation $f := f_{q+1}$ that looks like

$$
\left(\cdots\left(\left(\cdots\left(\left(y^{n_1}-x^{l_0^{(1)}}\right)^{n_2}-x^{l_0^{(2)}}y^{l_0^{(2)}}\right)^{n_3}-\cdots\right)^{n_i}x^{l_0^{(i)}}\cdots f_{i-1}^{l_{i-1}^{(i)}}\right)^{n_{i+1}}\cdots\right)^{n_{g-1}}x^{l_0^{(g)}}\cdots f_{g-1}^{l_{g-1}^{(g)}}.
$$

This proves the reverse implication in Proposition [3.2.](#page-9-3)

In the following sections, it will be essential to have a certain semigroup constant deformation of a plane branch equation, similar to f_{q+1} in Equation [\(3.6\)](#page-10-0), in such a way that any other plane branch with the same semigroup is analytically equivalent to some fiber of that deformation. From the previous discussion we obtain the following result.

Proposition 3.6. Let $\Gamma = \langle \beta_0, \beta_1, \dots, \beta_g \rangle$ be a plane branch semigroup. Consider, with the same notation as above,

(3.7)
$$
f_{i+1} = f_i^{n_i} - \lambda_i f_0^{l_0^{(i)}} f_1^{l_1^{(i)}} \cdots f_{i-1}^{l_{i-1}^{(i)}} + \sum_{\overline{\beta}_0 k_0 + \cdots + \overline{\beta}_i k_i > n_i \overline{\beta}_i} t_k^{(i)} f_0^{k_0} f_1^{k_1} \cdots f_i^{k_i},
$$

with $\lambda_i \neq 0, \lambda_1 = 1$ for $i = 1, \ldots, g$ and the sum being finite. Define,

$$
f_{\boldsymbol{t},\boldsymbol{\lambda}}(x,y) := f_{g+1}(x,y;\underline{t}^{(1)},\ldots,\underline{t}^{(g)};\lambda_2,\ldots,\lambda_g).
$$

Then, $\{f_{t,\lambda}(x,y)\}_{\lambda \in \mathbb{C}^{g-1}}$ is an infinite family of semigroup constant deformations, all having semigroup Γ, with the property that any other plane branch with semigroup Γ is analytically equivalent to a fiber of one element of the family.

Proof. Consider the semigroup constant deformation of the monomial curve $(C^{\Gamma}, 0)$ from Theorem [3.5.](#page-9-1) Since the semigroup Γ is of a plane branch, we can assume that the set of vectors $(u_2, 0, \ldots, 0), (0, u_3, 0, \ldots, 0), \ldots, (0, 0, \ldots, u_q, 0)$ are part of the semigroup constant deformation. X_{Γ} will have equations

$$
C: H_i = h_i - v_{i+1}u_{i+1} + \sum_{r=g+1}^{\tau_-} v_r \phi_{r,i}(u_0, \dots, u_g) = 0, \quad 1 \le i \le g.
$$

The embedding dimension of $(C, 0)$ is equal to $g + 1 - \text{rk } JH(0)$, see [\[27,](#page-33-23) 4.3]. Since all the monomials in H_i have (non-weighted) degree bigger than 2, except for those in the vectors $(u_2, 0, \ldots, 0), (0, u_3, 0, \ldots, 0), \ldots, (0, 0, \ldots, u_q, 0)$, the rank of the Jacobian is $g-1$, if and only if, $v_2 \cdots v_{q-1}$ is non-zero. Thus, the embedding dimension of $(C, 0)$ is 2, if and only if, all v_2, \ldots, v_{g-1} are different from zero.

Finally, performing elimination on the variables u_2, \ldots, u_g one obtains a plane branch equation similar to Equation [\(3.7\)](#page-10-1) with $\lambda_i = v_2^{n_2} \cdots v_i^{n_i} \neq 0$ and a finite number of deformation monomials with coefficients that are polynomials in the variables v_r . Therefore, there is an inclusion of the parameter space of X_{Γ} into the parameters of the family of deformations $f_{t,\lambda}$.

We will sometimes drop the dependency on the parameters $\lambda \in \mathbb{C}^{g-1}$, and denote just $f_t(x, y)$. Although we are considering a finite deformation of f_t , we can always assume that we have deformation terms of high enough order. Adding extra terms to the summation does not change the facts that the family contains all plane curves up to analytic isomorphism or that the deformation has constant semigroup.

4. Resolution of plane curve singularities

In this section, we will review some facts about resolution of singularities of plane curves. In the sequel, a plane curve will be a germ of a mapping $f : (\mathbb{C}^2, \mathbf{0}) \longrightarrow (\mathbb{C}, 0)$ with $f(0) = 0$. We will make a small abuse of notation and denote also by f an equation of the germ, assuming it is defined in a small neighborhood $U \subseteq \mathbb{C}^2$ of the origin. For irreducible plane curves f , we will present the notions of toric resolution, maximal contact elements, and their relation. We will also see how some resolution data can be described in terms of the semigroup and the characteristic sequence of f . In the last part of this section, we give equations for the resolution of the elements in the family f_t considered in Proposition [3.6](#page-10-2) at the so-called rupture divisors. The majority of the results in this section are well-known and can be found in the books of Casas-Alvero [\[22\]](#page-33-22) or Wall [\[70\]](#page-35-6).

4.1. Resolution of singularities. Let $\pi : X' \longrightarrow U$ be the minimal embedded resolution of the plane curve f. Here, π is birational proper morphism and X' is a smooth surface. We can assume that π is given as a composition of point blow-ups

$$
\pi: X' := X_{s+1} \xrightarrow{\pi_{p_s}} X_r \longrightarrow \cdots \longrightarrow X_1 \xrightarrow{\pi_{p_0}} X_0 := U \subseteq \mathbb{C}^2,
$$

with $X_{i+1} := \text{Bl}_{p_i} X_i, p_i \in \text{Exc}(\pi_{p_{i-1}} \circ \cdots \circ \pi_{p_0}) \subset X_i$ for $i = 1, \ldots, s$ and $p_0 := \mathbf{0}$. Denote by $E := \text{Exc}(\pi)$ the exceptional divisor of π . Let $K := \{p_0, p_1, \ldots, p_s\}$ be the set of points that have been blown-up. Then, ${E_p}_{p \in K}$ is the set of all the irreducible exceptional components of E. The total transform divisor F_{π} and relative canonical divisor K_{π} will have the following expressions

$$
F_{\pi} := \text{Div}(\pi^* f) = \sum_{p \in K} N_p E_p + \mu_1 \widetilde{C}_1 + \cdots + \mu_t \widetilde{C}_t, \quad K_{\pi} = \text{Div}(\text{Jac}(\pi)) := \sum_{p \in K} k_p E_p,
$$

where $C_1, \ldots, C_t \in \text{Div}_{\mathbb{Z}}(X')$ are the branches of the strict transform of f, that is to say, $C_1 + \cdots + C_t = Div(\pi^{-1}(f - \{\mathbf{0}\}))$. The divisor F_{π} is a simple normal crossing divisor.

We will distinguish between two types of exceptional divisors. An exceptional divisor is said to be of *rupture* type if it intersects three or more divisors in the support of F_{π} . It is said to be non-rupture otherwise. Those exceptional divisors that only intersect one divisor in the support of F_{π} will be called *dead-end* divisors. They are dead-end points in the dual graph of the resolution, hence the name.

4.2. Toric resolutions and maximal contact elements. We assume from now on that $f: (\mathbb{C}^2, \mathbf{0}) \longrightarrow (\mathbb{C}, 0)$ is irreducible with semigroup $\Gamma = \langle \overline{\beta}_0, \overline{\beta}_1, \ldots, \overline{\beta}_g \rangle$ and we use the notations from Section [3.1.](#page-8-4) All plane branches having the same semigroup Γ , or characteristic sequence, are *equisingular*, see $[22, 3.8]$. This means that the combinatorics of the resolution is the same for all of them and can be determined by the characteristic sequence.

A classical way to obtain the minimal resolution by point blow-ups of an irreducible plane curve f from its characteristic sequence $(n, \beta_1, \ldots, \beta_q)$ is using Enriques' theorem, see $[22, 5.5]$. We will however take the approach of Oka in $[59]$ and describe the minimal resolution of f as a composition of toric morphisms. Indeed, there exists a resolution map π of f that decomposes into $g \geq 1$ toric morphisms. For $i = 1, \ldots, g$,

$$
\pi^{(i)} := \pi_1 \circ \cdots \circ \pi_{i-1} \circ \pi_i : X^{(i)} \xrightarrow{\pi_i} X^{(i-1)} \xrightarrow{\pi_{i-1}} \cdots \xrightarrow{\pi_2} X^{(1)} \xrightarrow{\pi_1} X^{(0)} := U \subseteq \mathbb{C}^2,
$$

where π_i is a toric morphism for a suitable choice of coordinates on $X^{(i-1)}$ and $\pi := \pi^{(g)}$. Each π_i resolves one characteristic exponent of the plane branch f in the sense that the strict transform of f on $X^{(i)}$ has one characteristic exponent less than the strict transform on $X^{(i-1)}$. In this way, $X^{(i)}$ always contain one extra rupture divisor E_{p_i} than $X^{(i-1)}$. We will denote by U_i, V_i the affine open sets, and by $(x_i, y_i), (z_i, w_i)$ the coordinates, containing the *i*-th rupture divisor E_{p_i} on $X^{(i)}$ after the *i*-th toric modification π_i . In these coordinates, recalling the definitions of the integers n_i, q_i from Section [3.1,](#page-8-4) the toric morphism is given by

(4.1)
$$
\pi_i(x_i, y_i) = (x_i^{n_i} y_i^{a_i}, x_i^{q_i} y_i^{b_i}) \quad \text{and} \quad \pi_i(z_i, w_i) = (z_i^{c_i} w_i^{n_i}, z_i^{d_i} w_i^{q_i}),
$$

with $a_i, b_i, c_i, d_i \in \mathbb{Z}_{\geq 0}$ such that $n_i b_i - q_i a_i = 1, q_i c_i - n_i d_i = 1$ and $a_i q_i + d_i n_i = n_i q_i - 1$. These toric morphisms can be thought as a composition of point blow-ups. In the sequel, we will associate to each plane branch singularity these series of four integers a_i, b_i, c_i, d_i for every $i = 1, \ldots, g$. They are determined, although not explicitly, by the semigroup Γ of f since they depend on the continuous fraction expansion of q_i/n_i .

If the singularity is still not resolved at $X^{(i)}$, one needs to perform an analytic change of coordinates around the unique singular point of the strict transform of f on $X^{(i)}$ in

order for π_{i+1} to be toric. These new coordinates, let us say (\bar{x}_i, \bar{y}_i) and (\bar{z}_i, \bar{w}_i) , are such that $\pi^{(i)}_*\bar{y}_i = \pi^{(i)}_*\bar{z}_i$ is a germ $f_i: (\mathbb{C}^2, \mathbf{0}) \longrightarrow (\mathbb{C}, 0)$ that is a maximal contact element of f in the sense that the intersection of f_i and f is, precisely, β_i . By construction, each of these maximal contact elements f_i are resolved by the corresponding $\pi^{(i)}$. In the case of the plane curves constructed in Section [3.2,](#page-9-4) the maximal contact element f_i coincide with the elements f_i defined in Equation (3.6) and Equation (3.7) . For completeness, we will assume that the $(g + 1)$ –th maximal contact element f_{g+1} is the curve f itself.

It is easy to see that the semigroup Γ_{i+1} of the maximal contact element f_{i+1} is

(4.2)
$$
\Gamma_{i+1} = \langle n_1 n_2 \cdots n_i, n_2 \cdots n_i \overline{m}_1, \ldots, n_i \overline{m}_{i-1}, \overline{m}_i \rangle, \text{ for } i = 1, \ldots, g.
$$

Similarly, its characteristic sequence is given by $(n_1 n_2 \cdots n_i, n_2 \cdots n_i m_1, \ldots, n_i m_{i-1}, m_i)$. Apart from this numerical data of the maximal contact elements, we are interested in describing the multiplicities of the total transform and the relative canonical divisor along the rupture and dead-end divisors in terms of the semigroup Γ. Following the same notation as above, denote E_{p_i} , $i = 1, \ldots, g$ the rupture divisors of f. Similarly, denote E_{q_i} for $i = 0, \ldots, g$ the dead-end divisors. It is well-known, see [\[70,](#page-35-6) 8.5], that

$$
(4.3) N_{p_i} = n_i \overline{\beta}_i, \quad k_{p_i} + 1 = m_i + n_1 \cdots n_i, \quad N_{q_i} = \overline{\beta}_i, \quad k_{q_i} + 1 = \lceil (m_i + n_1 \cdots n_i)/n_i \rceil.
$$

We will end this section with a technical result about the resolution of the elements of the constant semigroup deformations ${f_{t,\lambda}}_{\lambda \in \mathbb{C}^{g-1}}$. Having constant semigroup means that all the fibers of all the elements of the family are equisingular. Hence, the toric resolution of the plane branches in the family f_t is the same modulo the coordinates needed at each $X^{(i)}$. The following proposition describes locally the equations of f_t around the rupture divisors after pulling back by $\pi^{(i)}$.

Proposition 4.1. Let E_{p_i} be the *i*-th rupture divisor on the surface $X^{(i)}$ and let U_i , V_i be the corresponding charts containing E_{p_i} with local coordinates (x_i, y_i) and (z_i, w_i) , respectively. Then,

• The equations of the total transform of $f_t(x, y)$ are given by

$$
(4.4) \t x_i^{n_i \overline{\beta}_i} y_i^{a_i \overline{\beta}_i} u_1(x_i, y_i) \tilde{f}_t(x_i, y_i), \t z_i^{(c_i n_{i-1} \overline{m}_{i-1} + d_i)e_{i-1}} w_i^{n_i \overline{\beta}_i} u_2(x_i, y_i) \tilde{f}_t(z_i, w_i),
$$

where u_1, u_2 are units at any point of E_{p_i} .

• The equations \hat{f}_t of the strict transform of f are

$$
\tilde{f}_{t}(x_i, y_i) = \tilde{f}_{g+1}(x_i, y_i; \underline{t}^{(1)}, \dots, \underline{t}^{(g)}), \quad \tilde{f}_{t}(z_i, w_i) = \tilde{f}_{g+1}(z_i, w_i; \underline{t}^{(1)}, \dots, \underline{t}^{(g)}).
$$

• The $(i + 1)$ –th maximal contact element has equations

(4.5)
\n
$$
\tilde{f}_{i+1}(x_i, y_i) = y_i - \lambda_i + \sum_{\overline{\beta}_0 k_0 + \dots + \overline{\beta}_i k_i > n_i \overline{\beta}_i} t_{\underline{k}}^{(i)} x_i^{a_{i+1}^{(i)}} y_i^{A_{i+1}^{(i)}} u_{\underline{k}}^{(i)}(x_i, y_i), \qquad (U_i \text{ chart}),
$$
\n
$$
\tilde{f}_{i+1}(z_i, w_i) = 1 - \lambda_i z_i + \sum_{\overline{\beta}_0 k_0 + \dots + \overline{\beta}_i k_i > n_i \overline{\beta}_i} t_{\underline{k}}^{(i)} z_i^{C_{i+1}^{(i)}} w_i^{a_{i+1}^{(i)}} u_{\underline{k}}^{(i)}(z_i, w_i), \qquad (V_i \text{ chart}),
$$

where $u_k^{(i)}$ $\frac{k}{k}$ are units at any point of E_{p_i} . • The remaining maximal contact elements $f_{j+1}, j > i$ have strict transforms given by (4.6)

$$
\tilde{f}_{j+1} = \tilde{f}_j^{n_j} - \lambda_j x_i^{\rho_{j+1}^{(i)}(l_j)} y_i^{A_{j+1}^{(i)}(l_j)} u_0^{(j)} \tilde{f}_{i+1}^{l_{i+1}^{(j)}} \cdots \tilde{f}_{j-1}^{l_{j-1}^{(j)}} + \sum_{\overline{\beta}_0 k_0 + \cdots + \overline{\beta}_j k_j > n_j \overline{\beta}_j} t_{\underline{k}}^{(j)} x_i^{\rho_{j+1}^{(i)}(k)} y_i^{A_{j+1}^{(i)}(k)} u_{\underline{k}}^{(j)} \tilde{f}_{i+1}^{k_{i+1}} \cdots \tilde{f}_j^{k_j}
$$

in the U_i chart, and similarly in V_i . As before, $u_0^{(j)}$ $\frac{(j)}{0}, u^{(j)}_k$ $\frac{dS}{dt}$ are units everywhere on E_{p_i} and we denote $\underline{l}_j := (l_0^{(j)})$ $\mathcal{L}_0^{(j)}, l_1^{(j)}, \ldots, l_{j-1}^{(j)}, 0$ the integers from Proposition [3.4.](#page-9-5)

Finally, $\rho_{j+1}^{(i)}, A_{j+1}^{(i)}, C_{j+1}^{(i)}, j \geq i$, are the following linear forms:

$$
\rho_{j+1}^{(i)}(\underline{k}) = \sum_{l=0}^{i} n_{l+1} \cdots n_i \overline{m}_l k_l + n_i \overline{m}_i \sum_{l=i+1}^{j} n_{i+1} \cdots n_{l-1} k_l - n_i \cdots n_j \overline{m}_i,
$$

\n
$$
A_{j+1}^{(i)}(\underline{k}) = a_i \sum_{l=0}^{i-1} n_{l+1} \cdots n_{i-1} \overline{m}_l k_l + (a_i n_{i-1} \overline{m}_{i-1} + b_i) k_i
$$

\n
$$
+ a_i \overline{m}_i \sum_{l=i+1}^{j} n_{i+1} \cdots n_{l-1} k_l - a_i \overline{m}_i n_{i+1} \cdots n_j,
$$

\n
$$
C_{j+1}^{(i)}(\underline{k}) = c_i \sum_{l=0}^{i-1} n_{l+1} \cdots n_{i-1} \overline{m}_l k_l + (c_i n_{i-1} \overline{m}_{i-1} + d_i) k_i
$$

\n
$$
+ n_i (c_i n_{i-1} \overline{m}_{i-1} + d_i) \sum_{l=i+1}^{j} n_{i+1} \cdots n_{l-1} k_l - (c_i n_{i-1} \overline{m}_{i-1} + d_i) n_i \cdots n_j.
$$

Proof. The results follow from the inductive procedure of applying the toric transforma-tions from Equation [\(4.1\)](#page-12-0) to the equations of f_t in Proposition [3.6.](#page-10-2) At each $X^{(i)}$ the analytic coordinates which make the morphism π_i toric are described by $\bar{y}_i = f_{i+1}, \bar{x}_i = x_i u_i$ in the U_i chart, for some unit u_i . The expressions for the linear forms $\rho_{j+1}^{(i)}, A_{j+1}^{(i)}, C_{j+1}^{(i)}$ follow, recursively, from the relations

$$
\rho_{j+1}^{(i)}(\underline{k}) = n_i \rho_{j+1}^{(i-1)}(\underline{k}) + q_i k_i + n_i q_i \sum_{l=i+1}^j n_{i+1} \cdots n_{l-1} k_l - n_i \cdots n_j q_i,
$$

\n
$$
A_{j+1}^{(i)}(\underline{k}) = a_i \rho_{j+1}^{(i-1)}(\underline{k}) + b_i k_i + a_i q_i \sum_{l=i+1}^j n_{i+1} \cdots n_{l-1} k_l - a_i q_i n_{i+1} \cdots n_j,
$$

\n
$$
C_{j+1}^{(i)}(\underline{k}) = c_i \rho_{j+1}^{(i-1)}(\underline{k}) + d_i k_i + n_i d_i \sum_{l=i+1}^j n_{i+1} \cdots n_{l-1} k_l - d_i n_i \cdots n_j.
$$

Following the notations from Proposition [4.1,](#page-13-0) we will fix, in the sequel, the index i to denote that we have resolved the singularity up to the *i*-th rupture divisor on $X^{(i)}$. On the other hand, the index j will make reference to the j -th maximal contact element. At any step of the resolution process, one has that $1 \leq i \leq j \leq g$.

 \Box

Lemma 4.2. Let $\rho_{j+1}^{(i)}(\underline{k}), A_{j+1}^{(i)}(\underline{k}), C_{j+1}^{(i)}(\underline{k})$ be the linear forms in Proposition [4.1.](#page-13-0) Then,

$$
A_{j+1}^{(i)}(\underline{k}) + C_{j+1}^{(i)}(\underline{k}) + \sum_{l=i+1}^{j} n_{i+1} \cdots n_{l-1} k_l = \rho_{j+1}^{(i)}(\underline{k}) + n_{i+1} \cdots n_j
$$

.

Proof. From the relations between a_i, b_i, c_i, d_i , one can deduce that $a_i + c_i = n_i, b_i + d_i = q_i$. The result follows from adding $A_{j+1}^{(i)}$ and $C_{j+1}^{(i)}$ and using these relations.

5. Poles and residues for plane curves

Let $f: (\mathbb{C}^2, \mathbf{0}) \longrightarrow (\mathbb{C}, 0)$ be a plane curve not necessarily reduced or irreducible. After fixing local coordinates x, y, and with a small abuse of notation, let $f \in \mathbb{C}\{x, y\}$ be an equation of the germ, assuming it is defined in a neighborhood $U \subseteq \mathbb{C}^2$ of the origin. Following Section [2.1,](#page-3-2) define the complex zeta function f^s of a local singularity as

(5.1)
$$
\langle f^s, \varphi \rangle := \int_U |f(x, y)|^{2s} \varphi(z) dz, \text{ for } \text{Re}(s) > 0,
$$

with $\varphi \in C_c^{\infty}(U)$ and $z := (x, y, \bar{x}, \bar{y})$. The poles of $\langle f^s, \varphi \rangle$ do not depend on the equation of the germ or the local coordinates x, y . As discussed in Section [2.1,](#page-3-2) f^s must be understood in the distributional sense. In this section, we will use the minimal resolution of singularities of f to study the structure of the residues of f^s at any candidate pole σ . The residue will be expressed as an improper integral along the exceptional divisor associated to σ . In order to do so, we first present the straightforward generalization of Proposition [2.1](#page-4-1) to the two dimensional case and see how poles of order two might arise. Finally, we will use the residue formula to prove that most non-rupture divisors do not contribute to poles of the complex zeta function f^s .

5.1. Regularization of monomials in two variables. The result from Proposition [2.1](#page-4-1) can be easily generalized to the two dimensional case, mimicking the proof in [\[32\]](#page-33-1), to see how poles of order two arise. Let $\varphi(z_1, z_2) \in C_c^{\infty}(\mathbb{C}^2)$ which is, in fact, a function of $z = (z_1, z_2, \overline{z}_1, \overline{z}_2)$, and consider

(5.2)
$$
\langle z_1^{s_1} z_2^{s_2}, \varphi \rangle = \int_{\mathbb{C}^2} |z_1|^{2s_1} |z_2|^{2s_2} \varphi(z) dz,
$$

which is absolutely convergent for $\text{Re}(s_1) > -1$, $\text{Re}(s_2) > -1$ since φ has compact support.

Let $\Delta_0 = D_1 \times D_2$ be the polydisc formed by the discs of radius one centered at the origin, i.e. $D_1 = \{|z_1| \leq 1\}$ and $D_2 = \{|z_2| \leq 1\}$. We can decompose \mathbb{C}^2 as the disjoint union

$$
\Delta_0 \cup (D_1 \times \mathbb{C} \setminus D_2) \cup (\mathbb{C} \setminus D_1 \times D_2) \cup (\mathbb{C} \setminus D_1 \times \mathbb{C} \setminus D_2).
$$

Using the notation $z^k = z_1^{k_1} z_2^{k_2} \bar{z}_1^{k_3} \bar{z}_2^{k_4}$, the integral in Equation [\(5.2\)](#page-15-1) on the region Δ_0 can be written as

$$
\int_{\Delta_0} |z_1|^{2s_1} |z_2|^{2s_2} \bigg(\varphi(z) - \sum_{|k| \le m} \frac{\partial^k \varphi(\mathbf{0})}{\partial z^k} \frac{z^k}{k!} \bigg) dz - \sum_{|k| \le m} \frac{\partial^k \varphi}{\partial z^k}(\mathbf{0}) \frac{4\pi^2}{k!(s_1 + k_1 + 1)(s_2 + k_2 + 1)},
$$

where in the second summation, we have $k_1 = k_3, k_2 = k_4$. The left-hand integral is holomorphic on the regions $\text{Re}(s_1) > -m-1, \text{Re}(s_2) > -m-1.$

With a small abuse of the notation, let $z_1 = (z_1, \bar{z}_1), z_2 = (z_2, \bar{z}_2)$ and $z_1^{k_1} = z_1^{k_1,1}$ $\frac{k_{1,1}}{2}\bar{z}_1^{k_{1,2}}$ $\frac{\kappa_{1,2}}{1}$. On the region $D_1 \times \mathbb{C} \setminus D_2$, the integral in Equation [\(5.2\)](#page-15-1) is

$$
\int_{D_1\times\mathbb{C}\backslash D_2} \hspace{-2mm} |z_1|^{2s_1} |z_2|^{2s_2} \bigg(\varphi(z)-\sum_{|k_1|\leq m} \frac{\partial^{k_1}\varphi}{\partial z_1^{k_1}}(0,z_2)\frac{z_1^{k_1}}{k_1!}\bigg) dz_1 dz_2 - 2\pi i \sum_{|k_1|\leq m} \frac{\displaystyle\int_{|z_2|>1} |z_2|^{2s_2} \frac{\partial^{k_1}\varphi}{\partial z_1^{k_1}}(0,z_2) dz_2}{(k_{1,1}!)^2(s_1+k_{1,1}+1)},
$$

where in the second sum, $k_{1,1} = k_{1,2}$. The left-hand integral is holomorphic in $\text{Re}(s_1)$ $-m-1$. By symmetry, a similar expression holds true in the other region, $\mathbb{C} \setminus D_1 \times D_2$. On the last region, $\mathbb{C}\setminus D_1\times \mathbb{C}\setminus D_2$, the integral in Equation [\(5.2\)](#page-15-1) is absolutely convergent for all $s_1, s_2 \in \mathbb{C}$.

From the regularization of $z_1^{s_1} z_2^{s_2}$ constructed above we can see that the residue of $z_1^{s_1} z_2^{s_2}$ at a simple pole $s_1 = -k - 1, k \in \mathbb{Z}_{\geq 0}$, i.e. the coefficient of $(s + k + 1)^{-1}$, is given by the following function of s_2

(5.3)
$$
\operatorname{Res}_{s_1=-k-1} \langle z_1^{s_1} z_2^{s_2}, \varphi \rangle = -\frac{2\pi i}{(k!)^2} \int_{\mathbb{C}} |z_2|^{2s_2} \frac{\partial^{2k} \varphi}{\partial z_1^k \partial \bar{z}_1^k} (0, s_2) dz_2 d\bar{z}_2.
$$

The residue in Equation [\(5.3\)](#page-16-0) being a function of s_2 implies that $\text{Res}_{s_1=-k-1} z_1^{s_1} z_2^{s_2}$ will have a simple pole, as a function of s_2 , on the poles of order two of $z_1^{s_1}z_2^{s_2}$. Conversely, if Equation [\(5.3\)](#page-16-0) is zero for certain $s_2 = \alpha$, the point $(s_1, s_2) = (-k-1, \alpha)$ is neither a pole of order one nor a pole of order two of $z_1^{s_1} z_2^{s_2}$.

5.2. The residue at the poles. Let $E_p, p \in K$ be an irreducible exceptional divisor. We will denote by $D_1, D_2, \ldots, D_r \in Div_{\mathbb{Z}}(X')$ the other prime components (exceptional or not) of F_{π} crossing E_p . By definition, dead-end divisors have only one divisor crossing them, which will be denoted D_1 . On the other hand, rupture divisors have at least three divisors crossing them, i.e. $r \geq 3$. In any other case, $r = 2$. We will denote by N_1, N_2, \ldots, N_r (resp. k_1, k_2, \ldots, k_r) the coefficients of D_1, D_2, \ldots, D_r in F_π (resp. in K_π). Since no confusion arises, we drop the explicit dependence on $p \in K$.

For each $E_p, p \in K$ we consider two affine charts U_p, V_p containing E_p that arise after the blow-up of a neighborhood of p in any chart containing p . The origin of the charts U_p, V_p are neighborhoods of opposite points in the projective line E_p . Usually, these points are the intersection points of E_p with two other components of F_π which we will assume to be D_1 and D_2 . For simplicity, if this is not the case, we will set N_1, k_1 (or N_2, k_2) to be zero. The only case in which both N_1D_1 and N_2D_2 are zero is when the minimal resolution π consists of a single blow-up, that is, an homogeneous singularity.

In order to define the complex zeta function f^s on X' , we need to work locally with coordinates. Accordingly, let (x_p, y_p) , (z_p, w_p) be the natural holomorphic coordinates of U_p, V_p centered at the origins of both charts which, by construction, are the origin and the infinity point on a $\mathbb{P}^1_{\mathbb{C}}$, or vice versa. The coordinates $(x_p, y_p), (z_p, w_p)$ are related at the intersection $U_p \cap V_p$ by $x_p = z_p^{\kappa_p} w_p, y_p z_p = 1$, and where the integer value $\kappa_p \in \mathbb{Z}_{>0}$ has a very precise geometric meaning, namely, $\kappa_p = -E_p \cdot E_p$.

Following the discussion in Section [2.1,](#page-3-2) each exceptional component E_p contributes with a sequence of candidate poles to the meromorphic continuation of f^s . Indeed, with the notations above,

$$
\left\{\sigma_{p,\nu} = -\frac{k_p + 1 + \nu}{N_p} \middle| \nu \in \mathbb{Z}_{\geq 0} \right\}, \quad p \in K.
$$

The set $\mathcal{W}_K := \{U_p, V_p\}_{p \in K}$ forms a finite affine open cover X', which applied to the construction presented in Section [2.1,](#page-3-2) Equation [\(2.3\)](#page-5-0), results in the following proposition. First, denote by $\{\eta_{1,q}, \eta_{2,q}\}_{q \in K}$ a partition of unity subordinated to the open cover \mathcal{W}_K .

Proposition 5.1. Using the affine open cover W_K , the part of the complex zeta function f^s on Equation [\(2.3\)](#page-5-0) involving the divisor E_p can be written as the sum of the integrals over just two affine charts $U_p, V_p \in \mathcal{W}_K$ containing E_p , namely

(5.4)
$$
\int_{U_p} |x_p|^{2(N_ps+k_p)} |y_p|^{2(N_1s+k_1)} \Phi_1(x_p, y_p; s) \eta_1 dx_p dy_p d\bar{x}_p d\bar{y}_p +
$$

$$
\int_{V_p} |z_p|^{2(N_2s+k_2)} |w_p|^{2(N_ps+k_p)} \Phi_2(z_p, w_p; s) \eta_2 dz_p dw_p d\bar{z}_p d\bar{w}_p,
$$

where $\Phi_1(x_i, y_i; s), \Phi_2(z_i, w_i; s)$ are infinitely many times differentiable at neighborhoods of the points $p_1 = E_p \cap D_1$ and $p_2 = E_p \cap D_2$. More precisely,

$$
\Phi_1 := |u_1|^{2s} |v_1|^2 (\pi^* \varphi)|_{U_p}, \quad \Phi_2 := |u_2|^{2s} |v_2|^2 (\pi^* \varphi)|_{V_p},
$$

with the elements u_1, v_1 (resp. u_2, v_2) being units in the local ring at the points p_1 (resp. p_2). Finally, η_1 and η_2 have compact support and $\eta_1|_{E_p} + \eta_2|_{E_p} \equiv 1$.

Proof. Let us denote by $\mathcal{W}_{p_1}, \mathcal{W}_{p_2}$ all the elements in \mathcal{W}_K that contain p_1 and p_2 . By construction, \mathcal{W}_{p_1} is disjoint from \mathcal{W}_{p_2} , since there can be no affine open set containing both p_1 and p_2 , and the union of \mathcal{W}_{p_1} and \mathcal{W}_{p_2} contains all the charts from \mathcal{W}_K containing E_p . Applying Equation [\(2.3\)](#page-5-0) from Section [2.1,](#page-3-2) the part of f^s on X' where the divisor E_p appears is a sum of the integrals over the affine open sets from \mathcal{W}_{p_1} and \mathcal{W}_{p_2} ,

$$
(5.5)\quad \sum_{U_q \in \mathcal{W}_{p_1}} \int_{U_q} |\pi^* f|^{2s} |_{U_q} (\pi^* \varphi)|_{U_q} |d\pi|_{U_q}|^2 \eta_{1,q} + \sum_{V_q \in \mathcal{W}_{p_2}} \int_{V_q} |\pi^* f|^{2s} |_{V_q} (\pi^* \varphi)|_{V_q} |d\pi|_{V_q}|^2 \eta_{2,q}.
$$

Let us see that we can reduce Equation (5.5) to Equation (5.4) . The proof is the same for both summations in Equation [\(5.5\)](#page-17-0). Since the elements of \mathcal{W}_{p_1} are blow-up charts, the difference between the union and the intersection of all the elements in \mathcal{W}_{p_1} is a finite number of lines. Given that a finite number of lines have measure zero, they do not affect the integral, and we can replace the left-hand summation of Equation [\(5.5\)](#page-17-0) by

$$
\int_{\cap U_q} |\pi^* f|^{2s} |\negthinspace\negthinspace\negthinspace\negthinspace\negthinspace v_q \; (\pi^* \varphi)|_{\cap U_q} |d\pi|_{\cap U_q} |^2 \eta_1
$$

This equality is true since $|\pi^* f|^{2s}\pi^* \varphi|d\pi|^2$ is a global section on X' and it coincides at the intersection of all the $U_q \in \mathcal{W}_{p_1}$. Concerning the partitions of unit, we just set $\eta_1 := \sum_{U_q \in \mathcal{W}_{p_1}} \eta_{1,q}$. Finally, by the same argument as before, we can replace $\bigcap_{U_q \in \mathcal{W}_{p_1}} U_q$ by any $U_p \in \mathcal{W}_{p_1}$ yielding Equation [\(5.4\)](#page-17-1). Notice that, by definition of \mathcal{W}_{p_1} , no other $\eta \in \{\eta_{1,q}, \eta_{2,q}\}_{q \in K}$, except for those in η_1 , has p_1 in its support.

Before presenting the formula for the residue, let us introduce the following rational numbers associated to a candidate pole $\sigma_{p,\nu}$ of an irreducible exceptional divisor $E_p, p \in K$. They will play an important role in the analysis of the residues.

Definition 5.2 (Residue numbers). Let $\sigma_{p,\nu}, \nu \in \mathbb{Z}_{\geq 0}$ be a candidate pole of f^s associated to an exceptional divisor $E_p, p \in K$ intersecting the divisors $D_1, D_2, \ldots, D_r \in Div_{\mathbb{Z}}(X')$. Define the residue numbers as

(5.6)
$$
\epsilon_{i,\nu} := N_i \sigma_{p,\nu} + k_i \in \mathbb{Q} \quad \text{for} \quad i = 1,\ldots,r.
$$

For the ease of notation, we will omit the dependence of $\epsilon_{1,\nu}, \epsilon_{2,\nu}$ on $p \in K$. The following relations between $\epsilon_{1,\nu}, \epsilon_{2,\nu}, \ldots, \epsilon_{r,\nu}$ holds.

Lemma 5.3. For any $\nu \in \mathbb{Z}_{\geq 0}$, we have

(5.7)
$$
\epsilon_{1,\nu} + \epsilon_{2,\nu} + \cdots + \epsilon_{r,\nu} + \kappa_p \nu + 2 = 0.
$$

Proof. Consider the Q-divisor $\sigma_{p,\nu}F_{\pi} + K_{\pi}$. Applying the adjunction formula for surfaces [\[34,](#page-33-24) §V.1], $(\sigma_{p,\nu}F_{\pi}+K_{\pi})\cdot E_p = \kappa_p-2$, recall $\kappa_p = -E_p\cdot E_p$. On the other hand,

$$
(\sigma_{p,\nu}F_{\pi} + K_{\pi}) \cdot E_p = \sum_{i=0}^r \epsilon_{i,\nu} - \kappa_p (N_p \sigma_{p,\nu} + k_p).
$$

Since $N_p \sigma_{p,\nu} + k_p = -\nu - 1$, the result follows.

A first instance of the numbers $\epsilon_{i,\nu}$ and of Equation [\(5.7\)](#page-18-1), in the case of rupture divisors of irreducible plane curves and $\nu = 0$, already appeared in an article of Lichtin [\[47\]](#page-34-5).

The formula for the residue at a candidate pole $\sigma_{p,\nu}$ is presented next. The residue is expressed as an improper integral, see Remark [5.6](#page-19-0) below, along the divisor E_p having singularities of orders $\epsilon_{1,\nu}, \epsilon_{2,\nu}, \ldots, \epsilon_{1,r}$ at the intersection points of E_p with D_1, D_2, \ldots, D_r .

Proposition 5.4. The residue of the complex zeta function f^s at a candidate pole $s = \sigma_{p,\nu}$ is given by

(5.8)
$$
\operatorname{Res}_{s=\sigma_{p,\nu}} \langle f^s, \varphi \rangle = \frac{-2\pi i}{(\nu!)^2} \int_{\mathbb{C}} |y_p|^{2\epsilon_{1,\nu}} \frac{\partial^{2\nu} \Phi_1}{\partial x_p^{\nu} \partial \bar{x}_p^{\nu}} (0, y_p; \sigma_{p,\nu}) dy_p d\bar{y}_p \qquad (U_p \text{ chart})
$$

$$
= \frac{-2\pi i}{(\nu!)^2} \int_{\mathbb{C}} |z_p|^{2\epsilon_{2,\nu}} \frac{\partial^{2\nu} \Phi_2}{\partial w_p^{\nu} \partial \bar{w}_p^{\nu}} (z_p, 0; \sigma_{p,\nu}) dz_p d\bar{z}_p, \qquad (V_p \text{ chart}).
$$

Proof. Applying Equation [\(5.3\)](#page-16-0) to Proposition [5.1](#page-17-2) with $s_2 = N_1 \sigma_{p,\nu} + k_1$ and $N_2 \sigma_{p,\nu} + k_2$, respectively, we obtain that the residue of f^s at $s = \sigma_{p,\nu}$ is

(5.9)
$$
\operatorname{Res}_{s=\sigma_{p,\nu}} \langle f^s, \varphi \rangle = \frac{-2\pi i}{(\nu!)^2} \bigg(\int_{\mathbb{C}} |y_p|^{2(N_1\sigma_{p,\nu}+k_1)} \frac{\partial^{2\nu} \Phi_1 \eta_1}{\partial x_p^{\nu} \partial \bar{x}_p^{\nu}} (0, y_p; \sigma_{p,\nu}) dy_p d\bar{y}_p + \int_{\mathbb{C}} |z_p|^{2(N_2\sigma_{p,\nu}+k_2)} \frac{\partial^{2\nu} \Phi_2 \eta_2}{\partial w_p^{\nu} \partial \bar{w}_p^{\nu}} (z_p, 0; \sigma_{p,\nu}) dz_p d\bar{z}_p \bigg)
$$

The global section $|\pi^* f|^{2s} (\pi^* \varphi) |d\pi|^2$ restricted to U_p , V_p contains Φ_1 , Φ_2 . The whole global section differs only from Φ_1, Φ_2 by the exceptional part in the total transform $|\pi^* f|^{2s}$. Thus, at the intersection $U_p \cap V_p$, having that $y_p z_p = 1, w_p = x_p^{\kappa_p} y_p$, one checks that

$$
\Phi_2(z_p, w_p; \sigma_{p,\nu}) = \Phi_2(y_p^{-1}, x_p^{\kappa_p} y_p; \sigma_{p,\nu})
$$

= $|y_p|^{-2(\epsilon_{3,\nu} + \cdots + \epsilon_{r,\nu})} \Phi_2(y_p, x_p^{\kappa_p} y_p; \sigma_{p,\nu}) = |y_p|^{-2(\epsilon_{3,\nu} + \cdots + \epsilon_{r,\nu})} \Phi_1(x_p, y_p; \sigma_{p,\nu}).$

Now, deriving both sides with respect to w_p and \overline{w}_p and setting $w_p, \overline{w}_p = 0$ yields,

$$
\frac{\partial^{2\nu}\Phi_2}{\partial w_p^{\nu}\partial \overline{w}_p^{\nu}}(z_p,0;\sigma_{p,\nu})=|y_p|^{-2(\epsilon_{3,\nu}+\cdots+\epsilon_{r,\nu}+\kappa_p\nu)}\frac{\partial^{2\nu}\Phi_1}{\partial x_p^{\nu}\partial \overline{x}_p^{\nu}}(0,y_p;\sigma_{p,\nu}).
$$

.

This, together with Lemma [5.3,](#page-18-2) shows that the differential forms

$$
|y_p|^{2\epsilon_{1,\nu}}\frac{\partial^{2\nu}\Phi_1}{\partial^\nu x_p\partial^\nu \bar{x}_p}(0, y_p; \sigma_{p,\nu}) dy_p \wedge d\bar{y}_p, \quad |z_p|^{2\epsilon_{2,\nu}}\frac{\partial^{2\nu}\Phi_2}{\partial w_p^{\nu}\partial \bar{w}_p}(z_p, 0; \sigma_{p,\nu}) dw_p \wedge d\bar{w}_p,
$$

define a global section on E_p . As a consequence, it suffices to use $z_p y_p = 1$ in either of the integrals in Equation [\(5.9\)](#page-18-3), together with the fact that $\eta_1|_{E_p} + \eta_2|_{E_p} \equiv 1$.

Corollary 5.5. The residue of the complex zeta function f^s at $s = \sigma_{p,\nu}$ is given, in the U_p chart, by

(5.10)
$$
\operatorname{Res}_{s=\sigma_{p,\nu}} \langle f^s, \varphi \rangle = \frac{-2\pi i}{(\nu!)^2} \int_{|y_p| \le R} |y_p|^{2\epsilon_{1,\nu}} \frac{\partial^{2\nu} \Phi_1}{\partial x_p^{\nu} \partial \bar{x}_p^{\nu}} (0, y_p; \sigma_{p,\nu}) dy_p d\bar{y}_p + \frac{-2\pi i}{(\nu!)^2} \int_{|y_p| > R} |y_p|^{2\epsilon_{1,\nu}} \frac{\partial^{2\nu} \Phi_1}{\partial x_p^{\nu} \partial \bar{x}_p^{\nu}} (0, y_p; \sigma_{p,\nu}) dy_p d\bar{y}_p, \quad R > 0,
$$

and analogously for the other chart V_p .

Proof. The function η_1 can be chosen continuous and such that its restriction to E_p is $\eta_1 |_{E_p} \equiv 1$ in $|y_p| \leq R$ and 0 in $|y_p| > R$. Because $z_p y_p = 1$ on the overlap of any two charts of E_p , $\eta_2 |_{E_p}$ must be identically 1 in $|z_p| < 1/R$, i.e. $|y_p| > R$, and zero in the complementary. The results follows now from the proof of the previous proposition. Indeed, substitute such η_1 and η_2 in Equation [\(5.9\)](#page-18-3) and use the fact that the integrand is a global section on E_p .

Remark 5.6. The value of the residue $\text{Res}_{s=\sigma_{p,\nu}}\langle f^s, \varphi \rangle$ must be understood as the analytic continuation of the functions,

$$
I_1(\alpha_1, \beta_3, \dots, \beta_r) = \int_{\mathbb{C}} |y_p|^{2\alpha_1} \frac{\partial^{2\nu} \Phi_1}{\partial x_p^{\nu} \partial \bar{x}_p^{\nu}} (0, y_p; \beta_3, \dots, \beta_r) dy_p d\bar{y}_p,
$$

$$
I_2(\alpha_2, \beta_3, \dots, \beta_r) = \int_{\mathbb{C}} |z_p|^{2\alpha_2} \frac{\partial^{2\nu} \Phi_2}{\partial w_p^{\nu} \partial \bar{w}_p^{\nu}} (z_p, 0; \beta_3, \dots, \beta_r) dz_p d\bar{z}_p,
$$

at the rational points $(\epsilon_{1,\nu}, \epsilon_{3,\nu}, \ldots, \epsilon_{r,\nu})$ and $(\epsilon_{2,\nu}, \epsilon_{3,\nu}, \ldots, \epsilon_{r,\nu})$, respectively, as these points will usually be outside the region of convergence of the integrals defining I_1, I_2 . For simplicity of the notation, we always present Φ_1 , Φ_2 depending only on a single variable $\sigma_{p,\nu}$, as $\epsilon_{3,r}, \ldots, \epsilon_{r,\nu}$ are in fact $N_3\sigma_{p,\nu} + k_3 = \epsilon_{3,\nu}, \ldots, N_r\sigma_{p,\nu} + k_r = \epsilon_{r,\nu}$.

Finally, we end this section with the following important observation. As in the monomial case $\langle z^s, \varphi \rangle$ considered in Proposition [2.1,](#page-4-1) where the residue is interpreted in terms of the derivatives of the test function φ , and consequently, in terms of the Dirac's delta function, the same holds true for any f^s . The derivatives of Φ_1, Φ_2 involve deriving $\pi^*\varphi$ which, by the product rule of differentiation and the fact that $(\pi^*\varphi)|_{E_p} = \varphi(0)$, imply that

$$
(5.11) \qquad \text{Res}_{s=\sigma_{p,\nu}} f^s \in \left\langle \delta_0^{(0,0,0,0)}, \delta_0^{(1,0,0,0)}, \delta_0^{(0,1,0,0)}, \dots, \delta_0^{(\nu,\nu,\nu,\nu)} \right\rangle_{\mathbb{C}}.
$$

Therefore, the residue of f^s at any candidate pole must be also understood as a distribution in this precise sense. As a consequence, the residue of f^s at a candidate pole will be zero when all the coefficients of the linear combination in Equation [\(5.11\)](#page-19-1) are zero. In a similar way, the residue will be non-zero when just one of the coefficients is non-zero.

5.3. Residues at non-rupture divisors. The exact expression of the residue is quite involved due to the presence of the ν -th derivative of Φ_1 or Φ_2 . However, from the study of the derivatives of the factors of Φ_1 or Φ_2 we will show when the residues at a candidate $\sigma_{p,\nu}$ is zero for a non-rupture exceptional divisor E_p . The proof uses the following technical results.

Lemma 5.7 (Faà di Bruno's formula, [\[26,](#page-33-25) III.3.4]). Let g, h be infinitely many times differentiable functions. Then,

(5.12)
$$
\frac{d^{\nu}}{dx^{\nu}}g(h(x)) = \sum_{k=1}^{\nu} \frac{d^k g}{dx^k}(h(x)) B_{\nu,k} \left(\frac{dh}{dx}(x), \frac{d^2 h}{dx^2}(x), \dots, \frac{d^{\nu-k+1} h}{dx^{\nu-k+1}}(x) \right),
$$

where $B_{\nu,k}$ are the partial exponential Bell polynomials

$$
B_{\nu,k}(x_1, x_2, \dots, x_{\nu-k+1}) := \sum \frac{\nu!}{j_1! j_2! \cdots j_{\nu-k+1}!} \left(\frac{x_1}{1!}\right)^{j_1} \left(\frac{x_2}{2!}\right)^{j_2} \cdots \left(\frac{x_{\nu-k+1}}{(\nu-k+1)!}\right)^{j_{\nu-k+1}},
$$

and where the summation takes places over all integers $j_1, j_2, j_3, \ldots, j_{\nu-k+1}$, such that

(5.13)
$$
j_1 + j_2 + j_3 + \cdots + j_{\nu-k+1} = k,
$$

$$
j_1 + 2j_2 + 3j_3 + \cdots + (\nu - k + 1)j_{\nu-k+1} = \nu.
$$

For instance, in the chart U_p around E_p , we are interested in the situation where $g(x) = x^s$ and $h(x_p)$ is equal to $u_1(x_p, y_p)$ from Proposition [5.1,](#page-17-2) and we set $x_p = 0$ after deriving. In this case, Equation [\(5.12\)](#page-20-0) reads as (5.14)

$$
\frac{\partial^{\nu} u_1^s}{\partial x_p^{\nu}}(0, y_p) = \sum_{k=1}^{\nu} (s)_k (u_1(0, y_p))^{s-k} B_{\nu,k} \left(\frac{du_1}{dx_p}(0, y_p), \frac{d^2 u_1}{dx_p^2}(0, y_p), \dots, \frac{d^{\nu-k+1} u_1}{dx_p^{\nu-k+1}}(0, y_p) \right),
$$

where $(s)_k := s(s-1)\cdots(s-k+1)$. And similarly in the other chart V_p .

Proposition 5.8 ([\[32,](#page-33-1) I.3.8]). For any $\alpha, \alpha' \in \mathbb{C}$ such that $\alpha' - \alpha = n \in \mathbb{Z}$, the analytic continuation of the sum

$$
I_n(\alpha) := \int_{|z| \le R} z^{\alpha'} \overline{z}^{\alpha} dz d\overline{z} + \int_{|z| > R} z^{\alpha'} \overline{z}^{\alpha} dz d\overline{z} \quad \text{for any} \quad R > 0.
$$

is zero everywhere, i.e. $I_n(\alpha) \equiv 0$.

Proof. Using polar coordinates^{[1](#page-20-1)}

$$
-2i\int_0^R \int_0^{2\pi} r^{2\alpha+n+1} e^{2\pi i n\theta} d\theta dr - 2i\int_R^{\infty} \int_0^{2\pi} r^{2\alpha+n+1} e^{2\pi i n\theta} d\theta dr.
$$

However,

$$
\int_0^{2\pi} e^{2\pi i n\theta} d\theta = \begin{cases} 0, & n \neq 0, \\ 2\pi, & n = 0. \end{cases}
$$

Hence, the result follows if $n \neq 0$. In the case that $n = 0$, the first integral defines an holomorphic function in $\text{Re}(\alpha) > -1$. It can be analytically continued by means of

$$
-4\pi i \int_0^R r^{2\alpha+1} dr = -2\pi i \frac{R^{2(\alpha+1)}}{\alpha+1} \quad \text{for} \quad \alpha \neq -1.
$$

¹By definition, $z^{\alpha'}\overline{z}^{\alpha} := |z|^{\alpha' + \alpha}e^{i(\alpha' - \alpha) \arg z}$, which, for integral $\alpha' - \alpha$, is a single valued function of z.

Similarly, the other integral defines an holomorphic functions in $\text{Re}(\alpha) < -1$, and the analytic continuation to the whole complex plane is

$$
-4\pi i \int_{R}^{\infty} r^{2\alpha+1} dr = 2\pi i \frac{R^{2(\alpha+1)}}{\alpha+1} \quad \text{for} \quad \alpha \neq -1.
$$

Finally, the sum of the analytic continuation of both integrals is identically zero. \Box

In the following proposition, we generalize a calculation attributed to Cohen appearing in an article of Barlet [\[5\]](#page-32-15) and used by Lichtin in [\[48\]](#page-34-6). The original result gives a closed form for the integral in Equation [\(5.15\)](#page-21-0) in the case $R_{0,0}(\alpha, \beta)$. We provide a formula for the general case $R_{n,m}(\alpha,\beta)$.

Proposition 5.9. For $\alpha, \alpha', \beta, \beta' \in \mathbb{C}$, such that $\alpha' - \alpha = n \in \mathbb{Z}$ and $\beta' - \beta = m \in \mathbb{Z}$, the integral

(5.15)
$$
R_{n,m}(\alpha,\beta) := \int_{\mathbb{C}} z^{\alpha'} \bar{z}^{\alpha} (1 - \lambda z)^{\beta'} (1 - \bar{\lambda} \bar{z})^{\beta} dz d\bar{z} = R_{-n,-m}(\alpha',\beta'), \quad \lambda \in \mathbb{C}^*,
$$

is absolutely convergent for $\text{Re}(\alpha' + \alpha) > -2$, $\text{Re}(\beta' + \beta) > -2$ and $\text{Re}(\alpha' + \alpha + \beta' + \beta) < -2$. It defines a meromorphic function on \mathbb{C}^2 equal to

(5.16)
$$
R_{n,m}(\alpha,\beta) = -2\pi i \lambda^{-\alpha'-1} \bar{\lambda}^{-\alpha-1} \frac{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(\gamma+1)}{\Gamma(-\alpha-n)\Gamma(-\beta-m)\Gamma(-\gamma-n-m)},
$$

where $\gamma := -\alpha - \beta - n - m - 2$.

Proof. Let us prove first the case $m = 0$. Since $R_{n,0}(\alpha, \beta) = R_{-n,0}(\alpha', \beta')$, we can assume $n \in \mathbb{Z}_{\geq 0}$. Using polar coordinates $\lambda z = re^{i\theta}$, we have that $(1-\lambda z)^{\beta'}(1-\bar{\lambda}\bar{z})^{\beta} = |1-\lambda z|^{2\beta} =$ $(1 - 2r \cos \theta + r^2)^{\beta}$ and

$$
R_{n,0}(\alpha,\beta) = -2i\lambda^{-\alpha'-1}\overline{\lambda}^{-\alpha-1}\int_0^{2\pi}\int_0^{\infty}r^{2\alpha+n+1}e^{in\theta}(1-2r\cos\theta+r^2)^{\beta}drd\theta.
$$

For simplicity, we can set $\lambda = 1$. We have, $(1 - 2r \cos \theta + r^2)^{\beta} = (1 + r^2)^{\beta}$ 1 − $2r\cos\theta$ $1 + r^2$ \setminus^{β} and since $|2r/(1+r^2)| \leq 1$, we may expand the binomial,

$$
\left(1 - \frac{2r\cos\theta}{1+r^2}\right)^{\beta} = \sum_{k=0}^{\infty} {\binom{\beta}{k}} \frac{(-2)^k r^k}{(1+r^2)^k} \cos^k\theta.
$$

The angular part of the integral is just

$$
\int_0^{2\pi} e^{in\theta} \cos^k \theta \, d\theta = \begin{cases} \frac{2\pi}{4^l} {2l \choose l-s}, & \text{if } k = 2l \ge n = 2s, \quad l \in \mathbb{Z}_{\ge 0}, s \in \mathbb{Z}_{\ge 0}, \\ \frac{\pi}{4^l} {2l+1 \choose l-s}, & \text{if } k = 2l+1 \ge n = 2s+1, \quad l \in \mathbb{Z}_{\ge 0}, s \in \mathbb{Z}_{\ge 0}, \\ 0, & \text{otherwise.} \end{cases}
$$

The integral then reads as

$$
R_{2s,0}(\alpha,\beta) = -4\pi i \int_0^\infty r^{2\alpha+2s+1} (1+r^2)^\beta \sum_{l=0}^\infty \binom{2l}{l-s} \binom{\beta}{2l} \frac{r^{2l}}{(1+r^2)^{2l}} dr,
$$

for n even, and

$$
R_{2s+1,0}(\alpha,\beta) = 4\pi i \int_0^\infty r^{2\alpha+2s+2} (1+r^2)^\beta \sum_{l=0}^\infty \binom{2l+1}{l-s} \binom{\beta}{2l+1} \frac{r^{2l+1}}{(1+r^2)^{2l+1}} dr,
$$

for *n* odd. Using that $\begin{pmatrix} \beta & \beta \\ 0 & \beta \end{pmatrix}$ k $= (-1)^k \frac{\Gamma(k-\beta)}{\Gamma(-\beta) \Gamma(k-\beta)}$ $\Gamma(-\beta)k!$ and interchanging the summation and integral signs,

$$
R_{2s,0}(\alpha,\beta) = \frac{-4\pi i}{\Gamma(-\beta)} \sum_{l=0}^{\infty} \frac{\Gamma(2l-\beta)}{(l-s)!(l+s)!} \int_0^{\infty} r^{2\alpha+2s+2l+1} (1+r^2)^{\beta-2l} dr,
$$

$$
R_{2s+1,0}(\alpha,\beta) = \frac{-4\pi i}{\Gamma(-\beta)} \sum_{l=0}^{\infty} \frac{\Gamma(2l+1-\beta)}{(l-s)!(l+s+1)!} \int_0^{\infty} r^{2\alpha+2s+2l+3} (1+r^2)^{\beta-2l-1} dr.
$$

Now, for $\text{Re}(\mu) > 0$ and $\text{Re}(2\nu + \mu) < 0$,

$$
\int_0^\infty x^{\mu-1} (1+x^2)^\nu dx = \frac{1}{2} \mathcal{B}\left(\frac{\mu}{2}, -\nu - \frac{\mu}{2}\right) = \frac{1}{2} \Gamma\left(\frac{\mu}{2}\right) \Gamma\left(-\nu - \frac{\mu}{2}\right) \Gamma(-\nu)^{-1}.
$$

See, for instance, [\[33,](#page-33-26) 3.251–2]. For $\mu_l = 2(\alpha + s + l + 1)$ and $\nu_l = \beta - 2l$,

$$
R_{2s}(\alpha,\beta) = \frac{-2\pi i}{\Gamma(-\beta)} \sum_{l=0}^{\infty} \frac{\Gamma(\alpha+s+l+1)\Gamma(l-\alpha-s-\beta-1)}{(l-s)!(l+s)!},
$$

since Re(μ_l) > 0 and Re($2\nu_l + \mu_l$) < 0 for all $l \in \mathbb{Z}_{\geq 0}$. Analogously, for $n \geq 0$ odd, $\mu_l = 2(\alpha + s + l + 2)$ and $\nu_l = \beta - 2l - 1$,

$$
R_{2s+1,0}(\alpha,\beta) = \frac{-2\pi i}{\Gamma(-\beta)} \sum_{l=0}^{\infty} \frac{\Gamma(\alpha+s+l+2)\Gamma(l-\alpha-s-\beta-1)}{(l-s)!(l+s+1)!}.
$$

Since $\Gamma(-k)^{-1} = 0$ for $k \in \mathbb{Z}_{\geq 0}$, we can write

$$
R_{2s,0}(\alpha,\beta) = \frac{-2\pi i}{\Gamma(-\beta)} \sum_{l=0}^{\infty} \frac{\Gamma(\alpha+l+1)\Gamma(l-\alpha-2s-\beta-1)}{\Gamma(l-2s+1)l!},
$$

$$
R_{2s+1,0}(\alpha,\beta) = \frac{-2\pi i}{\Gamma(-\beta)} \sum_{l=0}^{\infty} \frac{\Gamma(\alpha+l+1)\Gamma(l-\alpha-2s-\beta-2)}{\Gamma(l-2s)l!}.
$$

Finally, for $\text{Re}(c) > \text{Re}(a+b)$,

(5.17)
$$
\sum_{k=0}^{\infty} \frac{\Gamma(a+k)\Gamma(b+k)}{\Gamma(c+k)k!} = \frac{\Gamma(a)\Gamma(b)}{\Gamma(c)} {}_2F_1(a,b;c;1) = \frac{\Gamma(a)\Gamma(b)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)},
$$

where ${}_2F_1(a, b; c; z)$ is the hypergeometric function. For the last equality see, for instance, [\[33,](#page-33-26) 9.122–1]. For $n = 2s$, set $a = \alpha + 1$, $b = -\alpha - \beta - 2s - 1$ and $c = -2s + 1$. Similarly, for $n = 2s + 1$, $a = \alpha + 1$, $b = -\alpha - \beta - 2s - 2$ and $c = -2s$. Then,

$$
R_{2s,0}(\alpha,\beta) = R_{2s+1,0}(\alpha,\beta) = -2\pi i \frac{\Gamma(\alpha+1)\Gamma(-\alpha-\beta-n-1)\Gamma(\beta+1)}{\Gamma(-\beta)\Gamma(-n-\alpha)\Gamma(\alpha+\beta+2)}.
$$

For the case where $m \neq 0$, having proved the result for $n \in \mathbb{Z}, m = 0$, we can assume without loss of generality that $m \in \mathbb{Z}_{\geq 0}$. Keeping $\lambda = 1$, notice that

$$
R_{n,m}(\alpha,\beta) = \int_{\mathbb{C}} z^{\alpha'} \overline{z}^{\alpha} |1-z|^{2\beta} (1-z)^m dz d\overline{z} = \sum_{j=0}^m \binom{m}{j} (-1)^j R_{n+j,0}(\alpha,\beta).
$$

Hence,

$$
R_{n,m}(\alpha,\beta) = -2\pi i \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{\Gamma(-\beta)\Gamma(\alpha+\beta+2)} \sum_{j=0}^{m} {m \choose j} (-1)^j \frac{\Gamma(-\alpha-\beta-n-j-1)}{\Gamma(-n-j-\alpha)}.
$$

Since the binomial coefficient is zero if $j > m$, we can consider the infinite sum. Expanding the binomial coefficient in terms of the Gamma function

$$
R_{n,m}(\alpha,\beta) = -2\pi i \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{\Gamma(-\beta)\Gamma(\alpha+\beta+2)\Gamma(-m)} \sum_{j=0}^{\infty} \frac{\Gamma(j-m)\Gamma(-\alpha-\beta-n-j-1)}{\Gamma(-n-j-\alpha) j!}.
$$

Using the functional equation $\Gamma(z+1) = z\Gamma(z)$ at each term

$$
R_{n,m}(\alpha,\beta) = -2\pi i \frac{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(-\alpha-\beta-n-1)\Gamma(\alpha+\beta+n+2)}{\Gamma(-\beta)\Gamma(\alpha+\beta+2)\Gamma(-m)\Gamma(-\alpha-n)\Gamma(\alpha+n+1)}
$$

$$
\cdot \sum_{j=0}^{\infty} \frac{\Gamma(j-m)\Gamma(\alpha+n+j+1)}{\Gamma(\alpha+\beta+n+j+2)j!}.
$$

Applying Equation [\(5.17\)](#page-22-0) once again, since $\text{Re}(\beta' + \beta) > -2$ implies $\text{Re}(\beta + m) > -1$,

$$
R_{n,m}(\alpha,\beta) = -2\pi i \frac{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(-\alpha-\beta-n-1)\Gamma(\alpha+\beta+n+2)\Gamma(\beta+m+1)}{\Gamma(-\beta)\Gamma(\alpha+\beta+2)\Gamma(-\alpha-n)\Gamma(\alpha+\beta+n+m+2)\Gamma(\beta+1)}.
$$

And we get the desired result using the functional equation $\Gamma(z+1) = z\Gamma(z)$ once again. П

It is now possible to prove the following result regarding non-rupture divisors. Recall that with the notation above, the divisors crossing a non-rupture exceptional divisor E_p can only be D_1, D_2, D_3 , with at least one being non-zero and one being zero.

Theorem 5.10. Let $f : (\mathbb{C}^2, \mathbf{0}) \longrightarrow (\mathbb{C}, 0)$ be any plane branch. Let $E_p, p \in K$ be a non-rupture exceptional divisor with sequence of candidate poles $\sigma_{p,\nu}, \nu \in \mathbb{Z}_{\geq 0}$. Then,

\n- If
$$
D_3 = 0
$$
, $\underset{s = \sigma_{p,\nu}}{\text{Res }} f^s = 0$, for all $\nu \in \mathbb{Z}_{\geq 0}$.
\n- If $D_3 \neq 0$, $\underset{s = \sigma_{p,\nu}}{\text{Res }} f^s = 0$, if $\epsilon_{3,\nu} \notin \mathbb{Z}$.
\n

Proof. Let us first begin with D_3 non-zero. In this case, we must also have D_1 or D_2 non-zero. We can assume, for instance, D_1 non-zero.

By the definition of D_1, D_2, \ldots, D_r , if D_1 and D_3 are non-zero, D_3 is the only divisor crossing E_p in the V_p chart. In the (z_p, w_p) coordinates this means that $u_2^s(z_p, 0)v_2(z_p, 0)$ has the form $(1-\lambda y_p)^{N_3s+k_3}$ for some $\lambda \in \mathbb{C}^*$. By Faà di Bruno's formula in Lemma [5.7](#page-20-2) and Equation [\(5.14\)](#page-20-3), the *ν*-th holomorphic and antiholomorphic derivatives of Φ_2 at z_p , \overline{z}_p = $0, s = \sigma_{p,\nu}$ is an algebraic function involving the terms $z_p^{k'}$ $\frac{k'}{p} \bar{z}_p^k (1-\lambda z_p)^{\epsilon_{3,\nu}-l'}(1-\bar{\lambda}\bar{y}_p)^{\epsilon_{3,\nu}-l'}$

with $k, k', l, l' \in \mathbb{Z}_{\geq 0}$. Hence, the residue reduces to a finite sum involving the integrals $R_{n,m}(\alpha,\beta)$ from Equation [\(5.15\)](#page-21-0) which, by Proposition [5.9,](#page-21-1) are proportional to

$$
\frac{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(\gamma+1)}{\Gamma(-\alpha-n)\Gamma(-\beta-m)\Gamma(-\gamma-n-m)}, \quad n, m \in \mathbb{Z}.
$$

Applying Lemma [5.3](#page-18-2) to this situation implies that, $\epsilon_{1,\nu} + \epsilon_{3,\nu} + \kappa_{p} \nu + 2 = 0$. Notice that $\epsilon_{3,\nu} \notin \mathbb{Z}$ implies $\epsilon_{1,\nu} \notin \mathbb{Z}$. Since the poles of $\Gamma(z)$ are located at the negative integers, and $\Gamma(z)$ has no zeros, $\Gamma(\beta+1), \Gamma(\gamma+1), \Gamma(-\beta-m)^{-1}, \Gamma(-\gamma-n-m)^{-1} \in \mathbb{C}^*$. Finally, consider the non-negative integers k, k' in $z_p^{k'}$ $_{p}^{k'}\bar{z}_{p}^{k}$ above. The quotient $\Gamma(\alpha+1)/\Gamma(-\alpha-n)$ is zero since $\epsilon_{2,\nu} = 0$, and $\alpha + 1 = k + 1 > 0$, $-\alpha - n = -\alpha' = -k' \leq 0$.

If D_3 is zero and E_p is a non-rupture divisor, it may happen that D_2 is zero or not. If D_2 is not zero, it must cross E_p in the only point of V_p not in U_p . In both cases, in the chart U_p , none of the components u_1^s, v_1 of Φ_1 will cross E_p . In the (x_p, y_p) coordinates this means that, $u_1^s(0, y_p)v_1(0, y_p) \in \mathbb{C}^*$ for all $y_p \in \mathbb{C}$. Applying Faà di Bruno's formula again, the holomorphic and antiholomorphic derivatives of Φ_1 restricted to $x_p, \bar{x}_p = 0$ are just polynomials in y_p and \bar{y}_p . By Corollary [5.5](#page-19-2) and Proposition [5.8,](#page-20-4) the residue is zero. \Box

Example 1. There are example where $D_3 \neq 0$ and $\epsilon_{3,\nu} \in \mathbb{Z}$ and the corresponding $\sigma_{p,\nu}$ has non-zero residue. For instance, $f = (y^2 - x^3)(y - x^2)^3$. Its resolution is given by

$$
F_{\pi} = 5E_{p_0} + 9E_{p_1} + 15E_{p_2} + C_1 + 3C_2, \quad K_{\pi} = E_{p_0} + 2E_{p_1} + 4E_{p_2}.
$$

In this case, E_{p_1} is a non-rupture exceptional divisor with E_{p_2} and C_2 crossing E_{p_1} . It can be checked that $\sigma_{p_1,0} = -1/3$ is a pole of order two of f^s . Here, $D_1 = E_{p_2}, D_3 = C_2$ and $\epsilon_{1,0} = \epsilon_{3,0} = -1$.

6. The set of poles of the complex zeta function of a plane branch

In this section, we restrict our study of the poles of the complex zeta function to the case of plane branch singularities. Through the rest of this work we will fix a plane branch semigroup $\Gamma = \langle \beta_0, \beta_1, \ldots, \beta_g \rangle$ and we stick to the notations of Section [3.1.](#page-8-4) Instead of taking any germ $f: (\mathbb{C}^2, \mathbf{0}) \longrightarrow (\mathbb{C}, 0)$ with semigroup Γ , we will work with the family ${f_{t,\lambda}}_{\lambda \in \mathbb{C}^{g-1}}$ presented in Section [3.2.](#page-9-4) Since this family contains at least one representative for each analytic type in the equisingularity class of the semigroup Γ , we can give an optimal set of candidates for the poles of the complex zeta function f^s of all plane branches with semigroup Γ. Furthermore, we will prove that if f_{gen} is generic among all branches with semigroup Γ (in the sense that the coefficients of a μ -constant deformation are generic), all the candidates are indeed poles of f_{gen}^s . As a corollary, we prove Yano's conjecture under the assumption that eigenvalues of the monodromy are pairwise different.

6.1. Residues at rupture divisors. After Theorem [5.10](#page-23-0) in the preceding section, the only divisors that will contribute to poles of the complex zeta function of a plane branch will be rupture divisors and the divisor of the strict transform. Following the discussion in Section [4.2,](#page-12-1) a singular plane branch will have exactly $q \geq 1$ rupture divisors, denoted E_{p_1}, \ldots, E_{p_g} , where g is the number of characteristic exponents of f_t . This first observation reduces the list of candidate poles to $\sigma_{i,\nu}, \nu \in \mathbb{Z}$ for $i = 1, \ldots, g$ contributed by E_{p_i} , in addition to the negative integers corresponding to the strict transform \tilde{C} . With the notations from Equation [\(4.3\)](#page-13-1),

$$
\sigma_{i,\nu} = -\frac{m_i + n_1 \cdots n_i + \nu}{n_i \overline{\beta}_i}, \quad \nu \in \mathbb{Z}_{\geq 0}.
$$

We will basically use the same results about the residue presented in Section [5.2.](#page-16-1) However, in this case, a better understanding of the total transform around the rupture divisors is needed. To that end, we will make use of Proposition [4.1.](#page-13-0) The only thing that will differ from Section [5.2](#page-16-1) is that, instead of computing the residue on the minimal resolution surface X' , we proceed inductively and resolve up to the i -th rupture divisor, compute the residue on $X^{(i)}$ and blow-up up to the $(i + 1)$ –th rupture divisor. This process simplifies the notation and fits more naturally with the toric resolutions from Section [4.2.](#page-12-1)

First of all, we have to analyze the residue numbers from Equation [\(5.6\)](#page-18-4) in the case of a rupture divisor of a plane branch. In this case, Equation [\(5.7\)](#page-18-1) is $\epsilon_{1,\nu}+\epsilon_{2,\nu}+\epsilon_3+\kappa_{p_i}\nu+2=0$, since rupture divisors of plane branches only have three divisor, D_1, D_2, D_3 , crossing them. Since we will be working on $X^{(i)}$, the divisor previously written as D_3 is the strict transform \tilde{C} of f on $X^{(i)}$ and thus, $\epsilon_3 = e_i \sigma_{i,\nu}$, because $E_{p_i} \cdot \tilde{C} = e_i$. Similarly, on the surface $X^{(i)}$, $\kappa_{p_i} = -E_{p_i} \cdot E_{p_i} = 1$. Hence,

(6.1)
$$
\epsilon_{1,\nu} + \epsilon_{2,\nu} + e_i \sigma_{i,\nu} + \nu + 2 = 0.
$$

It is then enough to study the relation of $\epsilon_{1,\nu}, \epsilon_{2,\nu}, \nu \in \mathbb{Z}_{\geq 0}$ with the semigroup Γ. We point out that Lichtin [\[47\]](#page-34-5) studied the residue numbers $\epsilon_{1,\nu}$, $\epsilon_{2,\nu}$ for the case $\nu = 0$. Using the notations and definitions from Section [3.1,](#page-8-4)

Proposition 6.1 ([\[47,](#page-34-5) Prop. 2.12]). The residue numbers $\epsilon_{1,0}, \epsilon_{2,0}$ associated to a rupture divisor E_{p_i} of plane branch are given by

$$
\epsilon_{1,0} + 1 = \frac{1}{n_i}, \quad \epsilon_{2,0} + 1 = \frac{m_{i-1} - n_{i-1} \overline{m}_{i-1} + n_1 \cdots n_{i-1}}{\overline{m}_i}.
$$

Corollary 6.2. For any $\nu \in \mathbb{Z}_{\geq 0}$, the residue numbers are

$$
\epsilon_{1,\nu} + 1 = -\frac{a_i}{n_i} \nu + \frac{1}{n_i}, \quad \epsilon_{2,\nu} + 1 = -\frac{c_i n_{i-1} \overline{m}_{i-1} + d_i}{\overline{m}_i} \nu + \frac{m_{i-1} - n_{i-1} \overline{m}_{i-1} + n_1 \cdots n_{i-1}}{\overline{m}_i}.
$$

Proof. The result follows from the definition of $\epsilon_{1,\nu}$, $\epsilon_{2,\nu}$ in Equation [\(5.6\)](#page-18-4) and using Propo-sition [6.1](#page-25-0) together with the expression of N_1 and N_2 given in Proposition [4.1.](#page-13-0)

Secondly, we focus our attention on the functions Φ_1, Φ_2 from Proposition [5.1](#page-17-2) but now in the case of the family f_t . Around the rupture divisor E_{p_i} on $X^{(i)}$, we have that $\Phi_{1,t} = |u_1|^{2s} |v_1|^2 |\tilde{f}_t|^{2s} (\pi^{(i)})^* \varphi |_{U_i}$ and $\Phi_{2,t} = |u_2|^{2s} |v_2|^s |\tilde{f}_t|^{2s} (\pi^{(i)})^* \varphi |_{V_i}$. Since the only factor of $\Phi_{1,t}, \Phi_{2,t}$ crossing E_{p_i} is the strict transform \tilde{f}_t , by Faà di Bruno's formula in Lemma [5.7](#page-20-2) applied to the equations in Proposition [4.1,](#page-13-0) the ν -th derivatives of $\Phi_{1,t}$, $\Phi_{2,t}$ at $x_p, \bar{x}_p = 0, w_p, \bar{w}_p = 0$ with $s = \sigma_{i,\nu}$ are a finite sum with summands that look like

$$
y_p^{k'}\bar{y}_p^{k}(y_i-\lambda_i)^{e_i(\sigma_{i,\nu}-l')}(\bar{y}_i-\bar{\lambda}_i)^{e_i(\sigma_{i,\nu}-l)},\quad z_p^{k'}\bar{z}_p^{k}(1-\lambda_i z_p)^{e_i(\sigma_{i,\nu}-l')}(1-\bar{\lambda}_i\bar{z}_p)^{e_i(\sigma_{i,\nu}-l)},
$$

with $k', k, l', l \in \mathbb{Z}_{\geq 0}$. Therefore, it make sense to consider the order and the degree of y_i, z_i in $\Phi_{1,t}, \Phi_{2,t}$. On the U_i chart, they will be denoted by

$$
0 \leq \text{ord}_{y_i} \frac{\partial^{\nu} \Phi_{1,t}}{\partial x^{\nu} \partial \bar{x}^{\nu}} (0, y_i; \sigma_{i,\nu}) \leq \text{deg}_{y_i} \frac{\partial^{\nu} \Phi_{1,t}}{\partial x^{\nu} \partial \bar{x}^{\nu}} (0, y_i; \sigma_{i,\nu}),
$$

and respectively on the V_i chart. By the symmetry of the holomorphic and antiholomorphic parts, the orders and degrees are exactly the same if considered with respect to the conjugated variables \bar{y}_i, \bar{z}_i . By convention the order and degree of zero are $+\infty$ and 0, respectively. Let us first present some technical lemmas.

Lemma 6.3. Let $f_1, \ldots, f_n \in C^\infty(\mathbb{C}^2)$ be functions such that

$$
\alpha \nu \leq \text{ord}_y \frac{\partial^{\nu} f_i}{\partial x^{\nu}}(0, y), \quad \deg_y \frac{\partial^{\nu} f_i}{\partial x^{\nu}}(0, y) \leq \beta \nu, \quad \text{for} \quad i = 1, \dots, n,
$$

and $\alpha, \beta \in \mathbb{Q}_{>0}$. Then,

$$
\alpha \nu \leq \text{ord}_y \frac{\partial^{\nu} (f_1 \cdots f_n)}{\partial x^{\nu}}(0, y), \quad \deg_y \frac{\partial^{\nu} (f_1 \cdots f_n)}{\partial x^{\nu}}(0, y) \leq \beta \nu,
$$

and,

$$
\alpha\nu \leq \text{ord}_y \frac{\partial^{\nu} f_1^s}{\partial x^{\nu}}(0, y), \quad \deg_y \frac{\partial^{\nu} f_1^s}{\partial x^{\nu}}(0, y) \leq \beta \nu \quad \text{for all} \quad s \in \mathbb{C}.
$$

Proof. The first inequalities follow from the general Leibniz rule,

$$
\frac{\partial^{\nu}(f_1 \cdots f_n)}{\partial x^{\nu}}(0, y) = \sum_{k_1 + \cdots + k_n = \nu} {n \choose k_1, \ldots, k_n} \frac{\partial^{k_1} f_1}{\partial x^{k_1}}(0, y) \cdots \frac{\partial^{k_n} f_n}{\partial x^{k_n}}(0, y).
$$

Similarly, the second follow from Faà di Bruno's formula and the definition of the partial exponential Bell polynomials, see Lemma [5.7](#page-20-2) and Equation [\(5.14\)](#page-20-3). Specifically, they follow from the second part of Equation (5.13) .

Lemma 6.4. Let $f(x, y) \in C^{\infty}(\mathbb{C}^2)$ and let $\pi(x, y) = (x^n y^a, x^m y^b)$ with $n, m, a, b \in \mathbb{Z}_{\geq 0}$. Furthermore, assume that $nb - ma \geq 0$. Then,

$$
\frac{a}{n}\nu \leq \operatorname{ord}_{y} \frac{\partial^{\nu}(f \circ \pi)}{\partial x^{\nu}}(0, y), \quad \deg_{y} \frac{\partial^{\nu}(f \circ \pi)}{\partial x^{\nu}}(0, y) \leq \frac{b}{m}\nu \quad \text{for all} \quad \nu \in \mathbb{Z}_{\geq 0}.
$$

Proof. Consider the Taylor expansion of f at the origin of order $\tau > \nu$,

$$
f(x,y) = \sum_{i,j=0}^{\tau-1} \frac{\partial^{i+j} f}{\partial x^i \partial y^i} (0,0) \frac{x^i}{i!} \frac{y^j}{j!} + R_\tau(x,y) x^\tau y^\tau,
$$

where R_{τ} is the residual. Composing with π , we get

$$
f(\pi(x,y)) = \sum_{i,j=0}^{\tau-1} \frac{\partial^{i+j} f}{\partial x^i \partial y^i} (0,0) \frac{x^{ni+mj}}{i!} \frac{y^{ai+bj}}{j!} + R_\tau(x^n y^a, x^m y^b) x^{(n+m)\tau} y^{(a+b)\tau}.
$$

At the *v*-th derivative of this Taylor polynomial with respect to x and restricted to $x = 0$, we must have that $ni+mj = \nu$ for some integers $i, j \geq 0$. Notice that if there are no such $i, j \geq 0$ the derivative is zero, and the bounds are trivially fulfilled. Finally,

$$
\operatorname{ord}_y \frac{\partial^{\nu} (f \circ \pi)}{\partial x^{\nu}} (0, y) = \min_{ni + mj = \nu} \{ ai + bj \} \ge \min_{j \ge 0} \left\{ \frac{a}{n} \nu + \frac{(nb - ma)j}{n} \right\} \ge \frac{a}{n} \nu,
$$

$$
\deg_y \frac{\partial^{\nu} (f \circ \pi)}{\partial x^{\nu}} (0, y) = \max_{ni + mj = \nu} \{ ai + bj \} \le \max_{i \ge 0} \left\{ \frac{b}{m} \nu - \frac{(nb - ma)i}{m} \right\} \le \frac{b}{m} \nu,
$$

where in the first inequality of each equation used that $i = (\nu - mj)/n$ and $j = (\nu - ni)/m$, respectively. \Box

Recall the linear forms $\rho_{j+1}^{(i)}(\underline{k}), A_{j+1}^{(i)}(\underline{k}), C_{j+1}^{(i)}(\underline{k})$ from Proposition [4.1.](#page-13-0)

Lemma 6.5. For any $\nu \in \mathbb{Z}_{\geq 0}$, we have

$$
\frac{a_i}{n_i} \nu \le \min_{\rho_{j+1}^{(i)}(\underline{k})=\nu} A_{j+1}^{(i)}(\underline{k}), \qquad \frac{c_i n_{i-1} \overline{m}_{i-1} + d_i}{\overline{m}_i} \nu \le \min_{\rho_{j+1}^{(i)}(\underline{k})=\nu} C_{j+1}^{(i)}(\underline{k}),
$$

\n
$$
\max_{\rho_{j+1}^{(i)}(\underline{k})=\nu} A_{j+1}^{(i)}(\underline{k}) \le \frac{a_i n_{i-1} \overline{m}_{i-1} + b_i}{\overline{m}_i} \nu + n_{i+1} \cdots n_j, \quad \max_{\rho_{j+1}^{(i)}(\underline{k})=\nu} C_{j+1}^{(i)}(\underline{k}) \le \frac{c_i}{n_i} \nu + n_{i+1} \cdots n_j.
$$

Proof. For the first inequality, solve for k_0 in the constrain $\rho_{j+1}^{(i)}(k_0,\ldots,k_j) = \nu$ and substitute in $A_{j+1}^{(i)}(\underline{k})$. After some cancellations we get

(6.2)
$$
\frac{a_i}{n_i} \nu + a_i n_{i-1} \overline{m}_{i-1} k_i + b_i k_i - \frac{a_i}{n_i} \overline{m}_i k_i = \frac{a_i}{n_i} \nu + \frac{k_i}{n_i},
$$

where in the equality we applied $\overline{m}_i = n_i n_{i-1} \overline{m}_{i-1} + q_i$, from Equation [\(3.3\)](#page-8-2), and that $n_i b_i - a_i q_i = 1$. Since $k_i \geq 0$, we obtain the lower bound for the minimum of $A_{j+1}^{(i)}(\underline{k})$ restricted to $\rho_{j+1}^{(i)}(\underline{k}) = \nu$. The argument for the lower bound for the minimum of $C_{j+1}^{(i)}(\underline{k})$ works similarly. Instead, solve for k_i in the constrain $\rho_{j+1}^{(i)}(k_0,\ldots,k_i,\ldots,k_j) = \nu$ and substitute in $C_{j+1}^{(i)}(\underline{k})$. Applying Equation [\(3.3\)](#page-8-2) when necessary and that $q_i c_i - n_i d_i = 1$, we obtain

(6.3)
$$
\frac{c_i n_{i-1} \overline{m}_{i-1} + d_i}{\overline{m}_i} \nu + \frac{1}{\overline{m}_i} \sum_{l=0}^{i-1} n_{l+1} \cdots n_i \overline{m}_l k_l,
$$

which gives again the lower bound since $k_l \geq 0$. Having obtained the lower bounds for the minimums of $A_{j+1}^{(i)}$ and $C_{j+1}^{(i)}$ we can use Lemma [4.2](#page-15-2) to obtain the upper bounds for the maximums. Indeed,

$$
A_{j+1}^{(i)}(\underline{k})+C_{j+1}^{(i)}(\underline{k})\leq \rho_{j+1}^{(i)}(\underline{k})+n_{i+1}\cdots n_j,
$$

since $k_l \geq 0$. Hence,

$$
\max_{\rho_{j+1}^{(i)}(\underline{k})=\nu} A_{j+1}^{(i)}(\underline{k}) \leq \nu + n_{i+1} \cdots n_j - \min_{\rho_{j+1}^{(i)}(\underline{k})=\nu} C_{j+1}^{(i)}(\underline{k}) = \frac{a_i n_{i-1} \overline{m}_{i-1} + b_i}{\overline{m}_i} \nu + n_{i+1} \cdots n_j,
$$

since $a_i + c_i = n_i$ and $b_i + d_i = q_i$. We can argue similarly for the remaining bound. \square

All these technical lemmas are used in the proof of the following proposition.

Proposition 6.6. With the notations above, we have that

$$
\frac{a_i}{n_i} \nu \le \text{ord}_{y_i} \frac{\partial^{2\nu} \Phi_{1,t}}{\partial x_i^{\nu} \partial \bar{x}_i^{\nu}} (0, y_i; \sigma_{i,\nu}), \quad \frac{d_i}{q_i} \nu \le \text{ord}_{z_i} \frac{\partial^{2\nu} \Phi_{2,t}}{\partial w_i^{\nu} \partial \bar{w}_i^{\nu}} (z_i, 0; \sigma_{i,\nu}),
$$

$$
\deg_{y_i} \frac{\partial^{2\nu} \Phi_{1,t}}{\partial x_i^{\nu} \partial \bar{x}_i^{\nu}} (0, y_i; \sigma_{i,\nu}) \le \frac{b_i}{q_i} \nu + e_i, \quad \deg_{z_i} \frac{\partial^{2\nu} \Phi_{2,t}}{\partial w_i^{\nu} \partial \bar{w}_i^{\nu}} (z_i, 0; \sigma_{i,\nu}) \le \frac{c_i}{n_i} \nu + e_i,
$$

for all $\nu \in \mathbb{Z}_{\geq 0}$, and the same bounds hold for the conjugated variables \bar{y}_i, \bar{z}_i .

Proof. We will do the proof for $\Phi_{1,t}$ since the proof for $\Phi_{2,t}$ works similarly. Recall that $\Phi_{1,t} = |u_1|^{2s} |v_1|^2 |\tilde{f}_t|^{2s} (\pi^{(i)})^* \varphi|_{U_i}$. By Lemma [6.3,](#page-26-0) it is enough to prove the bounds for each holomorphic or antiholomorphic factor. The factors $u_1, v_1, (\pi^{(i)})^* \varphi|_{U_i}$ are the pull-back by the toric morphism π_i of some invertible elements in the unique point of the total transform of f_t that is not a simple normal crossing on $X^{(i-1)}$. Therefore, by Section [4.2](#page-12-1) and Lemma [6.4,](#page-26-1) their orders (resp. degrees) with respect to y_i are bounded by $a_i \nu / n_i$ (resp. $b_i \nu / q_i$). It remains to prove the bounds for the strict transform \tilde{f}_t .

For the strict transform, consider Proposition [4.1](#page-13-0) and proceed by induction from f_{i+1} . Analyzing Equation [\(4.5\)](#page-13-2), the only part depending on x_i is the summation. By Lemma [6.3,](#page-26-0) it is enough to show that each factor of each summand fulfills the bounds. By the same argument as before, the units $u_k^{(i)}$ $\frac{k}{k}$ satisfy the bounds. On the other hand, Lemma [6.5](#page-27-0) assures the bounds for the monomials in x_i, y_i . The lower-bound for the order is clear. For the upper-bound on the degree just notice that

$$
\frac{a_i n_{i-1} \overline{m}_{i-1} + b_i}{\overline{m}_i} = \frac{a_i n_{i-1} \overline{m}_{i-1} + b_i}{n_i n_{i-1} \overline{m}_{i-1} + q_i} < \frac{b_i}{q_i},
$$

since $\overline{m}_i = n_i n_{i-1} \overline{m}_{i-1} + q_i$ and $n_i b_i - q_i a_i = 1$. Therefore, we are done for \tilde{f}_{i+1} . By induction, if all f_{k+1} , $i \leq k < j$ satisfy the bounds, so does f_{i+1} . To see this, it is just a matter of applying Lemma [6.3,](#page-26-0) Lemma [6.4,](#page-26-1) Lemma [6.5,](#page-27-0) and the induction hypothesis to Equation [\(4.6\)](#page-14-0).

We are ready to present the main result of this section.

Theorem 6.7. For any plane branch singularity $f : (\mathbb{C}^2, \mathbf{0}) \longrightarrow (\mathbb{C}, 0)$ the poles of the $complex zeta function f^s are simple and contained in the sets$

(6.4)
$$
\left\{\sigma_{i,\nu}=-\frac{m_i+n_1\cdots n_i+\nu}{n_i\overline{\beta}_i}\;\middle|\;\nu\in\mathbb{Z}_{\geq0},\;\overline{\beta}_i\sigma_{i,\nu},e_{i-1}\sigma_{i,\nu}\notin\mathbb{Z}\right\},\quad i=1,\ldots,g,
$$

contributed by the rupture divisors E_{p_i} , together with the negative integers $\mathbb{Z}_{\leq 0}$, contributed by the strict transform \tilde{C} .

Proof. In order to prove this result for any plane branch, it is enough to restrict the study to the family of f_t from Proposition [3.6.](#page-10-2) By the previous discussion, we only have to show that the candidates $\sigma_{i,\nu}$ such that $\overline{\beta}_i \sigma_{i,\nu}, e_{i-1} \overline{\sigma}_{i,\nu} \in \mathbb{Z}$ have always residue zero. The first important observation is that $\overline{\beta}_i \sigma_{i,\nu}, e_{i-1} \sigma_{i,\nu} \in \mathbb{Z}$, if and only if, $\epsilon_{1,\nu}, \epsilon_{2,\nu} \in \mathbb{Z}$, respectively. To see this, consider the definitions $\epsilon_{1,\nu} = N_1 \sigma_{1,\nu} + k_1, \epsilon_{2,\nu} = N_2 \sigma_{2,\nu} + k_2$ from Equation [\(5.6\)](#page-18-4). Hence, $\epsilon_{1,\nu}, \epsilon_{2,\nu} \in \mathbb{Z}$, if and only if, $N_1\sigma_{i,\nu}, N_2\sigma_{i,\nu} \in \mathbb{Z}$, respectively. By Proposition [4.1,](#page-13-0) $N_1 = a_i\overline{\beta}_i$ and $N_2 = (c_i n_{i-1}\overline{m}_{i-1} + d_i)e_{i-1}$, and the remark follows because

$$
\gcd(n_i\overline{\beta}_i, a_i\overline{\beta}_i) = \overline{\beta}_i \gcd(n_i, a_i) = \overline{\beta}_i,
$$

since $n_i b_i - q_i a_i = 1$, and

$$
\gcd(n_i\overline{\beta}_i,(c_in_{i-1}\overline{m}_{i-1}+d_i)e_{i-1})=e_{i-1}\gcd(\overline{m}_i,c_in_{i-1}\overline{m}_{i-1}+d_i)=e_{i-1},
$$

since $\overline{m}_i = n_i n_{i-1} \overline{m}_{i-1} + q_i$ and $q_i c_i - n_i d_i = 1$. The argument to show that the residue is zero if $\epsilon_{1,\nu} \in \mathbb{Z}$ or $\epsilon_{2,\nu} \in \mathbb{Z}$ is fundamentally different for each case. Let us begin by the case where $\beta_i \sigma_{i,\nu} \in \mathbb{Z}$ and $e_{i-1} \sigma_{i,\nu} \notin \mathbb{Z}$.

In order to study the residues of the candidates $\sigma_{i,\nu}$ of the *i*-th rupture divisor E_{pi} such that $\bar{\beta}_i \sigma_{i,\nu} \in \mathbb{Z}$, we place ourselves in the chart U_i of $X^{(i)}$ with local coordinates (x_i, y_i) . The origin of this chart is the intersection point $E_{p_1} \cap D_1$. The only point of the total transform on $X^{(i)}$ that is not a simple normal crossing is the intersection of the strict transform of f_t with E_{p_i} , with E_{p_i} being the only exceptional divisor at that point. Therefore, we can apply the formula for the residue in Proposition [5.4](#page-18-0) for the U_i chart.

From the preceding discussion, the derivatives of $\Phi_{1,t}$ appearing in the residue formula are a finite sum of terms that look like $y_p^{k'}$ ${}_{p}^{k'}\bar{y}_{p}^{k}(y_{i}-\lambda_{i})^{e_{i}(\sigma_{i,\nu}-l')}\left(\bar{y}_{i}-\bar{\lambda}_{i}\right)^{e_{i}(\sigma_{i,\nu}-l)}$. Consequently, we can reduce the residue to a finite sum of the integrals from Equation [\(5.15\)](#page-21-0), which, by Proposition [5.9,](#page-21-1) are equal to

(6.5)
$$
-2\pi i \lambda^{-\alpha-n-1} \bar{\lambda}^{-\alpha-1} \frac{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(\gamma+1)}{\Gamma(-\alpha-n)\Gamma(-\beta-m)\Gamma(-\gamma-n-m)}, \quad n, m \in \mathbb{Z}.
$$

As noted earlier, for E_{p_i} on $X^{(i)}$ we have that $\epsilon_{1,\nu} + \epsilon_{2,\nu} + e_i \sigma_{i,\nu} + \nu + 2 = 0$. If we are assuming that $\epsilon_{1,\nu} \in \mathbb{Z}$ but $\epsilon_{2,\nu} \notin \mathbb{Z}$, i.e. $e_{i-1}\sigma_{i,\nu} \notin \mathbb{Z}$, it must happen that $e_i\sigma_{i,\nu} \notin \mathbb{Z}$. This implies that $\Gamma(\beta+1), \Gamma(-\beta-n), \Gamma(\gamma+1), \Gamma(-\gamma-n-m) \in \mathbb{C}^*$. However, the remaining factor, $\Gamma(\alpha+1)/\Gamma(-\alpha-n)$, is always zero, since $\epsilon_{1,\nu} \in \mathbb{Z}$ implies that $\alpha, \alpha' \in \mathbb{Z}$,

$$
\alpha + 1 = \epsilon_{1,\nu} + 1 + k \ge -\frac{a_i}{n_i}\nu + \frac{1}{n_i} + \frac{a_i}{n_i}\nu = \frac{1}{n_i} > 0,
$$

and $-\alpha - n = -\alpha' < 1$, by Corollary [6.2](#page-25-1) and Proposition [6.6.](#page-27-1) This proves that the residue at the candidate poles $\sigma_{i,\nu}$ such that $\overline{\beta}_i \sigma_{i,\nu} \in \mathbb{Z}$, $e_{i-1} \sigma_{i,\nu} \notin \mathbb{Z}$ is zero.

We move now to the case where $e_{i-1}\sigma_{i,\nu} \in \mathbb{Z}, \overline{\beta}_i \sigma_{i,\nu} \notin \mathbb{Z}$, i.e. $\epsilon_{2,\nu} \in \mathbb{Z}$ and $e_i \sigma_{i,\nu} \notin \mathbb{Z}$. For this case observe that the previous argument, applied to the residue in the V_i chart, only works for the first rupture divisor. For $i = 1$, $\epsilon_{2,\nu} + 1 = -d_1\nu/q_1 + 1/q_1$, since $n_0, m_0 = 0$ and $\overline{m}_1 = q_1$. Otherwise, the formula for $\epsilon_{2,\nu}$ from Corollary [6.2](#page-25-1) and the bound for the order of $\Phi_{2,t}$ in Proposition [6.6](#page-27-1) do not match. Thus, from now on, we will assume $i \geq 2$.

To study the residue at these poles, we consider $\text{Res}_{s=\sigma_{i,\nu}} f_{t}^{s}$ as a function of $\lambda_{i} \in \mathbb{C}$ and we place ourselves in the V_i chart. Since λ_i is the intersection coordinate of the strict transform with E_{p_i} , if we let $\lambda_i \to 0$ or $\lambda_i \to \infty$, we are in the situation of a non-rupture divisor and the residue is zero. By Proposition [5.9](#page-21-1) and since $\epsilon_2 \in \mathbb{Z}$, the residue is a Laurent series on λ_i , $\bar{\lambda}_i \neq 0$. Deriving under the integral sign in the formula for the residue from Corollary [5.5](#page-19-2) we increase the order in z_i, \bar{z}_i of $\Phi_{2,t}$ by one unit. Therefore, after deriving enough times we can assume that in Equation (6.5) , $\alpha+1 > 0$, $-\alpha' \leq 0$, $\alpha, \alpha' \in \mathbb{Z}$. This implies that the principal part in λ_i , $\bar{\lambda}_i$ of $\text{Res}_{s=\sigma_{i,\nu}} f^s_{t}(\lambda_i)$ must be zero. However, the same argument is true if we consider the residue as a function of λ_i^{-1} $\overline{i}^{-1}, \overline{\lambda}_i^{-1}$ i^{-1} . Hence, the residue is independent of λ_i , $\bar{\lambda}_i$. This implies that the residue is zero overall.

It remains to show that the residue is zero in the case that $\overline{\beta}_i \sigma_{i,\nu} \in \mathbb{Z}$ and $e_{i-1} \sigma_{i,\nu} \in \mathbb{Z}$. Both conditions imply that, $e_i \sigma_{i\nu} \in \mathbb{Z}$. To see that the residue is zero in this situation, it is just a matter of combining the previous arguments and recalling that the Gamma function has only simple poles. After deriving with respect to λ_i in the V_i chart, we can get Equation [\(6.5\)](#page-29-0) with $\alpha + 1 > 0$ and $-\alpha - n = -\alpha' \leq 0$, i.e., the factor $\Gamma(\alpha + 1)/\Gamma(-\alpha - n)$ is zero. Assume we have derived d' times with respect to λ_i , then

$$
\alpha' = \epsilon_{2,\nu} + d' + k' \qquad \beta' = e_i \sigma_{i,\nu} - e_i l' - e_i d'.
$$

Consequently, since $e_i \geq 1$,

$$
\gamma + 1 = -\alpha' - \beta' - 1 = -\epsilon_{2,\nu} + 1 - k' - \epsilon_i \sigma_{i,\nu} + \epsilon_i l' - 2 \ge \epsilon_{1,\nu} + \nu + 1 - k' + \epsilon_i
$$

$$
\ge -\frac{a_i}{n_i} \nu + \frac{1}{n_i} + \nu - \frac{c_i}{n_i} \nu - \epsilon_i + \epsilon_i = \frac{1}{n_i} > 0,
$$

by Corollary [6.2](#page-25-1) and Proposition [6.6.](#page-27-1) Similarly, $-\gamma' - n - m < 1$. Therefore, the factor $\Gamma(\gamma+1)/\Gamma(-\gamma-n-m)$ is also zero. Since $e_{i-1}\sigma_{i\nu}\in\mathbb{Z}$, the piece, $\Gamma(\beta+1)/\Gamma(-\beta-m)$, has a pole. However, Equation [\(6.5\)](#page-29-0) is zero because the poles of $\Gamma(z)$ are simple.

Finally, we need to see that the negative integers, the candidates coming from the strict transform \tilde{C} , are poles. We can argue directly from the definition of f^s given in Equation [\(5.1\)](#page-15-3). Take $0 \neq p \in f_t^{-1}(0) \cap U$ at which the equation f_t can be taken as one of the holomorphic coordinates. Thus, we reduce the problem to the monomial case and, by Proposition [2.1,](#page-4-1) the negative integers are simple poles. The poles contributed by rupture divisors are also simple because they must have $\epsilon_{1,\nu}, \epsilon_{2,\nu}, e_i \sigma_{i,\nu} \notin \mathbb{Z}$ for all $\nu \in \mathbb{Z}_{\geq 0}$. These conditions imply that Equation [\(6.5\)](#page-29-0) cannot have a pole and hence, the residue does not have a pole. By Section [5.1,](#page-15-4) all the poles of f^s are simple.

We point out that the candidate poles $\sigma_{1,0} > \sigma_{2,0} > \cdots > \sigma_{g,0}$ are always poles of f^s for any plane branch f as shown by Lichtin in [\[47,](#page-34-5) [48\]](#page-34-6).

Example 2. There are examples where the candidate poles of f^s that are in the sets from Equation [\(6.4\)](#page-28-1) vary in a μ -constant deformations of f. For instance, consider $f = y^4 - x^9$ and the μ -constant deformation $f_t = y^4 - x^9 + tx^5y^2$. For the unique rupture divisor, the sequence of candidate poles is $\sigma_{1,\nu} = -(13 + \nu)/36, \nu \in \mathbb{Z}_{\geq 0}$. Taking $\nu = 2$, $\sigma_{1,2} = -5/12, \epsilon_{1,2} = -9/4, \epsilon_{2,2} = -4/3, \text{ and}$

$$
\operatorname{Res}_{s=\sigma_{1,2}} f_t = -16\pi^2 \sigma_{1,2}^2 \frac{\Gamma(\epsilon_{1,2}+3)\Gamma(\sigma_{1,2})\Gamma(\epsilon_{2,2}+2)}{\Gamma(-\epsilon_{1,2}-2)\Gamma(-\sigma_{1,2}+1)\Gamma(-\epsilon_{2,2}-1)} |t|^2 \delta_0^{(0,0,0,0)}.
$$

Therefore, $\sigma_{1,2} = -5/12$ is a pole, if and only if, $t \neq 0$.

6.2. Generic poles. Studying the residue in terms of the deformation parameters of f_t , we can get open conditions in which a certain candidate pole is indeed a pole, as seen in Example [2.](#page-30-1) Recalling Equation [\(5.11\)](#page-19-1), the first observation is that, in terms of t ,

(6.6)
$$
\operatorname{Res}_{s=\sigma_{i,\nu}} f_t^s = \sum_{k',l',k,l=0}^{\nu} p_{k',l'}(\boldsymbol{t}) p_{k,l}(\bar{\boldsymbol{t}}) \, \delta_{\boldsymbol{0}}^{(k',l',k,l)},
$$

with $p_{k',l'}(t) = \overline{p_{k,l}(\bar{t})}$ if $k' = k$ and $l' = l$. The following theorem shows that, actually, any candidate is a pole in a certain Zariski open set in the deformation space of f_t .

Theorem 6.8. For any $M_1, \ldots, M_g \in \mathbb{Z}_{\geq 0}$, generic plane branches f_{gen} in the equisingularity class corresponding to the semigroup $\Gamma = \langle \beta_0, \beta_1, \ldots, \beta_g \rangle$ satisfy that

(6.7)
$$
\left\{\sigma_{i,\nu}=-\frac{m_i+n_1\cdots n_i+\nu}{n_i\overline{\beta}_i}\middle| 0\leq \nu < M_i, \overline{\beta}_i\sigma_{i,\nu}, e_{i-1}\sigma_{i,\nu}\notin\mathbb{Z}\right\}, \quad i=1,\ldots,g,
$$

are simple poles of the complex zeta function f_{gen}^s .

Proof. By Theorem [6.7,](#page-28-0) we only have to check the candidates in the sets given in Equa-tion [\(6.4\)](#page-28-1). In the case that $\nu = 0$, the residue for the candidates $\sigma_{1,0} > \sigma_{2,0} > \cdots > \sigma_{q,0}$ does not depend on t and consists only of one of the integrals from Proposition 5.9 which is non-zero by Proposition [6.1.](#page-25-0)

Therefore, fix a candidate pole $\sigma_{i,\nu}$ with $0 < \nu < M_i$. It is enough to show that one of the polynomials $p_{k',l'}(t)$ on the parameters t in the expression from Equation [\(6.6\)](#page-30-2) is not identically zero. Consider the residue formula from Proposition [5.4](#page-18-0) in, for instance, the U_i

chart and recall that $\Phi_{1,t} = |u_1|^{2s} |v_1|^2 |f_t|^{2s} (\pi^{(i)})^* \varphi|_{U_i}$. Now, considering Equation [\(4.5\)](#page-13-2), we claim that the strict transform has a deformation term

$$
t_{\underline{k}}^{(i)} x_i^{\rho_{i+1}^{(i)}(\underline{k})} y_i^{A_{i+1}^{(i)}(\underline{k})} u_{\underline{k}}^{(i)}(x_i, y_i)
$$

for a certain <u>k</u> such that $\rho_{i+1}^{(i)}(\underline{k}) = \nu$. Indeed, this happens since $\rho_{i+1}^{(i)}(\underline{k}) = \nu$ is equivalent, by Equation (4.7) , to

$$
n_1\cdots n_ik_0+n_2\cdots n_i\overline{m}_1k_1+\cdots+\overline{m}_ik_i=n_i\overline{m}_i+\nu.
$$

But this is an identity in the semigroup Γ_{i+1} of the maximal contact element f_{i+1} , see Equation [\(4.2\)](#page-13-3), and such a <u>k</u> always exist because $n_i\overline{m}_i + \nu$ is bigger than the conductor of Γ_{i+1} . The deformation parameter $t_k^{(i)}$ $\frac{k}{k}$ for such a <u>k</u> can only appear when f_t is derived ν times. By Faà di Bruno's formula, this implies that the polynomial $p_{0,0}(t)$ in Equation (6.6) is equal to $\zeta t_k^{(i)} + \cdots$, where the dots represents other terms on \boldsymbol{t} not containing $t_k^{(i)}$ ⁽ⁱ⁾. The coefficient $\zeta \in \mathbb{C}$ is different from zero since it has the form of Equation [\(5.16\)](#page-21-2) and we are assuming that $\overline{\beta}_i \sigma_{i,\nu}, e_{i-1} \sigma_{i,\nu} \notin \mathbb{Z}$. Hence, the condition $p_{0,0}(\boldsymbol{t}) \neq 0$ gives a non-empty Zariski open subset on the deformation space of f_t in which $\sigma_{i,\nu}$ is a pole.

In the case that there is a resonance between two poles, i.e. $\sigma_{i,\nu} = \sigma_{i',\nu'}$, $i \neq i'$, we can always add, if necessary, an extra condition, giving a Zariski open set, which ensures that the residues do not cancel out. The intersection of all the open sets defines generic plane branches f_{gen} .

Consider the sets from Equation [\(6.7\)](#page-30-3) with $M_i = n_i \beta_i$, namely

$$
\Pi_i := \left\{ \sigma_{i,\nu} = -\frac{m_i + n_1 \cdots n_i + \nu}{n_i \overline{\beta}_i} \; \middle| \; 0 \leq \nu < n_i \overline{\beta}_i, \; \overline{\beta}_i \sigma_{i,\nu}, e_{i-1} \sigma_{i,\nu} \notin \mathbb{Z} \right\},
$$

for $i = 1, \ldots, g$ and define $\Pi := \bigcup_{i=1}^g \Pi_i$. An easy computation shows that there are exactly μ elements in Π , counted with possible multiplicities,

$$
|\Pi| = \sum_{i=1}^{g} n_i \overline{\beta}_i - \overline{\beta}_i - n_i e_i + e_i = \sum_{i=1}^{g} (n_i - 1) \overline{\beta}_i + \sum_{i=1}^{g} e_i - e_{i-1} = \sum_{i=1}^{g} (n_i - 1) \overline{\beta}_i - n + 1 = \mu,
$$

using Equation [\(3.4\)](#page-9-0) in the last equality. The sets Π_i are precisely the b-exponents in Yano's conjecture from Section [2.3.](#page-7-0) Indeed, the relation between the notations in Equa-tion [\(2.7\)](#page-7-2) and the resolution data in Section [4.2](#page-12-1) is clear. Namely, $R_i = N_{p_i} = n_i \overline{\beta}_i$, $R'_i =$ $N_{q_i} = \overline{\beta}_i, r_i = k_{p_i} + 1 = m_i + n_1 \cdots n_i$ and $r'_i = k_{q_i} + 1 = [(m_i + n_1 \cdots n_i)/n_i]$. To see the equality between the exponents in Equation (2.8) and the set Π is enough to notice that $R_i = N_{p_i} = n_i N_{q_i} = n_i R'_i$ and $r_i = k_{p_i} + 1 = n_i (k_{q_i} + 1) = n_i r'_i$.

The results of Malgrange [\[52\]](#page-34-3) and Barlet [\[4\]](#page-32-7) imply that the elements of Π generate all the eigenvalues of the monodromy. The characteristic polynomial of the monodromy is a topological invariant of the singularity, see A'Campo's formula [\[1\]](#page-32-5). Consequently, it can be computed from the semigroup Γ of f, see [\[56\]](#page-34-8). The hypothesis that the eigenvalues of the monodromy are pairwise different is a condition on the equisingularity class, i.e. on the semigroup Γ, implying that there are exactly μ different elements in Π.

As a corollary of Theorem [6.8,](#page-30-0) we can deduce Yano's conjecture for any number of characteristic exponents if we assume that the eigenvalues of the monodromy are different. Yano's conjecture is proved by Cassou-Noguès for plane branches with one characteristic exponent in [\[25\]](#page-33-18). For two characteristic exponents with different monodromy eigenvalues, Yano's conjecture is proved by Artal Bartolo, Cassou-Noguès, Luengo and Melle Hernández in $|8|$.

Corollary 6.9. Let $\Gamma = \langle \beta_0, \beta_1, \dots, \beta_g \rangle$ be a semigroup defining an equisingularity class of plane branches. If the eigenvalues of the monodromy in the equisingularity class associated to the semigroup Γ are pairwise different, then Yano's conjecture holds.

Proof. We must check that all the μ different elements of Π are roots of the Bernstein-Sato polynomial $b_{gen,0}(s)$ of f_{gen} . If $\sigma_{i,\nu} \in \Pi_i$ is in between $-\mathrm{lct}(f) - 1$ and $-\mathrm{lct}(f)$, then $\sigma_{i,\nu}$ is automatically a roots of $b_{gen,0}(s)$ by Corollary [2.6.](#page-6-3) Otherwise, by Corollary [2.6,](#page-6-3) we must check that $\sigma_{i,\nu} + 1$ is not a root of $b_{gen,0}(s)$. By contradiction, assume $\sigma_{i,\nu} + 1$ is a root of $b_{gen,0}(s)$. By [\[49,](#page-34-4) Th. 1.9], the roots of the Bernstein-Sato polynomial of a reduced plane curve can only be of the form $\sigma - k$, with $k \in \mathbb{Z}_{\geq 0}$ and σ a pole of f^s . Hence, by Theorem [6.7,](#page-28-0) $\sigma_{i,\nu} + 1 = \sigma_{i',\nu'} \in \Pi_{i'}, i \neq i'$. But this is impossible since, as eigenvalues of the monodromy, they are both equal, contradicting the hypothesis. Finally, by the definitions in Section [2.3,](#page-7-0) if the Bernstein-Sato polynomial has exactly μ different roots they must coincide with the opposites in sign to the b-exponents. \Box

REFERENCES

- $[1]$ N. A'Campo, La fonction zêta d'une monodromie, Comment. Math. Helv. 50 (1975), 233–248. [2,](#page-1-0) [8,](#page-7-4) [32](#page-31-0)
- [2] V. I. Arnold, A. Varchenko, and S. M. Gusein-Zade, Singularities of differentiable maps, vol. 2, Monogr. Math., no. 8[3](#page-2-0), Birkhäuser, Boston, 1988. 3
- [3] M. F. Atiyah, *Resolution of singularities and division of distributions*, Comm. Pure Appl. Math. 23 (1970), no. 2, 145–150. [1,](#page-0-2) [5](#page-4-2)
- [4] D. Barlet, Contribution effective de la monodromie aux développements asymptotiques, Ann. Sci. Éc. Norm. Supér. (4) 17 (1984), no. [2,](#page-1-0) 293-315. 2, [32](#page-31-0)
- [5] ___, Le calcul de la forme hermitienne canonique pour $X^a + Y^b + Z^c = 0$, Séminaire d'analyse P. Lelong – P. Dolbeault – H. Skoda, années 1983/1984 (P. Lelong, P. Dolbeault, and H. Skoda, eds.), Lecture Notes in Math., vol. 1198, Springer, Berlin, 1986, pp. 35–46. [22](#page-21-3)
- [6] $______\$, Monodromie et pôles du prolongement méromorphe de $\int_X |f|^{2\lambda} \Box$, Bull. Soc. Math. France 114 (1986), 247–269. [2](#page-1-0)
- [7] E. Artal Bartolo, Pi. Cassou-Noguès, I. Luengo, and A. Melle Hernández, Quasi-ordinary power series and their zeta functions, Mem. Amer. Math. Soc. 178 (2005), no. 841. [3](#page-2-0)
- [8] , Yano's conjecture for two-Puiseux-pair irreducible plane curve singularities, Publ. Res. Inst. Math. Sci. 53 (2017), no. 1, 211–239. [4,](#page-3-3) [33](#page-32-16)
- [9] J. Bernstein, Modules over a ring of differential operators. Study of the fundamental solutions of equations with constant coefficients, Funct. Anal. Appl. 5 ([1](#page-0-2)971), no. 2, 1–16. 1
- [10] , Analytic continuation of distributions with respect to a parameter, Funct. Anal. Appl. 6 (1972), no. 4, 26–40. [1,](#page-0-2) [4,](#page-3-3) [6](#page-5-3)
- [11] J. Bernstein and S. I. Gel'fand, Meromorphic continuation of the function P^{λ} , Funct. Anal. Appl. 3 (1969), no. 1, 84–85. [1,](#page-0-2) [5](#page-4-2)
- [12] J.-E. Björk, *Dimensions of modules over algebras of differential operators*, Fonctions analytiques de plusieurs variables et analyse complexe (Colloq. Internat. CNRS, No. 208, Paris, 1972), Agora Mathematica, vol. 1, Gauthier-Villars, Paris, 1974, pp. 6–11. [1](#page-0-2)
- [13] , Rings of differential operators, North-Holland Math. Library, no. 21, North-Holland Pub. Co., Amsterdam, 1979. [6](#page-5-3)
- [14] B. Bories and W. Veys, Igusa's p-adic local zeta function and the monodromy conjecture for non-degenerate surface singularities, Mem. Amer. Math. Soc. 242 (2016), no. 1145. [3](#page-2-0)
- [15] J. Briançon, M. Granger, P. Maisonobe, and M. Miniconi, Algorithme de calcul du polynôme de *Bernstein : Cas non dégénéré*, Ann. Inst. Fourier (Grenoble) **[3](#page-2-0)9** (1989), no. 3, 553–610. 3

- [16] J. Briançon and P. Maisonobe, Caractérisation géométrique de l'existence du polynôme de Bernstein relatif, Algebraic Geometry and Singularities (A. Campillo and L. Narváez-Macarro, eds.), Progr. Math., vol. 1[3](#page-2-0)4, Birkhäuser, Basel, 1996, pp. 215–236. 3
- [17] J. Briançon, P. Maisonobe, and T. Torrelli, Matrice magique associée à un germe de courbe plane et division par l'idéal jacobien, Ann. Inst. Fourier (Grenoble) 57 (2007), no. [3](#page-2-0), 919–953. 3
- [1[8](#page-7-4)] E. Brieskorn, Die Monodromie der isolierten Singularitäten von Hyperflächen, 2 (1970), 103–161. 8
- [19] N. Budur, *On Hodge spectrum and multiplier ideals*, Math. Ann. **[3](#page-2-0)27** (2003), no. 2, 257–270. **3**
- [20] N. Budur, M. Mustată, and Z. Teitler, *The monodromy conjecture for hyperplane arrangements*, Geom. Dedicata 153 (2011), no. 1, 131–137. [3](#page-2-0)
- [21] N. Budur and M. Saito, Multiplier ideals, V -filtration, and spectrum, 14 (2005), no. 2, 269–282. [3](#page-2-0)
- [22] E. Casas-Alvero, Singularities of plane curves, London Math. Soc. Lecture Note Ser., no. 276, Cambridge Univ. Press, Cambridge, 2000. [10,](#page-9-6) [12,](#page-11-1) [13](#page-12-2)
- [23] Pi. Cassou-Noguès, Racines de polynômes de Bernstein, Ann. Inst. Fourier (Grenoble) 36 (1986), no. 4, 1–30. [3](#page-2-0)
- $[24]$ $_____\$ ftude du comportement du pôlynome de Bernstein lors d'une déformation à μ constant de $X^a + Y^b$ avec $(a, b) = 1$, Compos. Math. **6[3](#page-2-0)** (1987), no. 3, 291-313. 3
- [25] , Polynôme de Bernstein générique, Abh. Math. Semin. Univ. Hambg. 58 (1988), no. 1, 103–124. [3,](#page-2-0) [4,](#page-3-3) [33](#page-32-16)
- [26] L. Comtet, Advanced combinatorics. The art of finite and infinite expansions, D. Reidel Pub. Co., Dordrecht, 1974. [21](#page-20-6)
- [27] T. de Jong and G. Pfister, Local analytic geometry. Basic theory and applications., Adv. Lect. Math., Friedr. Vieweg & Sohn, Braunschweig, 2000. [12](#page-11-1)
- [28] J. Denef, Report on Igusa's local zeta function, Séminaire N. Bourbaki 33 (1991), no. 741, 359–386. [3](#page-2-0)
- [29] J. Denef and F. Loeser, Caractéristiques d'Euler-Poincaré, fonctions zêta locales et modifications analytiques, J. Amer. Math. Soc. 5 (1992), no. 4, 705–720. [3](#page-2-0)
- [30] L. Ein, R. Lazarsfeld, K. E. Smith, and D. Varolin, Jumping coefficients of multiplier ideals, Duke Math. J. **12[3](#page-2-0)** (2004), no. 3, 469–506. 3
- [31] I. M. Gel'fand, Some aspects of functional analysis and algebra, Proceedings of the International Congress of Mathematicians 1954 (Amsterdam) (J. C. H. Gerretsen and J. de Groot, eds.), vol. 1, Erven P. Noordhoff N. V., Groningen and North-Holland Pub. Co., Amsterdam, 1957, pp. 253–276. [1,](#page-0-2) [5,](#page-4-2) [6](#page-5-3)
- [32] I. M. Gel'fand and G. E. Shilov, Generalized functions, vol. 1, Academic Press, New York, 1964. [1,](#page-0-2) [4,](#page-3-3) [5,](#page-4-2) [16,](#page-15-5) [21](#page-20-6)
- [33] I. S. Gradshteyn and I. M. Ryzhik, Table of integrals, series, and products, eighth ed., Elsevier/Academic Press, Amsterdam, 2015. [23](#page-22-1)
- [34] R. Hartshorne, *Algebraic geometry*, Grad. Texts in Math, no. 52, Springer, Berlin, [19](#page-18-5)77. 19
- [35] C. Hertling and C. Stahlke, *Bernstein polynomial and Tjurina number*, Geom. Dedicata **75** (1999), no. 2, 137–176. [3](#page-2-0)
- [36] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero: I, Ann. of Math. 79 (1964), no. 1, 109–203. [1](#page-0-2)
- [37] , Resolution of singularities of an algebraic variety over a field of characteristic zero: II, Ann. of Math. 79 (1964), no. 2, 205–326. [1](#page-0-2)
- [38] J.-i. Igusa, Complex powers and asymptotic expansions. I. Functions of certain types, J. Reine Angew. Math. 268-269 (1974), 110–130. [2](#page-1-0)
- [39] Complex powers and asymptotic expansions. II, J. Reine Angew. Math. 278-279 (1975), 307–321. [2](#page-1-0)
- [40] , B-functions and p-adic integrals, Algebraic analysis, vol. 1, Academic Press, Boston, 1988, pp. 231–241. [2](#page-1-0)
- [41] $________________\._$ on local zeta functions, Selected papers on number theory, algebraic geometry, and differential geometry (K. Nomizu, ed.), Amer. Math. Soc. Transl. Ser. 2, vol. 160, Amer. Math. Soc., Providence, RI, 1996, pp. 1–20. [3](#page-2-0)
- [42] M. Kashiwara, B-functions and holonomic systems, Invent. Math. 38 (1976), no. 1, 33–53. [1,](#page-0-2) [3,](#page-2-0) [6](#page-5-3)
- [43] M. Kato, *The b-function of a* μ *-constant deformation of* $x^7 + y^5$, Bull. College Sci. Univ. Ryukyus 32 (1981), 5–10. [3](#page-2-0)
- [44] \ldots , The b-function of a μ -constant deformation of $x^9 + y^4$, Bull. College Sci. Univ. Ryukyus 33 $(1982), 5-8.3$ $(1982), 5-8.3$
- [45] J. Kollár, Singularities of pairs, Algebraic Geometry (Santa Cruz, 1995) (J. Kollár, R. Lazarsfeld, and D. R. Morrison, eds.), Proc. Sympos. Pure Math., vol. 62, part 2, Amer. Math. Soc., Providence, RI, 1997, pp. 221–287. [2,](#page-1-0) [3](#page-2-0)
- [46] V. Levandovskyy and J. Martín-Morales, Algorithms for checking rational roots of b-functions and *their applications*, J. Algebra 352 352 (2012), no. 1, 408–429. 3
- [47] B. Lichtin, Some algebro-geometric formulae for poles of $|f(x, y)|^s$, Amer. J. Math. **107** (1985), no. 1, 139–162. [2,](#page-1-0) [19,](#page-18-5) [26,](#page-25-2) [31](#page-30-4)
- [48] \ldots , Poles of $|f(z, w)|^{2s}$ and roots of the B-function, Ark. Mat. 27 (1989), no. 1-[2,](#page-1-0) 283-304. 2, [3,](#page-2-0) [6,](#page-5-3) [22,](#page-21-3) [31](#page-30-4)
- [49] F. Loeser, Quelques conséquences locales de la théorie de Hodge, Ann. Inst. Fourier (Grenoble) 35 (1985), no. 1, 75–92. [2,](#page-1-0) [3,](#page-2-0) [7,](#page-6-4) [33](#page-32-16)
- [50] , Fonctions d'Igusa p-adiques et polynomes de Bernstein, Amer. J. Math. 110 (1988), no. 1, 1–21. [2](#page-1-0)
- [51] , Fonctions d'Igusa p-adiques, polynomes de Bernstein, et polyedres de Newton, J. Reine Angew. Math. 412 (1990), 75–96. [3](#page-2-0)
- [52] B. Malgrange, Le polynôme de Bernstein d'une singularité, Lecture Notes in Math. 4 (1975), 98–119. [1,](#page-0-2) [2,](#page-1-0) [3,](#page-2-0) [4,](#page-3-3) [8,](#page-7-4) [32](#page-31-0)
- [53] , Polynôme de Bernstein-Sato et cohomologie évanescente, Analysis and topology on singular spaces, II, III (Luminy, 1981), Astérisque, vol. 101-102, Soc. Math. France, Paris, 1983, pp. 243–267. [2](#page-1-0)
- [54] Z. Mebkhout and L. Narváez-Macarro, La théorie du polynôme de Bernstein-Sato pour les algèbres de Tate et de Dwork-Monsky-Washnitzer, Ann. Sci. Éc. Norm. Supér. (4) 24 (1991), no. 2, 227–256. [1](#page-0-2)
- [55] D. Meuser, A survey of Igusa's local zeta function, Amer. J. Math. 138 (2016), no. 1, 149–179. [3](#page-2-0)
- [56] W. D. Neumann, Invariants of plane curve singularities, Knots, braids and singularities (Plans-sur-Bex, 1982), Monogr. Enseign. Math., Enseignement Math., Geneva, 1983, pp. 223–232. [2,](#page-1-0) [32](#page-31-0)
- [57] M. Noro, An efficient modular algorithm for computing the global b-function, Mathematical software (Beijing, 2002) (M. Cohen, ed.), Proceedings of the First International Congress of Mathematical Software, World Sci. Pub., 2002, pp. 147–157. [3](#page-2-0)
- [58] T. Oaku, An algorithm of computing b-functions, Duke Math. J. **87** (1997), no. 1, 115–1[3](#page-2-0)2. 3
- [59] M. Oka, Geometry of plane curves via toroidal resolution, Algebraic Geometry and Singularities (A. Campillo and L. Narváez-Macarro, eds.), Progr. Math., vol. 134, Birkhäuser, Basel, 1996, pp. 215–236. [13](#page-12-2)
- [60] M. Saito, On the structure of the Brieskon lattice, Ann. Inst. Fourier (Grenoble) 39 (1989), no. 1, 27–72. [3](#page-2-0)
- [61] $______\$, *On microlocal b-function*, Bull. Soc. Math. France **122** (1994), 163–184. [3,](#page-2-0) [7](#page-6-4)
- [62] , Bernstein-Sato polynomials of hyperplane arrangements, Selecta Math. (N.S.) 22 (2016), no. 4, 2017–2057. [3](#page-2-0)
- [63] M. Sato, M. Kashiwara, T. Kimura, and T. Oshima, Micro-local analysis of prehomogeneous vector spaces, Invent. Math. 62 ([1](#page-0-2)980), no. 1, 117–179. 1
- [64] M. Sato and T. Shintani, Theory of prehomogeneous vector spaces (algebraic part) – the English translation of Sato's lecture from Shintani's note, Nagoya Math. J. 120 (1990), 1–34. [1](#page-0-2)
- [65] K. E. Smith and H. M. Thompson, Irrelevant exceptional divisors for curves on a smooth surface, Algebra, geometry and their interactions (A. Corso, J. Migliore, and C. Polini, eds.), Contemp. Math., vol. 448, Amer. Math. Soc., Providence, RI, 2007, pp. 245–254. [2](#page-1-0)
- [66] J. H. M. Steenbrink, Mixed Hodge structure on the vanishing cohomology, Real and complex singularities (Oslo, 1976), Proc. Ninth Nordic Summer School/NAVF Sympos. Math., Sijthoff and Noordhoff, Alphen aan den Rijn, 1977, pp. 525–563. [3](#page-2-0)
- [67] , The spectrum of hypersurface singularities, Actes du Colloque de Théorie de Hodge (Luminy, 1987), Astérisque, vol. 179-180, Soc. Math. France, Paris, 1989, pp. 16[3](#page-2-0)–184. 3

- [68] B. Teissier, Appendix, in [\[74\]](#page-35-5), 1986. [2,](#page-1-0) [9,](#page-8-5) [10,](#page-9-6) [11](#page-10-3)
- [69] A. N. Varchenko, The complex exponent of a singularity does not change along strata $\mu = const$, Funct. Anal. Appl. 16 (1982), no. 1, 1–9. [3](#page-2-0)
- [70] C. T. C. Wall, Singular points of plane curves, London Math. Soc. Stud. Texts, no. 63, Cambridge, 2004. [12,](#page-11-1) [14](#page-13-4)
- [71] U. Walther, Bernstein-Sato polynomial versus cohomology of the milnor fiber for generic hyperplane arrangements, Compos. Math. 141 (2005), no. 1, 121–145. [3](#page-2-0)
- [72] T. Yano, On the theory of b-functions, Publ. Res. Inst. Math. Sci. 14 (1978), no. 1, 111–202. [3](#page-2-0)
- [73] , Exponents of singularities of plane irreducible curves, Sci. Rep. Saitama Univ. Ser. 10 (1982), no. 2, 21–28. [2,](#page-1-0) [3,](#page-2-0) [4,](#page-3-3) [8](#page-7-4)
- [74] O. Zariski, Le problème des modules pour les branches planes, Hermann, Paris, 1986. [9,](#page-8-5) [10,](#page-9-6) [35](#page-34-25)

DEPARTAMENT DE MATEMÀTIQUES, UNIV. POLITÈCNICA DE CATALUNYA, AV. DIAGONAL 647, Barcelona 08028, Spain.

E-mail address: Guillem.Blanco@upc.edu