

# Holographic Thermodynamics of Accelerating Black Holes

Andrés Anabalón,<sup>1</sup> Michael Appels,<sup>2</sup> Ruth Gregory,<sup>2,3</sup> David Kubizňák,<sup>3</sup> Robert B. Mann,<sup>4,3</sup> and Ali Övgün<sup>5,6</sup>

<sup>1</sup>*Universidad Adolfo Ibáñez; Dep. de Ciencias, Facultad de Artes Liberales,  
Av. Padre Hurtado 750, - Viña del Mar, Chile.*

<sup>2</sup>*Centre for Particle Theory, Durham University, South Road, Durham, DH1 3LE, UK*

<sup>3</sup>*Perimeter Institute, 31 Caroline St., Waterloo, Ontario, N2L 2Y5, Canada*

<sup>4</sup>*Department of Physics and Astronomy, University of Waterloo, Waterloo, Ontario, Canada, N2L 3G1*

<sup>5</sup>*Instituto de Física, Pontificia Universidad Católica de Valparaíso, Casilla 4950, Valparaíso, Chile*

<sup>6</sup>*Physics Department, Arts and Sciences Faculty, Eastern Mediterranean University,  
Famagusta, 99628, North Cyprus via Mersin 10, Turkey*

(Dated: November 11, 2018)

We present a careful study of accelerating black holes in anti-de Sitter spacetime, formulating the thermodynamics and resolving discrepancies that have appeared in previous investigations of the topic. We compute the dual stress-energy tensor for the spacetime and identify the energy density associated with a static observer at infinity. The dual energy-momentum tensor can be written as a three-dimensional perfect fluid plus a non-hydrodynamic contribution with a universal coefficient which is given in gauge theory variables. We demonstrate that both the holographic computation and the method of conformal completion yield the same result for the mass. We compare to previous work on black funnels and droplets, showing that the boundary region can be endowed with non-compact geometry, and comment on this novel holographic dual geometry.

PACS numbers: 04.20.Jb, 04.70.-s

The importance of black holes in advancing our understanding of physics cannot be underestimated. They provide a setting for testing our most fundamental ideas about gravity under extreme conditions and offer us insight into the underlying microscopic degrees of freedom that may be associated with quantum gravity. The subject of black hole thermodynamics [1–3] has proven to be an invaluable tool to this end, and broad classes of black holes have been shown to exhibit a rich and varied range of thermodynamic behaviour, particularly in anti-de Sitter spacetime [4].

Within this framework, accelerating black holes have presented a unique challenge. The idealized solution is described by the *C-metric* [5–8], whose spacetime has a string-like singularity along one polar axis attached to the black hole. We can think of this conical singularity as a cosmic string (indeed, the conical singularity can be replaced by a finite width topological defect [9], or a magnetic flux tube [10]) with the tension providing the force driving the acceleration. Surprisingly, even though these black holes are not isolated because of the “cosmic strings” it is possible to derive sensible looking thermodynamics, although recent studies have apparently conflicting results [11–14].

We consider here the interpretation of an accelerating black hole in anti-de Sitter (AdS) spacetime, with a focus on a holographic interpretation of the thermodynamics. We resolve conflicting issues that exist in the literature, obtain a distinct set of thermodynamic variables that are now consistent with the gravitational action, and agree with both the conformal and holographic methods for computing conserved charges. To this end, we focus our attention to black holes with no acceleration horizon [15], so that there is no ambiguity as to which horizon temper-

ature should be considered, or as to whether there is an equilibrium thermodynamics for the system. In addition, as we discuss, the holographic computation and interpretation are also unambiguous and straightforward. We also comment on the cases when the acceleration horizons appear and provide a novel interpretation of the boundary geometry.

An accelerating black hole in AdS can be described by the metric [8, 15, 16]

$$ds^2 = \frac{1}{\Omega^2} \left[ -f dt^2 + \frac{dr^2}{f} + r^2 \left( \frac{d\theta^2}{\Sigma} + \Sigma \sin^2 \theta \frac{d\phi^2}{K^2} \right) \right], \quad (1)$$

where

$$\begin{aligned} \Omega &= 1 + Ar \cos \theta, & \Sigma &= 1 + 2mA \cos \theta, \\ f(r) &= (1 - A^2 r^2) \left( 1 - \frac{2m}{r} \right) + \frac{r^2}{\ell^2}. \end{aligned} \quad (2)$$

The potential  $f(r)$  shows clearly the black hole nature of the solution, as well as the effects of acceleration ( $A$ ) and cosmological constant ( $\Lambda = -3/\ell^2$ ). Note that we require  $2mA < 1$  to preserve the metric signature. The parameter  $K$  encodes information about the conical deficits on the north and south poles that have tensions given by [12]

$$\mu_{\pm} = \frac{\delta_{\pm}}{8\pi} = \frac{1}{4} \left( 1 - \frac{\Sigma(\theta_{\pm})}{K} \right) = \frac{1}{4} \left( 1 - \frac{1 \pm 2mA}{K} \right). \quad (3)$$

The absence of an acceleration horizon yields the constraint  $f(-1/A \cos \theta) > 0$ , in turn constraining the parameter space  $(m, \ell)$  to the white region bounded by the blue and red lines in figure 1. It is straightforward to show via a linear transformation [16] on the coordinates  $(x = \cos \theta, y = -1/Ar)$  that the latter bound is equivalent to the absence of black droplets [17].




FIG. 1. **Parameter space.** The blue and red lines denote the boundaries in the parameter space  $(mA, A\ell)$  for which the holographic computation is valid. The hashed red region is where acceleration horizons are present and the hashed blue region is where the metric signature is not preserved, leaving the white region as the physical parameter space.

As discussed in [12, 13], setting  $m = 0$  removes the black hole horizon, and leaves pure AdS spacetime in Rindler-type coordinates. Performing the coordinate transformation [15]:

$$1 + \frac{R^2}{\ell^2} = \frac{1 + (1 - A^2\ell^2)r^2/\ell^2}{(1 - A^2\ell^2)\Omega^2}, \quad R \sin \vartheta = \frac{r \sin \theta}{\Omega}, \quad (4)$$

recovers AdS in global coordinates:

$$ds_{AdS}^2 = -\left(1 + \frac{R^2}{\ell^2}\right)\alpha^2 dt^2 + \frac{dR^2}{1 + \frac{R^2}{\ell^2}} + R^2\left(d\vartheta^2 + \sin^2 \vartheta \frac{d\phi^2}{K^2}\right), \quad (5)$$

but with the notable feature that the time coordinate is not the expected AdS time, but is rescaled by a factor of  $\alpha = \sqrt{1 - A^2\ell^2}$ . Conventionally, we choose the normalisation of our time coordinate so that it corresponds to the “time” of an asymptotic observer. While this is potentially a slightly slippery concept in AdS, taken together with the spherical asymptotic spatial coordinates, this scaling suggests that the correct time coordinate is not in fact  $t$ , but rather  $\tau = \alpha t$ , giving a rescaling of the time-coordinate in (1). If we now proceed with this metric, and compute the temperature associated with the black hole (also the temperature of the boundary field theory), then we obtain

$$T = \frac{f'(r_+)}{4\pi\alpha} = \frac{1 + 3\frac{r_+^2}{\ell^2} - A^2r_+^2 \left(2 + \frac{r_+^2}{\ell^2} - A^2r_+^2\right)}{4\pi\alpha r_+(1 - A^2r_+^2)}, \quad (6)$$

where  $f(r_+) = 0$ .

It is worth pausing to reflect on this result. In past work [11–13], the standard time coordinate appearing in the AdS C-metric was used to derive the temperature of the black hole horizon. This appeared to be a natural approach as the blackening factor of the metric was in its canonical form. However, as pointed out in [18],

normalising the time and timelike Killing vector is key to obtaining the correct thermodynamics, although the method of obtaining this correct normalisation was less transparent. Here, having uncovered this suggestive result, we now proceed carefully with considering thermodynamics of the accelerating black hole. As usual, we will take the entropy to be one quarter of the horizon area:

$$S = \frac{\mathcal{A}}{4} = \frac{\pi r_+^2}{K(1 - A^2r_+^2)}. \quad (7)$$

The remaining task is to correctly identify the black hole mass, often the biggest challenge in studying thermodynamics of black holes with non-trivial asymptotics. In what follows, we will provide two independent arguments, beginning with the conformal completion method [19, 20]. Although consistency of the thermodynamic relations is a common method for deriving thermodynamics (used for example in [14]), we do not consider this sufficient; hence we return to our theme of holography, computing the holographic stress tensor of the boundary theory, thereby confirming our result. As an ancillary argument, we finally check consistency with a computation of the free energy.

The first argument uses the Ashtekar–Das definition of conformal mass [19, 20], which extracts the mass via conformal regularisation of the AdS C-metric near the boundary. The idea is to perform a conformal transformation on (1),  $\bar{g}_{\mu\nu} = \bar{\Omega}^2 g_{\mu\nu}$ , to remove the divergence near the boundary, then obtain a conserved charge by integrating the conserved current

$$Q(\xi) = \frac{\ell}{8\pi} \lim_{\bar{\Omega} \rightarrow 0} \oint \frac{\ell^2}{\bar{\Omega}} N^\alpha N^\beta \bar{C}^\nu{}_{\alpha\mu\beta} \xi_\nu d\bar{S}^\mu \quad (8)$$

composed of the Weyl tensor of the conformal metric,  $\bar{C}^\mu{}_{\alpha\nu\beta}$ , the normal to the boundary,  $N_\mu = \partial_\mu \bar{\Omega}$ , and a suitable Killing vector for the mass,  $\xi = \partial_\tau$ . Even though the conformal completion is not unique, the charge thus obtained is independent of the choice of conformal completion. We pick  $\bar{\Omega} = \ell\Omega r^{-1}$ , which provides a smooth conformal completion in the limit  $A = 0$ . The spacelike surface element tangent to  $\bar{\Omega} = 0$  is

$$d\bar{S}_\mu = \delta_\mu^\tau \frac{\ell^2 (d \cos \theta) d\phi}{\alpha K}, \quad (9)$$

obtained by inserting  $Ar \cos \theta = -1$  into the metric  $\bar{g}_{\mu\nu}$  and computing the relevant determinant. This yields

$$M = Q(\partial_\tau) = \alpha \frac{m}{K} \quad (10)$$

for the mass, in agreement with the temperature (6), but in contrast to previous results [11, 14]. The absence of acceleration horizons ensures that  $M$  vanishes in the limit  $A\ell \rightarrow 1$  only for  $m = 0$  and is positive otherwise.

It is now straightforward to verify the first law and Smarr [21] relation

$$\begin{aligned} \delta M &= T\delta S + V\delta P - \lambda_+ \delta\mu_+ - \lambda_- \delta\mu_-, \\ M &= 2TS - 2PV, \end{aligned} \quad (11)$$

using (6), (7), and (10), provided

$$V = \frac{4}{3} \frac{\pi}{K\alpha} \left[ \frac{r_+^3}{(1 - A^2 r_+^2)^2} + mA^2 \ell^4 \right], \quad (12)$$

$$\lambda_{\pm} = \frac{1}{\alpha} \left[ \frac{r_+}{1 - A^2 r_+^2} - m \left( 1 \pm \frac{2A\ell^2}{r_+} \right) \right],$$

where  $P = 3/8\pi\ell^2$  is the thermodynamic pressure associated with the cosmological constant [4], and  $\lambda_{\pm}$  are the thermodynamic lengths introduced in [12, 13] that are conjugate to the tensions. We have included the possibility that the tensions vary, as otherwise the system is constrained and identification of the correct parameters can be misleading.

We now turn to another method for deriving the thermodynamic mass, by computing the holographic stress tensor. This provides an alternate and completely independent method of computation, and will reveal the dual interpretation of this system. The idea here is to perform a Fefferman–Graham expansion of the metric [22], identifying the fall-off of sub-leading terms in the metric at the boundary. These are then used to compute the dual stress-energy tensor that can be integrated to give the mass of the system.

The action, including boundary counterterms [23, 24], is

$$I[g] = \frac{1}{16\pi} \int_M d^4x \sqrt{-g} \left[ R + \frac{6}{\ell^2} \right] + \frac{1}{8\pi} \int_{\partial M} d^3x \sqrt{-h} \mathcal{K} - \frac{1}{8\pi} \int_{\partial M} d^3x \sqrt{-h} \left[ \frac{2}{\ell} + \frac{\ell}{2} \mathcal{R}(h) \right], \quad (13)$$

where  $\mathcal{K}_{ab}$  is the extrinsic curvature of the boundary metric, evaluated asymptotically in an appropriate coordinate system, defined presently.  $h_{ab}$  is the intrinsic metric on  $\partial\mathcal{M}$ , and  $\mathcal{R}$  its Ricci curvature. Varying the action gives the energy momentum tensor:

$$8\pi\mathcal{T}_{ab} = \ell\mathcal{G}_{ab}(h) - \frac{2}{\ell}h_{ab} - \mathcal{K}_{ab} + h_{ab}\mathcal{K}. \quad (14)$$

To compute these terms requires new coordinates near the boundary of AdS, typically parametrised by Fefferman–Graham coordinates, in which

$$ds^2 = \frac{\ell^2}{\rho^2} d\rho^2 + \frac{\rho^2}{\ell^2} \left( \gamma_{ab}^{(0)} + \frac{1}{\rho^2} \gamma_{ab}^{(2)} + \dots \right) dx^a dx^b. \quad (15)$$

Although often one identifies a  $\rho$  coordinate globally, due to the complexity of (1), we instead perform an asymptotic expansion for the coordinate transformation, writing

$$\frac{1}{Ar} = -x - \sum F_n(x) \rho^{-n}, \quad \cos\theta = x + \sum G_n(x) \rho^{-n}. \quad (16)$$

The functions  $F_n$  and  $G_n$  are fixed by the required fall-off properties of (15), apart from  $F_1$ , that we choose to write as

$$F_1(x) = -\frac{(1 - A^2\ell^2 X)^{3/2}}{A\omega(x)\alpha}, \quad (17)$$

in order to elucidate the conformal degree of freedom in the boundary metric,  $\omega$ , with  $X = (1 - x^2)(1 + 2mA\ell^2 X)$ . Computing this boundary metric,  $ds_{(0)}^2 = \gamma_{ab}^{(0)} dx^a dx^b$ , we find it sufficient to truncate the series (16) at  $n = 4$  and find:

$$ds_{(0)}^2 = -\omega^2 d\tau^2 + \frac{\omega^2 \alpha^2 \ell^2 dx^2}{X(1 - A^2\ell^2 X)^2} + \frac{X\omega^2 \alpha^2 \ell^2 d\phi^2}{K^2(1 - A^2\ell^2 X)}. \quad (18)$$

Note that the transformation (16) is valid in general only when  $A^2\ell^2 X < 1$ , which is precisely the constraint that acceleration horizons are absent.

The expectation value of the energy momentum of the  $CFT_3$  can then be calculated, yielding

$$\langle \mathcal{T}_a^b \rangle = \lim_{\rho \rightarrow \infty} \frac{\rho}{\ell} \mathcal{T}_a^b = \begin{pmatrix} -\rho_E & 0 & 0 \\ 0 & \frac{\rho_E}{2} + \Pi & 0 \\ 0 & 0 & \frac{\rho_E}{2} - \Pi \end{pmatrix}, \quad (19)$$

where

$$\rho_E = \frac{m}{8\pi\ell^2\alpha^3\omega^3} (1 - A^2\ell^2 X)^{3/2} (2 - 3A^2\ell^2 X), \quad (20)$$

$$\Pi = \frac{3mA^2 X}{16\pi\alpha^3\omega^3} (1 - A^2\ell^2 X)^{\frac{3}{2}}.$$

We can re-express this in the language of the fluid/gravity correspondence (for a review and references see [25]) as

$$\langle \mathcal{T}_{ab} \rangle = \frac{3}{2} d(x) U_a U_b + \frac{d(x)}{2} \gamma_{ab}^{(0)} + \xi \Theta_{ab}, \quad (21)$$

where the 4-velocity  $U$  and  $\Theta$  are defined as

$$U = \sqrt{\frac{3 - 2A^2\ell^2 X}{3\omega(1 - A^2\ell^2 X)}} \partial_\tau - \frac{AX}{\sqrt{3\omega\alpha}} \sqrt{(1 - A^2\ell^2 X)} \partial_x,$$

$$\Theta_{ab} = C_{abd} U^d + C_{bad} U^d, \quad (22)$$

with  $C_{abc} = \nabla_{[c} R_{b]a} - g_{a[b} \nabla_{c]} R/4$  the Cotton tensor [26] of the metric (18); boundary indices are raised and lowered with  $\gamma_{ab}^{(0)}$ .

The fact that the boundary is non-conformally flat leads to the inclusion of non-hydrodynamic corrections to the perfect fluid, with

$$d(x) = \frac{m\sqrt{(1 - A^2\ell^2 X)^5}}{4\pi\ell^2\omega^3\alpha^3}. \quad (23)$$

The universal (in the sense of independent of black hole mass and acceleration) coefficient is

$$\xi = \frac{\ell^2}{8\pi\sqrt{3}} = \sqrt{\frac{2}{3}} \frac{1}{12\pi} k^{1/2} N^{3/2}, \quad (24)$$

using the standard  $AdS_4/CFT_3$  dictionary to identify  $k$  with the level and  $N$  with the rank of the gauge groups of the ABJ(M) theory. All dissipative corrections enumerated in [25] are seen to vanish, ensuring the uniqueness of the decomposition given in (21).




FIG. 2. **Free energy.** The red curve is the Schwarzschild-AdS case, illustrating the well-known Hawking–Page transition, situated at a temperature given by the intersection of the red curve with  $F = 0$ . We do not know of any such interpretation for all other curves with  $\mu \neq 0$ . The upper parts of these curves do not continue to arbitrarily large  $M$  but terminate at the boundary given in figure 1; this is visible in the above plot only for  $4\mu_- = 0.9$ .

Integrating the energy density, measured with respect to a static geodesic observer  $\rho_E$  yields

$$M = \int \rho_E \sqrt{-\gamma^{(0)}} dx d\phi = \frac{\alpha m}{K}, \quad (25)$$

for the mass, in agreement with (10), and independent of the conformal frame (the choice of  $\omega$ ).

We see from (18) that the boundary metric does not satisfy Dirichlet boundary conditions. However for arbitrary variations of the parameters  $A$  and  $m$  we find that  $\delta I[g] = 0$  provided we set  $\alpha = \sqrt{1 - A^2 \ell^2}$ . Our analysis therefore points towards the possibility of generalizing the conditions under which the conformal and holographic methods coincide for the mass computation [27, 28].

Finally, let us return to the computation of the action (26). We find

$$I = \frac{\beta}{2\alpha K} \left( m - 2mA^2\ell^2 - \frac{r_+^3}{\ell^2(1 - A^2r_+^2)} \right), \quad (26)$$

using the time coordinate  $\tau$ . Some simple algebra then yields the expected result  $F = I/\beta = M - TS$  for the free energy, which we plot in figure 2.

Although similar in form, the behaviour of the free energy no longer indicates the presence of a standard Hawking–Page transition [29]. As the string tension is fixed for the curves in the plot, no transition to pure radiation (with zero tension) is possible. One may, however, speculate that a transition to a different type of spacetime (for example that of the expanding spherical wave with an attached semi-infinite string of given ten-

sion, similar to [30]) may still be possible—such an investigation, however, remains to be carried out.

We can also explore the isoperimetric ratio, or the ratio of volume to areal radius:  $\mathcal{R} = \left(\frac{3V}{\omega_2}\right)^{\frac{1}{3}} \left(\frac{\omega_2}{A}\right)^{\frac{1}{2}}$  (recall  $\omega_2 = 4\pi/K$  here). Using (7) and (12) we find  $\mathcal{R} \geq 1$ , indicating it satisfies the standard *reverse isoperimetric inequality* [31], not adding to the notable exceptions [32–34].

Our full and consistent description of the thermodynamics of an accelerating black hole reconciles discrepancies and conflicts that have appeared in previous investigations of this system [11, 12, 14]. For example, while a set of thermodynamic variables for charged accelerating black holes respecting the first law was obtained [11–13] the computations employed an incorrectly normalized Killing vector at infinity; furthermore the resultant free energy is not consistent with the standard Euclidean action calculation. Alternate expressions for mass and temperature have been posited [14], with the tension of one deficit held fixed to zero. The other tension, while allowed to vary, was not included in the first law, which was derived by assuming integrability of a scaling of mass and temperature. However no physical interpretation was given either for this scaling or for why the energy content of the tension was thermodynamically irrelevant. Furthermore, the vacuum accelerating black hole has an acceleration horizon, akin to a Rindler horizon, and the full structure of the spacetime is that of two accelerating black holes in two Rindler regions. Whether one should be considering a single thermodynamic mass and first law with an additional horizon and black hole, or whether, as suggested in [35], this should be considered as a single system with a mass dipole is an open question.

We also found a decomposition of the dual stress energy tensor for the accelerating black hole in terms of a perfect fluid plus conformal tensors. We obtained a new “universal” coefficient,  $\xi$  that is relevant for the fluid/gravity correspondence in non-conformally flat manifolds. It is natural to expect the existence of non-hydrodynamic corrections to the energy momentum tensor for an even-dimensional CFT due to the conformal anomalies. We have shown here that a similar picture arises in the odd-dimensional case by explicitly constructing the relevant non-hydrodynamic corrections necessary to provide a complete holographic description of the system, c.f. [36].

It would also be interesting to make a connection with the weak coupling calculation of stress tensors in the presence of conical deficits [37]. Future work will involve investigating accelerating black holes with rotation, scalar fields [38, 39], and charge. The latter system will be a challenge due to the asymptotic structure of the gauge field.

Since our computation is independent of the conformal frame, we can compare to investigations of holographic C-metrics with an acceleration horizon. For example, by choosing  $\omega^2 = (1 - A^2\ell^2 X)\alpha^{-2}$ , we recover the form of

the boundary metric employed in [17], and our coordinate transformation (16) is now valid throughout  $x \in [-1, 1]$ . However, if the condition  $A^2 \ell^2 X < 1$  is violated, then a black droplet/black funnel is present, and we no longer have an equilibrium temperature for the system in general. The boundary geometry corresponds to a black hole in a spatially compact universe, and so there is no spatial asymptotic region as pointed out in [17]. However, with the full conformal degree of freedom present in our expression, we can easily remedy this shortcoming by, for example, multiplying the  $\omega$  above by  $\frac{1}{\sqrt{1-x}}$ , giving an  $AdS_2 \times S^1$  asymptotic region at  $x = 1$  with the  $AdS_2$  and  $S^1$  radii being equal. If we multiply by  $\frac{1}{\sqrt{1-x^2}}$  then there are actually two  $AdS_2 \times S^1$  asymptotic regions at  $x = \pm 1$  and  $\gamma_{ab}^{(0)}$  yields the geometry of a wormhole when there are no horizons at the boundary. The  $AdS_2 \times S^1$  asymptotic geometry is supersymmetric and to our knowledge has been unnoticed so far in the literature.

## Acknowledgements

We thank Alex Buchel, Rob Myers, Jiří Podolský, Mukund Rangamani and Kostas Skenderis for valuable discussions. This work was supported in part by the Natural Sciences and Engineering Research Council of Canada, a Chilean FONDECYT [Grant No. 1181047 (AA), 1170279 (AA), 1161418 (AA), 3170035 (A.Ö.)], CONYCT-RCUK [Newton-Picarte Grants DPI20140053 (AA) and DPI20140115 (AA)], by the STFC [consolidated grant ST/P000371/1 (RG)]. MA and AO would like to thank Perimeter Institute and University of Waterloo for hospitality. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Research, Innovation and Science.

- 
- [1] J. D. Bekenstein, *Black holes and entropy*, Phys. Rev. D **7**, 2333 (1973).
- [2] J. D. Bekenstein, *Generalized second law of thermodynamics in black hole physics*, Phys. Rev. D **9**, 3292 (1974).
- [3] S. W. Hawking, *Particle Creation by Black Holes*, Commun. Math. Phys. **43**, 199 (1975) Erratum: [Commun. Math. Phys. **46**, 206 (1976)].
- [4] D. Kubizňák, R. B. Mann and M. Teo, *Black hole chemistry: thermodynamics with Lambda*, Class. Quant. Grav. **34**, 063001 (2017) [arXiv:1608.06147 [hep-th]].
- [5] W. Kinnersley and M. Walker, *Uniformly accelerating charged mass in general relativity*, Phys. Rev. D **2**, 1359 (1970).
- [6] J. F. Plebanski and M. Demianski, *Rotating, charged, and uniformly accelerating mass in general relativity*, Annals Phys. **98**, 98 (1976).
- [7] O. J. C. Dias and J. P. S. Lemos, *Pair of accelerated black holes in anti-de Sitter background: AdS C metric*, Phys. Rev. D **67**, 064001 (2003) [hep-th/0210065].
- [8] J. B. Griffiths and J. Podolsky, *A New look at the Plebanski-Demianski family of solutions*, Int. J. Mod. Phys. D **15**, 335 (2006) [gr-qc/0511091].
- [9] R. Gregory and M. Hindmarsh, *Smooth metrics for snapping strings*, Phys. Rev. D **52** (1995) 5598–5605, [gr-qc/9506054].
- [10] F. Dowker, J. P. Gauntlett, D. A. Kastor and J. H. Traschen, *Pair creation of dilaton black holes*, Phys. Rev. D **49**, 2909 (1994) [hep-th/9309075].
- [11] M. Appels, R. Gregory, and D. Kubizňák, *Thermodynamics of Accelerating Black Holes*, Phys. Rev. Lett. **117**, 131303 (2016) [arXiv:1604.08812 [hep-th]].
- [12] M. Appels, R. Gregory, and D. Kubizňák, *Black Hole Thermodynamics with Conical Defects*, JHEP **1705**, 116 (2017) [arXiv:1702.00490 [hep-th]].
- [13] R. Gregory, *Accelerating Black Holes*, J. Phys. Conf. Ser. **942**, no. 1, 012002 (2017) [arXiv:1712.04992 [hep-th]].
- [14] M. Astorino, *Thermodynamics of Regular Accelerating Black Holes*, Phys. Rev. D **95**, no. 6, 064007 (2017) [arXiv:1612.04387 [gr-qc]].
- [15] Podolsky J, *Accelerating black holes in anti-de Sitter universe*, Czech. J. Phys. **52**, 1 (2002) [gr-qc/0202033].
- [16] Kenneth Hong and Edward Teo. *A New form of the C metric*, Class. Quant. Grav. **20**, 3269 (2003) [gr-qc/0305089].
- [17] V. E. Hubeny, D. Marolf and M. Rangamani, *Black funnels and droplets from the AdS C-metrics*, Class. Quant. Grav. **27**, 025001 (2010) [arXiv:0909.0005 [hep-th]].
- [18] G. W. Gibbons, M. J. Perry and C. N. Pope, *The First law of thermodynamics for Kerr-anti-de Sitter black holes*, Class. Quant. Grav. **22**, 1503 (2005) [hep-th/0408217].
- [19] A. Ashtekar and S. Das, *Asymptotically Anti-de Sitter space-times: Conserved quantities*, Class. Quant. Grav. **17** (2000) L17–L30, [hep-th/9911230].
- [20] S. Das and R. Mann, *Conserved quantities in Kerr-anti-de Sitter space-times in various dimensions*, JHEP **08** (2000) 033, [hep-th/0008028].
- [21] L. Smarr, *Mass formula for Kerr black holes*, Phys. Rev. Lett. **30**, 71 (1973) Erratum: [Phys. Rev. Lett. **30**, 521 (1973)].
- [22] C. Fefferman and C. Robin Graham, *Conformal Invariants in Elie Cartan et les Mathématiques d'aujourd'hui* (Astérisque, 1985) Numero Hors Serie, 95-116, see also: [arXiv:0710.0919 [math.DG]].
- [23] V. Balasubramanian and P. Kraus, *A Stress tensor for Anti-de Sitter gravity*, Commun. Math. Phys. **208**, 413 (1999) [hep-th/9903238].
- [24] R. B. Mann, *Misner string entropy*, Phys. Rev. D **60**, 104047 (1999) [hep-th/9903229].
- [25] M. Rangamani, *Gravity and Hydrodynamics: Lectures on the fluid-gravity correspondence*, Class. Quant. Grav. **26** (2009) 224003 [arXiv:0905.4352 [hep-th]].
- [26] A. Mukhopadhyay, A. C. Petkou, P. M. Petropoulos, V. Pozzoli and K. Siampos, *Holographic perfect fluidity*,

- Cotton energy-momentum duality and transport properties*, JHEP **1404** (2014) 136 [[arXiv:1309.2310](#)] [[hep-th](#)].
- [27] S. Hollands, A. Ishibashi and D. Marolf, *Comparison between various notions of conserved charges in asymptotically AdS-spacetimes*, Class. Quant. Grav. **22**, 2881 (2005) [[hep-th/0503045](#)].
- [28] I. Papadimitriou and K. Skenderis, *Thermodynamics of asymptotically locally AdS spacetimes*, JHEP **0508**, 004 (2005) [[hep-th/0505190](#)].
- [29] S. W. Hawking and D. N. Page, *Thermodynamics of Black Holes in anti-De Sitter Space*, Commun. Math. Phys. **87**, 577 (1983).
- [30] J. Podolsky and J. B. Griffiths, *A Snapping cosmic string in a de Sitter or anti-de Sitter universe*, Class. Quant. Grav. **21**, 2537 (2004) [[gr-qc/0403089](#)].
- [31] M. Cvetič, G. W. Gibbons, D. Kubizňák and C. N. Pope, *Black Hole Enthalpy and an Entropy Inequality for the Thermodynamic Volume*, Phys. Rev. D **84**, 024037 (2011) [[arXiv:1012.2888](#)] [[hep-th](#)].
- [32] R. A. Hennigar, D. Kubizňák and R. B. Mann, *Entropy Inequality Violations from Ultraspinning Black Holes*, Phys. Rev. Lett. **115**, no. 3, 031101 (2015) [[arXiv:1411.4309](#)] [[hep-th](#)].
- [33] R. A. Hennigar, D. Kubizňák, R. B. Mann and N. Musoke, *Ultraspinning limits and super-entropic black holes*, JHEP **1506**, 096 (2015) [[arXiv:1504.07529](#)] [[hep-th](#)].
- [34] W. G. Brenna, R. B. Mann and M. Park, *Mass and Thermodynamic Volume in Lifshitz Spacetimes*, Phys. Rev. D **92**, no. 4, 044015 (2015) [[arXiv:1505.06331](#)] [[hep-th](#)].
- [35] K. Dutta, S. Ray and J. Traschen, *Boost mass and the mechanics of accelerated black holes*, Class. Quant. Grav. **23**, 335 (2006) [[hep-th/0508041](#)].
- [36] G. Bernardi de Freitas and H. S. Reall, *Algebraically special solutions in AdS/CFT*, JHEP **1406**, 148 (2014) [[arXiv:1403.3537](#)] [[hep-th](#)].
- [37] J. S. Dowker, *Quantum Field Theory on a Cone*, J. Phys. A **10**, 115 (1977).
- [38] A. Anabalón and H. Maeda, *New Charged Black Holes with Conformal Scalar Hair*, Phys. Rev. D **81**, 041501 (2010) [[arXiv:0907.0219](#)] [[hep-th](#)].
- [39] A. Anabalón, *Exact Black Holes and Universality in the Backreaction of non-linear Sigma Models with a potential in (A)dS<sub>4</sub>*, JHEP **1206**, 127 (2012) [[arXiv:1204.2720](#)] [[hep-th](#)].