

LEVEL CURVES OF RATIONAL FUNCTIONS AND UNIMODULAR POINTS ON RATIONAL CURVES

FEDOR PAKOVICH AND IGOR E. SHPARLINSKI

ABSTRACT. We obtain an improvement and broad generalisation of a result of N. Ailon and Z. Rudnick (2004) on common zeros of shifted powers of polynomials. Our approach is based on reducing this question to a more general question of counting intersections of level curves of complex functions. We treat this question via classical tools of complex analysis and algebraic geometry.

1. INTRODUCTION

Recall that Ailon and Rudnick [1, Theorem 1] have shown that for any multiplicatively independent polynomials $P_1(z)$ and $P_2(z)$ with complex coefficients there exists a polynomial $F(z) \in \mathbb{C}[z]$ such that for any positive integer k the greatest common divisor of $P_1(z)^k - 1$ and $P_2(z)^k - 1$ divides F , that is

$$\gcd(P_1(z)^k - 1, P_2(z)^k - 1) \mid F(z), \quad k = 1, 2, \dots$$

Since it is easy to see that for a non-trivial polynomial $P(z) \in \mathbb{C}[z]$ the multiplicity of any factor of $P(z)^k - 1$ does not exceed $\deg P$, the theorem of Ailon and Rudnick is equivalent to the following statement: if P_1 and P_2 are complex polynomials, then

$$(1.1) \quad \#\bigcup_{k=1}^{\infty} \{z \in \mathbb{C} : P_1(z)^k = P_2(z)^k = 1\} \leq C(P_1, P_2),$$

for some constant $C(P_1, P_2)$ that depends only on P_1 and P_2 , unless for some non-zero integers m_1 and m_2 we have

$$(1.2) \quad P_1^{m_1}(z)P_2^{m_2}(z) = 1$$

identically. Different versions and generalization Ailon-Rudnick result have been studied in many recent papers (see, for example, [5, 6, 9, 10] and the references therein).

2010 *Mathematics Subject Classification.* 11D61, 12D10, 30C15, 30J10.

Key words and phrases. Unimodular points, Ailon and Rudnick theorem, Blaschke product.

The method of Ailon and Rudnick [1] relies on a result conjectured by Lang and proved by Ihara, Serre and Tate, which states that the intersection of an irreducible curve \mathcal{C} in $\mathbb{C}^* \times \mathbb{C}^*$ with the roots of unity $\mu_\infty \times \mu_\infty$ is finite, unless \mathcal{C} is of the form $X^n Y^m - \eta = 0$ or $X^m - \eta Y^n = 0$, where $\eta \in \mu_\infty$, that is unless \mathcal{C} is a translate by a torsion point of an algebraic subgroup of $\mathbb{C}^* \times \mathbb{C}^*$ (see [7], [8], and also [2]).

Corvaja, Masser, and Zannier in the paper of [3] ask about a possible extension of the Lang statement [7], where instead of the intersection of \mathcal{C} with $\mu_\infty \times \mu_\infty$ the intersection with $S^1 \times S^1$ is considered (here S^1 is treated as the topological closure of torsion points). In particular, they proved that the system

$$(1.3) \quad |z| = |P(z)| = 1,$$

where $P(z)$ is a polynomial, has finitely many solutions, unless $P(z)$ is a monomial. They also remarked that if $P(z)$ is allowed to be a rational function, then the system (1.3) might have infinitely many solutions for non-monomial $P(z)$.

In this paper we consider the system of equations for the level curves

$$(1.4) \quad |P_1(z)| = |P_2(z)| = 1,$$

where $P_1(z)$ and $P_2(z)$ arbitrary rational functions, generalising the systems (1.1) and (1.3). Using classical tools of complex analysis and algebraic geometry, we describe P_1 and P_2 for which this system has infinitely many solutions and provide bounds for the number of solutions in the other cases. Thus, our results can be considered as extensions of the result of Ailon and Rudnick [1] as well as of the Lang statement [7] in the particular case concerning of curves of genus zero.

2. RESULTS

Recall that a *finite Blaschke product* is a rational function $B(z) \in \mathbb{C}(z)$ of the form

$$B(z) = \zeta \prod_{i=1}^n \left(\frac{z - a_i}{1 - \bar{a}_i z} \right)^{m_i},$$

where a_i are complex numbers in the open unit disc

$$\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\},$$

the exponents m_i , $i = 1, \dots, n$, are positive integers, and $|\zeta| = 1$. A rational function $Q(z)$ of the form $Q(z) = B_1(z)/B_2(z)$, where B_1 and B_2 are finite Blaschke products, is called a *quotient of finite Blaschke products*.

In the above notation, our first result is the following.

Theorem 2.1. *Let $\mathcal{C} : F(x, y) = 0$, where $F(x, y) \in \mathbb{C}[x, y]$, be an irreducible algebraic curve of genus zero and of degree $d = \deg F$. Then \mathcal{C} has at most d^2 unimodular points, unless it can be parametrised by some quotients of finite Blaschke products $x = Q_1(z)$ and $y = Q_2(z)$.*

Our second result is the following generalisation of the bound (1.1).

Theorem 2.2. *Let $P_1(z)$ and $P_2(z)$ be complex rational functions of degrees n_1 and n_2 . Then*

$$(2.1) \quad \#\{z \in \mathbb{C} : |P_1(z)| = |P_2(z)| = 1\} \leq (n_1 + n_2)^2,$$

unless

$$(2.2) \quad P_1 = B_1 \circ W \quad \text{and} \quad P_2 = B_2 \circ W$$

for some quotients of finite Blaschke products B_1 and B_2 and rational function W .

In order to see that Theorem 2.2 implies (1.1) it is enough to observe that if a quotient of finite Blaschke products is a polynomial, then this polynomial is necessarily a power. Thus, (2.2) reduces to

$$P_1 = W^{m_1} \quad \text{and} \quad P_2 = W^{m_2},$$

implying (1.2).

Notice that since any quotient of finite Blaschke products maps the unit circle

$$\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$$

to itself, if P_1 and P_2 satisfy (2.2), then the level curves $|P_1(z)| = 1$ and $|P_2(z)| = 1$ have a common component $W^{-1}\{\mathbb{T}\}$, so a bound like (2.1), or any other finiteness result, cannot exist. In particular, this happens if P_1 is a unimodular constant and P_2 is an arbitrary rational function (in this case (2.2) holds for $B_1 = P_1$, $B_2 = z$, and $W = P_2$).

3. PROOFS

Since the inverse Cayley transform

$$z \mapsto T(z) = i \frac{1+z}{1-z}$$

maps \mathbb{D} to the upper half-plane, and the unit circle \mathbb{T} maps under T to the extended real line, a rational function Q is a quotient of a finite Blaschke product if and only if the rational function

$$R = T \circ Q \circ T^{-1}$$

maps $\mathbb{R} \cup \infty$ to $\mathbb{R} \cup \infty$. In turn, the last condition is equivalent to the condition that R has real coefficients (since $R(z)$ and $\overline{R}(z)$ coincide for infinitely many values of z).

Thus, Theorem 2.1 is equivalent to the following statement.

Theorem 3.1. *If an irreducible algebraic curve $\mathcal{C} : F(x, y) = 0$ of genus zero and degree d has more than d^2 real points, then \mathcal{C} can be parametrised by rational functions with real coefficients.*

Proof. Observe that real points of \mathcal{C} belong to the intersection of the curve \mathcal{C} and the curve $\overline{\mathcal{C}} : \overline{F}(x, y) = 0$. Therefore, it follows from the Bézout theorem that whenever \mathcal{C} has more than d^2 real points there exists $c \in \mathbb{C}$ such that $\overline{F} = cF$. Such c must satisfy $c\overline{c} = 1$, implying that we can find a complex number λ such that $\lambda^2 = c$ and $\lambda\overline{\lambda} = 1$. Since

$$\overline{\lambda F} = \overline{\lambda}\lambda^2 F = \lambda F,$$

the polynomial λF has real coefficients, and hence \mathcal{C} can be defined over \mathbb{R} .

Since the maximal number of singular points of a plane curve of degree d does not exceed

$$\frac{(d-1)(d-2)}{2}$$

(see, for example, [4, Page 49]) and \mathcal{C} has more than d^2 real points, \mathcal{C} has a non-singular real point. Finally, an algebraic curve \mathcal{C} of genus zero defined over \mathbb{R} admits a parametrisation by rational functions defined over \mathbb{R} whenever \mathcal{C} has at least one non-singular \mathbb{R} -point (see, for example, [11, Theorem 7.6]). \square

In order to prove Theorem 2.2, recall that if a parametrisation $x = P_1(z)$, $y = P_2(z)$ of an algebraic curve \mathcal{C} of genus zero is proper, that is, if

$$\mathbb{C}(z) = \mathbb{C}(P_1(z), P_2(z)),$$

then

$$\deg P_1 = \deg_y F \quad \text{and} \quad \deg P_2 = \deg_x F,$$

(see, for example, [11, Theorem 4.21]).

Let now P_1 and P_2 be rational functions of degrees n_1 and n_2 . Then the Lüroth theorem implies that there exist a rational function W and rational functions Q_1 and Q_2 such that the equalities (2.2) hold, and

$$x = Q_1(z), \quad y = Q_2(z),$$

is a proper parametrisation of an algebraic curve \mathcal{C} of degree at most $n_1 + n_2$. Therefore, if (2.1) does not hold, then Q_1 and Q_2 are quotients of finite Blaschke products by Theorem 2.1.

Remark 3.2. We observe that the above argument provides a simple geometric criterion for a curve $\mathcal{C} : G(x, y) = 0$ to have infinitely many unimodular points. Namely, considering instead of the curve \mathcal{C} a curve $\widehat{\mathcal{C}} : \widehat{G}(x, y) = 0$, where

$$\widehat{G}(x, y) = G(T(x), T(y)),$$

we reduce the question to the question about real points of $\widehat{\mathcal{C}}$. On the other hand, it is easy to see that an algebraic curve has infinitely many real points if and only if it is defined over \mathbb{R} and has at least one simple \mathbb{R} -point. Indeed, the necessity has been proved above. In the other direction, if a curve defined over \mathbb{R} has a simple \mathbb{R} -point, then the implicit function theorem implies that it has infinitely many \mathbb{R} -points.

ACKNOWLEDGEMENT

The authors would like to thank Yuri Bilu and Umberto Zannier for their comments.

This work of I. E. Shparlinski. was supported in part by the Australian Research Council Grants DP170100786 and DP180100201.

REFERENCES

- [1] N. Ailon and Z. Rudnick, *Torsion points on curves and common divisors of $a^k - 1$ and $b^k - 1$* , Acta Arith., **113** (2004), 31–38. (pp. 1 and 2)
- [2] F. Beukers and C. J. Smyth, *Cyclotomic points on curves*, Number theory for the millenium (Urbana, Illinois, 2000), I, A K Peters, 2002, 67–85. (p. 2)
- [3] P. Corvaja, D. Masser and U. Zannier, *Sharpening ‘Manin–Mumford’ for certain algebraic groups of dimension 2*, Enseign. Math., **59** (2013), 1–45. (p. 2)
- [4] G. Fischer, *Plane algebraic curves*, Amer. Math. Soc., Providence, RI, 2001. (p. 4)
- [5] D. Ghioca, L.-C. Hsia and T. J. Tucker, *On a variant of the Ailon-Rudnick theorem in finite characteristic*, New York Journal of Math., **23** (2017), 213–225. (p. 1)
- [6] L.-C. Hsia and T. J. Tucker, *Greatest common divisors of iterates of polynomials*, Algebra and Number Theory, **11** (2017) 1437–1459. (p. 1)
- [7] S. Lang, *Division points on curves*, Ann. Mat. Pura Appl., **70** (1965), 229–234. (p. 2)
- [8] S. Lang, *Fundamentals of Diophantine geometry*, Springer-Verlag, New York, 1983. (p. 2)
- [9] A. Ostafe, *On some extensions of the Ailon-Rudnick theorem*, Monat. Math., **181** (2016), 451–471. (p. 1)
- [10] H. Pasten and J. T.-Y. Wang, *GCD bounds for analytic functions*, Intern. Math. Res. Not., **2017** (2017), 47–95. (p. 1)

- [11] J. Sendra, F. Winkler and S. Pérez-Díaz, *Rational algebraic curves. A computer algebra approach*, Algorithms and Computation in Mathematics, vol.22. Springer, Berlin, 2008. (p.4)

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV,
P.O.B. 653, BEER SHEVA, 8410501, ISRAEL
E-mail address: pakovich@math.bgu.ac.il

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF NEW SOUTH WALES,
SYDNEY, NSW 2052, AUSTRALIA
E-mail address: igor.shparlinski@unsw.edu.au