

MORSE-NOVIKOV COHOMOLOGY ON COMPLEX MANIFOLDS

LINGXU MENG

ABSTRACT. We view Dolbeault-Morse-Novikov cohomology $H_{\eta}^{p,q}(X)$ as the cohomology of the sheaf $\Omega_{X,\eta}^p$ of η -holomorphic p -forms and give several bimeromorphic invariants. Analogue to Dolbeault cohomology, we establish the Leray-Hirsch theorem and the blow-up formula for Dolbeault-Morse-Novikov cohomology. At last, we consider the relations between Morse-Novikov cohomology and Dolbeault-Morse-Novikov cohomology, moreover, investigate stabilities of their dimensions under the deformations of complex structures. In some aspects, Morse-Novikov and Dolbeault-Morse-Novikov cohomology behave similarly with de Rham and Dolbeault cohomology.

Keywords: Morse-Novikov cohomology, weight θ -sheaf, Dolbeault-Morse-Novikov cohomology, Leray-Hirsch theorem, blow-up formula, sheaf of η -holomorphic functions, bimeromorphic, θ -beti number, η -hodge number, stability.

AMSC: 32C35, 57R19.

1. INTRODUCTION

Let X be a smooth manifold and θ a real closed 1-form on X . Set $\mathcal{A}^p(X)$ the space of real smooth p -forms and define $d_{\theta} : \mathcal{A}^p(X) \rightarrow \mathcal{A}^{p+1}(X)$ as $d_{\theta}\alpha = d\alpha + \theta \wedge \alpha$ for $\alpha \in \mathcal{A}^p(X)$. Clearly, $d_{\theta} \circ d_{\theta} = 0$, so we have a complex

$$\cdots \longrightarrow \mathcal{A}^{p-1}(X) \xrightarrow{d_{\theta}} \mathcal{A}^p(X) \xrightarrow{d_{\theta}} \mathcal{A}^{p+1}(X) \cdots \longrightarrow \cdots,$$

whose cohomology $H_{\theta}^p(X) = H^p(\mathcal{A}^{\bullet}(X), d_{\theta})$ is called the p -th *Morse-Novikov cohomology*. For a complex closed 1-form θ on X , denote $H_{\theta}^p(X, \mathbb{C}) = H^p(\mathcal{A}_{\mathbb{C}}^{\bullet}(X), d_{\theta})$, where $\mathcal{A}_{\mathbb{C}}^{\bullet}(X) = \mathcal{A}^{\bullet}(X) \otimes_{\mathbb{R}} \mathbb{C}$. If θ is real, $H_{\theta}^p(X, \mathbb{C}) = H_{\theta}^p(X) \otimes_{\mathbb{R}} \mathbb{C}$. Similarly, we can define *Morse-Novikov cohomology with compact support* $H_{\theta,c}^p(X)$ and $H_{\theta,c}^p(X, \mathbb{C})$.

This cohomology was originally defined by A. Lichnerowicz ([13]) and D. Sullivan ([24]) in the context of Poisson geometry and infinitesimal computations in topology, respectively. It is well used to study the locally conformally Kählerian (l.c.K.) and locally conformally symplectic (l.c.s.) structures ([2, 3, 4, 10, 12, 26]). $H_{\theta}^*(X)$ can be viewed as the cohomology of a flat bundle (weight line bundle) or a local constant sheaf of \mathbb{R} -modules with finite rank, referring to [24, 14, 16, 17, 29]. As we know, the two viewpoints are equivalent, whereas the latter is much more convenient, seeing [14].

In his seminal paper [16], S. P. Novikov introduced a generalization of the classical Morse theory to the case of circle-valued Morse functions. A. Pajitnov [21] observed the relation of the circle-valued Morse theory to the homology with local coefficients and perturbed de Rham differential, see also [22], p. 414-416.

For smooth manifolds, the Mayer-Vietoris sequence and Poincaré duality theorem were generalized on Morse-Novikov cohomology by S. Haller and T. Rybicki [10]. M. León, B.

López, J. C. Marrero and E. Padrón [12] proved that a compact Riemannian manifold X endowed with a parallel one-form θ has trivial Morse-Novikov cohomology. By Atiyah-Singer index theorem, G. Bande and D. Kotschick [4] found that the Euler characteristic of Morse-Novikov cohomology coincides with the usual Euler characteristic. In [14], we proved several Künneth formulas and theorems of Leray-Hirsch type.

For complex manifolds, I. Vaisman [26] studied the classical operators twisted with a closed one-form on l.c.K. manifolds. In [14], we gave two explicit formulas of blow-ups of complex manifolds for Morse-Novikov cohomology. As we know, de Rham cohomology is closely related to Dolbeault cohomology on complex manifolds, such as Hodge decomposition theorem, hard Lefschetz theorem, Hodge's index theorem, etc.. Inspired by these, it is necessary to study Dolbeault-Morse-Novikov cohomology, which is a generalization of Dolbeault cohomology. Recently, L. Ornea, M. Verbitsky, and V. Vuletescu [20] showed that, for a locally conformally Kähler manifold X with proper potential, $H_{a\bar{\eta}}^{*,*}(X) = 0$ holds for all $a \in \mathbb{C}$ but a discrete countable subset, where η is the $(0, 1)$ -part of Lee form θ of X .

L. Ornea, M. Verbitsky, and V. Vuletescu [19] proved that the blow-up of an l.c.K. manifold along a submanifold is l.c.K. if and only if the submanifold is globally conformally equivalent to a Kähler submanifold. So, it is necessary to consider the variance of the Morse-Novikov ([14]) and Dolbeault-Morse-Novikov cohomology under blowing up.

Theorem 1.1. *Let $\pi : \tilde{X} \rightarrow X$ be the blow-up of a connected complex manifold X along a connected complex submanifold Z and $i_E : E = \pi^{-1}(Z) \rightarrow \tilde{X}$ the inclusion of the exceptional divisor E into \tilde{X} . Suppose that η is a $\bar{\partial}$ -closed $(0, 1)$ -form on X and $\tilde{\eta} = \pi^*\eta$. Then, for any p, q ,*

$$\pi^* + \sum_{i=0}^{r-2} (i_E)_* \circ (h^i \cup) \circ (\pi|_E)^*$$

gives an isomorphism

$$(1) \quad H_{\eta}^{p,q}(X) \oplus \bigoplus_{i=0}^{r-2} H_{\eta|_Z}^{p-1-i, q-1-i}(Z) \xrightarrow{\sim} H_{\tilde{\eta}}^{p,q}(\tilde{X}),$$

where $r = \text{codim}_{\mathbb{C}} Z$ and h is defined in (4).

For $\eta = 0$, S. Rao, S. Yang, and X.-D. Yang [23] first proved there exists an isomorphism (1) on a compact complex manifold X . It seems difficult to write out it explicitly using their method. In [15], we write out an isomorphism explicitly on any (possibly noncompact) base with a different way.

Deformations of complex structures play a significant role in studying Kählerian, balanced, strongly Gauduchon and $\partial\bar{\partial}$ -manifolds. For l.c.K. geometry, we have known the facts that a deformation of a l.c.K. manifold is generally not l.c.K. ([5]) and the class of compact l.c.K. manifolds with potential is stable under small deformations ([18]). These results inspire us to investigate behaviors of Dolbeault-Morse-Novikov cohomology under deformations.

Lemma 1.2. *Let $f : X \rightarrow Y$ be a proper surjective submersion of connected smooth manifolds and θ a real (resp. complex) closed 1-form on X . Then, for any k , the higher direct image $R^k f_* \underline{\mathbb{R}}_{X,\theta}$ (resp. $R^k f_* \underline{\mathbb{C}}_{X,\theta}$) is a local system of \mathbb{R} (resp. \mathbb{C})-modules with finite rank.*

Using above lemma and the relation between Morse-Novikov and Dolbeault-Morse-Novikov cohomologies, we get the theorem of stability of η -hodge numbers under the deformation.

Theorem 1.3. *Let $f : X \rightarrow Y$ be a family of complex manifolds and θ a complex closed 1-form on X . Assume $b_k(X_o, \theta|_{X_o}) = \sum_{p+q=k} h_{\eta|_{X_o}}^{p,q}(X_o)$ for some k and some point $o \in Y$, where η is the $(0, 1)$ -part of θ . Then, for any t near o , $h_{\eta|_{X_t}}^{p,q}(X_t) = h_{\eta|_{X_o}}^{p,q}(X_o)$, where η is the $(0, 1)$ -part of θ and $p + q = k$.*

In this article, we investigate the Dolbeault-Morse-Novikov cohomology via the theory of sheaves. In Sec. 2 and 3, we recall the Morse-Novikov cohomology and define the Dolbeault-Morse-Novikov cohomology, respectively. In Sec. 4, we study the properties of the sheaf $\mathcal{O}_{X,\eta}$ of η -holomorphic functions and show that $H_{\eta}^{p,0}(X)$, $H_{\eta,c}^{p,0}(X)$, $H_{\eta}^{0,p}(X)$ and $H_{\eta,c}^{0,p}(X)$ are all bimeromorphic invariants. In particular, we prove Leray-Hirsch theorem and Theorem 1.1. In Sec. 5, Lemma 1.2 and Theorem 1.3 are proved.

2. MORSE-NOVIKOV COHOMOLOGY

We first recall the weight θ -sheaf, referring to [14]. Let \mathcal{A}_X^k be the sheaf of germs of real smooth k -forms and $\mathbb{R}_X, \mathbb{C}_X$ be constant sheaves with coefficient \mathbb{R}, \mathbb{C} on X , respectively. Set $\mathcal{A}_{X,\mathbb{C}}^k = \mathcal{A}_X^k \otimes_{\mathbb{R}_X} \mathbb{C}_X$. Define $d_{\theta} : \mathcal{A}_{X,\mathbb{C}}^k \rightarrow \mathcal{A}_{X,\mathbb{C}}^{k+1}$ as $d_{\theta}\alpha = d\alpha + \theta \wedge \alpha$, for $\alpha \in \mathcal{A}_{X,\mathbb{C}}^k$.

Definition 2.1. *The kernel of $d_{\theta} : \mathcal{A}_{X,\mathbb{C}}^0 \rightarrow \mathcal{A}_{X,\mathbb{C}}^1$ is called a weight θ -sheaf, denoted by $\underline{\mathbb{C}}_{X,\theta}$.*

Locally, $\theta = du$ for a smooth complex-valued function u , so $d_{\theta} = e^{-u} \circ d \circ e^u$ and $\underline{\mathbb{C}}_{X,\theta} = \mathbb{C}e^{-u}$. Hence, the weight θ -sheaf $\underline{\mathbb{C}}_{X,\theta}$ is a local system of \mathbb{C} -modules with rank 1. We have a resolution of soft sheaves of $\underline{\mathbb{C}}_{X,\theta}$

$$0 \longrightarrow \underline{\mathbb{C}}_{X,\theta} \xrightarrow{i} \mathcal{A}_{X,\mathbb{C}}^0 \xrightarrow{d_{\theta}} \mathcal{A}_{X,\mathbb{C}}^1 \xrightarrow{d_{\theta}} \cdots \xrightarrow{d_{\theta}} \mathcal{A}_{X,\mathbb{C}}^n \longrightarrow 0,$$

where i is the natural inclusion. So

$$H_{\theta}^*(X, \mathbb{C}) \cong H^*(X, \underline{\mathbb{C}}_{X,\theta}), \quad H_{\theta,c}^*(X, \mathbb{C}) \cong H_c^*(X, \underline{\mathbb{C}}_{X,\theta}).$$

For d_{θ} -closed $\alpha \in \mathcal{A}_{\mathbb{C}}^*(X)$, denote by $[\alpha]_{\theta}$ (resp. $[\alpha]_{\theta,c}$) the class in $H_{\theta}^*(X, \mathbb{C})$ (resp. $H_{\theta,c}^*(X, \mathbb{C})$).

Assume X is also oriented. Let \mathcal{D}_X^k be the sheaf of germs of real k -currents and $\mathcal{D}_{X,\mathbb{C}}^k = \mathcal{D}_X^k \otimes_{\mathbb{R}_X} \mathbb{C}_X$. Similarly, define $d_{\theta} : \mathcal{D}_{X,\mathbb{C}}^k \rightarrow \mathcal{D}_{X,\mathbb{C}}^{k+1}$ as $d_{\theta}T = dT + \theta \wedge T$ for $T \in \mathcal{D}_{X,\mathbb{C}}^k$. We have another resolution

$$0 \longrightarrow \underline{\mathbb{C}}_{X,\theta} \xrightarrow{i} \mathcal{D}_{X,\mathbb{C}}^0 \xrightarrow{d_{\theta}} \mathcal{D}_{X,\mathbb{C}}^1 \xrightarrow{d_{\theta}} \cdots \xrightarrow{d_{\theta}} \mathcal{D}_{X,\mathbb{C}}^n \longrightarrow 0,$$

of soft sheaves of $\underline{\mathbb{C}}_{X,\theta}$, where i is the natural inclusion. By [6], p. 213 (6.3) (6.4) and p. 217 (7.8), the natural morphism $\mathcal{A}_{X,\mathbb{C}}^{\bullet} \hookrightarrow \mathcal{D}_{X,\mathbb{C}}^{\bullet}$ of resolutions induces isomorphisms

$$H_{\theta}^*(X, \mathbb{C}) \xrightarrow{\sim} H^*(\mathcal{D}_{\mathbb{C}}^{\bullet}(X), d_{\theta}), \quad H_{\theta,c}^*(X, \mathbb{C}) \xrightarrow{\sim} H^*(\mathcal{D}'_{\mathbb{C},c}{}^{\bullet}(X), d_{\theta}).$$

For d_{θ} -closed $T \in \mathcal{D}_{\mathbb{C}}^*(X)$, denote by $[T]_{\theta}$ (resp. $[T]_{\theta,c}$) the class in $H_{\theta}^*(X, \mathbb{C})$ (resp. $H_{\theta,c}^*(X, \mathbb{C})$).

Lemma 2.2 ([14]). *Let X be a connected smooth manifold and θ a complex closed 1-form on X .*

(1) $\underline{\mathbb{C}}_{X,\theta} \cong \underline{\mathbb{C}}_X$ if and only if θ is exact. More precisely, if $\theta = du$ for $u \in \mathcal{A}_{\mathbb{C}}^0(X)$, then $h \mapsto e^u \cdot h$ gives an isomorphism $\underline{\mathbb{C}}_{X,\theta} \xrightarrow{\sim} \underline{\mathbb{C}}_X$ of sheaves.

(2) If μ is a closed 1-form on X , then $\underline{\mathbb{C}}_{X,\theta} \otimes_{\underline{\mathbb{C}}_X} \underline{\mathbb{C}}_{X,\mu} = \underline{\mathbb{C}}_{X,\theta+\mu}$.

(3) Suppose $f : Y \rightarrow X$ is a smooth map between connected smooth manifolds. Then inverse image sheaf $f^{-1}\underline{\mathbb{C}}_{X,\theta} = \underline{\mathbb{C}}_{Y,f^*\theta}$.

Proof. (1) If $\underline{\mathbb{C}}_{X,\theta} \cong \underline{\mathbb{C}}_X$, $H_{\theta}^0(X) = H^0(X, \underline{\mathbb{C}}_{X,\theta}) = \mathbb{C}$. By [10], Example 1.6, θ is exact. Inversely, if $\theta = du$, $\underline{\mathbb{C}}_{X,\theta} = \underline{\mathbb{C}}e^{-u}$, which implies the conclusion.

(2) Locally, $\theta = du$ and $\mu = dv$. Then, $\underline{\mathbb{C}}_{X,\theta} = \mathbb{C}e^{-u}$, $\underline{\mathbb{C}}_{X,\mu} = \mathbb{C}e^{-v}$ and $\underline{\mathbb{C}}_{X,\theta+\mu} = \mathbb{C}e^{-u-v}$, locally. Clearly, the products of functions give an isomorphism $\underline{\mathbb{C}}_{X,\theta} \otimes_{\underline{\mathbb{C}}_X} \underline{\mathbb{C}}_{X,\mu} \rightarrow \underline{\mathbb{C}}_{X,\theta+\mu}$.

(3) Locally, $\theta = du$, $\underline{\mathbb{C}}_{X,\theta} = \mathbb{C}e^{-u}$ and $\underline{\mathbb{C}}_{Y,f^*\theta} = \mathbb{C}e^{-f^*u}$. The pullbacks of functions give an isomorphism $f^{-1}\underline{\mathbb{C}}_{X,\theta} \xrightarrow{\sim} \underline{\mathbb{C}}_{Y,f^*\theta}$. \square

Let X be a smooth manifold and θ, μ complex closed 1-forms on X . The wedge product $\alpha \wedge \beta$ defines a *cup product*

$$\cup : H_{\theta}^p(X, \mathbb{C}) \times H_{\mu}^q(X, \mathbb{C}) \rightarrow H_{\theta+\mu}^{p+q}(X, \mathbb{C}).$$

Similarly, we can define cup products between $H_{\theta}^p(X, \mathbb{C})$ or $H_{\theta,c}^p(X, \mathbb{C})$ and $H_{\mu}^q(X, \mathbb{C})$ or $H_{\mu,c}^q(X, \mathbb{C})$.

Let $f : X \rightarrow Y$ be a smooth map between connected smooth manifolds and θ a complex closed 1-form on Y . Set $\tilde{\theta} = f^*\theta$ and $r = \dim X - \dim Y$.

(i) Define *pullback* $f^* : H_{\theta}^*(Y, \mathbb{C}) \rightarrow H_{\tilde{\theta}}^*(X, \mathbb{C})$ as $[\alpha]_{\theta} \mapsto [f^*\alpha]_{\tilde{\theta}}$. If f is proper, we can also define $f^* : H_{\theta,c}^*(Y, \mathbb{C}) \rightarrow H_{\tilde{\theta},c}^*(X, \mathbb{C})$ in the same way.

(ii) If X and Y are oriented, define *pushout* $f_* : H_{\tilde{\theta},c}^*(X, \mathbb{C}) \rightarrow H_{\theta,c}^{*-r}(Y, \mathbb{C})$ as $[T]_{\tilde{\theta},c} \mapsto [f_*T]_{\theta,c}$. Moreover, if f is proper, $f_* : H_{\tilde{\theta}}^*(X, \mathbb{C}) \rightarrow H_{\theta}^{*-r}(Y, \mathbb{C})$ is defined well similarly.

Let $f : X \rightarrow Y$ be a proper smooth map between connected oriented smooth manifolds. If μ is a closed 1-forms on Y and $\tilde{\theta} = f^*\theta$, we have the *projection formula*

$$f_*(\sigma \cup f^*\tau) = f_*(\sigma) \cup \tau$$

for $\sigma \in H_{\tilde{\theta}}^*(X, \mathbb{C})$ or $H_{\tilde{\theta},c}^*(X, \mathbb{C})$ and $\tau \in H_{\mu}^*(Y, \mathbb{C})$ or $H_{\mu,c}^*(Y, \mathbb{C})$. We get it easily by $f_*(T \wedge f^*\beta) = f_*T \wedge \beta$, where $T \in \mathcal{D}'^*(X)$ and $\beta \in \mathcal{A}^*(Y)$.

Recall that a complex manifold X is called *p-Kählerian*, if it admits a closed strictly positive (p, p) -form Ω ([1], Definition 1.1, 1.2). For any p -dimensional connected complex submanifold Z of a p -Kähler manifold X , $\Omega|_Z$ is a volume form on Z . We have

Proposition 2.3. *Let $f : X \rightarrow Y$ be a proper surjective holomorphic map between connected complex manifolds, and θ a complex closed 1-form on Y . Set $r = \dim_{\mathbb{C}} X - \dim_{\mathbb{C}} Y$ and $\tilde{\theta} = f^*\theta$. Assume that X is r -Kählerian. Then, for any p , $f^* : H_{\theta}^p(Y, \mathbb{C}) \rightarrow H_{\tilde{\theta}}^p(X, \mathbb{C})$ is injective and $f_* : H_{\tilde{\theta}}^p(X, \mathbb{C}) \rightarrow H_{\theta}^{p-2r}(Y, \mathbb{C})$ is surjective. They also hold for the cases of compact supports.*

Proof. Let Ω be a strictly positive closed (r, r) -form on X . Then $c = f_*\Omega$ is a closed current of degree 0, hence a constant. By Sard's theorem, the set U of regular values of f is

nonempty. For any $y \in U$, $X_y = f^{-1}(y)$ is a r -dimensional compact complex submanifold, so $c = \int_{X_y} \Omega|_{X_y} > 0$ on U . By the projection formula, $f_*([\Omega] \cup f^*\tau) = c \cdot \tau$, where $[\Omega] \in H^{2r}(X, \mathbb{C})$ and $\tau \in H_\theta^p(Y, \mathbb{C})$ or $H_{\theta,c}^p(Y, \mathbb{C})$. It is easily to deduce the conclusion. \square

Clearly, any complex manifold is 0-Kählerian and any Kähler manifold X is p -Kählerian for every $p \leq \dim_{\mathbb{C}} X$, so we get

Corollary 2.4. *Let $f : X \rightarrow Y$ be a proper surjective holomorphic map between connected complex manifolds with the same dimensions. Let θ be a complex closed 1-form on Y and $\tilde{\theta} = f^*\theta$. Then, for any p , $f^* : H_\theta^p(Y, \mathbb{C}) \rightarrow H_{\tilde{\theta}}^p(X, \mathbb{C})$ is injective and $f_* : H_{\tilde{\theta}}^p(X, \mathbb{C}) \rightarrow H_\theta^p(Y, \mathbb{C})$ is surjective. They also hold for the cases of compact supports.*

Corollary 2.5. *Let $f : X \rightarrow Y$ be a proper surjective holomorphic map between connected complex manifolds and θ a complex closed 1-form on Y . Set $r = \dim_{\mathbb{C}} X - \dim_{\mathbb{C}} Y$ and $\tilde{\theta} = f^*\theta$. Assume that X is a Kähler manifold. Then, for any p , $f^* : H_\theta^p(Y, \mathbb{C}) \rightarrow H_{\tilde{\theta}}^p(X, \mathbb{C})$ is injective and $f_* : H_{\tilde{\theta}}^p(X, \mathbb{C}) \rightarrow H_\theta^{p-2r}(Y, \mathbb{C})$ is surjective. They also hold for the cases of compact supports.*

3. DOLBEAULT-MORSE-NOVIKOV COHOMOLOGY

Let X be a n -dimensional complex manifold and η a $\bar{\partial}$ -closed $(0, 1)$ -form on X . Suppose $\mathcal{A}^{p,q}(X)$ is the space of smooth (p, q) -forms on X . Define $\bar{\partial}_\eta : \mathcal{A}^{p,q}(X) \rightarrow \mathcal{A}^{p,q+1}(X)$ as follows:

$$\bar{\partial}_\eta \alpha = \bar{\partial} \alpha + \eta \wedge \alpha,$$

for every $\alpha \in \mathcal{A}^{p,q}(X)$. Clearly, $\bar{\partial}_\eta \circ \bar{\partial}_\eta = 0$, so we have a complex

$$\dots \longrightarrow \mathcal{A}^{p,q-1}(X) \xrightarrow{\bar{\partial}_\eta} \mathcal{A}^{p,q}(X) \xrightarrow{\bar{\partial}_\eta} \mathcal{A}^{p,q+1}(X) \longrightarrow \dots$$

We call its cohomology $H_\eta^{p,q}(X) = H^q(\mathcal{A}^{p,\bullet}(X), \bar{\partial}_\eta)$ *Dolbeault-Morse-Novikov cohomology*. Similarly, we can define *Dolbeault-Morse-Novikov cohomology with compact support* $H_{\eta,c}^{p,q}(X)$. If $\eta = 0$, $H_\eta^{p,q}(X)$ is the classical Dolbeault cohomology $H^{p,q}(X)$. Suppose $\mathcal{A}_X^{p,q}$ is the sheaf of germs of smooth (p, q) -forms on X . We naturally get a morphism $\bar{\partial}_\eta : \mathcal{A}_X^{p,q} \rightarrow \mathcal{A}_X^{p,q+1}$ of sheaves.

Definition 3.1. *We call the kernel of $\bar{\partial}_\eta : \mathcal{A}_X^{p,0} \rightarrow \mathcal{A}_X^{p,1}$ a weight η -sheaf of holomorphic p -forms, denoted by $\Omega_{X,\eta}^p$. In particular, $\mathcal{O}_{X,\eta} := \Omega_{X,\eta}^0$ is called a weight η -sheaf of holomorphic functions.*

Locally, by Grothendieck-Poincaré lemma, $\eta = \bar{\partial}u$ for a smooth complex-valued function u , and then, $\bar{\partial}_\eta = e^{-u} \circ \bar{\partial} \circ e^u$. Hence, locally, $\Omega_{X,\eta}^p = e^{-u} \Omega_X^p$, where Ω_X^p is the sheaf of germs of holomorphic p -forms. So $\mathcal{O}_{X,\eta}$ is a locally free sheaf of \mathcal{O}_X -modules with rank 1 and

$$(2) \quad \Omega_{X,\eta}^p = \Omega_X^p \otimes_{\mathcal{O}_X} \mathcal{O}_{X,\eta}.$$

Moreover, we have a soft resolution of $\Omega_{X,\eta}^p$

$$0 \longrightarrow \Omega_{X,\eta}^p \xrightarrow{i} \mathcal{A}_X^{p,0} \xrightarrow{\bar{\partial}_\eta} \mathcal{A}_X^{p,q} \xrightarrow{\bar{\partial}_\eta} \dots \xrightarrow{\bar{\partial}_\eta} \mathcal{A}_X^{p,n} \longrightarrow 0.$$

Similarly, we can define $\bar{\partial}_\eta$ on the sheaf $\mathcal{D}'_X{}^{p,q}$ of germs of (p, q) -currents and have a soft resolution

$$0 \longrightarrow \Omega_{X,\eta}^p \xrightarrow{i} \mathcal{D}'_X{}^{p,0} \xrightarrow{\bar{\partial}_\eta} \mathcal{D}'_X{}^{p,1} \xrightarrow{\bar{\partial}_\eta} \dots \xrightarrow{\bar{\partial}_\eta} \mathcal{D}'_X{}^{p,n} \longrightarrow 0.$$

So

$$H^q(\mathcal{D}'^{p,\bullet}(X), \bar{\partial}_\eta) \cong H_\eta^{p,q}(X) \cong H^q(X, \Omega_{X,\eta}^p)$$

and

$$H^q(\mathcal{D}'_c{}^{p,\bullet}(X), \bar{\partial}_\eta) \cong H_{\eta,c}^{p,q}(X) \cong H_c^q(X, \Omega_{X,\eta}^p).$$

Similarly with Morse-Novikov cohomology, we can define *pullback* f^* , *pushout* f_* , *cup product* \cup and have *projection formulas* on Dolbeault-Morse-Novikov cohomology. Moreover, by the similar proofs of Proposition 2.3, Corollary 2.4 and 2.5, we have

Proposition 3.2. *Let $f : X \rightarrow Y$ be a proper surjective holomorphic map between complex manifolds and η a $\bar{\partial}$ -closed $(0, 1)$ -forms on Y . Set $r = \dim_{\mathbb{C}} X - \dim_{\mathbb{C}} Y$ and $\tilde{\eta} = f^*\eta$. Assume that X is a r -Kähler manifold. Then, for any p, q , $f^* : H_{\tilde{\eta}}^{p,q}(Y) \rightarrow H_{\tilde{\eta}}^{p,q}(X)$ is injective and $f_* : H_{\tilde{\eta}}^{p,q}(X) \rightarrow H_{\tilde{\eta}}^{p-r,q-r}(Y)$ is surjective. They also hold for the cases of compact supports.*

Corollary 3.3. *Let $f : X \rightarrow Y$ be a proper surjective holomorphic map between complex manifolds with the same dimensions. Let η be a $\bar{\partial}$ -closed $(0, 1)$ -forms on Y and $\tilde{\eta} = f^*\eta$. Then, for any p, q , $f^* : H_{\tilde{\eta}}^{p,q}(Y) \rightarrow H_{\tilde{\eta}}^{p,q}(X)$ is injective and $f_* : H_{\tilde{\eta}}^{p,q}(X) \rightarrow H_{\tilde{\eta}}^{p,q}(Y)$ is surjective. They also hold for the cases of compact supports.*

Corollary 3.4. *Let $f : X \rightarrow Y$ be a proper surjective holomorphic map between complex manifolds and η a $\bar{\partial}$ -closed $(0, 1)$ -forms on Y . Set $r = \dim_{\mathbb{C}} X - \dim_{\mathbb{C}} Y$ and $\tilde{\eta} = f^*\eta$. If X is a Kähler manifold. Then, for any p, q , $f^* : H_{\tilde{\eta}}^{p,q}(Y) \rightarrow H_{\tilde{\eta}}^{p,q}(X)$ is injective and $f_* : H_{\tilde{\eta}}^{p,q}(X) \rightarrow H_{\tilde{\eta}}^{p-r,q-r}(Y)$ is surjective. They also hold for the cases of compact supports.*

Remark 3.5. *On de Rham and Dolbeault cohomologies, several particular cases were proved in [28].*

4. DOLBEAULT-MORSE-NOVIKOV COHOMOLOGY VIA SHEAF THEORY

4.1. weight η -sheaf. First, we give several properties of weight η -sheaves of holomorphic functions.

Lemma 4.1. *Let X be a complex manifold and θ a complex closed 1-form on X . Assume $\theta = \bar{\zeta} + \eta$, where ζ and η are the $(0, 1)$ -forms on X . Then*

$$(1) \mathcal{O}_{X,\eta} = \mathcal{O}_X \otimes_{\mathbb{C}_X} \underline{\mathbb{C}}_{X,\theta};$$

(2) $\mathcal{O}_{X,\eta}$, $\mathcal{O}_{X,\zeta}$ and $\underline{\mathbb{C}}_{X,\theta}$ are subsheaves of $\mathcal{A}_{X,\mathbb{C}}^0$. Moreover, $\mathcal{O}_{X,\eta} \cap \overline{\mathcal{O}_{X,\zeta}} = \underline{\mathbb{C}}_{X,\theta}$, where $\overline{\mathcal{O}_{X,\zeta}}$ is the sheaf of complex conjugation of $\mathcal{O}_{X,\zeta}$ in $\mathcal{A}_{X,\mathbb{C}}^0$.

Proof. Locally, $\theta = du$, $\zeta = \bar{\partial}u$, $\eta = \bar{\partial}u$, hence, $\underline{\mathbb{C}}_{X,\theta} = \mathbb{C}e^{-u}$, $\mathcal{O}_{X,\eta} = e^{-u} \cdot \mathcal{O}_X$ and $\mathcal{O}_{X,\zeta} = e^{-\bar{u}} \cdot \mathcal{O}_X$. Clearly, $\mathcal{O}_{X,\eta} \cap \overline{\mathcal{O}_{X,\zeta}} = \underline{\mathbb{C}}_{X,\theta}$, and the products of functions give an isomorphism $\mathcal{O}_X \otimes_{\mathbb{C}_X} \underline{\mathbb{C}}_{X,\theta} \rightarrow \mathcal{O}_{X,\eta}$. \square

Lemma 4.2. *Let X be a complex manifold and η a $\bar{\partial}$ -closed $(0,1)$ -form on X .*

(1) *Suppose η is $\bar{\partial}$ -exact, i.e., there exists $u \in \mathcal{A}_{\mathbb{C}}^0(X)$, such that $\eta = \bar{\partial}u$. Then*

$$\mathcal{O}_{X,\eta} \rightarrow \mathcal{O}_X, h \mapsto h \cdot e^u$$

is an isomorphism of sheaves of \mathcal{O}_X -modules.

(2) *Suppose ζ is a $\bar{\partial}$ -closed $(0,1)$ -form on X . Then $\mathcal{O}_{X,\zeta} \otimes_{\mathcal{O}_X} \mathcal{O}_{X,\eta} = \mathcal{O}_{X,\zeta+\eta}$. So $(\mathcal{O}_{X,\eta})^\vee = \mathcal{O}_{X,-\eta}$, where $(\mathcal{O}_{X,\eta})^\vee = \text{Hom}_{\mathcal{O}_X}(\mathcal{O}_{X,\eta}, \mathcal{O}_X)$ is the dual of $\mathcal{O}_{X,\eta}$ of \mathcal{O}_X -modules.*

(3) *If $f : Y \rightarrow X$ is a holomorphic map of complex manifolds, then*

$$f^* \mathcal{O}_{X,\eta} = \mathcal{O}_{Y,f^*\eta},$$

where $f^ \mathcal{O}_{X,\eta} = f^{-1} \mathcal{O}_{X,\eta} \otimes_{f^{-1} \mathcal{O}_X} \mathcal{O}_Y$ is the inverse image sheaf of \mathcal{O}_Y -modules.*

Proof. We can get (1) (2) immediately with the similar proof of Lemma 2.2.

(3) For any presheaf \mathcal{G} , denote by \mathcal{G}^+ the sheaf associated to \mathcal{G} . Define presheaves \mathcal{F} and \mathcal{R} on Y as

$$\mathcal{F}(U) = \varinjlim_{W \supseteq f(U)} \mathcal{O}_{X,\eta}(W)$$

and

$$\mathcal{R}(U) = \varinjlim_{W \supseteq f(U)} \mathcal{O}_X(W),$$

for any open subset U of Y . Then $\mathcal{F}^+ = f^{-1} \mathcal{O}_{X,\eta}$, $\mathcal{R}^+ = f^{-1} \mathcal{O}_X$ and $(\mathcal{F} \otimes_{\mathcal{R}} \mathcal{O}_Y)^+ = f^* \mathcal{O}_{X,\eta}$.

Define $\varphi(U) : \mathcal{F}(U) \otimes_{\mathcal{R}(U)} \mathcal{O}_Y(U) \rightarrow \mathcal{O}_{Y,f^*\eta}(U)$ as $[h] \otimes g \mapsto g \cdot (f^*h)|_U$, for every open subset U of Y , where $[h]$ is the class of the η -holomorphic function h under the direct limit. We get a morphism $\varphi : \mathcal{F} \otimes_{\mathcal{R}} \mathcal{O}_Y \rightarrow \mathcal{O}_{Y,f^*\eta}$ of presheaves, and moreover, induce a morphism $\varphi^+ : f^* \mathcal{O}_{X,\eta} \rightarrow \mathcal{O}_{Y,f^*\eta}$ of sheaves.

We claim that φ^+ is an isomorphism. Actually, for any $y \in Y$, choose a open ball V near $f(y)$, such that $\eta = \bar{\partial}u$ on V for some $u \in \mathcal{A}_{\mathbb{C}}^0(V)$. The elements of $\mathcal{F}_y = (\mathcal{O}_{X,\eta})_{f(y)}$ and $(\mathcal{O}_{Y,f^*\eta})_y$ can be written as $[pe^{-u}]$ and $[qe^{-f^*u}]$ respectively, where p, q are holomorphic functions near $f(y), y$ respectively, where $[a]$ denote the the class of a under direct limit. At the stalk over y , $\varphi_y^+([pe^{-u}] \otimes [g]) = [g \cdot f^*p \cdot e^{-f^*u}]$, which is isomorphic. We complete the proof. \square

Remark 4.3. If η is the $(0,1)$ -part of a closed 1-form, Lemma 4.2 (3) can be proved simply by Lemma 4.1 (1).

For a complex closed 1-form θ on a complex manifold X , we write $\theta = \bar{\zeta} + \eta$, where ζ and η are both $(0,1)$ -forms. Let $\partial_{\bar{\zeta}} = \partial + \bar{\zeta} \wedge$. Then $d_{\theta} = \partial_{\bar{\zeta}} + \bar{\partial}_{\eta}$, $\partial_{\bar{\zeta}}^2 = 0$, $\bar{\partial}_{\eta}^2 = 0$, and $\partial_{\bar{\zeta}} \bar{\partial}_{\eta} + \bar{\partial}_{\eta} \partial_{\bar{\zeta}} = 0$. Locally, $\theta = du$, for a smooth complex-valued function u . Then $\eta = \bar{\partial}u$, $\bar{\zeta} = \partial u$ and $\partial_{\bar{\zeta}} = e^{-u} \circ \partial \circ e^u$, locally. By the holomorphic de Rham resolution of \mathbb{C} , there exists a resolution of $\underline{\mathbb{C}}_{X,\theta}$

$$0 \longrightarrow \underline{\mathbb{C}}_{X,\theta} \xrightarrow{i} \mathcal{O}_{X,\eta} \xrightarrow{\partial_{\bar{\zeta}}} \Omega_{X,\eta}^1 \xrightarrow{\partial_{\bar{\zeta}}} \cdots \xrightarrow{\partial_{\bar{\zeta}}} \Omega_{X,\eta}^n \longrightarrow 0.$$

So we can compute Morse-Novikov cohomology by the hypercohomology $H_{\theta}^p(X, \mathbb{C}) = \mathbb{H}^p(X, \Omega_{X,\eta}^{\bullet})$. If X satisfies that $H_{\eta}^{p,q}(X) = 0$ for any $p \geq 1, q \geq 0$, then

$$H_{\theta}^p(X, \mathbb{C}) = H^p(\Gamma(X, \Omega_{X,\eta}^{\bullet}), \partial_{\bar{\zeta}}).$$

In this case, $H_{\bar{\partial}}^p(X, \mathbb{C}) = 0$ for $p > \dim_{\mathbb{C}} X$.

4.2. Künneth formula and Serre's duality. If \mathcal{F} and \mathcal{G} are sheaves of \mathcal{O}_X and \mathcal{O}_Y -modules on complex manifolds X and Y respectively. The *cartesian product sheaf* of \mathcal{F} and \mathcal{G} is defined as

$$\mathcal{F} \boxtimes \mathcal{G} = pr_1^* \mathcal{F} \otimes_{\mathcal{O}_{X \times Y}} pr_2^* \mathcal{G},$$

where pr_1 and pr_2 are projections from $X \times Y$ onto X , Y , respectively. Assume that ζ and η are $\bar{\partial}$ -closed forms on complex manifolds X and Y respectively. By the formula (2) and Lemma 4.2 (3),

$$pr_1^* \Omega_{X, \zeta}^p = pr_1^* \Omega_X^p \otimes_{\mathcal{O}_{X \times Y}} \mathcal{O}_{X \times Y, pr_1^* \zeta}$$

and

$$pr_2^* \Omega_{Y, \eta}^q = pr_2^* \Omega_Y^q \otimes_{\mathcal{O}_{X \times Y}} \mathcal{O}_{X \times Y, pr_2^* \eta},$$

hence $\Omega_{X, \zeta}^p \boxtimes \Omega_{Y, \eta}^q = (\Omega_X^p \boxtimes \Omega_Y^q) \otimes_{\mathcal{O}_{X \times Y}} \mathcal{O}_{X \times Y, \omega}$, where $\omega = pr_1^* \zeta + pr_2^* \eta$. So

$$\begin{aligned} \Omega_{X \times Y, \omega}^k &= \Omega_{X \times Y}^k \otimes_{\mathcal{O}_{X \times Y}} \mathcal{O}_{X \times Y, \omega} \\ (3) \quad &= \left(\bigoplus_{p+q=k} \Omega_X^p \boxtimes \Omega_Y^q \right) \otimes_{\mathcal{O}_{X \times Y}} \mathcal{O}_{X \times Y, \omega} \\ &= \bigoplus_{p+q=k} \Omega_{X, \zeta}^p \boxtimes \Omega_{Y, \eta}^q. \end{aligned}$$

If X or Y is compact, by (3) and [6], Chap. IX, (5.23) (5.24), we have an isomorphism

$$\bigoplus_{p+q=k, r+s=l} H_{\zeta}^{p,r}(X) \otimes_{\mathbb{C}} H_{\eta}^{q,s}(Y) \cong H_{\omega}^{k,l}(X \times Y)$$

for any k, l . We call it *Künneth formula* for Dolbeault-Morse-Novikov cohomology.

Let X be a connected compact complex manifold of dimension n and η a $\bar{\partial}$ -closed $(0, 1)$ -form on X . By Lemma 4.1, (2) and Serre duality theorem,

$$\cup : H_{\eta}^{p,q}(X) \times H_{-\eta}^{n-p, n-q}(X) \rightarrow \mathbb{C}$$

is a nondegenerate pair, for $0 \leq p, q \leq n$.

4.3. Bimeromorphic invariants. We give several bimeromorphic invariants by Dolbeault-Morse-Novikov cohomology.

Proposition 4.4. *Let $f : X \dashrightarrow Y$ be a bimeromorphic map of complex manifolds and η_X, η_Y $\bar{\partial}$ -closed $(0, 1)$ -forms on X, Y respectively. Assume that there exist nowhere dense analytic subsets $E \subseteq X$ and $F \subseteq Y$, such that $f : X - E \rightarrow Y - F$ is biholomorphic and $f^*(\eta_Y|_{Y-F}) = \eta_X|_{X-E}$. Then, for any p ,*

- (1) $H_{\eta_X}^{0,p}(X) \cong H_{\eta_Y}^{0,p}(Y)$ and $H_{\eta_X, c}^{0,p}(X) \cong H_{\eta_Y, c}^{0,p}(Y)$;
- (2) $H_{\eta_X}^{p,0}(X) \cong H_{\eta_Y}^{p,0}(Y)$ and $H_{\eta_X, c}^{p,0}(X) \cong H_{\eta_Y, c}^{p,0}(Y)$.

Proof. We choose two proper modifications $g : Z \rightarrow X$ and $h : Z \rightarrow Y$ such that there is nowhere dense analytic subset S in Z , $E \subseteq g(S)$ and $F \subseteq h(S)$, $g : Z - S \rightarrow X - g(S)$, $h : Z - S \rightarrow Y - h(S)$ are biholomorphic and $fg|_{Z-S} = h|_{Z-S}$. Obviously,

$$(g^* \eta_X - h^* \eta_Y)|_{Z-S} = g^*((\eta_X|_{X-E} - f^*(\eta_Y|_{Y-F}))|_{X-g(S)}) = 0.$$

By the continuity, $g^*\eta_X = h^*\eta_Y$. Hence, we need only to prove the proposition for the case that f is a proper modification and $f^*\eta_Y = \eta_X$. By [9], page 215, we assume $E = f^{-1}(F)$, $\text{codim}_Y F \geq 2$ and $\text{codim}_X E = 1$.

(1) By Lemma 4.2 (3) and [25], Proposition 1.13, 2.14,

$$R^q f_* \mathcal{O}_{X, \eta_X} = R^q f_* \mathcal{O}_X \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y, \eta_Y} = \begin{cases} \mathcal{O}_{Y, \eta_Y}, & q = 0; \\ 0, & \text{otherwise.} \end{cases}$$

Consider Leray spectral sequences,

$$E_2^{p,q} = H^p(Y, R^q f_* \mathcal{O}_{X, \eta_X}) \Rightarrow H^{p+q} = H^{p+q}(X, \mathcal{O}_{X, \eta_X})$$

and

$$E_2^{p,q} = H_c^p(Y, R^q f_* \mathcal{O}_{X, \eta_X}) \Rightarrow H^{p+q} = H_c^{p+q}(X, \mathcal{O}_{X, \eta_X}).$$

Then $E_2^{p,q} = 0$ for $q > 0$. Hence $E_2^{p,0} = H^p$. We get (1).

(2) Set $U = X - E$, $V = Y - F$ and $j_U : U \rightarrow X$, $j_V : V \rightarrow Y$ are inclusions. We have a commutative diagram

$$\begin{array}{ccc} H^0(Y, \Omega_{Y, \eta_Y}^p) & \xrightarrow{f^*} & H^0(X, \Omega_{X, \eta_X}^p) , \\ \downarrow j_V^* & & \downarrow j_U^* \\ H^0(V, \Omega_{Y, \eta_Y}^p) & \xrightarrow{(f|_U)^*} & H^0(U, \Omega_{X, \eta_X}^p) \end{array}$$

By the continuity, the restriction j_U^* is injective. By the second Riemann continuation theorem ([8], p. 133), j_V^* is isomorphic. Since $f|_U$ is biholomorphic, j_U^* is surjective, and then, an isomorphism. So f^* is an isomorphism.

Consider the commutative diagram

$$\begin{array}{ccc} H_c^0(X, \Omega_{X, \eta_X}^p) & \xrightarrow{f_*} & H_c^0(Y, \Omega_{Y, \eta_Y}^p) . \\ \downarrow & & \downarrow \\ H^0(X, \Omega_{X, \eta_X}^p) & \xrightarrow{f_*} & H^0(Y, \Omega_{Y, \eta_Y}^p) \end{array}$$

The two vertical maps are inclusions, hence are both injective. We have proven that $f^* : H_{\eta_Y}^{p,0}(Y) \rightarrow H_{\eta_X}^{p,0}(X)$ is an isomorphism. By the projection formula, $f_* f^* = \text{id}$ on $H_{\eta_Y}^{p,0}(Y)$. So the map at the bottom is an isomorphism. Then the map at the top is injective. By the projection formula again, $f_* f^* = \text{id}$ on $H_{\eta_Y, c}^{p,0}(Y)$, hence f_* is isomorphic on $H_{\eta_X, c}^{p,0}(X)$. \square

Remark 4.5. $H_\theta^1(X, \mathbb{C})$ and $H_{\theta, c}^{2n-1}(X, \mathbb{C})$ are also bimeromorphic invariants, referring to [14], Corollary 4.8.

4.4. Leray-Hirsch theorem. Now we establish the Leray-Hirsch theorem for the Dolbeault-Morse-Novikov cohomology.

Theorem 4.6. *Let $\pi : E \rightarrow X$ be a holomorphic fiber bundle over a connected complex manifold X whose general fiber F is compact and η a $\bar{\partial}$ -closed $(0, 1)$ -form on X . Assume there*

exist classes e_1, \dots, e_r of pure degrees in $H^{**}(E)$, such that, for every $x \in X$, their restrictions $e_1|_{E_x}, \dots, e_r|_{E_x}$ freely linearly generate $H^{**}(E_x)$. Then, $\pi^*(\bullet) \cup \bullet$ gives isomorphisms of bigraded vector spaces

$$H_{\tilde{\eta}}^{**}(X) \otimes_{\mathbb{C}} \text{span}_{\mathbb{C}}\{e_1, \dots, e_r\} \xrightarrow{\sim} H_{\tilde{\eta}}^{**}(E),$$

where $\tilde{\eta} = \pi^*\eta$.

Proof. If X is a Stein manifold, the theorem holds. Actually, since $H^{0,1}(X) = 0$, η is $\bar{\partial}$ -exact. By (2) and Lemma 4.2 (1), we may assume $\eta = 0$. It is exactly [15], Theorem 1.2.

Go back to the general case. Let t_1, \dots, t_r be forms of pure degrees in $\mathcal{A}^{**}(E)$, such that $e_i = [t_i]$ for $1 \leq i \leq r$. Set $L^{*,*} = \text{span}_{\mathbb{C}}\{t_1, \dots, t_r\}$, which is a bigraded vector spaces and isomorphic to $\text{span}_{\mathbb{C}}\{e_1, \dots, e_r\}$. For any open set U in X , set

$$B^{p,q}(U) = \bigoplus_{k+l=p, u+v=q} \mathcal{A}^{k,u}(U) \otimes_{\mathbb{C}} L^{l,v}$$

and $\bar{\partial}_B = \bar{\partial}_{\tilde{\eta}} \otimes 1$. For any p , $(B^{p,\bullet}(U), \bar{\partial}_B)$ is a complex, whose cohomology is

$$\begin{aligned} D^{p,q}(U) &= (H_{\tilde{\eta}}^{*,*}(U) \otimes_{\mathbb{C}} \text{span}_{\mathbb{C}}\{e_1, \dots, e_r\})^{p,q} \\ &= \bigoplus_{k+l=p, u+v=q} H_{\tilde{\eta}}^{k,u}(U) \otimes_{\mathbb{C}} (\text{span}_{\mathbb{C}}\{e_1, \dots, e_r\})^{l,v}. \end{aligned}$$

Clearly, the morphism $\pi^*(\bullet) \wedge \bullet : B^{p,\bullet}(U) \rightarrow C^{p,\bullet}(U) := \mathcal{A}^{p,\bullet}(E_U)$ of complexes induces a morphism on the cohomological level

$$\pi^*(\bullet) \cup \bullet : D^{p,q}(U) \rightarrow E^{p,q}(U) := H_{\tilde{\eta}}^{p,q}(E_U),$$

denoted by Φ_U . We need to prove Φ_X is an isomorphism.

Given p , for any open subsets U, V in X , there is a commutative diagram of complexes

$$\begin{array}{ccccccc} 0 & \longrightarrow & B^{p,\bullet}(U \cup V) & \xrightarrow{(\rho_U^{UV}, \rho_V^{UV})} & B^{p,\bullet}(U) \oplus B^{p,\bullet}(V) & \xrightarrow{\rho_{U \cap V}^U - \rho_{U \cap V}^V} & B^{p,\bullet}(U \cap V) \longrightarrow 0, \\ & & \downarrow \pi^*(\bullet) \wedge \bullet & & \downarrow (\pi^*(\bullet) \wedge \bullet, \pi^*(\bullet) \wedge \bullet) & & \downarrow \pi^*(\bullet) \wedge \bullet \\ 0 & \longrightarrow & C^{p,\bullet}(U \cup V) & \xrightarrow{(j_U^{UV}, j_V^{UV})} & C^{p,\bullet}(U) \oplus C^{p,\bullet}(V) & \xrightarrow{j_{U \cap V}^U - j_{U \cap V}^V} & C^{p,\bullet}(U \cap V) \longrightarrow 0 \end{array}$$

where ρ, j are restrictions and the differentials of complexes in the first, second rows are all $\bar{\partial}_B, \bar{\partial}$, respectively. The two rows are both exact sequences of complexes. Therefore, we have a commutative diagram of long exact sequences

$$\begin{array}{ccccccc} \dots & \longrightarrow & D^{p,q-1}(U \cap V) & \longrightarrow & D^{p,q}(U \cup V) & \longrightarrow & D^{p,q}(U) \oplus D^{p,q}(V) & \longrightarrow & D^{p,q}(U \cap V) & \longrightarrow & \dots \\ & & \downarrow \Phi_{U \cap V} & & \downarrow \Phi_{U \cup V} & & \downarrow (\Phi_U, \Phi_V) & & \downarrow \Phi_{U \cap V} & & \\ \dots & \longrightarrow & E^{p,q-1}(U \cap V) & \longrightarrow & E^{p,q}(U \cup V) & \longrightarrow & E^{p,q}(U) \oplus E^{p,q}(V) & \longrightarrow & E^{p,q}(U \cap V) & \longrightarrow & \dots \end{array}$$

If Φ_U, Φ_V and $\Phi_{U \cap V}$ are isomorphisms, then $\Phi_{U \cup V}$ is an isomorphism by Five Lemma (seeing [11], p. 6). We claim that:

(*) For open subsets $U_1, \dots, U_s \subseteq X$, if $\Phi_{U_{i_1} \cap \dots \cap U_{i_k}}$ is an isomorphism for any $1 \leq k \leq s$ and $1 \leq i_1 < \dots < i_k \leq s$, then $\Phi_{\bigcup_{i=1}^s U_i}$ is an isomorphism.

We prove this conclusion by induction. For $r = 1$, the conclusion holds clearly. Suppose it holds for s . For $s + 1$, set $U'_1 = U_1, \dots, U'_{s-1} = U_{s-1}, U'_s = U_s \cup U_{s+1}$. Then

$\Phi_{U'_{i_1} \cap \dots \cap U'_{i_k}} = \Phi_{U_{i_1} \cap \dots \cap U_{i_k}}$ is isomorphic for any $1 \leq i_1 < \dots < i_k \leq s-1$. Moreover, $\Phi_{U'_{i_1} \cap \dots \cap U'_{i_{k-1}} \cap U'_s}$ is also isomorphic for any $1 \leq i_1 < \dots < i_{k-1} \leq s-1$, since $\Phi_{U_{i_1} \cap \dots \cap U_{i_{k-1}} \cap U_s}$, $\Phi_{U_{i_1} \cap \dots \cap U_{i_{k-1}} \cap U_{s+1}}$ and $\Phi_{U_{i_1} \cap \dots \cap U_{i_{k-1}} \cap U_s \cap U_{s+1}}$ are isomorphic. By inductive hypothesis, $\Phi_{\bigcup_{i=1}^{s+1} U_i} = \Phi_{\bigcup_{i=1}^s U'_i}$ is an isomorphism. We proved (*).

For a disjoint union $U = \bigcup U_\alpha$ of open subsets U_α in X , Φ_U is exactly the direct product

$$\prod \Phi_{U_\alpha} : \prod D^{p,q}(U_\alpha) \rightarrow \prod H_{\tilde{\eta}}^{p,q}(E_{U_\alpha}).$$

If Φ_{U_α} are all isomorphic, then Φ_U is also an isomorphism.

Let \mathcal{U} be a basis for topology of X such that every $U \in \mathcal{U}$ is Stein and let \mathcal{U}_f be the collection of the finite unions of open sets in \mathcal{U} .

For any finite intersection V of open sets in \mathcal{U}_f , Φ_V is an isomorphism. Actually, $V = \bigcap_{i=1}^s U_i$, where $U_i = \bigcup_{j=1}^{r_i} U_{ij}$ and $U_{ij} \in \mathcal{U}$. Then $V = \bigcup_{J \in \Lambda} U_J$, where $\Lambda = \{J = (j_1, \dots, j_s) | 1 \leq j_1 \leq r_1, \dots, 1 \leq j_s \leq r_s\}$ and $U_J = U_{1j_1} \cap \dots \cap U_{sj_s}$. For any $J_1, \dots, J_t \in \Lambda$, $U_{J_1} \cap \dots \cap U_{J_t}$ is a Stein manifold, so $\Phi_{U_{J_1} \cap \dots \cap U_{J_t}}$ is isomorphic. By (*), $\Phi_V = \Phi_{\bigcup_{J \in \Lambda} U_J}$ is an isomorphism.

By [7], p. 16, Prop. II, $X = V_1 \cup \dots \cup V_l$, where V_i is a countable disjoint union of open sets in \mathcal{U}_f . For any $1 \leq i_1 < \dots < i_k \leq l$, $V_{i_1} \cap \dots \cap V_{i_k}$ is a disjoint union of the finite intersection of open sets in \mathcal{U}_f . Hence, $\Phi_{V_{i_1} \cap \dots \cap V_{i_k}}$ is isomorphic, so is Φ_X by (*). We complete the proof. \square

In particular, we can calculate the Dolbeault-Morse-Novikov cohomology of projectivized bundles.

Corollary 4.7. *Let $\pi : \mathbb{P}(E) \rightarrow X$ be the projectivization of a holomorphic vector bundle E on a connected complex manifold X . Assume η is a $\bar{\partial}$ -closed $(0,1)$ -form on X and $h = [\frac{i}{2\pi}\Theta(\mathcal{O}_{\mathbb{P}(E)}(-1))]$ is in $H^{1,1}(\mathbb{P}(E))$, where $\mathcal{O}_{\mathbb{P}(E)}(-1)$ is the universal line bundle on $\mathbb{P}(E)$ and $\Theta(\mathcal{O}_{\mathbb{P}(E)}(-1))$ is the Chern curvature of a hermitian metric on $\mathcal{O}_{\mathbb{P}(E)}(-1)$. Then $\pi^*(\bullet) \cup \bullet$ gives an isomorphism of graded vector spaces*

$$H_{\tilde{\eta}}^{*,*}(X) \otimes_{\mathbb{C}} \text{span}_{\mathbb{C}}\{1, \dots, h^{r-1}\} \xrightarrow{\sim} H_{\tilde{\eta}}^{*,*}(\mathbb{P}(E)),$$

where $\text{rank}_{\mathbb{C}} E = r$ and $\tilde{\eta} = \pi^* \eta$.

4.5. A blow-up formula. We have the following lemma by definition.

Lemma 4.8 ([15], Proposition 3.1). *Let X be a complex manifold and Z, U closed, open complex submanifolds of X , respectively. Assume $i : Z \rightarrow X$, $j : U \rightarrow X$, $i' : Z \cap U \rightarrow Z$ and $j' : Z \cap U \rightarrow U$ are inclusions. Then $i'_* j'^* = j^* i_*$ on $\mathcal{D}^{**}(Z)$.*

Let $\pi : \tilde{X} \rightarrow X$ be the blow-up of a connected complex manifold X along a connected complex submanifold Z . We know $\pi|_E : E = \pi^{-1}(Z) \rightarrow Z$ is the projectivization $E = \mathbb{P}(N_{Z/X})$ of the normal bundle $N_{Z/X}$. Set

$$(4) \quad h = [\frac{i}{2\pi}\Theta(\mathcal{O}_E(-1))]$$

in $H_{\bar{\partial}}^{1,1}(E)$, where $\Theta(\mathcal{O}_E(-1))$ is the curvature of the Chern connection of a hermitian metric of the universal line bundle $\mathcal{O}_E(-1)$ on E .

Theorem 4.9. *With above notations, let $i_E : E \rightarrow \tilde{X}$ be the inclusion and $r = \text{codim}_{\mathbb{C}} Z$. Suppose that η is a $\bar{\partial}$ -closed $(0, 1)$ -form on X and $\tilde{\eta} = \pi^*\eta$. Then, for any p, q ,*

$$(5) \quad \pi^* + \sum_{i=0}^{r-2} (i_E)_* \circ (h^i \cup) \circ (\pi|_E)^*$$

gives an isomorphism

$$H_{\tilde{\eta}}^{p,q}(X) \oplus \bigoplus_{i=0}^{r-2} H_{\eta|_Z}^{p-1-i, q-1-i}(Z) \xrightarrow{\sim} H_{\tilde{\eta}}^{p,q}(\tilde{X}).$$

Proof. For a Stein manifold X , we may assume $\eta = 0$ with the same reason with the proof of Theorem 4.6, so the theorem holds by [15], Theorem 1.3.

For the general complex manifold X , set

$$\mathcal{F}^{p,q} = \mathcal{A}_X^{p,q} \oplus \bigoplus_{i=0}^{r-2} i_{Z*} \mathcal{A}_Z^{p-1-i, q-1-i},$$

for any p, q . Define $\bar{\partial} : \mathcal{F}^{p,*} \rightarrow \mathcal{F}^{p,*+1}$ as $(\alpha, \beta_0, \dots, \beta_{r-2}) \mapsto (\bar{\partial}_{\eta}\alpha, \bar{\partial}_{\eta|_Z}\beta_0, \dots, \bar{\partial}_{\eta|_Z}\beta_{r-2})$. For any p , $(\mathcal{F}^{p,\bullet}, \bar{\partial})$ is a complex of sheaves. Let $t = \frac{i}{2\pi}\Theta(\mathcal{O}_E(-1)) \in \mathcal{A}^{1,1}(E)$. For any open subset U in X , define $\mathcal{F}^{p,q}(U) \rightarrow \mathcal{D}^{p,q}(\tilde{U})$ as

$$\varphi_U = \begin{cases} (\pi|_{\tilde{U}})^* + \sum_{i=0}^{r-2} (i_{E \cap \tilde{U}})_* \circ (t^i|_{E \cap \tilde{U}} \wedge) \circ (\pi|_{E \cap \tilde{U}})^*, & Z \cap U \neq \emptyset \\ (\pi|_{\tilde{U}})^*, & Z \cap U = \emptyset, \end{cases}$$

where $\tilde{U} = \pi^{-1}(U)$ and $i_{E \cap \tilde{U}} : E \cap \tilde{U} \rightarrow \tilde{U}$ is the inclusion. Clearly, $\bar{\partial}_{\tilde{\eta}} \circ \varphi_U = \varphi_U \circ \bar{\partial}$. Hence, φ_U induces a morphism of vector spaces

$$\Phi_U : H_{\tilde{\eta}}^{p,q}(U) \oplus \bigoplus_{i=0}^{r-2} H_{\eta|_Z}^{p-1-i, q-1-i}(Z \cap U) \rightarrow H_{\tilde{\eta}}^{p,q}(\tilde{U}).$$

We need to prove that Φ_X is an isomorphism.

For open sets $V \subseteq U$, denote by $\rho_V^U : \mathcal{F}^{p,q}(U) \rightarrow \mathcal{F}^{p,q}(V)$ the restriction of the sheaf $\mathcal{F}^{p,q}$ and $j_V^U : \mathcal{D}^{p,q}(\tilde{U}) \rightarrow \mathcal{D}^{p,q}(\tilde{V})$ the restriction of currents. By Lemma 4.8, $j_V^U \circ \varphi_U = \varphi_V \circ \rho_V^U$. Given p , for any open subsets U, V in X , there is a commutative diagram of complexes

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F}^{p,\bullet}(U \cup V) & \xrightarrow{(\rho_U^{UV}, \rho_V^{UV})} & \mathcal{F}^{p,\bullet}(U) \oplus \mathcal{F}^{p,\bullet}(V) & \xrightarrow{\rho_{U \cap V}^U - \rho_{U \cap V}^V} & \mathcal{F}^{p,\bullet}(U \cap V) \longrightarrow 0 \\ & & \downarrow \varphi_{U \cup V} & & \downarrow (\varphi_U, \varphi_V) & & \downarrow \varphi_{U \cap V} \\ 0 & \longrightarrow & \mathcal{D}^{p,\bullet}(\tilde{U} \cup \tilde{V}) & \xrightarrow{(j_U^{UV}, j_V^{UV})} & \mathcal{D}^{p,\bullet}(\tilde{U}) \oplus \mathcal{D}^{p,\bullet}(\tilde{V}) & \xrightarrow{j_{U \cap V}^U - j_{U \cap V}^V} & \mathcal{D}^{p,\bullet}(\tilde{U} \cap \tilde{V}) \longrightarrow 0 \end{array}$$

The two rows are both exact sequences of complexes. For convenience, denote

$$L^{p,q}(U) = H_{\tilde{\eta}}^{p,q}(U) \oplus \bigoplus_{i=0}^{r-2} H_{\eta|_Z}^{p-1-i, q-1-i}(Z \cap U).$$

Therefore, we have a commutative diagram of long exact sequences

$$\begin{array}{ccccccccc}
\cdots & \longrightarrow & L^{p,q-1}(U \cap V) & \longrightarrow & L^{p,q}(U \cup V) & \longrightarrow & L^{p,q}(U) \oplus L^{p,q}(V) & \longrightarrow & L^{p,q}(U \cap V) & \longrightarrow & L^{p,q+1}(U \cup V) & \longrightarrow & \cdots \\
& & \downarrow \Phi_{U \cap V} & & \downarrow \Phi_{U \cup V} & & \downarrow (\Phi_U, \Phi_V) & & \downarrow \Phi_{U \cap V} & & \downarrow \Phi_{U \cup V} & & \\
\cdots & \longrightarrow & H_{\bar{\eta}}^{p,q-1}(\bar{U} \cap \bar{V}) & \longrightarrow & H_{\bar{\eta}}^{p,q}(\bar{U} \cup \bar{V}) & \longrightarrow & H_{\bar{\eta}}^{p,q}(\bar{U}) \oplus H_{\bar{\eta}}^{p,q}(\bar{V}) & \longrightarrow & H_{\bar{\eta}}^{p,q}(\bar{U} \cap \bar{V}) & \longrightarrow & H_{\bar{\eta}}^{p,q+1}(\bar{U} \cup \bar{V}) & \longrightarrow & \cdots
\end{array}$$

Following the steps in the proof of Theorem 4.6, we proved that Φ_X is an isomorphism. \square

5. STABILITY OF θ -BETTI AND η -HODGE NUMBERS

For a compact smooth manifold X and a real (resp. complex) closed 1-form θ on X , $b_k(X, \theta) := \dim_{\mathbb{R}} H_{\theta}^k(X)$ (resp. $\dim_{\mathbb{C}} H_{\theta}^k(X, \mathbb{C})$) is called k -th θ -betti number of X . Similarly, for a compact complex manifold X and a $\bar{\partial}$ -closed $(0, 1)$ -form η on X , $h_{\eta}^{p,q}(X) := \dim_{\mathbb{C}} H_{\eta}^{p,q}(X)$ is called (p, q) -th η -hodge number of X .

Lemma 5.1. *Let $f : X \rightarrow Y$ be a proper surjective submersion of connected smooth manifolds and θ a real (resp. complex) closed 1-form on X . Then, for any k , the higher direct image $R^k f_* \mathbb{R}_{X,\theta}$ (resp. $R^k f_* \mathbb{C}_{X,\theta}$) is a local system of \mathbb{R} (resp. \mathbb{C})-modules with finite rank.*

In particular,

$$y \mapsto b_k(X_y, \theta|_{X_y})$$

is a constant function, where $X_y = f^{-1}(y)$ for any $y \in Y$.

Proof. We may assume Y is an open ball and only prove the real case.

Let o be the center of Y . By Ehresmann's trivialization theorem, there exists a diffeomorphism $T : X_o \times Y \rightarrow X$, such that $pr_2 = f \circ T$, where pr_2 is the projection from $X_o \times Y$ to Y . By Lemma 2.2 (3),

$$\begin{aligned}
(6) \quad R^k f_* \mathbb{R}_{X,\theta} &\cong R^k f_*(T_* \mathbb{R}_{X_o \times Y, T^* \theta}) \\
&\cong R^k (pr_2)_* \mathbb{R}_{X_o \times Y, T^* \theta}.
\end{aligned}$$

Set pr_2 the projection from $X_o \times Y$ to X_o . By Künneth formula, $pr_1^* : H^1(X_o) \rightarrow H^1(X_o \times Y)$ is an isomorphism, where we use the fact that $H^0(Y) = \mathbb{R}$ and $H^1(Y) = 0$. So, $T^* \theta$ can be written as $pr_1^* \theta_o + du$ for a closed 1-form θ_o on X_o and a smooth function u on $X_o \times Y$. Consider the cartesian diagram

$$\begin{array}{ccc}
X_o \times Y & \xrightarrow{pr_2} & Y \\
\downarrow pr_1 & & \downarrow p_Y \\
X_o & \xrightarrow{p_{X_o}} & \{pt\},
\end{array}$$

where $\{pt\}$ is a single point space and p_{X_o}, p_Y are constant map. Evidently, pr_2 and p_{X_o} are proper. By Lemma 2.2 and [11], p. 316, Corollary 1.5,

$$\begin{aligned}
(7) \quad R^k (pr_2)_* \mathbb{R}_{X_o \times Y, T^* \theta} &\cong R^k (pr_2)_* \mathbb{R}_{X_o \times Y, pr_1^* \theta_o} \\
&\cong R^k (pr_2)_* (pr_1^{-1} \mathbb{R}_{X_o, \theta_o}) \\
&\cong p_Y^{-1} R^k (p_{X_o})_* (\mathbb{R}_{X_o, \theta_o}) \\
&= \mathbb{R}_{X_o \times Y} \otimes_{\mathbb{R}} H_{\theta_o}^k(X_o).
\end{aligned}$$

Combined (6) and (7), $R^k f_* \mathbb{R}_{X,\theta}$ is constant on the open ball Y . Moreover, the stalk $(R^k f_* \mathbb{R}_{X,\theta})_y = H^k(X_y, \mathbb{R}_{X,\theta}|_{X_y}) = H_{\theta|_{X_y}}^k(X_y)$. We complete the proof. \square

Let X be a compact complex manifold and $\theta = \bar{\zeta} + \eta$ a complex closed 1-form on X , where ζ and η are both $(0,1)$ -forms. For the double complex $(\mathcal{A}^{*,*}(X), \partial_{\bar{\zeta}}, \bar{\partial}_{\eta})$, the associated simple complex is $(\mathcal{A}_{\mathbb{C}}^*(X), d_{\theta})$, which has a natural filtration

$$F^p \mathcal{A}_{\mathbb{C}}^k(X) = \bigoplus_{r \geq p, r+s=k} \mathcal{A}^{r,s}(X).$$

We get a spectral sequence $(E_r^{*,*}, d_r, H^*)$, where $E_1^{p,q} = H_{\eta}^{p,q}(X)$ and $H^k = H_{\theta}^k(X, \mathbb{C})$. If $\theta = 0$, this is *Frölicher spectral sequence*. Clearly, for $p < 0$, or $p > n$, or $q < 0$, or $q > n$, $E_r^{p,q} = 0$. So, for given p, q , if r is enough large,

$$E_r^{p,q} = E_{r+1}^{p,q} = \dots = E_{\infty}^{p,q} = F^p H_{\theta}^{p+q}(X, \mathbb{C}) / F^{p+1} H_{\theta}^{p+q}(X, \mathbb{C}).$$

Since $\dim_{\mathbb{C}} E_{r+1}^{p,q} \leq \dim_{\mathbb{C}} E_r^{p,q}$ for any r ,

$$b_k(X, \theta) = \sum_{p+q=k} E_{\infty}^{p,q} \leq \sum_{p+q=k} E_1^{p,q} = \sum_{p+q=k} h_{\eta}^{p,q}(X).$$

The degeneration of this spectral sequence at E_1 on compact locally conformally Kähler manifold is proved in some conditions in [20].

We say that $f : X \rightarrow Y$ is a *family of complex manifolds*, if f is a proper surjective holomorphic submersion.

Theorem 5.2. *Let $f : X \rightarrow Y$ be a family of complex manifolds and θ a complex closed 1-form on X . Assume $b_k(X_o, \theta|_{X_o}) = \sum_{p+q=k} h_{\eta|_{X_o}}^{p,q}(X_o)$ for some k and some point $o \in Y$, where η is the $(0,1)$ -part of θ . Then, for any t near o , $h_{\eta|_{X_t}}^{p,q}(X_t) = h_{\eta|_{X_o}}^{p,q}(X_o)$, where η is the $(0,1)$ -part of θ and $p+q=k$.*

Proof. Let $\Omega_{X/Y}^1 = \Omega_X^1 / f^* \Omega_Y^1$ be the sheaf of the relative holomorphic 1-forms and $\Omega_{X/Y}^p = \bigwedge^p \Omega_{X/Y}^1$. Set $i_t : X_t \rightarrow X$ the inclusion. Then $i_t^* \Omega_{X/Y}^p = \Omega_{X_t}^p$, seeing [27], p. 234-235. For the locally free sheaf $\Omega_{X/Y}^p \otimes_{\mathcal{O}_X} \mathcal{O}_{X,\eta}$, we have

$$i_t^*(\Omega_{X/Y}^p \otimes_{\mathcal{O}_X} \mathcal{O}_{X,\eta}) = i_t^* \Omega_{X/Y}^p \otimes_{\mathcal{O}_{X_t}} i_t^* \mathcal{O}_{X,\eta} = \Omega_{X_t, \eta|_{X_t}}^p.$$

By the semi-continuity theorem, $h_{\eta|_{X_t}}^{p,q}(X_t) \leq h_{\eta|_{X_o}}^{p,q}(X_o)$ for any t near o . So

$$b_k(X_o, \theta|_{X_o}) = \sum_{p+q=k} h_{\eta|_{X_o}}^{p,q}(X_o) \geq \sum_{p+q=k} h_{\eta|_{X_t}}^{p,q}(X_t) \geq b_k(X_t, \eta|_{X_t}).$$

By Lemma 5.1, $h_{\eta|_{X_t}}^{p,q}(X_t) = h_{\eta|_{X_o}}^{p,q}(X_o)$ for any $p+q=k$. \square

By Hodge decomposition of complex manifolds in Fujiki class \mathcal{C} , we get the following corollary immediately.

Corollary 5.3. *Let $f : X \rightarrow Y$ be a family of complex manifolds and θ a complex closed 1-form on X . Assume, for a point $o \in Y$, X_o is in the Fujiki class \mathcal{C} and $\theta|_{X_o} = 0$. Then, for any t near o , $h_{\eta|_{X_t}}^{p,q}(X_t) = h_{\eta|_{X_o}}^{p,q}(X_o)$, for any p, q , where η is the $(0,1)$ -part of θ .*

Acknowledgements. I would like to express my gratitude to referees for their helpful suggestions and careful reading of my manuscript.

References

1. Alessandrini, L., Bassanelli, G.: Compact p-Kähler manifolds, *Geometriae Dedicata* **38**, 199-210 (1991)
2. Banyaga, A.: Some properties of locally conformal symplectic structures. *Comment. Math. Helv.* **77**(2), 383-398 (2002)
3. Banyaga, A.: Examples of non $d\omega$ -exact locally conformal symplectic forms. *J. Geom.* **87**, 1-13 (2007)
4. Bande, G., Kotschick, D.: Moser stability for locally conformally symplectic structures. *Proc. Amer. Math. Soc.* **137**(7), 2419-2424 (2009)
5. Belgun, F. A.: On the metric structure of non-Kähler complex surfaces. *Math. Ann.* **317**, 1-40 (2000)
6. Demailly, J.-P.: *Complex Analytic and Differential Geometry*, <http://www-fourier.ujf-grenoble.fr/~demailly/documents.html>.
7. Greub, W., Halperin, S., Vanstone, R.: *Connections, curvature, and cohomology, Vol. I*. Academic Press, New York, 1972
8. Grauert, H., Remmert, R.: *Theory of Stein Spaces*. Classics in Math., Springer-Verlag, Berlin, 1979
9. Grauert, H., Remmert, R.: *Coherent analytic sheaves*. Grundlehren der Math. Wiss. **265**, Springer-Verlag, Berlin, 1984
10. Haller, S., Rybicki, T.: On the group of diffeomorphisms preserving a locally conformal symplectic structure. *Ann. Glob. Anal. Geom.* **17**, 475-502 (1999)
11. Iversen, B.: *Cohomology of Sheaves*. Universitext, Springer-Verlag, Berlin, 1986
12. León, M., López, B., Marrero, J. C., Padrón, E.: On the computation of the Lichnerowicz-Jacobi cohomology. *J. Geom. Phys.* **44**, 507-522 (2003)
13. Lichnerowicz, A.: Les variétés de Poisson et leurs algèbres de Lie associées. *J. Differ. Geom.* **12**(2), 253-300 (1977)
14. Meng, L.: Morse-Novikov cohomology for blow-ups of complex manifolds. arXiv:1806.06622v3 (22 October, 2018)
15. Meng, L.: An explicit formula of blow-ups for Dolbeault cohomology. arXiv:1806.11435v4 (12 October, 2018)
16. Novikov, S. P.: The Hamiltonian formalism and a multivalued analogue of Morse theory (Russian). *Uspekhi Mat. Nauk.* **37**, 3-43 (1982); English translation, *Russian Math. Surveys* **37** (5), 1-56 (1982).
17. Ornea, L., Verbitsky, M.: Morse-Novikov cohomology of locally conformally Kähler manifolds. *J. Geom. Phys.* **59**, 295-305 (2009)
18. Ornea, L., Verbitsky, M.: Locally conformal Kähler manifolds with potential. *Math. Ann.* **348**, 25-33 (2010)
19. Ornea, L., Verbitsky, M., Vuletescu, V.: Blow-ups of locally conformally Kähler manifolds. *Int. Math. Res. Not.* (12), 2809-2821 (2013)
20. Ornea, L., Verbitsky, M., Vuletescu, V.: Weighted Bott-Chern and Dolbeault cohomology for LCK-manifolds with potential. *J. Math. Soc. Japan* **70** (1), 409-422 (2018)
21. Pajitnov, A.: An analytic proof of the real part of the Novikov inequalities (in Russian). *Dokl. Akad. Nauk SSSR* **293** (6), 1305-1307 (1987); English translation, *Soviet Math. Dokl.* **35** (2), 456-457 (1987)
22. Pajitnov, A.: *Circle-valued Morse Theory*. De Gruyter Studies in Math. **32**, Walter de Gruyter, Berlin, 2006
23. Rao, S., Yang, S., Yang, X.-D.: Dolbeault cohomologies of blowing up complex manifolds. arXiv:1712.06749v4 (23 August, 2018) to appear in *J. Math. Pures Appl.*
24. Sullivan, D.: Infinitesimal computations in topology. *Publ. Math. Inst. Hautes Études Sci.* **47**, 269-331 (1977)
25. Ueno, K.: *Classification Theory of Algebraic Varieties and Compact Complex Spaces*. Lecture Notes in Math. **439**, Berlin-Heidelberg-New York, Springer, 1975
26. Vaisman, I.: Remarkable operators and commutation formulas on locally conformal Kähler manifolds. *Compos. Math.* **40**(3), 287-299 (1980)
27. Voisin, C.: *Hodge Theory and Complex Algebraic Geometry. Vol. I*. Camb. Stud. Adv. Math. **76**, Cambridge University Press, Cambridge, 2003
28. Wells, R. O.: Comparison of de Rham and Dolbeault cohomology for proper surjective mappings. *Pacific J. Math.* **53**, 281-300 (1974)

29. Yang, X., Zhao, G.: A note on the Morse-Novikov cohomology of blow-ups of locally conformal Kähler manifolds. *Bull. Aust. Math. Soc.* **91**, 155-166 (2015)

DEPARTMENT OF MATHEMATICS, NORTH UNIVERSITY OF CHINA, TAIYUAN, SHANXI 030051, P.R. CHINA
E-mail address: menglingxu@nuc.edu.cn