

# Prescribing Morse scalar curvatures: subcritical blowing-up solutions

Andrea Malchiodi & Martin Mayer

Published on Journal of Differential Equations

Scuola Normale Superiore, Pisa, ITALY  
andrea.malchiodi@sns.it , martin.mayer@sns.it

January 28, 2020

## Abstract

Prescribing conformally the scalar curvature of a Riemannian manifold as a given function consists in solving an elliptic PDE involving the critical Sobolev exponent. One way of attacking this problem consist in using subcritical approximations for the equation, gaining compactness properties. Together with the results in [30] we completely describe the blow-up phenomenon in case of uniformly bounded energy, zero weak limit and positive Yamabe invariant. In particular for dimension greater or equal to five and Morse functions with non-zero Laplacian at each critical point we show, that subsets of critical points with negative Laplacian are in one-to-one correspondence with such subcritical blowing-up solutions. *Key Words:* Conformal geometry, sub-critical approximation, blow-up analysis.

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Preliminaries</b>	<b>6</b>
<b>3</b>	<b>Existence of subcritical solutions</b>	<b>9</b>
<b>4</b>	<b>The second variation</b>	<b>16</b>
<b>5</b>	<b>Appendix: Some estimates and list of constants</b>	<b>33</b>

## 1 Introduction

Consider a compact manifold  $(M^n, g_0)$  with  $n \geq 3$  and a conformal metric

$$g = u^{\frac{4}{n-2}} g_0, \quad u > 0.$$

With this notation the scalar curvature transforms via

$$R_{g_u} u^{\frac{n+2}{n-2}} = L_{g_0} u = -c_n \Delta_{g_0} u + R_{g_0} u, \quad c_n = \frac{4(n-1)}{(n-2)}$$

with  $\Delta_{g_0}$  denoting the Laplace-Beltrami operator of  $g_0$ , cf. [4].  $L_{g_0}$  is called the *conformal Laplacian* and transforms according to

$$L_g(\phi) = u^{-\frac{n+2}{n-2}} L_{g_0}(u\phi).$$

In the 70's, Kazdan and Warner considered in [28] the problem of prescribing the scalar curvature of manifolds via conformal deformation of the metric, see also [26], [27]. By the above transformation law, if one wishes to prescribe  $R_g$  as a given function  $K$  then would need to solve

$$L_{g_0} u = K u^{\frac{n+2}{n-2}} \quad \text{on} \quad (M, g_0). \quad (1)$$

There are rather easy obstructions to the solvability of (1). For example, if the sign of  $K$  is constant, it has to coincide with that of the first eigenvalue of  $L_{g_0}$ . Depending on the latter sign, which is conformally invariant, a conformal class of metrics is said to be of *negative, zero or positive Yamabe class*. We will discuss for simplicity the case of function  $K$  with constant sign, despite in the literature there are many interesting papers dealing with changing-sign functions.

In [28] Kazdan and Warner proved existence results for zero or negative Yamabe classes using the sub- and super-solution method. For positive Yamabe class instead, they found a now well-known obstruction to existence on the sphere, namely that if  $u$  solves (1), then for  $f$  affine on  $S^n$  one must have

$$\int_{S^n} \langle \nabla K, \nabla f \rangle_{g_{S^n}} u^{\frac{2n}{n-2}} d\mu_{g_{S^n}} = 0, \quad (2)$$

and hence, for conformal curvatures  $K$ , the function  $\langle \nabla K, \nabla f \rangle_{g_{S^n}}$  must change sign.

Later on some existence results were found under conditions that would imply *topological richness* of the sub-levels of  $K$ , contrary to the above example. In two dimensions, where (1) is replaced by an equation in exponential form, J. Moser showed that the problem is solvable on the standard sphere if  $K$  is antipodally symmetric. In higher dimensions, existence results under the action of symmetry groups were proven in [20] and [21], [22]. A general difficulty in studying (1) is the lack of compactness due to the presence of the critical exponent. A typical phenomenon encountered here is that of *bubbling*. *Bubbles* are solutions of (1) on  $S^n$  with  $K \equiv 1$  and these arise as profiles of general diverging solutions and were classified in [11], see also [3], [37]. From the variational point of view bubbles generate diverging Palais-Smale sequences for the Euler-Lagrange energy of (1)

$$J(u) = J_K(u) = \frac{\int_M (c_n |\nabla u|_{g_0}^2 + R_{g_0} u^2) d\mu_{g_0}}{\left( \int_M K u^{\frac{2n}{n-2}} d\mu_{g_0} \right)^{\frac{n-2}{n}}}.$$

As seen from a formal expansion of  $J$  on a finite sum of bubbles, cf. the introduction in [30], the mutual interaction among bubbles becomes weaker as  $n$  increases. As a consequence in case  $n = 3$  at most one bubble can form. Exploiting this and after some work on  $S^2$  by A. Chang and P. Yang in [16], [17], A. Bahri and J.M. Coron proved an existence result in [6] on  $S^3$  assuming that  $K$  is a Morse function and

$$\{\nabla K = 0\} \cap \{\Delta K = 0\} = \emptyset; \quad (3)$$

$$\sum_{\{x \in M : \nabla K(x)=0, \Delta K(x)<0\}} (-1)^{m(x,K)} \neq -1, \quad (4)$$

where  $m(x, K)$  denotes the Morse index of  $K$  at  $x$ , cf. [12] and [36] for more general related results. The above existence statement was extended to arbitrary dimensions in [24] for functions satisfying a suitable flatness condition, and in [18], [1], [29] for functions  $K$  close to a positive constant in the  $C^2$ -sense.

In four dimensions, see [7] and [25], it was shown that even if multiple bubbles can form, they cannot be too close to each-other; such phenomenon is usually referred to as *isolated simple blow-up*. Results of different kind were also proven in [19] for  $n = 2$  and in [9][8], [10], see also Chapter 6 in [4].

Two main approaches have been used to understand the blow-up phenomenon, namely sub-critical approximations or the construction of pseudo-gradient flows. In this paper we focus on the former, while the other one will be the subject of [33], where a one-to-one correspondence of zero weak limit blowing-up solutions with bounded energy and *critical points at infinity* is shown, see also [34]. Consider the problem

$$-c_n \Delta_{g_0} u + R_{g_0} u = K u^{\frac{n+2}{n-2}-\tau}, \quad 0 < \tau \ll 1, \quad (5)$$

which upon rescaling is the Euler-Lagrange equation for the functional

$$J_\tau(u) = \frac{\int_M (c_n |\nabla u|_{g_0}^2 + R_{g_0} u^2) d\mu_{g_0}}{(\int_M K u^{p+1} d\mu_{g_0})^{\frac{2}{p+1}}}, \quad u \in \mathcal{A}. \quad (6)$$

Being now the exponent lower than critical, solutions can be easily found, even though one could lose uniform estimates as  $\tau$  tends to zero. In [12], [36], [24] the single-bubbling behaviour for diverging solutions of (5) was proved. Then by degree- or Morse-theoretical arguments it was shown that under (4) there must be families of solutions that stay uniformly bounded, therefore converging to solutions of (1). For this argument to work, one crucial step was to completely characterize blowing-up solutions of (5), showing that in three dimensions single blow-ups occur at any critical point of  $K$  with negative Laplacian and that they are unique. On four-dimensional spheres, a similar property was proved in [25] for multiple blow-ups, see also [7], assuming a suitable condition related to the multi-bubble interactions.

For Morse functions, if  $n \geq 5$  the situation is more involved, and blow-ups might be possibly of infinite energy, see e.g. [13], [14], [15], [38]. In [30] it

was however proved that if a sequence of blowing-up solutions has uniformly-bounded  $W^{1,2}$ -energy and zero weak limit, then blow-ups are still isolated simple. Although the result is similar to the case of dimensions three and four, the phenomenon is somehow opposite since it is *driven* by the function  $K$  rather than from the mutual bubble interactions. Both assumptions, i.e. zero weak limit and bounded energy, are indeed natural. If the former fails then problem (1) would have a solution; the second one instead is usually found when using min-max or Morse-theoretical arguments, as it will be done in [31]. However, differently from  $n = 3, 4$ , in [30] no restriction is proven on the number or location of blow-up points, provided they occur at critical points of  $K$  with negative Laplacian.

In this paper reshown, that the characterization of the above blow-ups in [30] is sharp, namely that they can occur at arbitrary subsets of

$$\{\nabla K = 0\} \cap \{\Delta K < 0\}.$$

Furthermore, we prove uniqueness of such solutions, their non-degeneracy and determine their Morse index. Our main result is the following one, that follows from Theorem 1 in [30] and from Proposition 3.1, Corollary 4.1.

**Theorem 1.** *Let  $(M, g)$  be a compact manifold of dimension  $n \geq 5$  of positive Yamabe invariant and let  $K : M \rightarrow \mathbb{R}$  be a positive Morse function satisfying (3). Let  $x_1, \dots, x_q$  be distinct critical points of  $K$  with negative Laplacian. Then there exists, as  $\tau \rightarrow 0$  and up to scaling, a unique solution  $u_{\tau, x_1, \dots, x_q}$  developing a simple bubble at each point  $x_i$  converging weakly to zero in  $W^{1,2}(M, g)$  as  $\tau \rightarrow 0$ . Moreover and up to scaling  $u_{\tau, x_1, \dots, x_q}$  is non-degenerate for  $J_\tau$  and*

$$m(J_\tau, u_{\tau, x_1, \dots, x_q}) = (q - 1) + \sum_{i=1}^q (n - m(K, x_i)).$$

*Conversely all blow-ups of uniformly bounded energy and zero weak limit type are as above.*

As it will be shown in [31], for  $n \geq 5$  there cannot be a direct counterpart of (4), which is an index-counting condition. However, existence results of different type will be derived there.

**Remark 1.1.** (i) *More precise expressions for  $u_{\tau, x_1, \dots, x_q}$  are given by*

$$\left\| u_m - \sum_{j=1}^q \alpha_{j,m} \delta_{\lambda_{j,m}, a_{j,m}} \right\|_{W^{1,2}(M, g_0)} \rightarrow 0 \quad \text{as} \quad m \rightarrow \infty,$$

and

$$\alpha_{j,m} = \frac{\Theta}{K(x_j)^{\frac{n-2}{4}}} + o(1), \quad a_{j,m} \rightarrow x_j \quad \text{and} \quad \lambda_{j,m} \simeq \lambda_{\tau_m} = \tau_m^{-\frac{1}{2}}.$$

Here the multiplicative constant  $\Theta$  depends on the blowing-up solutions but it is independent of  $j$ . For this and more precise formulae we refer to Section 3 and Theorem 2 in the Appendix. If  $n = 4$ , the same conclusions hold replacing  $\Delta K(a_j) < 0$  for all  $j$  with (iv) of Theorem 2 in [30].

- (ii) Although upon scaling the above solutions  $u_{\tau, x_1, \dots, x_q}$  are non-degenerate, they Hessian of  $J_\tau$  there has  $\sum_{i=1}^q (n - m(K, x_i))$  eigenvalues approaching zero as  $\tau \rightarrow 0$ , see Section 4.
- (iii) Theorem 1 gives a one-to-one correspondence of zero weak limit subcritical blow-up solutions to subsets of critical points of  $K$  with negative Laplacian, while in [33] this correspondence is shown with zero weak limits, i.e. pure critical points at infinity of energy decreasing type, cf. [5], [32].

The proof of Theorem 1 relies on the estimates in [30] and a finite dimensional reduction, see e.g. [2], with a careful asymptotic analysis. In dimension four this approach was used in Section 2 of [25]. Here we show that in higher dimensions blow-up might occur at arbitrary critical points of  $K$  with negative Laplacian, which affects the global structure of the solutions of problem (1). Via careful expansions, we also determine the Hessian of the Euler-Lagrange functional and the Morse index of these solutions, which we prove to be non-degenerate.

The solutions we consider here lie in a set  $V(q, \varepsilon) \subset W^{1,2}(M, g_0)$ , which contains a manifold of approximate solutions for (5), namely

$$\sum_{i=1}^q \alpha^i \varphi_{a_i, \lambda_i},$$

and is transversally non-degenerate, Section 2 for notation. This allows to solve (5) orthogonally to this manifold via a proper transversal correction to the approximate solutions, see Definition 3.1 and Lemma 3.1, and reduce to the study of the tangent component. By Theorem 2 from [30] we can reduce ourselves to a smaller set  $\bar{V}(q, \varepsilon)$ , see (17), where more precise estimates hold for the gradient of  $J_\tau$ . These allow us to use an orthogonal correction  $\bar{v}$  small in size, solve also for the tangent component and to estimate the second differential of  $J_\tau$  at  $\sum_{i=1}^q \alpha^i \varphi_{a_i, \lambda_i} + \bar{v}$ , see Section 4. Finally this allows in turn to compute the Morse index of the solutions  $u_{\tau, x_1, \dots, x_q}$  and to prove their uniqueness. In this step we show that even though the correction  $\bar{v}$  is of the same order as some eigenvalues of  $\partial^2 J_\tau$  some cancellation occur in the corresponding estimates.

The plan of the paper is the following. In Section 2 we collect some preliminary material concerning approximate solutions and the finite-dimensional reduction of the problem, which is then worked-out in detail in Section 3. In Section 4 we study the Hessian of the Euler-Lagrange functional  $J_\tau$  in  $\bar{V}(q, \varepsilon)$ , finding a proper base with respect to which the Hessian nearly diagonalizes. Finally we collect in an Appendix some useful and technical estimates from [30] and a table of constants.

**Acknowledgements.** A.M. has been supported by the project *Geometric Variational Problems* and *Finanziamento a supporto della ricerca di base* from Scuola Normale Superiore and by MIUR Bando PRIN 2015 2015KB9WPT 001. He is also member of GNAMPA as part of INdAM.

## 2 Preliminaries

In this section we collect some background and preliminary material, concerning the variational properties of the problem and some estimates on highly-concentrated approximate solutions of bubble type.

We consider a smooth, closed Riemannian manifold  $M = (M^n, g_0)$  with volume measure  $\mu_{g_0}$  and scalar curvature  $R_{g_0}$ . Letting

$$\mathcal{A} = \{u \in W^{1,2}(M, g_0) \mid u \geq 0, u \not\equiv 0\}$$

the *Yamabe invariant* is defined as

$$Y(M, g_0) = \inf_{\mathcal{A}} \frac{\int (c_n |\nabla u|_{g_0}^2 + R_{g_0} u^2) d\mu_{g_0}}{\left(\int u^{\frac{2n}{n-2}} d\mu_{g_0}\right)^{\frac{n-2}{n}}} \quad \text{with} \quad c_n = 4 \frac{n-1}{n-2}$$

and it turns out to depend only on the conformal class of  $g_0$ . We will assume that this invariant is positive, i.e.  $(M, g_0)$  to be of *positive Yamabe class*. As a consequence the *conformal Laplacian*

$$L_{g_0} = -c_n \Delta_{g_0} + R_{g_0}$$

is a positive and self appointed operator. Without loss of generality we assume  $R_{g_0} > 0$  and denote by

$$G_{g_0} : M \times M \setminus \Delta \longrightarrow \mathbb{R}_+$$

the Green function of  $L_{g_0}$ . Considering a conformal metric

$$g = g_u = u^{\frac{4}{n-2}} g_0$$

there holds

$$d\mu_{g_u} = u^{\frac{2n}{n-2}} d\mu_{g_0} \quad \text{and} \quad R = R_{g_u} = u^{-\frac{n+2}{n-2}} (-c_n \Delta_{g_0} u + R_{g_0} u) = u^{-\frac{n+2}{n-2}} L_{g_0} u.$$

Note that

$$c \|u\|_{W^{1,2}(M, g_0)} \leq \int u L_{g_0} u d\mu_{g_0} \leq C \|u\|_{W^{1,2}(M, g_0)}.$$

In particular we may define

$$\|u\|^2 = \|u\|_{L_{g_0}}^2 = \int u L_{g_0} u d\mu_{g_0}$$

and use  $\|\cdot\|$  as an equivalent norm on  $W^{1,2}(M, g_0)$ . Setting

$$R = R_u \quad \text{for} \quad g = g_u = u^{\frac{4}{n-2}} g_0$$

we have

$$r = r_u = \int R d\mu_{g_u} = \int u L_{g_0} u d\mu_{g_0}, \quad (7)$$

and hence

$$J_\tau(u) = \frac{r}{k_\tau^{\frac{2}{p+1}}} \quad \text{with} \quad k_\tau = \int K u^{p+1} d\mu_{g_0}. \quad (8)$$

The first- and second-order derivatives of the functional  $J_\tau$  are given by

$$\partial J_\tau(u)v = \frac{2}{k_\tau^{\frac{2}{p+1}}} \left[ \int L_{g_0} u v d\mu_{g_0} - \frac{r}{k_\tau} \int K u^p v d\mu_{g_0} \right]; \quad (9)$$

and

$$\begin{aligned} \partial^2 J_\tau(u)vw &= \frac{2}{k_\tau^{\frac{2}{p+1}}} \left[ \int L_{g_0} v w d\mu_{g_0} - p \frac{r}{k_\tau} \int K u^{p-1} v w d\mu_{g_0} \right] \\ &\quad - \frac{4}{k_\tau^{\frac{2}{p+1}+1}} \left[ \int L_{g_0} u v d\mu_{g_0} \int K u^p w d\mu_{g_0} \right. \\ &\quad \quad \quad \left. + \int L_{g_0} u w d\mu_{g_0} \int K u^p v d\mu_{g_0} \right] \\ &\quad + \frac{2(p+3)r}{k_\tau^{\frac{2}{p+1}+2}} \int K u^p v d\mu_{g_0} \int K u^p w d\mu_{g_0}. \end{aligned} \quad (10)$$

In particular  $J_\tau$  is of class  $C_{loc}^{2,\alpha}(\mathcal{A})$  and for  $\varepsilon > 0$  uniformly Hölder continuous on each set of the form

$$U_\varepsilon = \{u \in \mathcal{A} \mid \varepsilon < \|u\|, J_\tau(u) \leq \varepsilon^{-1}\}.$$

To understand the blow-up phenomenon, it is convenient to consider some highly concentrated approximate solutions to (1). Let us first recall the construction of *conformal normal coordinates* from [23]. Given  $a \in M$  these are defined as geodesic normal coordinates for a suitable conformal metric  $g_a \in [g_0]$ . Let  $r_a$  be the geodesic distance from  $a$  with respect to the metric  $g_a$ . With this choice, the expression of the Green function  $G_{g_a}$  for the conformal Laplacian  $L_{g_a}$  with pole at  $a \in M$ , denoted by  $G_a = G_{g_a}(a, \cdot)$ , simplifies considerably. In Section 6 of [23] one can find the expansion

$$G_a = \frac{1}{4n(n-1)\omega_n} (r_a^{2-n} + H_a), \quad H_a = H_{r,a} + H_{s,a} \quad \text{for} \quad g_a = u_a^{\frac{4}{n-2}} g_0. \quad (11)$$

Here  $r_a = d_{g_a}(a, \cdot)$  and  $H_{r,a} \in C_{loc}^{2,\alpha}$ , while the *singular* error term is of type

$$H_{s,a} = O \left( \begin{array}{ll} r_a & \text{for } n = 5 \\ \ln r_a & \text{for } n = 6 \\ r_a^{6-n} & \text{for } n \geq 7 \end{array} \right).$$

The leading term in  $H_{s,a}$  for  $n = 6$  is  $\mathbb{W}$  the Weyl tensor. Define

$$\varphi_{a,\lambda} = u_a \left( \frac{\lambda}{1 + \lambda^2 \gamma_n G_a^{\frac{2}{2-n}}} \right)^{\frac{n-2}{2}}, \quad G_a = G_{g_a}(a, \cdot) \quad (12)$$

for  $\lambda > 0$  large, where  $\gamma_n = (4n(n-1)\omega_n)^{\frac{2}{2-n}}$  is chosen so that

$$\gamma_n G_a^{\frac{2}{2-n}}(x) = d_{g_a}^2(a, x) + o(d_{g_a}^2(a, x)) \text{ as } x \rightarrow a. \quad (13)$$

Such functions are approximate solutions of (1), see Lemma 5.1, and for suitable values of  $\lambda$  depending on  $\tau$  these are also approximate solutions of (5), see Lemma 5.7 for a multi-bubble version.

**Notation.** For  $p \geq 1$ ,  $L_{g_0}^p$  will stand for the family of functions of class  $L^p$  with respect to the measure  $d\mu_{g_0}$ . Recall also that for  $u \in W^{1,2}(M, g_0)$  we have set  $r_u = \int u L_{g_0} u d\mu_{g_0}$ , while for  $a \in M$  we denote by  $r_a$  the geodesic distance from  $a$  with respect to the conformal metric  $g_a$  introduced before. For a finite set of points  $\{a_i\}_i$  of  $M$  we will denote by  $K_i, \nabla K_i, W_i$  the quantities  $K(a_i), \nabla K(a_i), |\mathbb{W}(a_i)|^2$  etc..

For  $k, l = 1, 2, 3$  and  $\lambda_i > 0$ ,  $a_i \in M$ ,  $i = 1, \dots, q$  let

- (i)  $\varphi_i = \varphi_{a_i, \lambda_i}$  and  $(d_{1,i}, d_{2,i}, d_{3,i}) = (1, -\lambda_i \partial_{\lambda_i}, \frac{1}{\lambda_i} \nabla_{a_i})$ ;
- (ii)  $\phi_{1,i} = \varphi_i$ ,  $\phi_{2,i} = -\lambda_i \partial_{\lambda_i} \varphi_i$ ,  $\phi_{3,i} = \frac{1}{\lambda_i} \nabla_{a_i} \varphi_i$ , so  $\phi_{k,i} = d_{k,i} \varphi_i$ .

Note, that the  $\phi_{k,i}$  are uniformly bounded in  $W^{1,2}(M, g_0)$  for any  $\lambda_i > 0$ .

We next recall a standard finite-dimensional reduction for functions that are close in  $W^{1,2}$  to a finite sum of bubbles, wherefore we define

$$\varepsilon_{i,j} = \left( \frac{\lambda_j}{\lambda_i} + \frac{\lambda_i}{\lambda_j} + \lambda_i \lambda_j \gamma_n G_{g_0}^{\frac{2}{2-n}}(a_i, a_j) \right)^{\frac{2-n}{2}}. \quad (14)$$

Given  $\varepsilon > 0$ ,  $q \in \mathbb{N}$ ,  $u \in W^{1,2}(M, g_0)$  and  $(\alpha^i, \lambda_i, a_i) \in (\mathbb{R}_+^q, \mathbb{R}_+^q, M^q)$ , we set

- (i)  $A_u(q, \varepsilon) = \{(\alpha^i, \lambda_i, a_i) \mid \forall_{i \neq j} \lambda_i^{-1}, \lambda_j^{-1}, \varepsilon_{i,j}, \left| 1 - \frac{r \alpha_i^{\frac{4}{n-2}} K(a_i)}{4n(n-1)k_\tau} \right|, \|u - \alpha^i \varphi_{a_i, \lambda_i}\| < \varepsilon, \lambda_i^\tau < 1 + \varepsilon\}$ ;

- (ii)  $V(q, \varepsilon) = \{u \in W^{1,2}(M, g_0) \mid A_u(q, \varepsilon) \neq \emptyset\}$ ,

see(7), (8) and (12). For  $A_u(q, \varepsilon)$  to be non-empty we will always assume that  $\tau \ll \varepsilon$ . Under the above conditions on the parameters  $\alpha_i, a_i$  and  $\lambda_i$  the functions  $\sum_{i=1}^q \alpha^i \varphi_{a_i, \lambda_i}$  constitute a smooth manifold in  $W^{1,2}(M, g_0)$ , which implies the following well known result, cf. [5].



**Proposition 2.1.** *Given  $\varepsilon_0 > 0$  there exists  $\varepsilon_1 > 0$  such that for  $u \in V(q, \varepsilon)$  with  $\varepsilon < \varepsilon_1$ , the problem*

$$\inf_{(\tilde{\alpha}_i, \tilde{a}_i, \tilde{\lambda}_i) \in A_u(q, 2\varepsilon_0)} \int (u - \tilde{\alpha}^i \varphi_{\tilde{a}_i, \tilde{\lambda}_i}) L_{g_0} (u - \tilde{\alpha}^i \varphi_{\tilde{a}_i, \tilde{\lambda}_i}) d\mu_{g_0}$$

*admits an unique minimizer  $(\alpha_i, a_i, \lambda_i) \in A_u(q, \varepsilon_0)$  and we set*

$$\varphi_i = \varphi_{a_i, \lambda_i}, \quad v = u - \alpha^i \varphi_i, \quad K_i = K(a_i). \quad (15)$$

*Moreover  $(\alpha_i, a_i, \lambda_i)$  depends smoothly on  $u$ .*

The term  $v = u - \alpha^i \varphi_i$  is orthogonal to all  $\varphi_i, -\lambda_i \partial_{\lambda_i} \varphi_i, \frac{1}{\lambda_i} \nabla_{a_i} \varphi_i$ , with respect to the product

$$\langle \cdot, \cdot \rangle_{L_{g_0}} = \langle L_{g_0} \cdot, \cdot \rangle_{L_{g_0}^2}.$$

Finally for  $u \in V(q, \varepsilon)$  let

$$H_u = H_u(q, \varepsilon) = \langle \varphi_i, \lambda_i \partial_{\lambda_i} \varphi_i, \frac{1}{\lambda_i} \nabla_{a_i} \varphi_i \rangle^{\perp_{L_{g_0}}}. \quad (16)$$

### 3 Existence of subcritical solutions

Theorem 2, from [30], describes in detail the behaviour as  $\tau \rightarrow 0$  of blowing-up solutions to (5) with uniformly bounded energy and zero weak limit in  $V(q, \varepsilon)$ , providing positive lower bounds on  $\|\partial J_\tau\|$  in a suitable subset of the functional space. In view of this, we can restrict our attention to *centers*  $a_1, \dots, a_q$  close to distinct critical points  $x_1, \dots, x_q$  of  $K$  with negative Laplacian. More precisely for  $n \geq 6$  we can assume the following conditions, which for  $n = 5$  are slightly modified,

- (i)  $|\alpha_j - \Theta \sqrt[p-1]{\frac{\lambda_j^\theta}{K(a_j)}}| < \frac{\varepsilon}{\lambda_j^3};$
- (ii)  $|\frac{\tilde{a}_j}{\lambda_j} + c_1(\nabla^2 K(x_j)) - \frac{\nabla \Delta K(x_j)}{\lambda_j^3}| \leq \frac{\varepsilon}{\lambda_j^3};$
- (iii)  $|\lambda_j^2 + c_2 \frac{\Delta K(x_j)}{K(x_j)\tau}| \leq \frac{\varepsilon}{\lambda_j^2},$

for  $\lambda^2 = \frac{1}{\tau}$  and some

$$x_j \in \{\nabla K = 0\} \cap \{\Delta K < 0\} \quad \text{with} \quad x_i \neq x_j \quad \text{for} \quad i \neq j.$$

Here  $\Theta > 0$  is uniformly bounded and bounded away from zero and depends on the function in  $V(q, \varepsilon)$ , determined in Remark 6.2 of [30]. We then define a neighbourhood of potential subcritical blowing-up solutions as

$$\bar{V}(q, \varepsilon) = \{u \in V(q, \varepsilon) \mid \text{(i), (ii) and (iii) above hold true}\}. \quad (17)$$

Indeed from Lemmata 5.4, 5.5 and 5.6 it follows, that there exists  $\tilde{\varepsilon} > 0$ , tending to zero as  $\varepsilon \rightarrow 0$ , such that

$$|\partial J_\tau(u)| \gtrsim \frac{\tilde{\varepsilon}}{\lambda^3} \quad \text{for } u \in V(q, \varepsilon) \setminus \bar{V}(q, \varepsilon) \quad \text{with } k_\tau = 1,$$

so this justifies to look for solutions in  $\bar{V}(q, \varepsilon)$  only. Moreover for

$$\alpha^i \varphi_i \in \bar{V}(q, \varepsilon) \quad \text{with } c < \alpha_i < C$$

we have the expansion

$$J_\tau(\alpha^i \varphi_i + v) = J_\tau(\alpha^i \varphi_i) + \partial J_\tau(\alpha^i \varphi_i)v + \frac{1}{2} \partial^2 J_\tau(\alpha^i \varphi_i)v^2 + O(\|v\|^3). \quad (18)$$

Recall the uniform positivity of  $\partial^2 J_\tau(\alpha^i \varphi_i)$  on  $H_u(q, \varepsilon)$ , cf. (16) and [5], which justifies the following

**Definition 3.1.** For  $\alpha^i \varphi_i \in V(q, \varepsilon)$  we define  $\bar{v}$  as the unique solution of the minimization problem

$$J_\tau(\alpha^i \varphi_i + \bar{v}) = \min_{v \in H_{\alpha^i \varphi_i}, \|v\| < \varepsilon} J_\tau(\alpha^i \varphi_i + v). \quad (19)$$

**Lemma 3.1.** Let  $\bar{v}$  be as in the above definition. Then

- (i) for  $\alpha^i \varphi_i \in \bar{V}(q, \varepsilon)$  there holds  $\|\bar{v}\| \lesssim \frac{1}{\lambda^2} \simeq \tau$  ;
- (ii) if  $u \in V(q, \varepsilon)$  is such that  $\partial J_\tau(u) = 0$ , then  $\alpha^i \varphi_i \in \bar{V}(q, \varepsilon)$  and  $u = \alpha^i \varphi_i + \bar{v}$ .

Moreover for  $\alpha^i \varphi_i \in \bar{V}(q, \varepsilon)$  we have

$$\partial J_\tau(\alpha^i \varphi_i + \bar{v}) = O\left(\frac{\tilde{\varepsilon}}{\lambda^3}\right), \quad \text{where } \tilde{\varepsilon} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad (20)$$

*Proof.* We first justify Definition 3.1, which amounts to solving in a unique way

$$\Pi_{H_{\alpha^i \varphi_i}} \partial J_\tau(\alpha^i \varphi_i + \bar{v}) = 0 \quad (21)$$

denoting by

$$\Pi_{H_{\alpha^i \varphi_i}} \hat{=} \quad \text{the projection onto } H_{\alpha^i \varphi_i}.$$

Equivalently, as  $\partial^2 J_\tau$  is invertible on the latter subspace,

$$\begin{aligned} \bar{v} = & -(H_{\alpha^i \varphi_i} \partial^2 J_\tau(\alpha^i \varphi_i))^{-1} \\ & [\partial J_\tau(\alpha^i \varphi_i) + (\partial J_\tau(\alpha^i \varphi_i + \bar{v}) - \partial J_\tau(\alpha^i \varphi_i) - \partial^2 J_\tau(\alpha^i \varphi_i)\bar{v})]. \end{aligned}$$

Note, that by Lemma 5.7 and for  $\alpha^i \varphi_i \in \bar{V}(q, \varepsilon)$  we have

$$\|\partial J_\tau(\alpha^i \varphi_i)\| \lesssim \frac{1}{\lambda^2}.$$

Moreover

$$J_\tau(\alpha^i \varphi_i + \bar{v}) - \partial J_\tau(\alpha^i \varphi_i) - \partial^2 J_\tau(\alpha^i \varphi_i) \bar{v} = o(\|\bar{v}\|)$$

by Hölder continuity. Hence we may use a contraction argument in a ball

$$B_{\frac{C}{\lambda^2}}(0) \subset H_{\alpha^i \varphi_i}$$

to obtain existence of some  $\bar{v}$  solving (21) and satisfying estimate (i). Uniqueness follows from the aforementioned invertibility. Hence we have justified Definition 3.1. We are left with proving (ii).

By the definition of  $\bar{v}$  and the above contraction argument we have that

$$\partial^2 J_\tau(\alpha^i \varphi_i) \bar{v} = -\partial J_\tau(\alpha^i \varphi_i) + o\left(\frac{1}{\lambda^2}\right) \text{ on } \langle \phi_{k,i} \rangle^{\perp L_{g_0}}. \quad (22)$$

Testing thus  $\partial J_\tau(\alpha^i \varphi_i)$  on  $\langle \phi_{k,i} \rangle$ , we find from Lemmata 5.4, 5.5 and 5.6

$$|\partial J_\tau(\alpha^i \varphi_i) \phi_{k,i}| \leq \frac{\tilde{\epsilon}}{\lambda^3} \quad \text{for } \alpha^i \varphi_i \in \bar{V}(q, \varepsilon).$$

It is easy to see from (10) and Lemma 5.1 that  $\partial^2 J_\tau \phi_{k,i} = o(\frac{1}{\lambda})$ , and since  $\|\bar{v}\| \lesssim \frac{1}{\lambda^2}$  we have that

$$\partial^2 J_\tau(\alpha^i \varphi_i) \bar{v} \phi_{k,i} = o\left(\frac{1}{\lambda^3}\right), \quad (23)$$

More generally we also find, that for any  $\theta \in (0, 1)$

$$\partial^2 J(\alpha^i \varphi_i + \theta \bar{v}) \bar{v} \phi_{k,j} = o\left(\frac{1}{\lambda^3}\right).$$

To see this, since  $\bar{v} \in \langle \phi_{k,i} \rangle^{\perp L_{g_0}}$ , recalling (10) it is sufficient to show that

$$\int K(\alpha^i \varphi_i + \theta \bar{v})^{p-1} \bar{v} \varphi_j d\mu_{g_0} - \int K(\alpha^i \varphi_i)^{p-1} \bar{v} \varphi_j d\mu_{g_0} = O\left(\frac{1}{\lambda^3}\right).$$

This in return can be verified by dividing the domain of integration into  $\{|\bar{v}| \leq \alpha^i \varphi_i\}$  and its complementary set, using Hölder inequality and the fact that  $\|\bar{v}\| \lesssim \frac{1}{\lambda^2}$ . Consequently

$$\partial J_\tau(\alpha^i \varphi_i + \bar{v}) = \partial J_\tau(\alpha^i \varphi_i + \bar{v})|_{\langle \phi_{k,i} \rangle} = \partial J_\tau(\alpha^i \varphi_i)|_{\langle \phi_{k,i} \rangle} + o\left(\frac{1}{\lambda^3}\right) = O\left(\frac{\tilde{\epsilon}}{\lambda^3}\right),$$

where  $\tilde{\epsilon}$  tends to zero as  $\varepsilon$  does. Finally, if a solution  $\partial J_\tau(u) = 0$  exists on  $V(q, \varepsilon)$ , then we may write

$$u = \alpha^i \varphi_i + \bar{v} + \tilde{v} \quad \text{with } \tilde{v} \perp_{L_{g_0}} \langle \phi_{k,i} \rangle.$$

But then

$$0 = \partial J_\tau(\alpha^i \varphi_i + \bar{v} + \tilde{v}) \tilde{v} = \partial J_\tau(\alpha^i \varphi_i + \bar{v}) \tilde{v} + \partial^2 J_\tau(\alpha^i \varphi_i + \bar{v}) \tilde{v} \tilde{v} + o(|\tilde{v}|^2),$$

whence necessarily  $\bar{v} = 0$  by uniform positivity

$$\partial^2 J_\tau(\alpha^i \varphi_i) \quad \text{on} \quad \langle \phi_{k,i} \rangle^{\perp L_{g_0}}.$$

Thus

$$\partial J_\tau(u) = 0 \quad \text{with} \quad u \in \bar{V}(q, \varepsilon) \quad \implies \quad u = \alpha^i \varphi_i + \bar{v}$$

where  $\bar{v} = \bar{v}_{\alpha, a, \lambda}$  is the unique solution to (19), for which  $\alpha^i \varphi_i + \bar{v} \in \bar{V}(q, \varepsilon)$ .  $\square$

**Remark 3.1.** For  $\alpha^i \varphi_i \in \bar{V}(q, \varepsilon)$  and  $\|\nu\| = 1$  we have

$$\begin{aligned} & \frac{(k_\tau)^{\frac{2}{p+1}}}{8n(n-1)} \alpha^i \varphi_i \partial J_\tau(\alpha^i \varphi_i) \nu \\ &= -\alpha^i \tau \int_{B_\varepsilon(a_i)} \left( \varphi_i^{\frac{n+2}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1} \varphi_i^{\frac{n+2}{n-2}} + \frac{2}{n-2} \frac{\tilde{c}_1}{c_2} \varphi_i^{\frac{4}{n-2}} \lambda_i \partial_{\lambda_i} \varphi_i \right) \nu d\mu_{g_0} \\ &+ \alpha^i \tau \int_{B_\varepsilon(a_i)} \left( \frac{\tilde{c}_1}{\tilde{c}_2} \frac{\lambda_i^2 r^2}{2n} \varphi_i^{\frac{n+2}{n-2}} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 c_1} \varphi_i^{\frac{n+2}{n-2}} + \frac{2}{n-2} \frac{\tilde{c}_1}{c_2} \varphi_i^{\frac{4}{n-2}} \lambda_i \partial_{\lambda_i} \varphi_i \right) \nu d\mu_{g_0} \\ &- \alpha^i \int_{B_\varepsilon(a_i)} \left( \frac{\nabla_{k,l}^2 K_i}{2K_i} x^k x^l - \frac{\Delta K_i}{2nK_i} r^2 \right) \varphi_i^{\frac{n+2}{n-2}} \nu d\mu_{g_0} + o\left(\frac{1}{\lambda^2}\right), \end{aligned}$$

referring to the table at the end of the paper for the definition of the constants. As a consequence of these formulae one can prove that  $\bar{v}$  is indeed of order  $\frac{1}{\lambda^2}$  and not smaller, as well as determine the leading order in its expansion. In any case due to some cancellation properties this will not substantially affect the eigenvalues of the Hessian of  $J_\tau$  at  $\alpha^i \varphi_i + \bar{v}$ , which we estimate in the next section.

Let us set  $(d_{1,i}, d_{2,i}, d_{3,i}) = (1, -\lambda_i \partial_{\lambda_i}, \frac{1}{\lambda_i} \nabla_{a_i})$  for  $i = 1, \dots, q$ .

**Lemma 3.2.** For  $u = \alpha^i \varphi_i + \bar{v} \in \bar{V}(q, \varepsilon)$  there holds

$$\|\bar{v}\|, \|d_{l,j} \bar{v}\| = O\left(\frac{1}{\lambda^2}\right).$$

*Proof.* The bound on  $\|\bar{v}\|$  follows from Lemma 3.1. Differentiating

$$\langle \phi_{k,i}, \bar{v} \rangle_{L_{g_0}} = 0$$

we obtain

$$\langle \phi_{k,i}, d_{l,j} \bar{v} \rangle_{L_{g_0}} = -\langle d_{l,j} \phi_{k,i}, \bar{v} \rangle_{L_{g_0}} = O(\|\bar{v}\|),$$

whence denoting by  $\Pi_{\langle \phi_{k,i} \rangle}$  the orthogonal projection onto  $\Pi_{\langle \phi_{k,i} \rangle}$  we have

$$\|\Pi_{\langle \phi_{k,i} \rangle} \bar{v}\| \simeq \frac{1}{\lambda^2} \quad \text{due to} \quad \|\bar{v}\| \lesssim \frac{1}{\lambda^2}.$$

Moreover, since  $\partial J_\tau(\alpha^i \varphi_i + \bar{v})v = 0$  for every smoothly varying vector field  $v \in \langle \phi_{k,i} \rangle^{\perp L_{g_0}}$  of unit norm we have

$$0 = d_{l,j}(\partial J_\tau(\alpha^i \varphi_i + \bar{v})v) = \partial^2 J_\tau(\alpha^i \varphi_i + \bar{v})d_{l,j}(\alpha^i \varphi_i + \bar{v})v + \partial J_\tau(\alpha^i \varphi_i + \bar{v})d_{l,j}v$$

and we can estimate the last summand above as

$$\partial J_\tau(\alpha^i \varphi_i + \bar{v})d_{l,j}v = \partial J_\tau(\alpha^i \varphi_i + \bar{v})\Pi_{\langle \phi_{k,i} \rangle}(d_{l,j}v) = O(\|\partial J_\tau(\alpha^i \varphi_i + \bar{v})\| \|v\|),$$

since

$$\langle \phi_{k,i}, d_{l,j}v \rangle = \langle d_{l,j}\phi_{k,i}, v \rangle = O(\|v\|).$$

Thence  $\partial J_\tau(\alpha^i \varphi_i + \bar{v}) = O(\frac{1}{\lambda^2})$  implies

$$\partial^2 J_\tau(\alpha^i \varphi_i + \bar{v})v d_{l,j}\bar{v} = -\partial^2 J_\tau(\alpha^i \varphi_i + \bar{v})v d_{l,j}(\alpha^i \varphi_i) + O(\frac{1}{\lambda^2}).$$

Then the claim would follow from

$$\|\Pi_{\langle \phi_{k,i} \rangle}(d_{l,j}\bar{v})\| \simeq \frac{1}{\lambda^2},$$

which we had seen before, and the uniform positivity

$$\partial^2 J_\tau(\alpha^i \varphi_i) > 0 \quad \text{on} \quad \langle \phi_{k,i} \rangle^{\perp L_{g_0}},$$

provided we show

$$\partial^2 J_\tau(\alpha^i \varphi_i + \bar{v})\phi_{l,j}v = O(\frac{1}{\lambda^2}), \tag{24}$$

cf. (25), (31) for weaker statements. Let us prove (24) for  $l = 1$ . We claim

$$\partial^2 J_\tau(\alpha^i \varphi_i + \bar{v})\varphi_j v = \partial^2 J_\tau(\alpha^i \varphi_i)\varphi_j v + O(\frac{1}{\lambda^2}).$$

From (10), since  $v \in \langle \phi_{k,i} \rangle^{\perp L_{g_0}}$ , it is sufficient to show that we must show, cf. the proof of Lemma 3.1,

$$\int K(\alpha^i \varphi_i + \bar{v})^{p-1} v \varphi_j d\mu_{g_0} - \int K(\alpha^i \varphi_i)^{p-1} v \varphi_j d\mu_{g_0} = O(\frac{1}{\lambda^2}).$$

Again this can be seen considering the set  $\{|\bar{v}| \leq \alpha^i \varphi_i\}$  and its complementary, using Hölder inequality and  $\|\bar{v}\| \lesssim \frac{1}{\lambda^2}$ . Thus from the above claim and (10) we find, due to the orthogonalities  $\langle \phi_{k,i}, v \rangle_{L_{g_0}} = 0$ ,

$$\begin{aligned} \partial^2 J_\tau(\alpha^i \varphi_i)\varphi_i v &= -\frac{2p}{(k_\tau)_{\alpha^i \varphi_i}^{\frac{2}{p+1}}} \frac{r_{\alpha^i \varphi_i}}{(k_\tau)_{\alpha^i \varphi_i}} \int K(\alpha^i \varphi_i)^{p-1} \varphi_j v d\mu_{g_0} \\ &\quad - \frac{4}{(k_\tau)_{\alpha^i \varphi_i}^{\frac{2}{p+1}+1}} \int L_{g_0}(\alpha^i \varphi_i)\varphi_j d\mu_{g_0} \int K(\alpha^i \varphi_i)^p v d\mu_{g_0} \\ &\quad + \frac{2(p+3)r_{\alpha^i \varphi_i}}{(k_\tau)_{\alpha^i \varphi_i}^{\frac{2}{p+1}+2}} \int K(\alpha^i \varphi_i)^p \varphi_j d\mu_{g_0} \int K(\alpha^i \varphi_i)^p v d\mu_{g_0}. \end{aligned}$$

By definition of  $\bar{V}(q, \varepsilon)$  we have  $\tau \simeq \frac{1}{\lambda^2}$  and recalling (37) and (40) we may simplify this to

$$\begin{aligned} \partial^2 J_\tau(\alpha^i \varphi_i) \varphi_j v &\simeq -4n(n-1) \frac{n+2}{n-2} \frac{\alpha^2}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} \int K(\alpha^i \varphi_i)^{\frac{4}{n-2}} \varphi_j v d\mu_{g_0} \\ &\quad - \frac{2}{\bar{c}_0 \alpha_{K,\tau}^{\frac{2n}{n-2}}} \int L_{g_0}(\alpha^i \varphi_i) \varphi_j d\mu_{g_0} \int K(\alpha^i \varphi_i)^{\frac{n+2}{n-2}} v d\mu_{g_0} \\ &\quad + 4n(n-1) \frac{(\frac{n+2}{n-2} + 3)\alpha^2}{\bar{c}_0 (\alpha_{K,\tau}^{\frac{2n}{n-2}})^2} \int K(\alpha^i \varphi_i)^{\frac{n+2}{n-2}} \varphi_j d\mu_{g_0} \int K(\alpha^i \varphi_i)^{\frac{n+2}{n-2}} v d\mu_{g_0} \end{aligned}$$

up to error  $O(\frac{1}{\lambda^2})$ . Moreover from (38) and (39) we have

$$\int L_{g_0}(\alpha^i \varphi_i) \varphi_j d\mu_{g_0} = 4n(n-1) \bar{c}_0 \alpha_j + O(\frac{1}{\lambda^2})$$

and since  $d(a_i, a_j) \simeq 1$ , we find by expanding and using Lemma 5.2

- (i)  $\int K(\alpha^i \varphi_i)^{\frac{4}{n-2}} \varphi_j v d\mu_{g_0} = \alpha_j^{\frac{4}{n-2}} \int K \varphi_j^{\frac{n+2}{n-2}} v d\mu_{g_0};$
- (ii)  $\int K(\alpha^i \varphi_i)^{\frac{n+2}{n-2}} v d\mu_{g_0} = \sum_i \alpha_i^{\frac{n+2}{n-2}} \int K \varphi_i^{\frac{n+2}{n-2}} v d\mu_{g_0};$
- (iii)  $\int K(\alpha^i \varphi_i)^{\frac{n+2}{n-2}} \varphi_j d\mu_{g_0} = \alpha_j^{\frac{n+2}{n-2}} \int K \varphi_j^{\frac{2n}{n-2}} d\mu_{g_0};$
- (iv)  $\int K(\alpha^i \varphi_i)^{\frac{n+2}{n-2}} v d\mu_{g_0} = \sum_i \alpha_i^{\frac{n+2}{n-2}} \int K \varphi_i^{\frac{n+2}{n-2}} v d\mu_{g_0},$

up to some  $O(\frac{1}{\lambda^2})$ . Therefore, since  $\frac{|\nabla K_i|}{\lambda_i} = O(\frac{1}{\lambda^2})$  due to (17), we obtain

$$\begin{aligned} \partial^2 J_\tau(\alpha^i \varphi_i) \varphi_j v &\simeq -4n(n-1) \frac{n+2}{n-2} \frac{\alpha^2}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} K_i \alpha_j^{\frac{4}{n-2}} \int \varphi_j^{\frac{n+2}{n-2}} v d\mu_{g_0} \\ &\quad - \frac{8n(n-1)\alpha_j}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} \sum_i K_i \alpha_i^{\frac{n+2}{n-2}} \int \varphi_i^{\frac{n+2}{n-2}} v d\mu_{g_0} \\ &\quad + 4n(n-1) \frac{(\frac{n+2}{n-2} + 3)\alpha^2}{(\alpha_{K,\tau}^{\frac{2n}{n-2}})^2} \alpha_j^{\frac{n+2}{n-2}} K_j \sum_i K_i \alpha_i^{\frac{n+2}{n-2}} \int \varphi_i^{\frac{n+2}{n-2}} v d\mu_{g_0} \end{aligned}$$

up to an error  $O(\frac{1}{\lambda^2})$ . Therefore using again (17) we have

$$\begin{aligned} \partial^2 J_\tau(\alpha^i \varphi_i) \varphi_j v &\simeq -\frac{n+2}{n-2} \int \varphi_j^{\frac{n+2}{n-2}} v d\mu_{g_0} - 2 \sum_i \frac{\alpha_i \alpha_j}{\alpha^2} \int \varphi_i^{\frac{n+2}{n-2}} v d\mu_{g_0} \\ &\quad + \left(\frac{n+2}{n-2} + 3\right) \sum_i \frac{\alpha_i \alpha_j}{\alpha^2} \int \varphi_i^{\frac{n+2}{n-2}} v d\mu_{g_0} \end{aligned}$$

up to the same error. Thus

$$\partial^2 J_\tau(\alpha^i \varphi_i) \varphi_j v = O\left(\frac{1}{\lambda^2}\right)$$

using (41), obtaining (24) for  $l = 1$ . For  $l = 2, 3$  the reasoning is analogous.  $\square$

Theorem 1 follows from the next proposition, based on the analysis of Section 4, and Corollary 4.1.

**Proposition 3.1.** *Let  $n \geq 5$  and let  $K : M \rightarrow \mathbb{R}$  be a positive Morse function satisfying (3). Then for every subset*

$$\{x_1, \dots, x_q\} \subseteq \{\nabla K = 0\} \cap \{\Delta K < 0\}$$

and, as  $\tau \rightarrow 0$ , there exists a unique  $u = \alpha^i \varphi_{a_i, \lambda_i} + \bar{v} \in V(q, \varepsilon)$  with

$$\|u\|_{L_{g_0}}^2 = 1, \quad d(a_i, x_i) = o(1) \quad \text{and} \quad \partial J_\tau(u) = 0.$$

*Proof.* Due to (20) we have

$$|\partial J| \leq \frac{\tilde{\varepsilon}}{\lambda^3} \quad \text{on} \quad \bar{V}(q, \varepsilon) \quad \text{and} \quad |\partial J| \geq \frac{\hat{\varepsilon}}{\lambda^3} \quad \text{on} \quad \partial \bar{V}(q, \varepsilon)$$

as long as  $c < \alpha_j < C$ . Thus by (ii) in Lemma 3.1 it is sufficient to look for critical points in the set

$$\tilde{\mathcal{C}} = \{\tilde{u}(\alpha, \lambda, a) = \alpha^i \varphi_i + \bar{v}(\alpha, \lambda, a) \in \bar{V}(q, \varepsilon) \mid \|\tilde{u}\|_{L_{g_0}}^2 = 1\},$$

which is a smooth  $(3(n+2) - 1)$ -dimensional manifold in  $W^{1,2}(M, g_0)$ .

Vice-versa we claim that a critical point of  $J_\tau|_{\tilde{\mathcal{C}}}$  is indeed a critical point of  $J_\tau$ . In fact by Lagrange multiplier rule the gradient of  $J_\tau$  at a constrained critical point  $\tilde{u}_0$  must be orthogonal to  $\tilde{\mathcal{C}}$ . Since  $J_\tau$  is scaling invariant, its gradient on  $\mathcal{C}$  must be tangent to the unit sphere in the  $\|\cdot\|_{L_{g_0}}$  norm. On the other hand, by construction of  $\bar{v}$ , the gradient of  $J_\tau$  at  $\tilde{u}_0$  is tangent to

$$\mathcal{C} = \{\alpha^i \varphi_i \in \bar{V}(q, \varepsilon) \mid \|u\|_{L_{g_0}}^2 = 1\}$$

at the point  $u_0$  such that  $\tilde{u}_0 = u_0 + \bar{v}_0$ . By the estimate on the derivatives of  $\bar{v}$  in Lemma 3.2,  $T_{\tilde{u}_0} \tilde{\mathcal{C}}$  is nearly parallel to  $T_{u_0} \mathcal{C}$ , which implies that  $\partial J_\tau(\tilde{u}_0) = 0$ , as desired.

It remains to prove existence and uniqueness of critical points of  $J_\tau|_{\tilde{\mathcal{C}}}$ . For the existence part we may use the expansions in Lemmas 5.4, 5.5 and 5.6 together with the definition of  $\bar{V}(q, \varepsilon)$  to show, that  $\partial J_\tau$  is non-vanishing on the boundary of  $\tilde{\mathcal{C}}$ . For example, cf. (iii) in the definition of  $\bar{V}(q, \varepsilon)$ , suppose

$$\lambda_j^2 = -c_2 \frac{\Delta K(x_j)}{K(x_j)\tau} + \frac{\varepsilon}{\lambda^2}; \quad \frac{1}{\lambda^2} = \tau.$$

From Lemma 5.5 we deduce, that there exists  $\tilde{\varepsilon} > 0$ , tending to zero as  $\varepsilon \rightarrow 0$ , such that

$$\lambda_j \partial_{\lambda_j} J_\tau(\alpha^i \varphi_i) > \frac{\tilde{\varepsilon}}{\lambda^3}.$$

From Lemmas 3.1 and 3.2 we also have

$$\lambda_j \partial_{\lambda_j} J_\tau(u(\alpha, \lambda, a)) > \frac{1}{2} \frac{\tilde{\varepsilon}}{\lambda^3}$$

and a similar reversed inequality with opposite sign, if

$$\lambda_j^2 = -c_2 \frac{\Delta K(x_j)}{K(x_j)\tau} - \frac{\varepsilon}{\lambda^2}.$$

Analogous estimates are derived for the  $\alpha$ - and  $a$ - derivatives, yielding that the degree of  $\partial J_\tau$  on  $\tilde{\mathcal{C}}$  is well-defined and non-zero. This shows the existence of a critical point for  $J_\tau|_{\tilde{\mathcal{C}}}$ , which is (freely) critical for  $J_\tau$  by the above discussion. Since by construction the negative part of the above solutions is small in  $W^{1,2}$  norm, it is possible to show from Sobolev inequality that it has to vanish identically, so full positivity follows then from the maximum principle.

Uniqueness follows from Lemma 3.2 and Proposition 4.1, implying the strict convexity or concavity of  $J_\tau|_{\tilde{\mathcal{C}}}$  with respect to all parameters  $\alpha$ ,  $\lambda$  and the coordinates of the points  $a_i$ , provided they are chosen so that  $\nabla^2 K(x_i)$  is diagonal.  $\square$

## 4 The second variation

Let  $\bar{V}(q, \varepsilon)$  be the open set defined in (17). The aim of this section is to find there a nearly diagonal form of the second differential of  $J_\tau$ . We recall our notation from Section 2, in particular that of the orthogonal space  $H_u$  in (16).

**Proposition 4.1.** *For  $\alpha^i \varphi_i + \bar{v} \in \bar{V}(q, \varepsilon)$ , consider the decomposition*

$$\begin{aligned} W^{1,2}(M, g_0) &= H_{\alpha^i \varphi_i} \oplus \langle \varphi_i \rangle_{1 \leq i \leq q} \oplus \langle \lambda_i \partial_{\lambda_i} \varphi_i \rangle_{1 \leq i \leq q} \oplus \langle \frac{\nabla_{a_i}}{\lambda_i} \varphi_i \rangle_{1 \leq i \leq q} \\ &=: \mathcal{V} \oplus X_\alpha \oplus X_\lambda \oplus X_a. \end{aligned}$$

Then there exists a basis  $\mathbb{B}$  of  $W^{1,2}(M, g_0)$  with elements in the subspaces of the above decomposition, such that the coefficients of the the second differential of  $J_\tau$  with respect to  $\mathbb{B}$  have the form

$$[\partial^2 J_\tau(\alpha^k \varphi_k + \bar{v})]_{\mathbb{B}} = \frac{1}{\lambda^2} \begin{pmatrix} \mathbb{V}_+ & 0 & 0 & 0 \\ 0 & \mathbb{A}_{q-1,0} & 0 & 0 \\ 0 & 0 & \mathbb{A}_+ & 0 \\ 0 & 0 & 0 & -\frac{\nabla^2 K}{K} \end{pmatrix} + o\left(\frac{1}{\lambda^2}\right), \quad \text{where}$$

- (i)  $\mathbb{A}_+$  represents the coefficients of a symmetric, positive-definite operator on  $\mathcal{V}$  with eigenvalues uniformly bounded away from zero;



- (ii)  $\mathbb{A}_{q-1,0}$  has  $q - 1$  negative eigenvalues uniformly bounded away from zero and one-dimensional kernel;
- (iii)  $\mathbb{A}_+$  is positive-definite with eigenvalues uniformly bounded away from zero;
- (iv)  $-\frac{\nabla^2 K}{K}$  stands for the diagonal matrix  $-(\frac{\nabla^2 K_i}{K_i})_{i=1,\dots,q}$ .

**Remark 4.1.** The basis elements in  $\mathbb{B}$  corresponding to the first two blocks have norms of order  $\frac{1}{\lambda^2}$ , while the ones corresponding to the last two blocks have norm of order 1. We made this choice to guarantee the off-diagonal terms in the above matrix to be of order  $o(\frac{1}{\lambda^2})$ .

*Proof.* We will analyse (10) for  $u = \alpha^i \varphi_i + \bar{v} \in \bar{V}(q, \varepsilon)$ . Recall from Section 2

$$W^{1,2}(M, g_0) = \langle \phi_{k,i} \rangle_{k,i} \oplus H_{\alpha^i \varphi_i},$$

We then choose a  $\langle \cdot, \cdot \rangle^{L_{g_0}}$ -orthonormal basis  $\{\nu_0, \nu_1, \nu_2, \dots\}$  for  $H_{\alpha^i \varphi_i}$  and for some  $\lambda \simeq \lambda_i \simeq \frac{1}{\sqrt{\tau}}$  define

$$\mathbb{B} = \{\tilde{\phi}_{k,i}, \tilde{\nu}_j\} = \left\{ \frac{\varphi_i}{\lambda_i}, \lambda_i \partial_{\lambda_i} \varphi_i, \frac{\nabla_{a_i} \varphi_i}{\lambda_i}, \frac{\nu_j}{\lambda} \right\}; \quad k = 1, 2, 3, \quad i = 1, \dots, q.$$

With this choice it is not hard to see that the coefficients  $[\partial^2 J_\tau(\alpha^k \varphi_k + \bar{v})]_{\mathbb{B}}$  are all of order  $O(\frac{1}{\lambda^2})$ , and our goal is to make their estimates more precise, considering different matrix blocks.

**First block.** The fact that  $\partial^2 J_\tau(\alpha^i \varphi_i)$  is (uniformly) positive-definite on  $H_{\alpha^i \varphi_i}$  is well-known, see e.g. [5]. The positivity of  $\partial^2 J_\tau(\alpha^i \varphi_i + \varepsilon_v)$  on the same subspace follows from the Hölder continuity of the second differential and the fact that  $\|\bar{v}\| = O(\frac{1}{\lambda^2})$ .

**First two blocks.** Testing the second differential with  $\tilde{\nu}_i$  and  $\tilde{\phi}_{1,j} = \frac{\varphi_j}{\lambda}$  we get

$$\partial^2 J_\tau(\alpha^i \varphi_i + \bar{v}) \tilde{\nu}_i \tilde{\phi}_{1,j} = o\left(\frac{1}{\lambda^2}\right) \quad (25)$$

using Lemma 5.1,  $\|\bar{v}\| \lesssim \frac{1}{\lambda^2}$  and the orthogonality

$$\langle \tilde{\nu}_i, \tilde{\phi}_{1,j} \rangle_{L_{g_0}} = 0.$$

Moreover from (10) and  $\|\tilde{\phi}_{1,i}\| = O(\frac{1}{\lambda})$  we find with  $c_0 = \int_{\mathbb{R}^n} \frac{dx}{(1+r^2)^n}$

$$\partial^2 J_\tau(\alpha^k \varphi_k + \bar{v}) \tilde{\phi}_{1,i} \tilde{\phi}_{1,j} = \frac{16n(n-1)c_0^{\frac{2}{n}}}{(n-2)(\alpha_{K,\tau}^{\frac{2n}{n-2}})^{\frac{n-2}{n}} \lambda^2} (-\delta_{k,l} + \frac{\alpha_k \alpha_l}{\alpha^2}) = \mathbb{A}_{i,j} \quad (26)$$

up to an error of order  $o(\frac{1}{\lambda^2})$ . Let us compare the above expression to

$$f(\alpha) = \frac{\alpha^2}{(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}}}; \quad \alpha := \sum_{i=1}^q \alpha_i^2, \quad \alpha_K^{\frac{2n}{n-2}} := \sum_{i=1}^q K_i \alpha_i^{\frac{2n}{n-2}}$$

with first- and second-order derivatives given by

$$\frac{1}{2} \partial_{\alpha_i} f(\alpha) = \frac{\alpha_i}{(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}}} - \frac{\alpha^2 K_i \alpha_i^{\frac{n+2}{n-2}}}{(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}+1}} = \frac{\alpha_i}{(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}}} \left(1 - \frac{\alpha^2}{\alpha_K^{\frac{2n}{n-2}}} K_i \alpha_i^{\frac{4}{n-2}}\right)$$

and

$$\begin{aligned} \frac{1}{2} \partial_{\alpha_i} \partial_{\alpha_j} f(\alpha) &= \delta_{i,j} \frac{1}{(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}}} \left(1 - \frac{n+2}{n-2} \frac{\alpha^2}{\alpha_K^{\frac{2n}{n-2}}} K_i \alpha_i^{\frac{4}{n-2}}\right) \\ &\quad + 2 \frac{\alpha_i \alpha_j}{(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}+1}} \frac{\alpha^2}{\alpha_K^{\frac{2n}{n-2}}} K_i \alpha_i^{\frac{4}{n-2}} K_j \alpha_j^{\frac{4}{n-2}} \\ &\quad - 2 \frac{\alpha_i \alpha_j}{(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}+1}} (K_i \alpha_i^{\frac{4}{n-2}} + K_j \alpha_j^{\frac{4}{n-2}}) \\ &\quad + \frac{2n}{n-2} \frac{\alpha^2}{(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}+2}} K_j \alpha_j^{\frac{n+2}{n-2}} K_i \alpha_i^{\frac{n+2}{n-2}}. \end{aligned}$$

The function  $f$  is scaling invariant and restricted to

$$\{\alpha_K^{\frac{2n}{n-2}} = 1\}$$

attains its maximum at  $(\alpha_i)_i$  satisfying

$$\frac{\alpha^2}{\alpha_K^{\frac{2n}{n-2}}} K_i \alpha_i^{\frac{4}{n-2}} = 1 \quad \text{for all } i = 1, \dots, q,$$

where we have

$$\frac{1}{2} \partial_{\alpha_i} \partial_{\alpha_j} f(\alpha) = \frac{4}{(n-2)(\alpha_K^{\frac{2n}{n-2}})^{\frac{n-2}{n}}} \left(-\delta_{i,j} + \frac{\alpha_i \alpha_j}{\alpha^2}\right). \quad (27)$$

Comparing (26) and (27) we conclude with obvious notation

$$[\partial^2 J_\tau(\alpha^k \varphi_k + \tilde{v})]_{\mathbb{B}} = \begin{pmatrix} \frac{1}{\lambda^2} \mathbb{V}_+ & 0 & \partial^2 J_\tau \tilde{\nu} \tilde{\phi}_2 & \partial^2 J_\tau \tilde{\nu} \tilde{\phi}_3 \\ 0 & \frac{1}{\lambda^2} \mathbb{A}_{q-1,0} & \partial^2 J_\tau \tilde{\phi}_1 \tilde{\phi}_2 & \partial^2 J_\tau \tilde{\phi}_1 \tilde{\phi}_3 \\ \partial^2 J_\tau \tilde{\phi}_2 \tilde{\nu} & \partial^2 J_\tau \tilde{\phi}_2 \tilde{\phi}_1 & \partial^2 J_\tau \tilde{\phi}_2 \tilde{\phi}_2 & \partial^2 J_\tau \tilde{\phi}_2 \tilde{\phi}_3 \\ \partial^2 J_\tau \tilde{\phi}_3 \tilde{\nu} & \partial^2 J_\tau \tilde{\phi}_3 \tilde{\phi}_1 & \partial^2 J_\tau \tilde{\phi}_3, \tilde{\phi}_2 & \partial^2 J_\tau \tilde{\phi}_3 \tilde{\phi}_3 \end{pmatrix}$$

up to some  $o(\frac{1}{\lambda^2})$ .

**Terms off 2x2 blocks.** Let us consider next the interaction of  $\tilde{\nu}_i$  with

$$\tilde{\phi}_{k,j} = \phi_{k,j} \quad \text{for } k = 2, 3.$$

Since

$$\bar{v} = O\left(\frac{1}{\lambda^2}\right), \quad \tilde{\nu}_i = O\left(\frac{1}{\lambda}\right), \quad \langle \varphi_k, \phi_{k,j} \rangle_{L_{g_0}} = O\left(\frac{1}{\lambda^2}\right) \quad \text{and} \quad \langle \nu_i, \phi_{k,j} \rangle_{L_{g_0}} = 0,$$

we simply find for (10)

$$\begin{aligned} \partial^2 J_\tau(\alpha^l \varphi_l + \bar{v}) \tilde{\nu}_i \tilde{\phi}_{j,k} &= \partial^2 J_\tau(\alpha^l \varphi_l) \tilde{\nu}_i \tilde{\phi}_{j,k} \\ &= -\frac{2pr\alpha^l \varphi_l}{k^{\frac{p+1}{2}+1}} \int K(\alpha^l \varphi_l)^{p-1} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} \end{aligned} \quad (28)$$

up to some  $o(\frac{1}{\lambda^2})$ . Indeed by (10) the crucial estimates to verify (28) are

$$\int K(\alpha^l \varphi_l)^p \tilde{\nu}_i d\mu_{g_0} = o\left(\frac{1}{\lambda^2}\right) = \int K(\alpha^l \varphi_l)^p \tilde{\phi}_{k,j} d\mu_{g_0}. \quad (29)$$

These however follow easily by expansion and interaction estimates using

$$\langle \varphi_l, \phi_{k,j} \rangle_{L_{g_0}} = O\left(\frac{1}{\lambda^2}\right), \quad \langle \nu_i, \phi_{k,j} \rangle_{L_{g_0}} = 0, \quad L_{g_0} \varphi_l = 4n(n-1)\varphi_l^{\frac{n+2}{n-2}} + o(1)$$

in  $W^{-1,2}$  and Lemma 5.3. For the remaining integral in (28) we then have

$$\begin{aligned} \int K(\alpha^l \varphi_l)^{p-1} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} &= K_j \int (\alpha^l \varphi_l)^{p-1} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} + o\left(\frac{1}{\lambda^2}\right) \\ &= K_j \int_{\{\varphi_j > \sum_{j \neq l} \alpha^l \varphi_l\}} (\alpha^l \varphi_l)^{p-1} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} \\ &\quad + O\left(\frac{1}{\lambda} \sum_{j \neq l} \|\varphi_l^{p-1} \varphi_j\|_{L^{\frac{p+1}{p}}}\right) + o\left(\frac{1}{\lambda^2}\right) \\ &= K_j \alpha_j^{p-1} \int_{\{\varphi_j > \sum_{j \neq l} \alpha^l \varphi_l\}} \varphi_j^{p-1} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} \\ &\quad + O\left(\frac{1}{\lambda} \sum_{j \neq l} \|\varphi_l^{p-1} \varphi_j + \varphi_l \varphi_j^{p-1}\|_{L^{\frac{p+1}{p}}}\right) + o\left(\frac{1}{\lambda^2}\right) \end{aligned} \quad (30)$$

and therefore using Lemma 5.2 with  $p = \frac{n+2}{n-2} - \tau$

$$\int K(\alpha^l \varphi_l)^{p-1} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} = K_j \alpha_j^{p-1} \int \varphi_j^{p-1} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} + o\left(\frac{1}{\lambda^2}\right).$$

Then, since  $\|\tilde{\nu}_i\| = O(\frac{1}{\lambda})$ ,  $\tau = O(\frac{1}{\lambda^2})$  and  $\varepsilon_{r,s} = O(\frac{1}{\lambda^{n-2}})$ , we find

$$\int K(\alpha^l \varphi_l)^{p-1} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} = K_j \alpha_j^{\frac{4}{n-2}} \int \varphi_j^{\frac{4}{n-2}} \tilde{\nu}_i \tilde{\phi}_{j,k} d\mu_{g_0} + o\left(\frac{1}{\lambda^2}\right) = o\left(\frac{1}{\lambda^2}\right),$$

where the last inequality follows from Lemma 5.1 and  $\langle \phi_{k,j}, \tilde{\nu}_i \rangle_{L_{g_0}} = 0$ . Thus

$$\partial^2 J_\tau(\alpha^l \varphi_l + \bar{v}) \tilde{\nu}_i \tilde{\phi}_{k,j} = o\left(\frac{1}{\lambda^2}\right) \text{ for } k = 2, 3. \quad (31)$$

By exactly the same arguments with  $\tilde{\phi}_{1,i} = O\left(\frac{1}{\lambda}\right)$  as for (29) there holds

$$\partial^2 J_\tau(\alpha^l \varphi_l + \bar{v}) \tilde{\phi}_{1,i} \tilde{\phi}_{k,j} = \partial^2 J_\tau(\alpha^l \varphi_l + \bar{v}) \frac{\phi_{1,i}}{\lambda} \phi_{k,j} = \frac{1}{\lambda} \partial^2 J_\tau(\alpha^l \varphi_l) \varphi_i \phi_{k,j} = o\left(\frac{1}{\lambda^2}\right)$$

for  $k = 2, 3$ . Thus we arrive at

$$[\partial^2 J_\tau(\alpha^l \varphi_l + \bar{v})]_{\mathbb{B}} = \begin{pmatrix} \frac{1}{\lambda^2} \mathbb{V}_+ & 0 & 0 & 0 \\ 0 & \frac{1}{\lambda^2} \mathbb{A}_{q-1,0} & 0 & 0 \\ 0 & 0 & \partial^2 J_\tau \tilde{\phi}_2 \tilde{\phi}_2 & \partial^2 J_\tau \tilde{\phi}_2 \tilde{\phi}_3 \\ 0 & 0 & \partial^2 J_\tau \tilde{\phi}_3 \tilde{\phi}_2 & \partial^2 J_\tau \tilde{\phi}_3 \tilde{\phi}_3 \end{pmatrix} + o\left(\frac{1}{\lambda^2}\right).$$

**Last 2x2 block.** We are left with the estimate of

$$\partial^2 J_\tau(\alpha^k \varphi_k + \bar{v}) \tilde{\phi}_{k,i} \tilde{\phi}_{l,j} = \partial^2 J_\tau(\alpha^k \varphi_k + \bar{v}) \phi_{k,i} \phi_{l,j}$$

for  $k, l = 2, 3$ . Using the fact that

$$\int \phi_{k,i} L_{g_0}(\alpha^m \varphi_m + \bar{v}) d\mu_{g_0} = o\left(\frac{1}{\lambda}\right) = \int \phi_{k,i} K(\alpha^m \varphi_m + \bar{v})^p d\mu_{g_0} \text{ for } k = 2, 3,$$

which follows from  $\|\bar{v}\| = O\left(\frac{1}{\lambda^2}\right)$ , Lemma 5.1 and Lemma 5.2, we find for (10)

$$\begin{aligned} & \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v}) \phi_{k,i} \phi_{l,j} \\ &= \frac{2}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}^{\frac{2}{p+1}}} \int [\phi_{k,i} L_{g_0} \phi_{l,j} - p \frac{r_{\alpha^m \varphi_m + \bar{v}}}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}} K(\alpha^m \varphi_m + \bar{v})^{p-1} \phi_{k,i} \phi_{l,j}] d\mu_{g_0} \\ &=: \frac{2I}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}^{\frac{2}{p+1}}} =: \frac{2(I_1 - I_2)}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}^{\frac{2}{p+1}}} = \frac{2}{(c_0 \alpha_{K,\tau}^{\frac{2n}{n-2}})^{\frac{n-2}{n}}} (I_1 - I_2) + o\left(\frac{1}{\lambda^2}\right). \end{aligned} \quad (32)$$

In the latter formula, recalling (8) and (17), we have used the fact that

$$(k_\tau)_{\alpha^m \varphi_m + \bar{v}}^{\frac{2}{p+1}} = (c_0 \alpha_{K,\tau}^{\frac{2n}{n-2}})^{\frac{n-2}{n}} + o(1)$$

and that both  $I_1, I_2 = O\left(\frac{1}{\lambda^2}\right)$ . Let us first compute  $I_2$ , for which we clearly have

$$\begin{aligned} I_2 &= p \frac{r_{\alpha^m \varphi_m + \bar{v}}}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}} \int K(\alpha^m \varphi_m)^{p-1} \phi_{k,i} \phi_{l,j} d\mu_{g_0} \\ &\quad + p(p-1) \frac{r_{\alpha^m \varphi_m + \bar{v}}}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}} \int K(\alpha^m \varphi_m)^{p-2} \phi_{k,i} \phi_{l,j} \bar{v} d\mu_{g_0} \end{aligned}$$

up to an error  $o(\frac{1}{\lambda^2})$ , as  $\|\bar{v}\| = O(\frac{1}{\lambda^2})$ , and therefore still up to an error  $o(\frac{1}{\lambda^2})$

$$\begin{aligned} I_2 = & p \frac{r_{\alpha^m \varphi_m + \bar{v}}}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}} \int K(\alpha^m \varphi_m)^{p-1} \phi_{k,i} \phi_{l,j} d\mu_{g_0} \\ & + 4n(n-1) \frac{n+2}{n-2} \frac{4}{n-2} \frac{\alpha^2}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} \int K(\alpha^m \varphi_m)^{\frac{6-n}{n-2}} \phi_{k,i} \phi_{l,j} \bar{v} d\mu_{g_0}. \end{aligned}$$

As due to  $d(a_i, a_j) \simeq 1$  for  $i \neq j$  the interactions terms in (14) are of order

$$\varepsilon_{i,j} = O\left(\frac{1}{\lambda^{n-2}}\right) = o\left(\frac{1}{\lambda^2}\right),$$

we find

$$\begin{aligned} I_2 = & p \frac{r_{\alpha^m \varphi_m + \bar{v}}}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}} \delta_{i,j} \alpha_i^{p-1} \int K \varphi_i^{p-1} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \\ & + 4n(n-1) \frac{n+2}{n-2} \frac{4}{n-2} \frac{\alpha^2}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} \delta_{i,j} \alpha_i^{\frac{6-n}{n-2}} \int K \varphi_i^{\frac{6-n}{n-2}} \phi_{k,i} \phi_{l,i} \bar{v} d\mu_{g_0} \end{aligned}$$

up to an error  $o(\frac{1}{\lambda^2})$ . Up to the same error we may simplify this using(17) to

$$\begin{aligned} I_2 = & p \frac{r_{\alpha^m \varphi_m + \bar{v}}}{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}} \delta_{i,j} K_i \alpha_i^{p-1} \int \varphi_i^{p-1} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \\ & + 4n(n-1) \frac{n+2}{n-2} \delta_{i,j} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \\ & + 4n(n-1) \frac{n+2}{n-2} \frac{4}{n-2} \delta_{i,j} \alpha_i^{-1} \int \varphi_i^{\frac{6-n}{n-2}} \phi_{k,i} \phi_{l,i} \bar{v} d\mu_{g_0} \end{aligned}$$

for some  $\varepsilon > 0$  small and fixed. Moreover by orthogonality and (47)

$$\frac{r_{\alpha^i \varphi_i + \bar{v}}}{(k_\tau)_{\alpha^i \varphi_i + \bar{v}}} = \frac{r_{\alpha^i \varphi_i}}{(k_\tau)_{\alpha^i \varphi_i}} = 4n(n-1) \frac{\alpha^2}{\alpha_{K,\theta}^{p+1}} \left(1 - \left(\frac{\bar{c}_1}{\bar{c}_0} - \frac{\bar{c}_1 \bar{c}_2}{\bar{c}_2 \bar{c}_0}\right) \tau\right) + o\left(\frac{1}{\lambda^2}\right),$$

whence by(17) and the fact that  $p = \frac{n+2}{n-2} - \tau$  we arrive at

$$\begin{aligned} I_2 = & 4n(n-1) \frac{n+2}{n-2} \left[ \left(1 - \left(\frac{n-2}{n+2} + \frac{\bar{c}_1}{\bar{c}_0} - \frac{\bar{c}_1 \bar{c}_2}{\bar{c}_2 \bar{c}_0}\right) \tau\right) \lambda_i^\theta \delta_{i,j} \int \varphi_i^{p-1} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \right. \\ & + 4n(n-1) \frac{n+2}{n-2} \delta_{i,j} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \\ & \left. + 4n(n-1) \frac{n+2}{n-2} \frac{4}{n-2} \delta_{i,j} \alpha_i^{-1} \int \varphi_i^{\frac{6-n}{n-2}} \phi_{k,i} \phi_{l,i} \bar{v} d\mu_{g_0} \right]. \end{aligned}$$

Let us compute the last integral above, which is of order  $O(\frac{1}{\lambda^2})$  as  $\|\bar{v}\|$ . Clearly

$$\begin{aligned} \frac{4}{n-2} \int \varphi_i^{\frac{6-n}{n-2}} \phi_{k,i} \phi_{l,i} \bar{v} d\mu_{g_0} & = \int d_{k,i} \varphi_i^{\frac{4}{n-2}} \phi_{l,i} \bar{v} d\mu_{g_0} \\ & = d_{k,i} \int \varphi_i^{\frac{4}{n-2}} \phi_{l,i} \bar{v} d\mu_{g_0} - \int \varphi_i^{\frac{4}{n-2}} d_{k,i} \phi_{l,i} \bar{v} d\mu_{g_0} - \int \varphi_i^{\frac{4}{n-2}} \phi_{l,i} d_{k,i} \bar{v} d\mu_{g_0}. \end{aligned}$$

Due to orthogonality the first integral above is of order  $o(\frac{1}{\lambda^2})$  and denoting by

$$\widehat{w} = \Pi_{\langle \phi_{k,i} \rangle^{\perp L_{g_0}}} w \text{ for } w \in W^{1,2}(M, g_0) \quad (33)$$

the orthogonal projection onto  $\langle \phi_{k,i} \rangle^{\perp L_{g_0}}$  we have up to an error  $o(\frac{1}{\lambda^2})$

$$\int \varphi_i^{\frac{4}{n-2}} d_{k,i} \phi_{l,i} \bar{v} d\mu_{g_0} = \int \varphi_i^{\frac{4}{n-2}} \widehat{d_{k,i} \phi_{l,i} \bar{v}} d\mu_{g_0}$$

due to the orthogonalities  $\langle \bar{v}, \phi_{k,i} \rangle_{L_{g_0}} = 0$  and  $\|\bar{v}\| = O(\frac{1}{\lambda^2})$ . Hence, using the notation in (33), we arrive at

$$\begin{aligned} I_2 &= 4n(n-1) \frac{n+2}{n-2} \left[ \left( 1 - \left( \frac{n-2}{n+2} + \frac{\bar{c}_1}{\bar{c}_0} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 \bar{c}_0} \right) \tau \right) \lambda_i^\theta \delta_{i,j} \int \varphi_i^{p-1} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \right. \\ &\quad + 4n(n-1) \frac{n+2}{n-2} \delta_{i,j} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \\ &\quad \left. - 4n(n-1) \frac{n+2}{n-2} \delta_{i,j} \alpha_i^{-1} \left( \int \varphi_i^{\frac{4}{n-2}} \widehat{d_{k,i} \phi_{l,i} \bar{v}} d\mu_{g_0} + \int \varphi_i^{\frac{4}{n-2}} \phi_{l,i} d_{k,i} \bar{v} d\mu_{g_0} \right) \right]. \end{aligned}$$

Due to  $\|\bar{v}\| = O(\frac{1}{\lambda^2})$  we havestill up to a  $o(\frac{1}{\lambda^2})$

$$\partial^2 J_\tau(\alpha^m \varphi_m) \bar{v} = \frac{8n(n-1)}{(\bar{c}_0 \alpha_{K,\tau}^{p+1})^{\frac{n-2}{n}}} \left( \frac{L_{g_0}}{4n(n-1)} \bar{v} - \frac{n+2}{n-2} \sum_m \varphi_m^{\frac{4}{n-2}} \bar{v} \right)$$

and we recall from (22) that

$$\partial^2 J_\tau(\alpha^m \varphi_m) \bar{v} = -\partial J_\tau(\alpha^m \varphi_m) + o\left(\frac{1}{\lambda^2}\right) \text{ on } \langle \phi_{l,j} \rangle^{\perp L_{g_0}}.$$

From this we deduce again by smallness of interactions terms  $\varepsilon_{i,j}$

$$\frac{n+2}{n-2} \int \varphi_i^{\frac{4}{n-2}} \widehat{d_{k,i} \phi_{l,i} \bar{v}} d\mu_{g_0} = \frac{(\bar{c}_0 \alpha_{K,\tau}^{p+1})^{\frac{n-2}{n}}}{8n(n-1)} \partial J_\tau(\alpha^m \varphi_m) \widehat{d_{k,i} \phi_{l,i}} + \frac{\langle \bar{v}, \widehat{d_{k,i} \phi_{l,i}} \rangle_{L_{g_0}}}{4n(n-1)}$$

and by orthogonality and Lemma 5.1 there holds up to an error  $o(\frac{1}{\lambda^2})$

$$\begin{aligned} \langle \bar{v}, \widehat{d_{k,i} \phi_{l,i}} \rangle_{L_{g_0}} &= -\langle d_{k,i} \bar{v}, \phi_{l,i} \rangle_{L_{g_0}} = -4n(n-1) \int \bar{d}_{k,i} v d_{l,i} \varphi_i^{\frac{n+2}{n-2}} d\mu_{g_0} \\ &= -4n(n-1) \frac{n+2}{n-2} \int \bar{\varphi}_i^{\frac{4}{n-2}} d_{k,i} v \phi_{l,i} d\mu_{g_0}. \end{aligned}$$

We therefore conclude that up to an error  $o(\frac{1}{\lambda^2})$

$$\begin{aligned} I_2 &= 4n(n-1) \frac{n+2}{n-2} \left[ \left( 1 - \left( \frac{n-2}{n+2} + \frac{\bar{c}_1}{\bar{c}_0} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 \bar{c}_0} \right) \tau \right) \lambda_i^\theta \delta_{i,j} \int \varphi_i^{p-1} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \right. \\ &\quad + 4n(n-1) \frac{n+2}{n-2} \delta_{i,j} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{l,i} d\mu_{g_0} \\ &\quad \left. - 4n(n-1) \delta_{i,j} \alpha_i^{-1} \frac{(\bar{c}_0 \alpha_{K,\tau}^{p+1})^{\frac{n-2}{n}}}{8n(n-1)} \partial J_\tau(\alpha^m \varphi_m) \widehat{d_{k,i} \phi_{l,i}}, \right. \end{aligned}$$

at which point  $\bar{v}$  has been eliminated from the main terms in the expansion. By Lemma 3.1 we then have

$$\partial J_\tau(\alpha^m \varphi_m)|_{\langle \phi_{k,i} \rangle} = o\left(\frac{1}{\lambda^2}\right),$$

so we may pass from  $\widehat{d_{k,i}\phi_{l,i}}$  to  $d_{k,i}\phi_{l,i}$  in the above formulae and, as

$$\partial J_\tau(\alpha^m \varphi_m) = O\left(\frac{1}{\lambda^2}\right),$$

we obtain

$$\begin{aligned} & \frac{(\bar{c}_0 \alpha_{K,\tau}^{p+1})^{\frac{n-2}{n}}}{8n(n-1)} \partial J_\tau(\alpha^m \varphi_m) d_{k,i}\phi_{l,i} \\ &= -\alpha^m \tau \int_{B_\varepsilon(a_m)} \left( \varphi_m^{\frac{n+2}{n-2}} \ln(1 + \lambda_m^2 r^2)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1} \varphi_m^{\frac{n+2}{n-2}} \right. \\ & \quad \left. + \frac{2}{n-2} \frac{\tilde{c}_1}{c_2} \varphi_m^{\frac{4}{n-2}} \lambda_m \partial_{\lambda_m} \varphi_m \right) d_{k,i}\phi_{l,i} d\mu_{g_0} \\ &+ \alpha^m \tau \int_{B_\varepsilon(a_m)} \left( \frac{\tilde{c}_1}{\tilde{c}_2} \frac{\lambda_m^2 r^2}{2n} \varphi_m^{\frac{n+2}{n-2}} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 c_1} \varphi_m^{\frac{n+2}{n-2}} \right. \\ & \quad \left. + \frac{2}{n-2} \frac{\tilde{c}_1}{c_2} \varphi_m^{\frac{4}{n-2}} \lambda_m \partial_{\lambda_m} \varphi_m \right) d_{k,i}\phi_{l,i} d\mu_{g_0} \\ &- \alpha^m \int_{B_\varepsilon(a_m)} \left( \frac{\nabla^2 K_m}{2K_m} x^2 - \frac{\Delta K_m}{2nK_m} r^2 \right) \varphi_m^{\frac{n+2}{n-2}} d_{k,i}\phi_{l,i} d\mu_{g_0}. \end{aligned}$$

Still  $\varepsilon_{i,j} = o(\frac{1}{\lambda^2})$  we therefore arrive at

$$\begin{aligned}
I_2 = & 4n(n-1)\frac{n+2}{n-2}\left[1 - \left(\frac{n-2}{n+2} + \frac{\bar{c}_1}{\bar{c}_0} - \frac{\tilde{c}_1\bar{c}_2}{\bar{c}_2\bar{c}_0}\right)\tau\right]\lambda_i^\theta\delta_{i,j}\int\varphi_i^{p-1}\phi_{k,i}\phi_{l,i}d\mu_{g_0} \\
& + 4n(n-1)\frac{n+2}{n-2}\delta_{i,j}\int_{B_\varepsilon(a_i)}\frac{\nabla^2K_i}{2K_i}x^2\varphi_i^{\frac{4}{n-2}}\phi_{k,i}\phi_{l,i}d\mu_{g_0} \\
& - 4n(n-1)\delta_{i,j}\left(-\tau\int_{B_\varepsilon(a_i)}\left(\varphi_i^{\frac{n+2}{n-2}}\ln(1+\lambda_i^2r^2)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1}\varphi_i^{\frac{n+2}{n-2}}\right.\right. \\
& \qquad \qquad \qquad \left.\left. + \frac{2}{n-2}\frac{\tilde{c}_1}{c_2}\varphi_i^{\frac{4}{n-2}}\lambda_i\partial_{\lambda_i}\varphi_i\right)d_{k,i}\phi_{l,i}d\mu_{g_0}\right. \\
& \qquad \qquad \qquad \left. + \tau\int_{B_\varepsilon(a_i)}\left(\frac{\tilde{c}_1}{\tilde{c}_2}\frac{\lambda_i^2r^2}{2n}\varphi_i^{\frac{n+2}{n-2}} - \frac{\tilde{c}_1\bar{c}_2}{\bar{c}_2c_1}\varphi_i^{\frac{n+2}{n-2}}\right.\right. \\
& \qquad \qquad \qquad \left.\left. + \frac{2}{n-2}\frac{\tilde{c}_1}{c_2}\varphi_i^{\frac{4}{n-2}}\lambda_i\partial_{\lambda_i}\varphi_i\right)d_{k,i}\phi_{l,i}d\mu_{g_0}\right. \\
& \qquad \qquad \qquad \left. - \int_{B_\varepsilon(a_i)}\left(\frac{\nabla^2K_i}{2K_i}x^2 - \frac{\Delta K_i}{2nK_i}r^2\right)\varphi_i^{\frac{n+2}{n-2}}d_{k,i}\phi_{l,i}d\mu_{g_0}\right),
\end{aligned}$$

up to some  $o(\frac{1}{\lambda^2})$ . By oddness we may simplify this to

$$\begin{aligned}
I_2 = & 4n(n-1)\frac{n+2}{n-2}\left[1 - \left(\frac{n-2}{n+2} + \frac{\bar{c}_1}{\bar{c}_0} - \frac{\tilde{c}_1\bar{c}_2}{\bar{c}_2\bar{c}_0}\right)\tau\right] \\
& \lambda_i^\theta\delta_{i,j}\delta_{k,l}\int\varphi_i^{p-1}\phi_{k,i}\phi_{k,i}d\mu_{g_0} \\
& + 4n(n-1)\frac{n+2}{n-2}\delta_{i,j}\delta_{k,l}\int_{B_\varepsilon(a_i)}\frac{\nabla^2K_i}{2K_i}x^2\varphi_i^{\frac{4}{n-2}}\phi_{k,i}\phi_{k,i}d\mu_{g_0} \\
& - 4n(n-1)\delta_{i,j}\delta_{k,l}\left(-\tau\int_{B_\varepsilon(a_i)}\left(\varphi_i^{\frac{n+2}{n-2}}\ln(1+\lambda_i^2r^2)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1}\varphi_i^{\frac{n+2}{n-2}}\right.\right. \\
& \qquad \qquad \qquad \left.\left. + \frac{2}{n-2}\frac{\tilde{c}_1}{c_2}\varphi_i^{\frac{4}{n-2}}\lambda_i\partial_{\lambda_i}\varphi_i\right)d_{k,i}\phi_{k,i}d\mu_{g_0}\right. \\
& \qquad \qquad \qquad \left. + \tau\int_{B_\varepsilon(a_i)}\left(\frac{\tilde{c}_1}{\tilde{c}_2}\frac{\lambda_i^2r^2}{2n}\varphi_i^{\frac{n+2}{n-2}} - \frac{\tilde{c}_1\bar{c}_2}{\bar{c}_2c_1}\varphi_i^{\frac{n+2}{n-2}}\right.\right. \\
& \qquad \qquad \qquad \left.\left. + \frac{2}{n-2}\frac{\tilde{c}_1}{c_2}\varphi_i^{\frac{4}{n-2}}\lambda_i\partial_{\lambda_i}\varphi_i\right)d_{k,i}\phi_{k,i}d\mu_{g_0}\right. \\
& \qquad \qquad \qquad \left. - \int_{B_\varepsilon(a_i)}\left(\frac{\nabla^2K_i}{2K_i}x^2 - \frac{\Delta K_i}{2nK_i}r^2\right)\varphi_i^{\frac{n+2}{n-2}}d_{k,i}\phi_{k,i}d\mu_{g_0}\right)
\end{aligned}$$



By Lemma 5.1it follows that for  $k = 2, 3$  and up to some  $o(\frac{1}{\lambda^2})$

$$\begin{aligned}
4n(n-1)\frac{n+2}{n-2} \int \varphi_i^{\frac{4}{n-2}} \lambda_i \partial_{\lambda_i} \varphi_i d_{k,i} \phi_{k,i} d\mu_{g_0} &= \int L_{g_0}(\lambda_i \partial_{\lambda_i} \varphi_i) d_{k,i} \phi_{k,i} d\mu_{g_0} \\
&= \langle \lambda_i \partial_{\lambda_i} \varphi_i, (d_{k,i})^2 \varphi_i \rangle_{L_{g_0}} = d_{k,i} \langle \lambda_i \partial_{\lambda_i} \varphi_i, d_{k,i} \varphi_i \rangle_{L_{g_0}} - \langle \lambda_i \partial_{\lambda_i} d_{k,i} \varphi_i, d_{k,i} \varphi_i \rangle_{L_{g_0}} \\
&= d_{k,i} \langle \phi_{2,i}, \phi_{k,i} \rangle_{L_{g_0}} - \frac{1}{2} \lambda_i \partial_{\lambda_i} \|\phi_{k,i}^2\|_{L_{g_0}} = o(1),
\end{aligned}$$

as  $\langle \phi_{2,i}, \phi_{k,i} \rangle_{L_{g_0}}$  and  $\|\phi_{k,i}^2\|_{L_{g_0}}$  are almost constant in  $a_i$  and  $\lambda_i$ . Hence

$$\begin{aligned}
\frac{I_2}{4n(n-1)} &= \frac{n+2}{n-2} \left[ \left( 1 - \left( \frac{n-2}{n+2} + \frac{\bar{c}_1}{\bar{c}_0} - \frac{\bar{c}_1 \bar{c}_2}{\bar{c}_2 \bar{c}_0} \right) \tau \right) \lambda_i^\theta \delta_{i,j} \delta_{k,l} \int \varphi_i^{p-1} \phi_{k,i} \phi_{k,i} d\mu_{g_0} \right. \\
&\quad + \frac{n+2}{n-2} \delta_{i,j} \delta_{k,l} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{k,i} d\mu_{g_0} \\
&\quad - \delta_{i,j} \delta_{k,l} \left( -\tau \int_{B_\varepsilon(a_i)} \left( \ln(1 + \lambda_i^2 r^2) \right)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1} \right) \varphi_i^{\frac{n+2}{n-2}} d_{k,i} \phi_{k,i} d\mu_{g_0} \\
&\quad + \tau \int_{B_\varepsilon(a_i)} \left( \frac{\bar{c}_1 \lambda_i^2 r^2}{\bar{c}_2 2n} - \frac{\bar{c}_1 \bar{c}_2}{\bar{c}_2 c_1} \right) \varphi_i^{\frac{n+2}{n-2}} d_{k,i} \phi_{k,i} d\mu_{g_0} \\
&\quad \left. - \int_{B_\varepsilon(a_i)} \left( \frac{\nabla^2 K_i}{2K_i} x^2 - \frac{\Delta K_i}{2nK_i} r^2 \right) \varphi_i^{\frac{n+2}{n-2}} d_{k,i} \phi_{k,i} d\mu_{g_0} \right).
\end{aligned}$$

Next for the first summand above we find that up to an error  $o(\frac{1}{\lambda^2})$

$$\begin{aligned}
&\lambda_i^\theta \int \varphi_i^{p-1} \phi_{k,i} \phi_{k,i} d\mu_{g_0} \\
&= \int \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{k,i} d\mu_{g_0} + \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} (\lambda_i^\theta \varphi_i^{-\tau} - 1) \phi_{k,i} \phi_{k,i} d\mu_{g_0} \\
&= \frac{n-2}{n+2} \int d_{k,i} \varphi_i^{\frac{n+2}{n-2}} \phi_{k,i} d\mu_{g_0} + \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} ((1 + \lambda_i^2 r^2)^\theta - 1) \phi_{k,i} \phi_{k,i} d\mu_{g_0} \\
&= \frac{1}{4n(n-1)} \frac{n-2}{n+2} \langle \phi_{k,i}, \phi_{k,i} \rangle_{L_{g_0}} + \theta \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2) \phi_{k,i} \phi_{k,i} d\mu_{g_0}
\end{aligned}$$

using Lemma 5.1 and properly expanding. Recalling (32) we thus conclude

$$\begin{aligned}
& \frac{(k_\tau)^{\frac{2}{p+1}}}{8n(n-1)} \alpha^m \varphi_{m+\bar{v}} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v}) \phi_{k,i} \phi_{l,j} \\
&= \int \frac{L_{g_0}}{4n(n-1)} \phi_{k,i} \phi_{l,j} d\mu_{g_0} - \frac{I_2}{4n(n-1)} \\
&= \delta_{i,j} \delta_{k,l} \left( \left( 1 + \frac{n+2}{n-2} \left( \frac{\bar{c}_1}{\bar{c}_0} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 \bar{c}_0} \right) \right) \tau \int \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{k,i} d\mu_{g_0} \right. \\
&\quad - \frac{n+2}{n-2} \tau \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \phi_{k,i} \phi_{k,i} d\mu_{g_0} \\
&\quad - \tau \int_{B_\varepsilon(a_i)} \left( \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1} \right) \varphi_i^{\frac{n+2}{n-2}} d_{k,i} \phi_{k,i} d\mu_{g_0} \quad (34) \\
&\quad + \tau \int_{B_\varepsilon(a_i)} \left( \frac{\tilde{c}_1 \lambda_i^2 r^2}{\tilde{c}_2} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 c_1} \right) \varphi_i^{\frac{n+2}{n-2}} d_{k,i} \phi_{k,i} d\mu_{g_0} \\
&\quad - \int_{B_\varepsilon(a_i)} \left( \frac{\nabla^2 K_i}{2K_i} x^2 - \frac{\Delta K_i}{2nK_i} r^2 \right) \varphi_i^{\frac{n+2}{n-2}} d_{k,i} \phi_{k,i} d\mu_{g_0} \\
&\quad \left. - \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{k,i} d\mu_{g_0} \right)
\end{aligned}$$

and in particular for  $i = 1, \dots, q$ , and  $k, l = 1, \dots, n$  we have up some  $o(\frac{1}{\lambda^2})$

$$[\partial^2 J_\tau(\alpha^k \varphi_k + \bar{v})]_{\mathbb{B}} = \begin{pmatrix} \frac{1}{\lambda^2} \mathbb{V}_+ & 0 & 0 & 0 \\ 0 & \frac{1}{\lambda^2} \mathbb{A}_{q-1,0} & 0 & 0 \\ 0 & 0 & \partial^2 J_\tau \lambda_i \partial_{\lambda_i} \varphi_i \lambda_i \partial_{\lambda_i} \varphi_i & 0 \\ 0 & 0 & 0 & \partial^2 J_\tau \frac{(\nabla_{a_i})_k}{\lambda_i} \varphi_i \frac{(\nabla_{a_i})_l}{\lambda_i} \varphi_i \end{pmatrix}.$$

**Last diagonal terms.** Concerning  $\lambda$ -derivatives we note, that mixed derivatives in different  $\lambda_i$  are of order  $\lambda^{2-n} = o(\lambda^{-2})$ , since  $n \geq 5$ . Therefore it is sufficient to compute second derivatives with respect to the same  $\lambda_i$ . This

corresponds to

$$\begin{aligned}
& \frac{(k\tau)^{\frac{2}{p+1}}}{8n(n-1)} \alpha^{m\varphi_m + \bar{v}} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v}) (\lambda_i \partial_{\lambda_i} \varphi_i)^2 \\
&= \left(1 + \frac{n+2}{n-2} \left(\frac{\bar{c}_1}{\bar{c}_0} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 \bar{c}_0}\right)\right) \tau \int \varphi_i^{\frac{4}{n-2}} \phi_{k,i} \phi_{k,i} d\mu_{g_0} \\
&\quad - \frac{n+2}{n-2} \tau \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0} \\
&\quad - \tau \int_{B_\varepsilon(a_i)} \left(\ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1}\right) \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0} \\
&\quad + \tau \int_{B_\varepsilon(a_i)} \left(\frac{\tilde{c}_1 \lambda_i^2 r^2}{\tilde{c}_2 2n} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 c_1}\right) \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0} \\
&\quad - \int_{B_\varepsilon(a_i)} \left(\frac{\nabla^2 K_i}{2K_i} x^2 - \frac{\Delta K_i}{2nK_i} r^2\right) \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0} \\
&\quad - \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0}.
\end{aligned}$$

The second-last summand vanishes and

$$\int \varphi_i^{p-1} \phi_{k,i} \phi_{k,i} d\mu_{g_0} = c_k + o(1),$$

cf. Lemma 5.2, whence

$$\begin{aligned}
& \frac{(k\tau)^{\frac{2}{p+1}}}{8n(n-1)} \alpha^{m\varphi_m + \bar{v}} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v}) (\lambda_i \partial_{\lambda_i} \varphi_i)^2 = c_2 \left(1 + \frac{n+2}{n-2} \left(\frac{\bar{c}_1}{\bar{c}_0} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 \bar{c}_0}\right)\right) \tau \\
&\quad - \frac{n+2}{n-2} \tau \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0} \\
&\quad - \tau \int_{B_\varepsilon(a_i)} \left(\ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1}\right) \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0} \\
&\quad + \tau \int_{B_\varepsilon(a_i)} \left(\frac{\tilde{c}_1 \lambda_i^2 r^2}{\tilde{c}_2 2n} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 c_1}\right) \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0} \\
&\quad - \frac{n+2}{n-2} \frac{\Delta K_i}{2nK_i} \int_{B_\varepsilon(a_i)} r^2 \varphi_i^{\frac{4}{n-2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0}.
\end{aligned}$$

Moreover

$$\begin{aligned} \int \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0} &= \lambda_i \partial_{\lambda_i} \int \varphi_i^{\frac{n+2}{n-2}} \lambda_i \partial_{\lambda_i} \varphi_i d\mu_{g_0} \\ &- \frac{n+2}{n-2} \int \varphi_i^{\frac{4}{n-2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0} = -\frac{n+2}{n-2} c_2 + o(1), \end{aligned}$$

and

$$\begin{aligned} \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} r^2 \varphi_i^{\frac{4}{n-2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0} &= \int_{B_\varepsilon(a_i)} r^2 \lambda_i \partial_{\lambda_i} \varphi_i \lambda_i \partial_{\lambda_i} \varphi_i^{\frac{n+2}{n-2}} d\mu_{g_0} \\ &= \lambda_i \partial_{\lambda_i} \int_{B_\varepsilon(a_i)} r^2 \lambda_i \partial_{\lambda_i} \varphi_i \varphi_i^{\frac{n+2}{n-2}} d\mu_{g_0} - \int_{B_\varepsilon(a_i)} r^2 \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0}. \end{aligned}$$

Thus recalling(17), in particular  $\tilde{c}_1 \tau + \tilde{c}_2 \frac{\Delta K_i}{K_i \lambda_i^2} = o(\frac{1}{\lambda_i^2})$ , we arrive at

$$\begin{aligned} \frac{(k_\tau)^{\frac{2}{p+1}}}{8n(n-1)} \alpha^m \varphi_m + \bar{v} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v})(\lambda_i \partial_{\lambda_i} \varphi_i)^2 &= c_2 \tau \\ &- \frac{n+2}{n-2} \tau \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0} \\ &- \tau \int_{B_\varepsilon(a_i)} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0} \\ &+ \frac{\tilde{c}_1}{\tilde{c}_2} \frac{\tau}{2n} \lambda_i^3 \partial_{\lambda_i} \int_{B_\varepsilon(a_i)} r^2 \lambda_i \partial_{\lambda_i} \varphi_i \varphi_i^{\frac{n+2}{n-2}} d\mu_{g_0}, \end{aligned}$$

and for the last integral above we find passing to integration over  $\mathbb{R}^n$

$$\begin{aligned} \lambda_i \partial_{\lambda_i} \int_{B_\varepsilon(a_i)} r^2 \lambda_i \partial_{\lambda_i} \varphi_i \varphi_i^{\frac{n+2}{n-2}} d\mu_{g_0} &= \lambda_i \partial_{\lambda_i} \int_{\mathbb{R}^n} r^2 \lambda_i \partial_{\lambda_i} \delta_{0, \lambda_i} \delta_{0, \lambda_i}^{\frac{n+2}{n-2}} dx \\ &= \frac{n-2}{2n} (\lambda_i \partial_{\lambda_i})^2 \int r^2 \delta_{0, \lambda_i}^{\frac{2n}{n-2}} dx = \frac{n-2}{2n} (\lambda_i \partial_{\lambda_i})^2 (\lambda_i^{-2} \int_{\mathbb{R}^n} \frac{r^2}{(1+r^2)^n} dx) \\ &= \frac{n-2}{8n} \frac{\tilde{c}_2}{\lambda_i^2} \end{aligned}$$

up to some error of order  $o(1)$ . Consequently

$$\begin{aligned} \frac{(k_\tau)^{\frac{2}{p+1}}}{8n(n-1)} \alpha^m \varphi_m + \bar{v} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v})(\lambda_i \partial_{\lambda_i} \varphi_i)^2 &= c_2 \left(1 + \frac{n-2}{16n^2} \frac{\tilde{c}_1 \tilde{c}_2}{\tilde{c}_2 c_2}\right) \tau \\ &- \frac{n+2}{n-2} \tau \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0} \\ &- \tau \int_{B_\varepsilon(a_i)} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \varphi_i^{\frac{n+2}{n-2}} (\lambda_i \partial_{\lambda_i})^2 \varphi_i d\mu_{g_0}. \end{aligned}$$

Finally we calculate passing to integration over  $\mathbb{R}^n$  and up to a  $o(1)$

$$\begin{aligned}
& \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} |\lambda_i \partial_{\lambda_i} \varphi_i|^2 d\mu_{g_0} \\
&= \int_{\mathbb{R}^n} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \lambda_i \partial_{\lambda_i} \delta_{0, \lambda_i} \lambda_i \partial_{\lambda_i} \delta_{0, \lambda_i}^{\frac{n+2}{n-2}} dx \\
&= \lambda_i \partial_{\lambda_i} \int_{\mathbb{R}^n} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \lambda_i \partial_{\lambda_i} \delta_{0, \lambda_i} \delta_{0, \lambda_i}^{\frac{n+2}{n-2}} dx \\
&\quad - (n-2) \int_{\mathbb{R}^n} \frac{\lambda_i^2 r^2}{1 + \lambda_i^2 r^2} \lambda_i \partial_{\lambda_i} \delta_{0, \lambda_i} \delta_{0, \lambda_i}^{\frac{n+2}{n-2}} dx \\
&\quad - \int_{\mathbb{R}^n} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} (\lambda_i \partial_{\lambda_i})^2 \delta_{0, \lambda_i} \delta_{0, \lambda_i}^{\frac{n+2}{n-2}} dx,
\end{aligned}$$

where the first summand above vanishes by rescaling, and we are reduced to

$$\begin{aligned}
& \frac{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}^{\frac{2}{p+1}}}{8n(n-1)} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v})(\lambda_i \partial_{\lambda_i} \varphi_i)^2 \\
&= c_2 \left(1 + \frac{n-2}{16n^2} \frac{\tilde{c}_1 \tilde{c}_2}{\tilde{c}_2 c_2}\right) \tau + (n-2) \tau \int_{\mathbb{R}^n} \frac{\lambda_i^2 r^2}{1 + \lambda_i^2 r^2} \lambda_i \partial_{\lambda_i} \delta_{0, \lambda_i} \delta_{0, \lambda_i}^{\frac{n+2}{n-2}} dx,
\end{aligned}$$

where up to some  $o(1)$  and with  $\hat{c}_3 = -\int_{\mathbb{R}^n} \frac{r^2(1-r^2)}{(1+r^2)^{n+2}} dx$

$$\int_{\mathbb{R}^n} \frac{\lambda_i^2 r^2}{1 + \lambda_i^2 r^2} \lambda_i \partial_{\lambda_i} \delta_{0, \lambda_i} \delta_{0, \lambda_i}^{\frac{n+2}{n-2}} dx = \frac{n-2}{2} \int_{\mathbb{R}^n} \frac{r^2(1-r^2)}{(1+r^2)^{n+2}} dx = -\frac{n-2}{2} \hat{c}_3. \quad (35)$$

By an explicit computation of the above constants we conclude that

$$\begin{aligned}
& \frac{(k_\tau)_{\alpha^m \varphi_m + \bar{v}}^{\frac{2}{p+1}}}{8n(n-1)} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v})(\lambda_i \partial_{\lambda_i} \varphi_i)^2 \\
&= \left( c_2 \left(1 + \frac{n-2}{16n^2} \frac{\tilde{c}_1 \tilde{c}_2}{\tilde{c}_2 c_2}\right) - \frac{(n-2)^2}{2} \hat{c}_3 \right) \tau = \frac{(n-2)^2 \Gamma^2\left(\frac{n}{2}\right)}{128n \Gamma(n+1)} \tau
\end{aligned}$$

up to an error  $o\left(\frac{1}{\lambda^2}\right)$ . Thence with  $i = 1, \dots, q$  and  $k, l = 1, \dots, n$  we derive

$$[\partial^2 J_\tau(\alpha^k \varphi_k + \bar{v})]_{\mathbb{B}} = \begin{pmatrix} \frac{1}{\lambda^2} \mathbb{V}_+ & 0 & 0 & 0 \\ 0 & \frac{1}{\lambda^2} \mathbb{A}_{q-1,0} & 0 & 0 \\ 0 & 0 & \frac{1}{\lambda^2} \mathbb{A}_+ & 0 \\ 0 & 0 & 0 & \partial^2 J_\tau \frac{(\nabla_{a_i})_k}{\lambda_i} \varphi_i \frac{(\nabla_{a_i})_l}{\lambda_i} \varphi_i \end{pmatrix}$$

up to  $o(\frac{1}{\lambda^2})$ , where  $\mathbb{A}_+ > 0$  is as in the statement. For instance consider

$$\begin{aligned}
& \frac{(k\tau)^{\frac{2}{p+1}}}{8n(n-1)} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v}) \left( \frac{(\nabla_{a_i})_1}{\lambda_i} \varphi_i \right)^2 \\
&= \left( 1 + \frac{n+2}{n-2} \left( \frac{\bar{c}_1}{\bar{c}_0} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 \bar{c}_0} \right) \right) \tau \int \varphi_i^{\frac{4}{n-2}} \left| \frac{(\nabla_{a_i})_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0} \\
&\quad - \frac{n+2}{n-2} \tau \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \left| \frac{(\nabla_{a_i})_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0} \\
&\quad - \tau \int_{B_\varepsilon(a_i)} \left( \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} - \frac{\bar{c}_1}{c_1} \right) \varphi_i^{\frac{n+2}{n-2}} \left( \frac{(\nabla_{a_i})_1}{\lambda_i} \right)^2 \varphi_i d\mu_{g_0} \\
&\quad + \tau \int_{B_\varepsilon(a_i)} \left( \frac{\tilde{c}_1 \lambda_i^2 r^2}{\tilde{c}_2 2n} - \frac{\tilde{c}_1 \bar{c}_2}{\tilde{c}_2 c_1} \right) \varphi_i^{\frac{n+2}{n-2}} \left( \frac{(\nabla_{a_i})_1}{\lambda_i} \right)^2 \varphi_i d\mu_{g_0} \\
&\quad - \int_{B_\varepsilon(a_i)} \left( \frac{\nabla^2 K_i}{2K_i} x^2 - \frac{\Delta K_i}{2nK_i} r^2 \right) \varphi_i^{\frac{n+2}{n-2}} \left( \frac{(\nabla_{a_i})_1}{\lambda_i} \right)^2 \varphi_i d\mu_{g_0} \\
&\quad - \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \left| \frac{(\nabla_{a_i})_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0}.
\end{aligned}$$

At this point some simplifications occur. From the relation

$$\tilde{c}_1 \tau + \tilde{c}_2 \frac{\Delta K_i}{K_i \lambda_i^2} = o\left(\frac{1}{\lambda^2}\right)$$

we obtain cancellation of the terms involving  $\Delta K_i$  and  $\frac{\tilde{c}_1 \lambda_i^2 r^2}{\tilde{c}_2 2n}$ . Using

$$\int \varphi_i^{\frac{4}{n-2}} \left| \frac{(\nabla_{a_i})_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0} = \frac{c_3}{n} + o(1)$$

and

$$\int \varphi_i^{\frac{n+2}{n-2}} \left( \frac{(\nabla_{a_i})_1}{\lambda_i} \right)^2 \varphi_i d\mu_{g_0} = -\frac{n+2}{n-2} \frac{c_3}{n} + o(1)$$

as well as  $(\frac{\bar{c}_1}{\bar{c}_0} - \frac{\bar{c}_1 \bar{c}_2}{\bar{c}_2 \bar{c}_0}) \frac{c_3}{n} = (\frac{\bar{c}_1}{c_1} - \frac{\bar{c}_1 \bar{c}_2}{\bar{c}_2 c_1}) \frac{c_2}{n}$  due  $\bar{c}_0 = c_1$  and  $c_2 = c_3$ , we find

$$\begin{aligned}
& \frac{(k_\tau)^{\frac{2}{p+1}}}{8n(n-1)} \alpha^m \varphi_m + \bar{v} \partial^2 J_\tau (\alpha^m \varphi_m + \bar{v}) \left( \frac{(\nabla a_i)_1}{\lambda_i} \varphi_i \right)^2 \\
&= \frac{c_3}{n} \tau - \frac{n+2}{n-2} \tau \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \left| \frac{(\nabla a_i)_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0} \\
&\quad - \tau \int_{B_\varepsilon(a_i)} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \varphi_i^{\frac{n+2}{n-2}} \left( \frac{(\nabla a_i)_1}{\lambda_i} \right)^2 \varphi_i d\mu_{g_0} \\
&\quad - \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{n+2}{n-2}} \left( \frac{(\nabla a_i)_1}{\lambda_i} \right)^2 \varphi_i d\mu_{g_0} \\
&\quad - \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \left| \frac{(\nabla a_i)_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0}.
\end{aligned}$$

Moreover we have, passing to integration over  $\mathbb{R}^n$ , up to an error  $o(1)$

$$\begin{aligned}
& \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} \varphi_i^{\frac{4}{n-2}} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \left| \frac{(\nabla a_i)_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0} \\
&= \int_{\mathbb{R}^n} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i}^{\frac{n+2}{n-2}} \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i} dx \\
&= -(n-2) \int_{\mathbb{R}^n} \frac{\lambda_i x_1}{1 + \lambda_i^2 r^2} \delta_{0,\lambda_i}^{\frac{n+2}{n-2}} \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i} dx \\
&\quad - \int_{\mathbb{R}^n} \ln(1 + \lambda_i^2 r^2)^{\frac{n-2}{2}} \delta_{0,\lambda_i}^{\frac{n+2}{n-2}} \left( \frac{(\nabla a_i)_1}{\lambda_i} \right) \delta_{0,\lambda_i} dx
\end{aligned}$$

and find for the first summand

$$(n-2) \int_{\mathbb{R}^n} \frac{\lambda_i x_1}{1 + \lambda_i^2 r^2} \delta_{0,\lambda_i}^{\frac{n+2}{n-2}} \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i} dx = - \int_{\mathbb{R}^n} \delta_{0,\lambda_i}^{\frac{4}{n-2}} \left| \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i} \right|^2 dx = -\frac{c_3}{n}.$$

We therefore are left with

$$\begin{aligned}
& \frac{(k_\tau)^{\frac{2}{p+1}}}{8n(n-1)} \alpha^m \varphi_m + \bar{v} \partial^2 J_\tau (\alpha^m \varphi_m + \bar{v}) \left( \frac{(\nabla a_i)_1}{\lambda_i} \varphi_i \right)^2 \\
&= - \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{n+2}{n-2}} \left( \frac{(\nabla a_i)_1}{\lambda_i} \right)^2 \varphi_i d\mu_{g_0} \\
&\quad - \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} \frac{\nabla^2 K_i}{2K_i} x^2 \varphi_i^{\frac{4}{n-2}} \left| \frac{(\nabla a_i)_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0}.
\end{aligned}$$

Finally passing to integration over  $\mathbb{R}^n$  up to some  $o(1)$  there holds

$$\begin{aligned} \frac{n+2}{n-2} \int_{B_\varepsilon(a_i)} x_l^2 \varphi_i^{\frac{4}{n-2}} \left| \frac{(\nabla a_i)_1}{\lambda_i} \varphi_i \right|^2 d\mu_{g_0} &= \int_{\mathbb{R}^n} x_l^2 \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i}^{\frac{n+2}{n-2}} \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i} dx \\ &= -2\delta_{1,l} \int_{\mathbb{R}^n} \frac{x_1}{\lambda_i} \delta_{0,\lambda_i}^{\frac{n+2}{n-2}} \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i} dx - \int_{\mathbb{R}^n} x_l^2 \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i}^{\frac{n+2}{n-2}} \left( \frac{(\nabla a_i)_1}{\lambda_i} \right)^2 \delta_{0,\lambda_i} dx, \end{aligned}$$

and similarly for  $j = 2, \dots, n$ . Diagonalizing the Hessian we have

$$\nabla^2 K_i x^2 = \sum_{l=1}^n \partial_l^2 K_i x_l^2$$

and

$$\begin{aligned} \int_{\mathbb{R}^n} \frac{x_1}{\lambda_i} \delta_{0,\lambda_i}^{\frac{n+2}{n-2}} \frac{(\nabla a_i)_1}{\lambda_i} \delta_{0,\lambda_i} dx &= -(n-2) \int_{\mathbb{R}^n} \delta_{0,\lambda_i}^{\frac{2n}{n-2}} \frac{x_1^2}{1 + \lambda_i^2 r^2} dx \\ &= -\frac{n-2}{n\lambda_i^2} \int_{\mathbb{R}^n} \frac{r^2}{(1+r^2)^{n+1}} dx, \end{aligned}$$

so we conclude that

$$\frac{(k_\tau)^{\frac{2}{p+1}}}{8n(n-1)} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v}) \left( \frac{(\nabla a_i)_1}{\lambda_i} \varphi_i \right)^2 = -c \frac{\partial_1^2 K_i}{K_i \lambda_i^2}.$$

Similarly one can show analogous formula for any couple of indices

$$\frac{(k_\tau)^{\frac{2}{p+1}}}{8n(n-1)} \partial^2 J_\tau(\alpha^m \varphi_m + \bar{v}) \frac{(\nabla a_i)_k}{\lambda_i} \varphi_i \frac{(\nabla a_i)_l}{\lambda_i} \varphi_i = -c \frac{\partial_{k,l}^2 K_i}{K_i \lambda_i^2}.$$

The proof is thereby complete.  $\square$

From Proposition 4.1 we deduce that the kernel of  $\partial^2 J_\tau$  is potentially one-dimensional. On the other hand the presence of an at one dimensional kernel at a solution is necessary due to the scaling invariance of  $J_\tau$ . Hence it is natural to impose some homogeneous constraint.

**Corollary 4.1.** *Let  $u \in \bar{V}(q, \varepsilon)$  be a solution of (5) and*

$$I_\tau = J_\tau|_{[\|\cdot\|_{L_{g_0}}=1]} \quad \text{or} \quad I_\tau = J_\tau|_{[\|\cdot\|_{k_\tau}=1]}.$$

*Then, if  $\tilde{u}$  denoted the corresponding normalization of  $u$ , we have*

$$m(I_\tau, \tilde{u}) = q - 1 + \sum_{i=1}^q (n - m(K, a_i)).$$



## 5 Appendix: Some estimates and list of constants

In this appendix, recalling our notation, we collect some useful statements and formulae proved in [30].

**Lemma 5.1.** *There holds  $L_{g_0}\varphi_{a,\lambda} = O(\varphi_{a,\lambda}^{\frac{n+2}{n-2}})$ . More precisely on a geodesic ball  $B_\alpha(a)$  for  $\alpha > 0$  small*

$$\begin{aligned} L_{g_0}\varphi_{a,\lambda} = & 4n(n-1)\varphi_{a,\lambda}^{\frac{n+2}{n-2}} - 2nc_n r_a^{n-2}((n-1)H_a + r_a \partial_{r_a} H_a)\varphi_{a,\lambda}^{\frac{n+2}{n-2}} \\ & + \frac{R_{g_a}}{\lambda} u_a^{\frac{2}{n-2}} \varphi_{a,\lambda}^{\frac{n}{n-2}} + o(r_a^{n-2})\varphi_{a,\lambda}^{\frac{n+2}{n-2}}, \quad r_a = d_{g_a}(a, \cdot). \end{aligned}$$

Since in conformal normal coordinates

$$R_{g_a} = O(r_a^2),$$

cf. [23], we obtain

(i) for  $n = 5$

$$L_{g_0}\varphi_{a,\lambda} = 4n(n-1)\left[1 - \frac{c_n}{2}r_a^{n-2}(H_a(a) + n\nabla H_a(a)x)\right]\varphi_{a,\lambda}^{\frac{n+2}{n-2}} + O(\lambda^{-2}\varphi_{a,\lambda});$$

(ii) for  $n = 6$  and with  $W(a) = |\mathbb{W}(a)|^2$

$$L_{g_0}\varphi_{a,\lambda} = 4n(n-1)\varphi_{a,\lambda}^{\frac{n+2}{n-2}} = 4n(n-1)\left[1 + \frac{c_n}{2}W(a)\ln r\right]\varphi_{a,\lambda}^{\frac{n+2}{n-2}} + O(\lambda^{-2}\varphi_{a,\lambda});$$

(iii) for  $n = 7$

$$L_{g_0}\varphi_{a,\lambda} = 4n(n-1)\varphi_{a,\lambda}^{\frac{n+2}{n-2}} + O(\lambda^{-2}\varphi_{a,\lambda}).$$

These expansions persist upon taking  $\lambda\partial\lambda$  and  $\frac{\nabla_a}{\lambda}$  derivatives.

**Lemma 5.2.** *Let  $\theta = \frac{n-2}{2}\tau$  and  $k, l = 1, 2, 3$  and  $i, j = 1, \dots, q$ . There holds uniformly as  $0 \leq \tau \rightarrow 0$*

(i)  $|\phi_{k,i}|, |\lambda_i \partial_{\lambda_i} \phi_{k,i}|, |\frac{1}{\lambda_i} \nabla_{a_i} \phi_{k,i}| \leq C\varphi_i$ ;

(ii)  $\lambda_i^\theta \int \varphi_i^{\frac{4}{n-2}-\tau} \phi_{k,i} \phi_{k,i} d\mu_{g_0} = c_k \cdot id + O(\tau + \frac{1}{\lambda_i^{2+\theta}})$ ,  $c_k > 0$ ;

(iii) for  $i \neq j$  up to some error of order  $O(\tau^2 + \sum_{i \neq j} (\frac{1}{\lambda_i^4} + \varepsilon_{i,j}^{\frac{n+2}{n}}))$

$$\lambda_i^\theta \int \varphi_i^{\frac{n+2}{n-2}-\tau} \phi_{k,j} d\mu_{g_0} = b_k d_{k,i} \varepsilon_{i,j} = \int \varphi_i^{1-\tau} d_{k,j} \varphi_j^{\frac{n+2}{n-2}} d\mu_{g_0};$$

$$(iv) \quad \lambda_i^\theta \int \varphi_i^{\frac{4}{n-2}-\tau} \phi_{k,i} \phi_{l,i} d\mu_{g_0} = O\left(\frac{1}{\lambda_i^\tau}\right) \text{ for } k \neq l \text{ and for } k = 2, 3$$

$$\lambda_i^\theta \int \varphi_i^{\frac{n+2}{n-2}-\tau} \phi_{k,i} d\mu_{g_0} = O\left(\tau + \begin{pmatrix} \lambda_i^{2-n} & \text{for } n = 5 \\ \frac{\ln \lambda_i}{\lambda_i^4} & \text{for } n = 6 \\ \lambda_i^4 & \text{for } n \geq 7 \end{pmatrix}\right);$$

$$(v) \quad \text{for } i \neq j, \alpha + \beta = \frac{2n}{n-2}, \alpha - \tau > \frac{n}{n-2} > \beta \geq 1$$

$$\lambda_i^\theta \int \varphi_i^{\alpha-\tau} \varphi_j^\beta d\mu_{g_0} = O(\varepsilon_{i,j}^\beta);$$

$$(vi) \quad \int \varphi_i^{\frac{n}{n-2}} \varphi_j^{\frac{n}{n-2}} d\mu_{g_0} = O(\varepsilon_{i,j}^{\frac{n}{n-2}} \ln \varepsilon_{i,j}), \quad i \neq j;$$

$$(vii) \quad (1, \lambda_i \partial_{\lambda_i}, \frac{1}{\lambda_i} \nabla_{a_i}) \varepsilon_{i,j} = O(\varepsilon_{i,j}), \quad i \neq j,$$

cf. (14), with constants

- $b_k = \int_{\mathbb{R}^n} \frac{dx}{(1+r^2)^{\frac{n+2}{2}}} \text{ for } k = 1, 2, 3;$
- $c_1 = \int_{\mathbb{R}^n} \frac{dx}{(1+r^2)^n};$
- $c_2 = \frac{(n-2)^2}{4} \int_{\mathbb{R}^n} \frac{|r^2-1|^2 dx}{(1+r^2)^{n+2}};$
- $c_3 = \frac{(n-2)^2}{n} \int_{\mathbb{R}^n} \frac{r^2 dx}{(1+r^2)^{n+2}}.$

**Lemma 5.3.** For  $u \in V(q, \varepsilon)$  with  $k_\tau = 1$  and  $\nu \in H_u(q, \varepsilon)$  there holds

$$\partial J_\tau(\alpha^i \varphi_i) \nu = O\left(\left[\sum_r \frac{\tau}{\lambda_r^\theta} + \sum_r \frac{|\nabla K_r|}{\lambda_r^{1+\theta}} + \sum_r \frac{1}{\lambda_r^{2+\theta}} + \sum_{r \neq s} \frac{\varepsilon_{r,s}^{\frac{n+2}{2n}}}{\lambda_r^\theta}\right] \|\nu\|\right).$$

**Lemma 5.4.** For  $u \in V(q, \varepsilon)$  and  $\varepsilon > 0$  sufficiently small the three quantities  $\partial J_\tau(u) \phi_{1,j}$ ,  $\partial J_\tau(\alpha^i \varphi_i) \phi_{1,j}$ ,  $\partial_{\alpha_j} J_\tau(\alpha^i \varphi_i)$  can be written as

$$\begin{aligned} & \frac{\alpha_j}{(\alpha_{K,\tau}^{\frac{2n}{n-2}})^{\frac{n-2}{n}}} \left( \dot{c}_0 \left(1 - \frac{\alpha^2}{\alpha_{K,\tau}^{p+1}} \frac{K_j}{\lambda_j^\theta} \alpha_j^{p-1}\right) - \dot{c}_2 \left(\frac{\Delta K_j}{K_j \lambda_j^2} - \sum_k \frac{\Delta K_k}{K_k \lambda_k^2} \frac{\alpha_k^2}{\alpha^2}\right) \right. \\ & \quad \left. + \dot{b}_1 \left(\sum_{k \neq l} \frac{\alpha_k \alpha_l}{\alpha^2} \varepsilon_{k,l} - \sum_{j \neq i} \frac{\alpha_i}{\alpha_j} \varepsilon_{i,j}\right) \right. \\ & \quad \left. - \dot{d}_1 \begin{pmatrix} \frac{H_j}{\lambda_j^3} - \sum_k \frac{\alpha_k^2}{\alpha^2} \frac{H_k}{\lambda_k^3} & \text{for } n = 5 \\ \frac{W_j \ln \lambda_j}{\lambda_j^4} - \sum_k \frac{\alpha_k^2}{\alpha^2} \frac{W_k \ln \lambda_k}{\lambda_k^4} & \text{for } n = 6 \\ 0 & \text{for } n \geq 7 \end{pmatrix} \right) \end{aligned}$$

up to an error of order

$$O(\tau^2 + \sum_{r \neq s} \frac{|\nabla K_r|^2}{\lambda_r^2} + \frac{1}{\lambda_r^4} + \varepsilon_{r,s}^{\frac{n+2}{n}} + |\partial J_\tau(u)|^2),$$

with positive constants

$$\begin{aligned} \bullet \quad \dot{b}_1 &= \frac{8n(n-1)(n+2)}{\bar{c}_0^{\frac{n-2}{n}}(n-2)} b_1; \\ \bullet \quad \dot{c}_2 &= \frac{8n(n-1)}{\bar{c}_0^{\frac{n-2}{n}}} \bar{c}_2; \\ \bullet \quad \dot{d}_1 &= \frac{8n(n-1)}{\bar{c}_0^{\frac{n-2}{n}}} \bar{d}_1; \\ \bullet \quad \dot{c}_0 &= 8n(n-1) \bar{c}_0^{\frac{2}{n}}. \end{aligned} \tag{36}$$

In particular for all  $j$

$$\frac{\alpha^2}{\alpha_{K,\tau}^{p+1}} \frac{K_j}{\lambda_j^\theta} \alpha_j^{p-1} = 1 + O(\tau + \sum_{r \neq s} \frac{1}{\lambda_r^2} + \varepsilon_{r,s} + |\partial J_\tau(u)|).$$

**Lemma 5.5.** For  $u \in V(q, \varepsilon)$  and  $\varepsilon > 0$  sufficiently small the three quantities  $\partial J_\tau(u) \phi_{2,j}$ ,  $\partial J_\tau(\alpha^i \varphi_i) \phi_{2,j}$  and  $\frac{\lambda_i}{\alpha_j} \partial_{\lambda_j} J_\tau(\alpha^i \varphi_i)$  can be written as

$$\frac{\alpha_j}{(\alpha_{K,\tau}^{\frac{2n}{n-2}})^{\frac{n-2}{n}}} \left( \tilde{c}_1 \tau + \tilde{c}_2 \frac{\Delta K_j}{K_j \lambda_j^2} - \tilde{b}_2 \sum_{j \neq i} \frac{\alpha_i}{\alpha_j} \lambda_j \partial_{\lambda_j} \varepsilon_{i,j} + \tilde{d}_1 \begin{pmatrix} \frac{H_j}{\lambda_j^3} & \text{for } n = 5 \\ \frac{W_j \ln \lambda_j}{\lambda_j^4} & \text{for } n = 6 \\ 0 & \text{for } n \geq 7 \end{pmatrix} \right),$$

with positive constants  $\tilde{c}_1, \tilde{c}_2, \tilde{d}_1, \tilde{b}_2$  up to some error

$$O(\tau^2 + \sum_{r \neq s} \frac{|\nabla K_r|^2}{\lambda_r^2} + \frac{1}{\lambda_r^4} + \varepsilon_{r,s}^{\frac{n+2}{n}} + |\partial J_\tau(u)|^2).$$

**Lemma 5.6.** For  $u \in V(q, \varepsilon)$  and  $\varepsilon > 0$  sufficiently small the three quantities  $\partial J_\tau(u) \phi_{3,j}$ ,  $\partial J_\tau(\alpha^i \varphi_i) \phi_{3,j}$  and  $\frac{\nabla_{a_j}}{\alpha_j \lambda_j} J_\tau(\alpha^i \varphi_i)$  can be written as

$$-\frac{\alpha_j}{(\alpha_{K,\tau}^{\frac{2n}{n-2}})^{\frac{n-2}{n}}} \left( \check{c}_3 \frac{\nabla K_j}{K_j \lambda_j} + \check{c}_4 \frac{\nabla \Delta K_j}{K_j \lambda_j^3} + \check{b}_3 \sum_{j \neq i} \frac{\alpha_i}{\alpha_j} \frac{\nabla_{a_j}}{\lambda_j} \varepsilon_{i,j} \right),$$

with positive constants  $\check{c}_3, \check{c}_4, \check{b}_3$  up to some error

$$O(\tau^2 + \sum_{r \neq s} \frac{|\nabla K_r|^2}{\lambda_r^2} + \frac{1}{\lambda_r^4} + \varepsilon_{r,s}^{\frac{n+2}{n}} + |\partial J_\tau(u)|^2).$$

**Lemma 5.7.** For every  $u \in V(q, \varepsilon)$  there holds

$$|\partial J_\tau(u)| \lesssim \tau + \sum_{r \neq s} \frac{|\nabla K_r|}{\lambda_r} + \frac{1}{\lambda_r^2} + \left| 1 - \frac{\alpha^2}{\alpha_{K, \tau}^{p+1}} \frac{K_r}{\lambda_r^\theta} \alpha_r^{p-1} \right| + \varepsilon_{r,s}^{\frac{n+2}{2n}} + \|v\|.$$

**Theorem 2.** Suppose that  $n \geq 5$ ,  $K : M \rightarrow \mathbb{R}_+$  is Morse and satisfies (3). Then for  $\varepsilon > 0$  sufficiently small there exists  $c > 0$  such that for any

$$u \in V(q, \varepsilon) \quad \text{with} \quad k_\tau = 1$$

there holds

$$|\partial J(u)| \geq c \left( \tau + \sum_{r \neq s} \frac{|\nabla K_r|}{\lambda_r} + \frac{1}{\lambda_r^2} + \left| 1 - \frac{\alpha^2}{\alpha_{K, \tau}^{p+1}} \frac{K_r}{\lambda_r^\theta} \alpha_r^{p-1} \right| + \varepsilon_{r,s} \right),$$

unless there is a violation of at least one of the four conditions

- (i)  $\tau > 0$  ;
- (ii)  $\exists x_i \neq x_j \in \{\nabla K = 0\} \cap \{\Delta K < 0\}$  and  $d(a_i, x_i) = O(\frac{1}{\lambda_i})$  ;
- (iii)  $\alpha_j = \Theta \cdot \left(\frac{\lambda_j^\theta}{K_j}\right)^{\frac{1}{p-1}} + o(\frac{1}{\lambda_j^2})$ ;
- (iv)  $\tilde{c}_1 \tau = -\tilde{c}_2 \frac{\Delta K_k}{K_k \lambda_k^{\frac{3}{2}}} + o(\frac{1}{\lambda_k^2})$

where  $\Theta$  is a positive constant, uniformly bounded and bounded away from zero, that depends on  $u$ , cf. Remark 6.2 in [30]. In the latter case there holds

$$\lambda_1 \simeq \dots \lambda_q \simeq \lambda = \frac{1}{\sqrt{\tau}}$$

and setting  $a_j = \exp_{g_{x_j}}(\bar{a}_j)$ , we still have up to an error  $o(\frac{1}{\lambda^3})$  the lower bound

$$\begin{aligned} |\partial J(u)| &\gtrsim \sum_j \left| \tau + \frac{2}{9} \frac{\Delta K(x_j)}{K(x_j) \lambda_j^2} + \frac{512}{9\pi} \left[ \frac{H(x_j)}{\lambda_j^3} + \sum_{j \neq i} \sqrt{\frac{K(x_j)}{K(x_i)}} \frac{G_{g_0}(x_i, x_j)}{\gamma_n(\lambda_i \lambda_j)^{\frac{3}{2}}} \right] \right| \\ &\quad + \sum_j \left| \frac{\bar{a}_j}{\lambda_j} + \frac{\check{c}_4}{\check{c}_3} (\nabla^2 K(x_j))^{-1} \frac{\nabla \Delta K(x_j)}{\lambda_j^3} \right| \\ &\quad + \sum_j \left| \alpha_j - \Theta \cdot \sqrt[p-1]{\frac{\lambda_j^\theta}{K(a_j)} \left( 1 - \frac{1}{90} \left( \frac{\Delta K(x_j)}{K(x_j) \lambda_j^2} + \frac{2816}{\pi} \frac{H(x_j)}{\lambda_j^3} - \frac{\sum_k \left( \frac{\Delta K(x_k)}{K(x_k)^2 \lambda_k^2} + \frac{2816}{\pi} \frac{H(x_k)}{K(x_k) \lambda_k^3} \right)}{\sum_k \frac{1}{K(x_k)}} \right) \right)} \right| \end{aligned}$$

in case  $n = 5$  and

$$\begin{aligned} |\partial J(u)| &\gtrsim \sum_j \left( \left| \tau + \frac{\check{c}_2}{\check{c}_1} \frac{\Delta K(x_j)}{K(x_j) \lambda_j^2} \right| \right. \\ &\quad \left. + \left| \frac{\bar{a}_j}{\lambda_j} + \frac{\check{c}_4}{\check{c}_3} (\nabla^2 K(x_j))^{-1} \frac{\nabla \Delta K(x_j)}{\lambda_j^3} \right| + \left| \alpha_j - \Theta \cdot \sqrt[p-1]{\frac{\lambda_j^\theta}{K(a_j)}} \right| \right) \end{aligned}$$

in case  $n \geq 6$ . The constants appearing above are defined by

- $\bar{c}_0 = \int_{\mathbb{R}^n} \frac{dx}{(1+r^2)^n}$  ;
- $\tilde{c}_1 = \frac{n(n-1)(n-2)^2}{\bar{c}_0^{\frac{n-2}{n}}} \int_{\mathbb{R}^n} \frac{1-r^2}{(1+r^2)^{n+1}} \ln \frac{1}{1+r^2} dx$ ;
- $\tilde{c}_2 = -\frac{(n-1)(n-2)}{\bar{c}_0^{\frac{n-2}{n}}} \int_{\mathbb{R}^n} \frac{r^2(1-r^2)}{(1+r^2)^{n+1}} dx$ ;
- $\check{c}_3 = \int_{\mathbb{R}^n} \frac{4(n-1)(n-2)}{(1+r^2)^n} dx$ ;
- $\tilde{b}_2 = \frac{4n(n-1)}{\bar{c}_0^{\frac{n-2}{n}}} \int_{\mathbb{R}^n} \frac{dx}{(1+r^2)^{\frac{n+2}{2}}}$  ;
- $\check{c}_4 = \int_{\mathbb{R}^n} \frac{2(n-1)r^2}{(1+r^2)^n} dx$  ;
- $\tilde{d}_1 = \frac{4n(n-1)}{\bar{c}_0^{\frac{n-2}{n}}} \int_{\mathbb{R}^n} r^n \frac{(n+2-nr^2)}{(1+r^2)^{n+2}} dx$ .

From the proof of Proposition 5.1 and Sections 4, 5 and 6 in [30] we will need the estimates

- (i) up to an error of order  $O\left(\tau^2 + \sum_r \frac{1}{\lambda_r^4} + \sum_{r \neq s} \varepsilon_{r,s}^{\frac{n+2}{n}}\right)$  there holds

$$\begin{aligned}
& \int K(\alpha^i \varphi_i)^{p+1} d\mu_{g_0} \\
&= \sum_i \left( \bar{c}_0 \frac{K_i}{\lambda_i^\theta} \alpha_i^{p+1} + \bar{c}_1 \frac{K_i}{\lambda_i^\theta} \alpha_i^{\frac{2n}{n-2}} \tau + \bar{c}_2 \frac{\Delta K_i}{\lambda_i^{2+\theta}} \alpha_i^{\frac{2n}{n-2}} \right) \\
& \quad + \bar{d}_1 \sum_i \frac{K_i}{\lambda_i^\theta} \alpha_i^{\frac{2n}{n-2}} \left( \frac{\frac{H_i}{\lambda_i^3}}{\frac{W_i \ln \lambda_i}{\lambda_i^4}} \right) + \bar{b}_1 \sum_{i \neq j} \alpha_i^{\frac{n+2}{n-2}} \alpha_j \frac{K_i}{\lambda_i^\theta} \varepsilon_{i,j}
\end{aligned} \tag{37}$$

with  $(\bar{b}_1 = \frac{2n}{n-2} b_1)$  and  $\bar{d}_1 = \int_{\mathbb{R}^n} \frac{r^n dx}{(1+r^2)^{n+1}}$ ;

- (ii) recalling (14) we have

$$\int \varphi_i L_{g_0} \varphi_j d\mu_{g_0} = \tilde{b}_1 \varepsilon_{i,j} + O\left(\sum_{r \neq s} \frac{1}{\lambda_r^4} + \varepsilon_{r,s}^{\frac{n+2}{n}}\right), \quad \tilde{b}_1 = 4n(n-1)b_1; \tag{38}$$

- (iii) up to an error  $O(\tau^2 + \frac{1}{\lambda_i^4})$ , there holds

$$\int \frac{\varphi_i L_{g_0} \varphi_i}{4n(n-1)} d\mu_{g_0} = \bar{c}_0; \tag{39}$$

(iv) up to an error of order  $O(\tau^2 + \sum_r \frac{1}{\lambda_r^4} + \sum_{r \neq s} \varepsilon_{r,s}^{\frac{n+2}{n}})$  we have

$$\alpha^i \alpha^j \int \varphi_i L_{g_0} \varphi_j d\mu_{g_0} = 4n(n-1) \bar{c}_0 \sum_i \alpha_i^2 + \tilde{b}_1 \sum_{i \neq j} \alpha_i \alpha_j \varepsilon_{i,j}. \quad (40)$$

(v) If  $\varphi_i$  is as in (12), then

$$\begin{aligned} \left| \int \varphi_i^{\frac{n+2}{n-2}} \nu d\mu_{g_0} \right| &\leq \|v\| \left\| \frac{L_{g_0} \varphi_i}{4n(n-1)} - \varphi_i^{\frac{n+2}{n-2}} \right\|_{L_{g_0}^{\frac{2n}{n+2}}} \\ &= O \left( \begin{array}{ll} \lambda_i^{-3} & \text{for } n=5 \\ \ln^{\frac{2}{3}} \lambda_i \lambda_i^{-\frac{10}{3}} & \text{for } n=6 \\ \lambda_i^{-4} & \text{for } n \geq 7 \end{array} \right) \|v\|; \end{aligned} \quad (41)$$

(vi) up to an error  $O(\tau^2 + \frac{1}{\lambda_i^4})$  we have with  $\bar{c}_2 = \frac{1}{2n} \int_{\mathbb{R}^n} \frac{r^2 dx}{(1+r^2)^n}$ ;

$$\int K \varphi_i^{p+1} d\mu_{g_0} = \frac{\bar{c}_0 K_i}{\lambda_i^\theta} + \bar{c}_1 \frac{K_i \tau}{\lambda_i^\theta} + \bar{c}_2 \frac{\Delta K_i}{\lambda_i^{2+\theta}} + \bar{d}_1 K_i \begin{pmatrix} \frac{H_i}{\lambda_i^{3+\theta}} \\ \frac{W_i \ln \lambda_i}{\lambda_i^{4+\theta}} \\ 0 \end{pmatrix}; \quad (42)$$

(vii) up to an error or order  $O(\tau^2 + \sum_{r \neq s} \frac{|\nabla K_r|^2}{\lambda_r^2} + \frac{1}{\lambda_r^4} + \varepsilon_{r,s}^{\frac{n+2}{n}})$  there holds

$$\begin{aligned} J_\tau(\alpha^i \varphi_i) &= \frac{\alpha^i \alpha^j \int \varphi_i L_{g_0} \varphi_j d\mu_{g_0}}{(\int K(\sum_i \alpha_i \varphi_i)^{p+1})^{\frac{2}{p+1}}} \\ &= \frac{\alpha^i \alpha^j \int \varphi_i L_{g_0} \varphi_j d\mu_{g_0}}{(\bar{c}_0 \sum_i \frac{K_i}{\lambda_i^\theta} \alpha_i^{p+1})^{\frac{2}{p+1}}} \left( 1 - \bar{c}_1 \sum_i \frac{K_i}{\lambda_i^\theta} \frac{\alpha_i^{\frac{2n}{n-2}}}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} \tau \right. \\ &\quad - \bar{c}_2 \sum_i \frac{\Delta K_i}{\lambda_i^{2+\theta}} \frac{\alpha_i^{\frac{2n}{n-2}}}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} \\ &\quad - \bar{d}_1 \sum_i \frac{K_i}{\lambda_i^\theta} \begin{pmatrix} \frac{H_i}{\lambda_i^3} \\ \frac{W_i \ln \lambda_i}{\lambda_i^4} \\ 0 \end{pmatrix} \frac{\alpha_i^{\frac{2n}{n-2}}}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} \\ &\quad \left. - \bar{b}_1 \sum_{i \neq j} \frac{\alpha_i^{\frac{n+2}{n-2}} \alpha_j}{\alpha_{K,\tau}^{\frac{2n}{n-2}}} \frac{K_i}{\lambda_i^\theta} \varepsilon_{i,j} \right); \end{aligned} \quad (43)$$

(viii) if  $\varepsilon_{i,j}$  is as in (14), then in case  $j < i$  or  $d_{g_0}(a_i, a_j) \neq o(1)$

$$\lambda_j \partial_{\lambda_j} \varepsilon_{i,j} = \frac{2-n}{2} \varepsilon_{i,j} + O\left(\frac{1}{\lambda_j^4} + \varepsilon_{i,j}^{\frac{n+2}{n}}\right). \quad (44)$$

Finally we derive one last technical estimate. Recalling (7), from (40) we have up to an error  $o(\frac{1}{\lambda^2})$ ,

$$r_{\alpha^i \varphi_i} = \alpha^i \alpha^j \int L_{g_0} \varphi_i \varphi_j d\mu_{g_0} = 4n(n-1) \bar{c}_0 \sum_i \alpha_i^2 = 4n(n-1) \bar{c}_0 \alpha^2 \quad (45)$$

with  $\bar{c}_0 = \int_{\mathbb{R}^n} \frac{dx}{(1+r^2)^n}$ . From (37) instead we get

$$\begin{aligned} \int K(\alpha^i \varphi_i)^{p+1} d\mu_{g_0} &= \sum_i \left( \bar{c}_0 \frac{K_i}{\lambda_i^\theta} \alpha_i^{p+1} + \bar{c}_1 \frac{K_i}{\lambda_i^\theta} \alpha_i^{\frac{2n}{n-2}} \tau + \bar{c}_2 \frac{\Delta K_i}{\lambda_i^{2+\theta}} \alpha_i^{\frac{2n}{n-2}} \right) \\ &= \bar{c}_0 \alpha_{K,\theta}^{p+1} + \sum_i \frac{K_i \alpha_i^{\frac{2n}{n-2}}}{\lambda_i^\theta} \left( \bar{c}_1 \tau + \bar{c}_2 \frac{\Delta K_i}{K_i \lambda_i^2} \right) \end{aligned}$$

up to an error  $o(\frac{1}{\lambda^2})$  and with constants given by

$$\bar{c}_1 = \frac{2}{n-2} \int_{\mathbb{R}^n} \frac{\ln(1+r^2)}{(1+r^2)^n} dx, \quad \text{and} \quad \bar{c}_2 = \frac{1}{2n} \int_{\mathbb{R}^n} \frac{r^2}{(1+r^2)^n} dx. \quad (46)$$

Therefore

$$\begin{aligned} \frac{r_{\alpha^i \varphi_i}}{(k_\tau)_{\alpha^i \varphi_i}} &= 4n(n-1) \frac{\alpha^2}{\alpha_{K,\theta}^{p+1}} \\ &\quad - 4n(n-1) \frac{\alpha^2}{(\alpha_{K,\theta}^{p+1})^2} \sum_i \frac{K_i \alpha_i^{\frac{2n}{n-2}}}{\lambda_i^\theta} \left( \frac{\bar{c}_1}{\bar{c}_0} \tau + \frac{\bar{c}_2}{\bar{c}_0} \frac{\Delta K_i}{K_i \lambda_i^2} \right) + o\left(\frac{1}{\lambda^2}\right) \end{aligned}$$

and we conclude again from (17) that

$$\frac{r_{\alpha^i \varphi_i}}{(k_\tau)_{\alpha^i \varphi_i}} = 4n(n-1) \frac{\alpha^2}{\alpha_{K,\theta}^{p+1}} \left( 1 - \left( \frac{\bar{c}_1}{\bar{c}_0} - \frac{\bar{c}_1 \bar{c}_2}{\bar{c}_2 \bar{c}_0} \right) \tau \right) + o\left(\frac{1}{\lambda^2}\right). \quad (47)$$

At last we display for the reader convenience the equations where some dimensional constants appear.

$c_0$		(45)		(36)		
$c_1$	Lemma 5.2	(46)			Theorem 2	
$c_2$	Lemma 5.2	(46)		(36)	Theorem 2	
$c_3$	Lemma 5.2		(35)			Theorem 2
$c_4$						Theorem 2
$d_1$		(37)		(36)	Theorem 2	
$b_1$	Lemma 5.2	(37)		(36)	(38)	
$b_2$	Lemma 5.2				Theorem 2	
$b_3$	Lemma 5.2					

## References

- [1] Ambrosetti A., Garcia Azorero J., Peral A., *Perturbation of  $\Delta u + u^{\frac{(N+2)}{(N-2)}} = 0$ , the Scalar Curvature Problem in  $\mathbb{R}^N$  and related topics*, Journal of Functional Analysis, 165 (1999), 117-149.
- [2] Ambrosetti A., Malchiodi A., *Perturbation Methods and Semilinear Elliptic Problems on  $R^n$* , Birkhäuser, 2006.
- [3] Aubin T., *Equations différentielles non linéaires et Problème de Yamabe concernant la courbure scalaire*, J. Math. Pures et Appl. 55 (1976), 269-296.
- [4] Aubin T., *Some Nonlinear Problems in Differential Geometry*, Springer-Verlag, 1998.
- [5] Bahri A., *Critical points at infinity in some variational problems*, Research Notes in Mathematics, 182, Longman-Pitman, London, 1989.
- [6] Bahri A., Coron J.M., *The Scalar-Curvature problem on the standard three-dimensional sphere*, Journal of Functional Analysis, 95 (1991), 106-172.
- [7] Ben Ayed M., Chen Y., Chtioui H., Hammami M., *On the prescribed scalar curvature problem on 4-manifolds*, Duke Mathematical Journal, 84 (1996), 633-677.
- [8] Ben Ayed M., Chtioui H., Hammami M., *The scalar-curvature problem on higher-dimensional spheres*, Duke Math. J. 93 (1998), no. 2, 379-424.
- [9] Bianchi G., *The scalar curvature equation on  $\mathbb{R}^n$  and on  $S^n$* , Adv. Diff. Eq. 1 (1996), 857-880.
- [10] Bianchi G., Egnell H., *A variational approach to the equation  $\Delta u + K u^{\frac{n+2}{n-2}} = 0$  in  $\mathbb{R}^n$* , Arch. Rat. Mech. Anal. 122 (1993), 159-182.
- [11] Caffarelli L., Gidas B., Spruck J., *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math. 42 (1989), no. 3, 271-297.
- [12] Chang S. A., Gursky M. J., Yang P., *The scalar curvature equation on 2- and 3- spheres*, Calc. Var. 1 (1993), 205-229.
- [13] Chen C.C., Lin C.S., *Estimates of the conformal scalar curvature equation via the method of moving planes*, Comm. Pure Appl. Math. 50 (1997), no. 10, 971-1017.
- [14] Chen C.C., Lin C.S., *Estimate of the conformal scalar curvature equation via the method of moving planes. II*, J. Diff. Geom. 49 (1998), no. 1, 115-178.
- [15] Chen C.C., Lin C.S., *Blowing up with infinite energy of conformal metrics on  $S^N$* , Comm. Partial Differential Equations 24 (1999), no. 5-6, 785-799.



- [16] Chang S. A., Yang P., *Prescribing Gaussian curvature on  $S^2$* , Acta Math. 159 (1987), 215-259.
- [17] Chang S. A., Yang P., *Conformal deformation of metrics on  $S^2$* , J. Diff. Geom. 27 (1988), 256-296.
- [18] Chang S. A., Yang P., *A perturbation result in prescribing scalar curvature on  $S^n$* , Duke Math. J. 64 (1991), 27-69.
- [19] Chen W. X., Ding W., *Scalar curvature on  $S^2$* , Trans. Amer. Math. Soc. 303 (1987), 365-382.
- [20] Escobar J., Schoen R., *Conformal metrics with prescribed scalar curvature*, Inventiones Mathematicae, 86 (1986), 243-254.
- [21] Hebey E., *Changements de métriques conformes sur la sphère - Le problème de Nirenberg*, Bull. Sci. Math. 114 (1990), 215-242.
- [22] Hebey E., Vaugon M., *Le probleme de Yamabe equivariant. [The equivariant Yamabe problem]*, Bull. Sci. Math. 117 (1993), no. 2, 241-286.
- [23] Lee J., Parker T., *The Yamabe problem*, Bull. Amer. Math. Soc. (N.S.) 17 (1987), no. 1, 37-91.
- [24] Li Y.Y., *Prescribing scalar curvature on  $S^n$  and related topics, Part I*, J.Diff.Equat.120 (1995), 319-410.
- [25] Li Y.Y., *Prescribing scalar curvature on  $S^n$  and related topics, Part II, Existence and compactness*, Comm.Pure Appl.Math.49 (1996), 437-477.
- [26] Kazdan J.L., Warner F., *Curvature functions for compact 2-manifolds*, Ann. of Math. (2) 99 (1974), 14-47.
- [27] Kazdan J.L., Warner F., *Existence and conformal deformation of metrics with prescribed Gaussian and scalar curvature*, Ann. of Math. 101 (1975), 317-331.
- [28] Kazdan J.L., Warner F., *Scalar curvature and conformal deformation of Riemannian structure*, J. Differential Geometry 10 (1975), 113-134.
- [29] Malchiodi A., *The Scalar Curvature problem on  $S^n$  : an approach via Morse Theory*, Calc. Var., 2002.
- [30] Malchiodi A., Mayer M., *Prescribing Morse scalar curvatures: blow-up analysis*, arxiv.org/1812.09457, to appear on International Mathematical Research Notes
- [31] Malchiodi A., Mayer M., *Prescribing Morse scalar curvatures: pinching and Morse theory*, arxiv.org/1909.03190
- [32] Mayer M., *A scalar curvature flow in low dimensions*, Calc. Var. Partial Differential Equations 56 (2017), no. 2, Art. 24, 41 pp.

- [33] Mayer M., *Prescribing Morse scalar curvatures: critical points at infinity*, arxiv.org/1901.06409
- [34] Mayer M. *Prescribing scalar curvatures: non compactness versus critical points at infinity*, Geometric Flows, 2019 Issue 4(1), pp. 51-82
- [35] Moser J., *On a nonlinear problem in differential geometry*, Dynamical Systems (M. Peixoto ed.), Academic Press, New York, 1973, 273-280.
- [36] Schoen R., Zhang D., *Prescribed scalar curvature on the  $n$ -sphere*, Calculus of Variations and Partial Differential Equations, 4 (1996), 1-25.
- [37] Talenti G., *Best constant in Sobolev Inequality*, Ann. Mat. Pura Appl. 110 (1976) 353-372.
- [38] Wei J., Yan S., *Infinitely many solutions for the prescribed scalar curvature problem on  $S^N$* , J. Funct. Anal. 258 (2010), no. 9, 3048-3081.