

Twisted de Rham Complex on Line and Singular Vectors in $\widehat{\mathfrak{sl}}_2$ Verma Modules

Alexey SLINKIN[†] and Alexander VARCHENKO[‡]

[†] Department of Mathematics, University of North Carolina at Chapel Hill,
Chapel Hill, NC 27599-3250, USA

E-mail: slinalex@live.unc.edu, anv@email.unc.edu

URL: <http://varchenko.web.unc.edu/>

[‡] Faculty of Mathematics and Mechanics, Lomonosov Moscow State University,
Leninskiye Gory 1, 119991 Moscow GSP-1, Russia

Received May 30, 2019, in final form September 21, 2019; Published online September 26, 2019

<https://doi.org/10.3842/SIGMA.2019.075>

Abstract. We consider two complexes. The first complex is the twisted de Rham complex of scalar meromorphic differential forms on projective line, holomorphic on the complement to a finite set of points. The second complex is the chain complex of the Lie algebra of \mathfrak{sl}_2 -valued algebraic functions on the same complement, with coefficients in a tensor product of contragredient Verma modules over the affine Lie algebra $\widehat{\mathfrak{sl}}_2$. In [Schechtman V., Varchenko A., *Mosc. Math. J.* **17** (2017), 787–802] a construction of a monomorphism of the first complex to the second was suggested and it was indicated that under this monomorphism the existence of singular vectors in the Verma modules (the Malikov–Feigin–Fuchs singular vectors) is reflected in the relations between the cohomology classes of the de Rham complex. In this paper we prove these results.

Key words: twisted de Rham complex; logarithmic differential forms; $\widehat{\mathfrak{sl}}_2$ -modules; Lie algebra chain complexes

2010 Mathematics Subject Classification: 17B56; 17B67; 33C80

*Dedicated to Dmitry Borisovich Fuchs
on the occasion of his 80-th birthday*

1 Introduction

We consider two complexes. The first complex is the twisted de Rham complex of scalar meromorphic differential forms on projective line, that are holomorphic on the complement to a finite set of points. The second complex is the chain complex of the Lie algebra of \mathfrak{sl}_2 -valued algebraic functions on the same complement, with coefficients in a tensor product of contragredient Verma modules over the affine Lie algebra $\widehat{\mathfrak{sl}}_2$. In [9] a construction of a monomorphism of the first complex to the second was suggested. That construction gives a relation between the singular vectors in the Verma modules and resonance relations in the de Rham complex.

That construction of the homomorphism was invented in the middle of 90s, while the paper [9] was prepared for publication 20 years later, when the proofs were forgotten, if they existed. The paper [9] provides supporting evidence to the results formulated in [9], but not the proofs. The goal of this paper is to give the proofs to the results formulated in [9], namely, the proofs that the construction in [9] indeed gives a homomorphism of complexes and relates the resonances in the de Rham complex and the $\widehat{\mathfrak{sl}}_2$ singular vectors.

This paper is a contribution to the Special Issue on Algebra, Topology, and Dynamics in Interaction in honor of Dmitry Fuchs. The full collection is available at <https://www.emis.de/journals/SIGMA/Fuchs.html>

The construction in [9] has two motivations.

The first motivation was to generalize the principal construction of [8]. In [8], the tensor products of contragredient Verma modules over a semisimple Lie algebra were identified with the spaces of the top degree logarithmic differential forms over certain configuration spaces. Also the logarithmic parts of the de Rham complexes over the configuration spaces were identified with some standard Lie algebra chain complexes having coefficients in these tensor products, cf. in [4, 5] a \mathcal{D} -module explanation of this correspondence.

The second idea was that the appearance of singular vectors in Verma modules over affine Lie algebras is reflected in the relations between the cohomology classes of logarithmic differential forms. This was proved in an important particular case in [1, 2], and in [7] a one-to-one correspondence was established “on the level of parameters”. In [9] and in the present paper this correspondence is developed for another non-trivial class of singular vectors, namely for (a part of) Malikov–Feigin–Fuchs singular vectors, cf. [6].

The paper has the following structure. In Section 2 we introduce the de Rham complex of a master function and resonance relations. In Section 3 we discuss $\widehat{\mathfrak{sl}}_2$ Verma modules, the Kac–Kazhdan reducibility conditions. We formulate Theorem 3.2 which describes certain relations in a contragredient Verma module. The proof of Theorem 3.2 is the main new result of this paper. In Theorem 3.3 we describe the connection between the relations, described in Theorem 3.2, and the Malikov–Feigin–Fuchs singular vectors. In Section 4 we construct a map of the de Rham complex of the master function to the chain complex of the Lie algebra of \mathfrak{sl}_2 -valued algebraic functions. Theorem 4.1 says that the map is a monomorphism of complexes. The proof of Theorem 4.1 is the second new result of this paper. Section 5 is devoted to the proof of Theorem 3.2. The proof is straightforward but rather nontrivial and lengthy.

2 The de Rham complex of master function

2.1 Twisted de Rham complex

Consider \mathbb{C} with coordinate t . Define the *master function* by the formula

$$\Phi(t) = \prod_{i=1}^n (t - z_i)^{-m_i/\kappa},$$

where $z_1, \dots, z_n, m_1, \dots, m_n, \kappa \in \mathbb{C}$ are parameters. Fix these parameters and assume that z_1, \dots, z_n are distinct. Set

$$z_{n+1} = \infty, \quad m_{n+1} = m_1 + \dots + m_n - 2.$$

Denote $U = \mathbb{C} - \{z_1, \dots, z_n\}$.

Consider the *twisted de Rham complex* associated with Φ ,

$$0 \longrightarrow \Omega^0(U) \xrightarrow{\partial} \Omega^1(U) \longrightarrow 0. \quad (2.1)$$

Here $\Omega^p(U)$ is the space of rational differential p -forms on \mathbb{C} regular on U . The differential ∂ is given by the formula

$$\partial = d + \alpha \wedge \cdot, \quad (2.2)$$

where d is the standard de Rham differential and the second summand is the left exterior multiplication by the form

$$\alpha = -\frac{1}{\kappa} \sum_{i=1}^n m_i \frac{dt}{t - z_i} = \frac{d\Phi}{\Phi}.$$

Formula (2.2) is motivated by the computation

$$d(\Phi\omega) = \Phi d\omega + d\Phi \wedge \omega = \Phi(d\omega + \alpha \wedge \omega).$$

The complex $\Omega^\bullet(U)$ is the complex of global *algebraic* sections of the de Rham complex of $(\mathcal{O}_U^{\text{an}}, \partial)$, where $\partial = d + \alpha \wedge \cdot$ is considered as the integrable connection on the sheaf $\mathcal{O}_U^{\text{an}}$ of holomorphic functions on U .

If the monodromy of Φ is non-trivial, that is, if at least one of the numbers $m_1/\kappa, \dots, m_n/\kappa$ is not an integer, then

$$H^0(\Omega^\bullet(U)) = 0, \quad \dim H^1(\Omega^\bullet(U)) = n - 1,$$

see for example [7].

2.2 Basis of $\Omega^\bullet(U)$

The functions

$$\frac{1}{(t - z_i)^a} \text{ for } a \in \mathbb{Z}_{>0} \quad \text{and} \quad t^a \text{ for } a \in \mathbb{Z}_{\geq 0}$$

form a basis of $\Omega^0(U)$. The differential forms

$$\frac{dt}{(t - z_i)^a} \text{ for } a \in \mathbb{Z}_{>0} \quad \text{and} \quad t^a dt \text{ for } a \in \mathbb{Z}_{\geq 0}$$

form a basis of $\Omega^1(U)$. The differential ∂ is given by the formulas

$$\begin{aligned} \kappa \partial \left(\frac{1}{(t - z_i)^a} \right) &= -(m_i + a\kappa) \frac{dt}{(t - z_i)^{a+1}} + \sum_{k=1}^a \sum_{j \neq i} \frac{m_j}{(z_j - z_i)^k} \frac{dt}{(t - z_i)^{a+1-k}} \\ &\quad - \sum_{j \neq i} \frac{m_j}{(z_j - z_i)^a} \frac{dt}{t - z_j}, \end{aligned} \quad (2.3)$$

$$\kappa \partial(t^a) = \left(a\kappa - \sum_{j=1}^n m_j \right) t^{a-1} dt - \sum_{k=1}^{a-1} \sum_{j=1}^n m_j z_j^k t^{a-1-k} dt - \sum_{j=1}^n m_j z_j^a \frac{dt}{t - z_j}. \quad (2.4)$$

2.3 Resonances

The equations

- (i) $m_i + (a - 1)\kappa = 0$ for some $a \in \mathbb{Z}_{>0}$, $i \in \{1, \dots, n\}$,
- (ii) $m_{n+1} + 2 - a\kappa = 0$ for some $a \in \mathbb{Z}_{>0}$,
- (iii) $\kappa = 0$,

are called the *resonance relations* for the parameters m_1, \dots, m_{n+1} , κ of the de Rham complex.

If $\kappa = 0$, then the twisted de Rham complex is not defined. If the resonance relation $m_i + a\kappa = 0$ is satisfied for some a , then the first term in the right-hand side of (2.3) equals zero. Similarly, if the resonance relation $m_{n+1} + 2 - a\kappa = 0$ is satisfied for some a , then the first term in the right-hand side of (2.4) equals zero.

2.4 Logarithmic subcomplex

Let $\Omega_{\log}^0(U) \subset \Omega^0(U)$ be the subspace generated over \mathbb{C} by function 1. Let $\Omega_{\log}^1(U) \subset \Omega^1(U)$ be the subspace generated over \mathbb{C} by the differential forms

$$\omega_j = \frac{dt}{t - z_j}, \quad j = 1, \dots, n.$$

These subspaces form the *logarithmic* subcomplex $(\Omega_{\log}^\bullet(U), \partial)$ of the de Rham complex $(\Omega^\bullet(U), \partial)$. We have

$$\partial: 1 \mapsto \alpha.$$

For generic m_1, \dots, m_n, κ , the embedding $(\Omega_{\log}^\bullet(U), \partial) \hookrightarrow (\Omega^\bullet(U), \partial)$ is a quasi-isomorphism, the logarithmic forms $\omega_1, \dots, \omega_n$ generate the space $H^1(\Omega^\bullet(U))$, and the cohomological relation $\sum_{i=1}^n m_i \omega_i \sim 0$ is the only one, see for example [7].

Each resonance relation implies a new cohomological relation between the forms $\omega_1, \dots, \omega_n$, see [9, Corollary 6.4]. For example, if $m_{n+1} + 2 - \kappa = 0$, then $\sum_{j=1}^n z_j m_j \omega_j \sim 0$, and if $m_{n+1} + 2 - 2\kappa = 0$, then

$$\sum_{j=1}^n z_j^2 m_j \omega_j - \frac{1}{\kappa} \left(\sum_{j=1}^n z_j m_j \right) \left(\sum_{i=1}^n z_i m_i \omega_i \right) \sim 0.$$

3 $\widehat{\mathfrak{sl}}_2$ -modules

3.1 Lie algebra $\widehat{\mathfrak{sl}}_2$

Let \mathfrak{sl}_2 be the Lie algebra of complex (2×2) -matrices with zero trace. Let e, f, h be standard generators subject to the relations

$$[e, f] = h, \quad [h, e] = 2e, \quad [h, f] = -2f.$$

Let $\widehat{\mathfrak{sl}}_2$ be the affine Lie algebra $\widehat{\mathfrak{sl}}_2 = \mathfrak{sl}_2[T, T^{-1}] \oplus \mathbb{C}c$ with the bracket

$$[aT^i, bT^j] = [a, b]T^{i+j} + i\langle a, b \rangle \delta_{i+j, 0} c,$$

where c is central element, $\langle a, b \rangle = \text{tr}(ab)$. Set

$$\begin{aligned} e_1 &= e, & f_1 &= f, & h_1 &= h, \\ e_2 &= fT, & f_2 &= eT^{-1}, & h_2 &= c - h. \end{aligned}$$

These are the standard Chevalley generators defining $\widehat{\mathfrak{sl}}_2$ as the Kac–Moody algebra corresponding to the Cartan matrix $\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$.

3.2 Automorphism π

The Lie algebra $\widehat{\mathfrak{sl}}_2$ has an automorphism π ,

$$\pi: c \mapsto c, \quad eT^i \mapsto fT^i, \quad fT^i \mapsto eT^i, \quad hT^i \mapsto -hT^i.$$

3.3 Verma modules

We fix $k \in \mathbb{C}$ and assume that the central element c acts on all our representations by multiplication by k .

For $m \in \mathbb{C}$, let $V(m, k - m)$ be the $\widehat{\mathfrak{sl}}_2$ Verma module with generating vector v . The Verma module is generated by v subject to the relations

$$e_1 v = 0, \quad e_2 v = 0, \quad h_1 v = m v, \quad h_2 v = (k - m) v.$$

Let $\hat{\mathfrak{n}}_- \subset \widehat{\mathfrak{sl}}_2$ be the Lie subalgebra generated by f_1, f_2 and $U\hat{\mathfrak{n}}_-$ its enveloping algebra. The map $U\hat{\mathfrak{n}}_- \rightarrow V(m, k - m)$, $F \mapsto Fv$, is an isomorphism of $U\hat{\mathfrak{n}}_-$ -modules.

The space $V(m, k - m)$ has a $\mathbb{Z}_{\geq 0}^2$ -grading: a vector $f_{i_1} \cdots f_{i_p} v$ with $i_j \in \{1, 2\}$ has degree (p_1, p_2) , if p_i is the number of i 's in the sequence i_1, \dots, i_p . For $\gamma \in \mathbb{Z}_{\geq 0}^2$, denote by $V(m, k - m)_\gamma \subset V(m, k - m)$ the corresponding γ -homogeneous component.

A homogeneous nonzero vector ω in $V(m, k - m)$, non-proportional to v , is called a *singular vector* if $e_1 \omega = e_2 \omega = 0$. The Verma module $V(m, k - m)$ is reducible, if and only if it contains a singular vector.

3.4 Reducibility conditions

See Kac–Kazhdan [3]. Set

$$\kappa = k + 2.$$

The Verma module $V(m, k - m)$ is reducible if and only if at least one of the following relations holds:

- (a) $m - l + 1 + (a - 1)\kappa = 0$,
- (b) $m + l + 1 - a\kappa = 0$,
- (c) $\kappa = 0$,

where $l, a \in \mathbb{Z}_{>0}$. If (m, κ) satisfies exactly one of the conditions (a), (b), then $V(m, k - m)$ contains a unique proper submodule, and this submodule is generated by a singular vector of degree $(la, l(a - 1))$ for condition (a) and of degree $(l(a - 1), la)$ for condition (b).

These singular vectors are highly nontrivial and are given by the following theorem.

Theorem 3.1 (Malikov–Feigin–Fuchs, [6]). *For $a, l \in \mathbb{Z}_{>0}$ and $\kappa \in \mathbb{C}$, the monomials*

$$\begin{aligned} F_{12}(l, a, \kappa) &= f_1^{l+(a-1)\kappa} f_2^{l+(a-2)\kappa} f_1^{l+(a-3)\kappa} \cdots f_2^{l-(a-2)\kappa} f_1^{l-(a-1)\kappa}, \\ F_{21}(l, a, \kappa) &= f_2^{l+(a-1)\kappa} f_1^{l+(a-2)\kappa} f_2^{l+(a-3)\kappa} \cdots f_1^{l-(a-2)\kappa} f_2^{l-(a-1)\kappa} \end{aligned}$$

are well-defined as elements of $U\hat{\mathfrak{n}}_-$. If $m = l - 1 - (a - 1)\kappa$, then $F_{12}(l, a, \kappa)v \in V(m, k - m)$ is a singular vector of degree $(la, l(a - 1))$ and if $m = -l - 1 + a\kappa$, then $F_{21}(l, a, \kappa)v \in V(m, k - m)$ is a singular vector of degree $(l(a - 1), la)$.

An explanation of the meaning of complex powers in these formulas see in [6].

For example for $m = -2 + \kappa$, we have

$$F_{21}(1, 1, \kappa)v = f_2 v = \frac{e}{T} v,$$

and for $m = -2 + 2\kappa$, we have

$$F_{21}(1, 2, \kappa)v = f_2^{1+\kappa} f_1 f_2^{1-\kappa} v = f \left(\frac{e}{T} \right)^2 v + (1 + \kappa) \frac{h}{T} \frac{e}{T} v - (1 + \kappa) \kappa \frac{e}{T^2} v.$$

3.5 Shapovalov form

The *Shapovalov form* on an $\widehat{\mathfrak{sl}}_2$ Verma module V with generating vector v is the unique symmetric bilinear form $S(\cdot, \cdot)$ on V such that

$$S(v, v) = 1, \quad S(f_i x, y) = S(x, e_i y) \quad \text{for } i = 1, 2; \ x, y \in V.$$

For $\gamma \in \mathbb{Z}_{\geq 0}^2$, let V_γ^* be the vector space dual to V_γ . Define $V^* = \bigoplus_\gamma V_\gamma^*$. The space V^* is an $\widehat{\mathfrak{sl}}_2$ -module with the $\widehat{\mathfrak{sl}}_2$ -action defined by the formulas:

$$\langle f_i \phi, x \rangle = \langle \phi, e_i x \rangle, \quad \langle e_i \phi, x \rangle = \langle \phi, f_i x \rangle,$$

where $\phi \in V^*$, $x \in V$, $i = 1, 2$. The $\widehat{\mathfrak{sl}}_2$ -module V^* is called the *contragredient* Verma module.

The Shapovalov form S considered as a map $S: V \rightarrow V^*$ is a morphism of $\widehat{\mathfrak{sl}}_2$ -modules.

3.6 Bases in V and V^*

Let V be an $\widehat{\mathfrak{sl}}_2$ Verma module V . For every $\gamma = (p_1, p_2) \in \mathbb{Z}_{\geq 0}^2$ with $p_1 \neq p_2$, we fix a basis in the homogeneous component $V_\gamma \subset V$.

For $p_1 > p_2$, we fix the basis

$$\left\{ \frac{f}{T^{i_1}} \cdots \frac{f}{T^{i_a}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_b}} \frac{e}{T^{k_1}} \cdots \frac{e}{T^{k_c}} v \right\},$$

where

$$\begin{aligned} 0 \leq i_a \leq i_{a-1} \leq \cdots \leq i_1, \quad 1 \leq j_b \leq j_{b-1} \leq \cdots \leq j_1, \quad 1 \leq k_c \leq k_{c-1} \leq \cdots \leq k_1; \\ \sum_{s=1}^a i_s + \sum_{s=1}^b j_s + \sum_{s=1}^c k_s + a - c = p_1, \quad \sum_{s=1}^a i_s + \sum_{s=1}^b j_s + \sum_{s=1}^c k_s = p_2. \end{aligned} \quad (3.1)$$

For $p_1 < p_2$, we fix the basis

$$\left\{ \frac{e}{T^{k_1}} \cdots \frac{e}{T^{k_c}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_b}} \frac{f}{T^{i_1}} \cdots \frac{f}{T^{i_a}} v \right\},$$

with the indices satisfying (3.1). Notice that for any $x \in \mathfrak{sl}_2$ the elements $\frac{x}{T^i}$ and $\frac{x}{T^j}$ commute.

These collections of vectors are bases by the Poincaré–Birkhoff–Witt theorem.

For any γ , we fix a basis in the γ -homogeneous component $V_\gamma^* \subset V^*$ as the basis dual of the basis in V_γ specified above. If $\{w_i\}$ is a basis in V_γ , then we denote by $\{(w_i)^*\}$ the dual basis in V_γ^* .

3.7 Main formula

Theorem 3.2 ([9, Theorem 5.12]). *For $m, k \in \mathbb{C}$ and $a \in \mathbb{Z}_{>0}$, the following identities hold in the contragredient Verma module $V(m, k - m)^*$,*

$$\begin{aligned} \frac{f}{T^{a-1}}(v)^* &= (m + (a-1)(k+2)) \left(\frac{f}{T^{a-1}} v \right)^* \\ &+ \sum_{\ell=1}^{a-1} \left[\frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^* + 2 \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^* \right], \\ \frac{e}{T^a}(v)^* &= (a(k+2) - m - 2) \left(\frac{e}{T^a} v \right)^* \end{aligned} \quad (3.2)$$

$$+ \sum_{\ell=0}^{a-2} \left[-\frac{h}{T^{\ell+1}} \left(\frac{e}{T^{a-\ell-1}} v \right)^* + 2\frac{f}{T^\ell} \sum_{\substack{i+j=a-\ell \\ i \geq j \geq 1}} \left(\frac{e}{T^i} \frac{e}{T^j} v \right)^* \right], \quad (3.3)$$

where v is the generating vector of the Verma module $V(m, k - m)$.

Theorem 3.2 was announced in [9]. The proof of Theorem 3.2 is the main result of this paper. The theorem is proved in Section 5.

Remark. The right-hand sides of formulas (3.2) and (3.3) have the factors $m + (a - 1)(k + 2)$ and $a(k + 2) - m - 2$. The vanishing of these factors corresponds to the resonance conditions $m_i + (a - 1)\kappa = 0$ and $m_{n+1} + 2 - a\kappa = 0$ for the de Rham complex in Section 2.3, if we recall that $\kappa = k + 2$.

Remark. Theorem 3.2 says that the action of the element $\frac{f}{T^{a-1}}$ of degree $(a, a - 1)$ on the covector $(v)^*$ can be expressed in terms of the actions of the elements $\frac{h}{T^\ell}$ and $\frac{e}{T^\ell}$ of smaller degree on some other covectors. Similarly the action of the element $\frac{e}{T^a}$ of degree $(a - 1, a)$ on the covector $(v)^*$ can be expressed in terms of the actions of the elements $\frac{h}{T^\ell}$, $\frac{f}{T^\ell}$ of smaller degree on some other covectors.

3.8 Relation to Malikov–Feigin–Fuchs vectors

Let

$$S: V(m, k - m) \rightarrow V(m, k - m)^*$$

be the Shapovalov form. Denote

$$X_a(m, k - m) = S^{-1} \left((m + (a - 1)(k + 2)) \left(\frac{f}{T^{a-1}} v \right)^* \right),$$

$$Y_a(m, k - m) = S^{-1} \left((m + 2 - a(k + 2)) \left(\frac{e}{T^a} v \right)^* \right).$$

For generic values of m and k , the Shapovalov form S is non-degenerate and X_a and Y_a are well defined elements of $V(m, k - m)$. The chosen basis in $V(m, k - m)$ allows us to compare these vectors for different values of k , m . The vectors $X_a(m, k - m)$, $Y_a(m, k - m)$ are holomorphic functions of k , m for generic k , m .

Recall the resonance lines in the (m, k) -plane, given by the equations

$$m - l + 1 + (a - 1)(k + 2) = 0, \quad m + l + 1 - a(k + 2) = 0, \quad k + 2 = 0,$$

for some $a, l \in \mathbb{Z}_{>0}$, see Section 3.4.

Theorem 3.3 ([9, Theorem 6.2]). *For $a \in \mathbb{Z}_{>0}$ let (m_0, k_0) be a point of the line $m + (a - 1)(k + 2) = 0$, which does not belong to other resonance lines. Then the function $X_a(m, k - m)$ can be analytically continued to the point (m_0, k_0) , and $X_a(m_0, k_0 - m_0)$ is a (nonzero) singular vector of $V(m_0, k_0 - m_0)$, hence it is proportional to the Malikov–Feigin–Fuchs vector $F_{12}(1, a, k_0 + 2)$.*

Similarly, for $a \in \mathbb{Z}_{>0}$ let (m_0, k_0) be a point of the line $m + 2 - a(k + 2) = 0$, which does not belong to other resonance lines. Then the function $Y_a(m, k - m)$ can be analytically continued to the point (m_0, k_0) , and $Y_a(m_0, k_0 - m_0)$ is a (nonzero) singular vector of $V(m_0, k_0 - m_0)$, hence it is proportional to the Malikov–Feigin–Fuchs vector $F_{21}(1, a, k_0 + 2)$.

4 Homomorphism of complexes

4.1 Lie algebra $\mathfrak{sl}_2(U)$

Recall that $\{z_1, \dots, z_n, z_{n+1} = \infty\}$ are pairwise distinct points of the complex projective line \mathbb{P}^1 and $U = \mathbb{P}^1 - \{z_1, \dots, z_n, z_{n+1}\}$. Fix local coordinates $t - z_1, \dots, t - z_n, 1/t$ on \mathbb{P}^1 at these points, respectively. Let $\mathfrak{sl}_2(U)$ be the Lie algebra of \mathfrak{sl}_2 -valued rational functions on \mathbb{P}^1 regular on U , with the pointwise bracket. Thus, an element of $\mathfrak{sl}_2(U)$ has the form $e \otimes u_1 + h \otimes u_2 + f \otimes u_3$ with $u_i \in \Omega^0(U)$, and the bracket is defined by the formula $[x \otimes u_1, y \otimes u_2] = [x, y] \otimes (u_1 u_2)$.

4.2 $\widehat{\mathfrak{sl}}_2(U)$ -modules

We say that an $\widehat{\mathfrak{sl}}_2$ -module W has the finiteness property, if for any $w \in W$ and $x \in \mathfrak{sl}_2$, we have $xT^j \cdot w = 0$ for all $j \gg 0$. For example, the contragradient Verma module has the finiteness property.

Let W_1, \dots, W_{n+1} be $\widehat{\mathfrak{sl}}_2$ -modules with the finiteness property. Then the Lie algebra $\mathfrak{sl}_2(U)$ acts on $W_1 \otimes \dots \otimes W_{n+1}$ by the formula

$$\begin{aligned} x \otimes u \cdot (w_1 \otimes \dots \otimes w_{n+1}) &= ([x \otimes u(t)]^{(z_1)} w_1) \otimes w_2 \otimes \dots \otimes w_{n+1} + \dots \\ &\quad + w_1 \otimes \dots \otimes w_{n-1} \otimes ([x \otimes u(t)]^{(z_n)} w_n) \otimes w_{n+1} \\ &\quad + w_1 \otimes \dots \otimes w_n \otimes (\pi([x \otimes u(t)]^{(\infty)}) w_{n+1}), \end{aligned}$$

where for $x \otimes u \in \mathfrak{sl}_2(U)$ the symbol $[x \otimes u(t)]^{(z_j)}$ denotes the Laurent expansion of $x \otimes u$ at $t = z_j$ and $[x \otimes u(t)]^{(\infty)}$ denotes the Laurent expansion at $t = \infty$; the symbol π in the last term denotes the $\widehat{\mathfrak{sl}}_2$ -automorphism defined in Section 3.2.

The finiteness property of the tensor factors ensures that the actions of the Laurent series are well-defined.

The $\widehat{\mathfrak{sl}}_2$ -action gives us a map

$$\mu: \mathfrak{sl}_2(U) \otimes \left(\otimes_{j=1}^{n+1} W_j \right) \rightarrow \otimes_{j=1}^{n+1} W_j. \quad (4.1)$$

4.3 Chain complex

For a Lie algebra \mathfrak{g} and a \mathfrak{g} -module W we denote by $C_\bullet(\mathfrak{g}, W)$ the standard chain complex of \mathfrak{g} with coefficients in W , where

$$\begin{aligned} C_p(\mathfrak{g}, W) &= \wedge^p \mathfrak{g} \otimes W, \\ d(g_p \wedge \dots \wedge g_1 \otimes w) &= \sum_{i=1}^p (-1)^{i-1} g_p \wedge \dots \wedge \widehat{g}_i \wedge \dots \wedge g_1 \otimes g_i w \\ &\quad + \sum_{1 \leq i < j \leq p} (-1)^{i+j} g_p \wedge \dots \wedge \widehat{g}_j \wedge \dots \wedge \widehat{g}_i \wedge \dots \wedge g_1 \otimes [g_j, g_i] w. \end{aligned}$$

4.4 Two complexes

4.4.1

Let $m_1, \dots, m_n, k \in \mathbb{C}$, $k + 2 \neq 0$. Define $m_{n+1} = m_1 + \dots + m_n - 2$. For $j = 1, \dots, n + 1$, let V_j be the \mathfrak{sl}_2 Verma module $V(m_j, k - m_j)$ and V_j^* the corresponding contragradient Verma module. Consider the chain complex $C_\bullet(\mathfrak{sl}_2(U), \otimes_{j=1}^{n+1} V_j^*)$ and its last two terms

$$\rightarrow \mathfrak{sl}_2(U) \otimes \left(\otimes_{j=1}^{n+1} V_j^* \right) \xrightarrow{d} \otimes_{j=1}^{n+1} V_j^* \rightarrow 0,$$

where $d = \mu$, see formula (4.1).

We assign degree 0 to the term $\otimes_{j=1}^{n+1} V_j^*$ of this complex and assign degree 1 to the differential d , so that the whole complex sits in the non-positive area.

4.4.2

Consider the twisted de Rham complex in (2.1) corresponding to $\kappa = k + 2$ with degrees shifted by 1, namely, the complex $\Omega^\bullet(U)[1]$,

$$0 \rightarrow \Omega^0(U) \xrightarrow{\partial} \Omega^1(U) \rightarrow 0,$$

where the shift [1] means that we assign degree $p - 1$ to the term $\Omega^p(U)$.

4.5 Construction

Define a linear map

$$\eta^1: \Omega^1(U) \longrightarrow \otimes_{j=1}^{n+1} V_j^*$$

by the formulas

$$\frac{dt}{(t - z_m)^a} \mapsto -\kappa(v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^{a-1}} v_m \right)^* \otimes \cdots \otimes (v_{n+1})^*, \quad (4.2)$$

$$t^{a-1} dt \mapsto \kappa(v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left(\frac{e}{T^a} v_{n+1} \right)^*, \quad (4.3)$$

for $a > 0$. Define a linear map

$$\eta^0: \Omega^0(U) \longrightarrow \mathfrak{sl}_2(U) \otimes \left(\otimes_{j=1}^{n+1} V_j^* \right)$$

by the formulas

$$\begin{aligned} \frac{1}{(t - z_m)^a} \mapsto & \frac{f}{(t - z_m)^a} \otimes (v_1)^* \otimes \cdots \otimes (v_{n+1})^* \\ & - \sum_{l=1}^a \left[\frac{e}{(t - z_m)^l} \otimes (v_1)^* \otimes \cdots \otimes 2 \sum_{\substack{i+j=a-l \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v_m \right)^* \otimes \cdots \otimes (v_{n+1})^* \right. \\ & \left. + \frac{h}{(t - z_m)^l} \otimes (v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^{a-l}} v_m \right)^* \otimes \cdots \otimes (v_{n+1})^* \right], \end{aligned} \quad (4.4)$$

for $a > 0$;

$$\begin{aligned} t^a \mapsto & ft^a \otimes (v_1)^* \otimes \cdots \otimes (v_{n+1})^* \\ & - \sum_{l=0}^{a-2} \left[et^l \otimes (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes 2 \sum_{\substack{i+j=a-l \\ i \geq j \geq 1}} \left(\frac{e}{T^i} \frac{e}{T^j} v_{n+1} \right)^* \right. \\ & \left. + ht^{l+1} \otimes (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left(\frac{e}{T^{a-l-1}} v_{n+1} \right)^* \right], \end{aligned} \quad (4.5)$$

for $a \geq 0$.

Theorem 4.1 ([9, Theorem 5.12]). *Formulas (4.2)–(4.5) define a homomorphism of complexes $\eta: \Omega^\bullet(U)[1] \rightarrow C_\bullet(\mathfrak{sl}_2(U); \otimes_{j=1}^{n+1} V_j^*)$, namely we have*

$$d\eta^0 = \eta^1 \partial.$$

The homomorphism is injective.

Theorem 4.1 was announced in [9]. Here is a proof of the theorem.

Proof. First we calculate $\eta^1(\partial((t - z_p)^{-a}))$,

$$\begin{aligned} \frac{1}{(t - z_p)^a} &\xrightarrow{\partial} -\frac{1}{\kappa}(m_p + a\kappa)\frac{dt}{(t - z_p)^{a+1}} + \frac{1}{\kappa}\sum_{k=1}^a\sum_{j \neq p} \frac{m_j}{(z_j - z_p)^k} \frac{dt}{(t - z_p)^{a+1-k}} \\ &\quad - \frac{1}{\kappa}\sum_{j \neq p} \frac{m_j}{(z_j - z_p)^a} \frac{dt}{t - z_j} \\ &\xrightarrow{\eta^1} (m_p + \kappa a)(v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^a}v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \\ &\quad - \sum_{k=1}^a\sum_{j \neq p} \frac{m_j}{(z_j - z_p)^k} (v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^{a-k}}v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \\ &\quad + \sum_{j \neq p} \frac{m_j}{(z_j - z_p)^a} (v_1)^* \otimes \cdots \otimes (fv_j)^* \otimes \cdots \otimes (v_{n+1})^*. \end{aligned}$$

Then we calculate $d(\eta^0((t - z_p)^{-a}))$,

$$\begin{aligned} \frac{1}{(t - z_p)^a} &\xrightarrow{\eta^0} \frac{f}{(t - z_p)^a} \otimes (v_1)^* \otimes \cdots \otimes (v_{n+1})^* \\ &\quad - \sum_{l=1}^a \left[\frac{h}{(t - z_p)^l} \otimes (v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^{a-l}}v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \right. \\ &\quad \left. + \frac{e}{(t - z_p)^l} \otimes 2 \sum_{\substack{i+j=a-l \\ i \geq j \geq 0}} (v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^i} \frac{f}{T^j} v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \right] \\ &\xrightarrow{\mu} (v_1)^* \otimes \cdots \otimes \left[(m_p + a(k+2)) \left(\frac{f}{T^a}v_p\right)^* \right. \\ &\quad \left. + \sum_{l=1}^a \left[\frac{h}{T^l} \left(\frac{f}{T^{a-l}}v_p\right)^* + 2\frac{e}{T^l} \sum_{\substack{i+j=a-l \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v_p\right)^* \right] \right] \otimes \cdots \otimes (v_{n+1})^* \\ &\quad + \sum_{j \neq p} \frac{m_j}{(z_j - z_p)^a} (v_1)^* \otimes \cdots \otimes (fv_j)^* \otimes \cdots \otimes (v_{n+1})^* \\ &\quad - \sum_{l=1}^a \left[(v_1)^* \otimes \cdots \otimes \frac{h}{T^l} \left(\frac{f}{T^{a-l}}v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \right. \\ &\quad \left. + (v_1)^* \otimes \cdots \otimes \sum_{\substack{i+j=a-l \\ i \geq j \geq 0}} 2\frac{e}{T^l} \left(\frac{f}{T^i} \frac{f}{T^j} v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \right. \\ &\quad \left. + \sum_{j \neq p} \frac{m_j}{(z_j - z_p)^l} (v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^{a-l}}v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \right] \\ &= (\kappa a + m_p)(v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^a}v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \\ &\quad - \sum_{l=1}^a \sum_{j \neq p} \frac{m_j}{(z_j - z_p)^l} (v_1)^* \otimes \cdots \otimes \left(\frac{f}{T^{a-l}}v_p\right)^* \otimes \cdots \otimes (v_{n+1})^* \\ &\quad + \sum_{j \neq p} \frac{m_j}{(z_j - z_p)^a} (v_1)^* \otimes \cdots \otimes (fv_j)^* \otimes \cdots \otimes (v_{n+1})^*. \end{aligned}$$

In this calculation we use formula (3.2) to express the action of $\frac{f}{T^a}$ on $(v_p)^*$. These formulas show that $d(\eta^0((t-z_p)^{-a})) = \eta^1(\partial((t-z_p)^{-a}))$.

Now we calculate $\eta^1(\partial(t^a))$,

$$\begin{aligned} t^a &\xrightarrow{\partial} \frac{1}{\kappa} \left(a\kappa - \sum_{j=1}^n m_j \right) t^{a-1} dt - \frac{1}{\kappa} \sum_{s=1}^{a-1} \sum_{j=1}^n m_j z_j^s t^{a-s-1} dt - \frac{1}{\kappa} \sum_{j=1}^n m_j z_j^a \frac{dt}{t-z_j} \\ &\xrightarrow{\eta^1} \left(a\kappa - \sum_{j=1}^n m_j \right) (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left(\frac{e}{T^a} v_{n+1} \right)^* \\ &\quad - \sum_{s=1}^{a-1} \sum_{j=1}^n m_j z_j^s (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left(\frac{e}{T^{a-s}} v_{n+1} \right)^* \\ &\quad + \sum_{j=1}^n m_j z_j^a (v_1)^* \otimes \cdots \otimes (fv_j)^* \otimes \cdots \otimes (v_{n+1})^*. \end{aligned}$$

Then we calculate $d(\eta^0(t^a))$,

$$\begin{aligned} t^a &\xrightarrow{\eta^0} ft^a \otimes (v_1)^* \otimes \cdots \otimes (v_{n+1})^* - \sum_{l=0}^{a-2} \left[ht^{l+1} \otimes (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left(\frac{e}{T^{a-l-1}} v_{n+1} \right)^* \right. \\ &\quad \left. + et^l \otimes (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes 2 \sum_{\substack{i+j=a-l \\ i \geq j \geq 1}} \left(\frac{e}{T^i} \frac{e}{T^j} v_{n+1} \right)^* \right] \\ &\xrightarrow{\mu} \sum_{j=1}^n m_j z_j^a (v_1)^* \otimes \cdots \otimes (fv_j)^* \otimes \cdots \otimes (v_{n+1})^* \\ &\quad + (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left[(-m_{n+1} - 2 + a(k+2)) \left(\frac{e}{T^a} v_{n+1} \right)^* \right. \\ &\quad \left. + \sum_{l=0}^{a-2} \left[-\frac{h}{T^{l+1}} \left(\frac{e}{T^{a-l-1}} v_{n+1} \right)^* + 2\frac{f}{T^l} \sum_{\substack{i+j=a-l \\ i \geq j \geq 1}} \left(\frac{e}{T^i} \frac{e}{T^j} v_{n+1} \right)^* \right] \right] \\ &\quad - (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \sum_{l=0}^{a-2} \left[2\frac{f}{T^l} \sum_{\substack{i+j=a-l \\ i \geq j \geq 1}} \left(\frac{e}{T^i} \frac{e}{T^j} v_{n+1} \right)^* - \frac{h}{T^{l+1}} \left(\frac{e}{T^{a-l-1}} v_{n+1} \right)^* \right] \\ &\quad - \sum_{s=1}^{a-1} \sum_{j=1}^n m_j z_j^s (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left(\frac{e}{T^{a-s}} v_{n+1} \right)^* \\ &= \left(a\kappa - \sum_{j=1}^n m_j \right) (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left(\frac{e}{T^a} v_{n+1} \right)^* \\ &\quad - \sum_{s=1}^{a-1} \sum_{j=1}^n m_j z_j^s (v_1)^* \otimes \cdots \otimes (v_n)^* \otimes \left(\frac{e}{T^{a-s}} v_{n+1} \right)^* \\ &\quad + \sum_{j=1}^n m_j z_j^a (v_1)^* \otimes \cdots \otimes (fv_j)^* \otimes \cdots \otimes (v_{n+1})^*. \end{aligned}$$

In this calculation we use formula (3.3) to express the action of $\frac{e}{T^a}$ on $(v_{n+1})^*$. Notice also that calculating the action on V_{n+1}^* we use the automorphism π , see Section 3.2. These formulas show that $d(\eta^0(t^a)) = \eta^1(\partial(t^a))$.

Clearly the maps η^1, η^2 are injective. Theorem 4.1 is proved. \blacksquare

4.6 Image of logarithmic subcomplex

Under the monomorphism η of Theorem 4.1 the image of the logarithmic subcomplex $(\Omega_{\log}^\bullet(U), \partial)$ is the chain complex $C_\bullet(\mathfrak{n}_-, \otimes_{j=1}^{n+1} V_j^*)$ of the nilpotent subalgebra $\mathfrak{n}_- \subset \mathfrak{sl}_2$ generated by f . More precisely, we have

$$\eta: 1 \mapsto f \otimes (v_1)^* \otimes \cdots \otimes (v_{n+1})^*, \quad \frac{dt}{x-t_j} \mapsto -\kappa(v_1)^* \otimes \cdots \otimes (fv_j)^* \otimes \cdots \otimes (v_{n+1})^*,$$

$j = 1, \dots, n$, and

$$\mu: f \otimes (v_1)^* \otimes \cdots \otimes (v_{n+1})^* \mapsto \sum_{j=1}^n m_j (v_1)^* \otimes \cdots \otimes (fv_j)^* \otimes \cdots \otimes (v_{n+1})^*.$$

Far-reaching generalizations of this identification of the logarithmic subcomplex with the chain complex of the nilpotent Lie algebra \mathfrak{n}_- see in [8].

5 Proof of Theorem 3.2

5.1 Formula (3.3) follows from formula (3.2)

The Lie algebra $\widehat{\mathfrak{sl}}_2$ has an automorphism ρ , corresponding to the involution of the Dynkin diagram:

$$\rho(e_i) = e_{3-i}, \quad \rho(f_i) = f_{3-i}, \quad \rho(h_i) = h_{3-i}, \quad i = 1, 2.$$

We have $\rho^2 = \text{id}$. In other words, ρ acts by the formulas

$$e \leftrightarrow fT, \quad f \leftrightarrow eT^{-1}, \quad h \leftrightarrow c - h.$$

Lemma 5.1. *For $i \in \mathbb{Z}_{>0}$, we have*

$$\rho: \frac{f}{T^i} \mapsto \frac{e}{T^{i+1}}, \quad \frac{e}{T^i} \mapsto \frac{f}{T^{i-1}}, \quad \frac{h}{T^i} \mapsto -\frac{h}{T^i}.$$

Proof. We have

$$\begin{aligned} \frac{f}{T} &= \frac{1}{2} \left[f, \frac{h}{T} \right] = \frac{1}{2} \left[f, \left[\frac{e}{T}, f \right] \right] \xrightarrow{\delta} \frac{1}{2} \left[\frac{e}{T}, \left[f, \frac{e}{T} \right] \right] = \frac{1}{2} \left[\frac{e}{T}, -\frac{h}{T} \right] = \frac{e}{T^2}, \\ \frac{f}{T^i} &= \frac{1}{2} \left[\frac{f}{T^{i-1}}, \left[\frac{e}{T}, f \right] \right] \xrightarrow{\delta} \frac{1}{2} \left[\frac{e}{T^i}, \left[f, \frac{e}{T} \right] \right] = \frac{1}{2} \left[\frac{e}{T^i}, -\frac{h}{T} \right] = \frac{e}{T^{i+1}}. \end{aligned}$$

Similarly we prove that $\rho\left(\frac{e}{T^i}\right) = \frac{f}{T^{i-1}}$, $\rho\left(\frac{h}{T^i}\right) = -\frac{h}{T^i}$. ■

For $m \in \mathbb{C}$, let $\sigma_m: \widehat{\mathfrak{sl}}_2 \rightarrow \text{End}(V(m, k-m))$ be the Verma module structure. Let $\sigma_m \circ \rho: \widehat{\mathfrak{sl}}_2 \rightarrow \text{End}(V(m, k-m))$ be the twisted module structure.

Clearly the $\widehat{\mathfrak{sl}}_2$ -modules $(\sigma_m \circ \rho, V(m, k-m))$ and $(\sigma_{m-k}, V(m-k, m))$ are isomorphic. If $v_m \in V(m, k-m)$ and $v_{k-m} \in V(k-m, m)$ are generating vectors, then an isomorphism $\chi: (\sigma_m \circ \rho, V(m, k-m)) \rightarrow (\sigma_{m-k}, V(m-k, m))$ is defined by the formula,

$$f_{i_1} \cdots f_{i_l} v_{k-m} \mapsto f_{3-i_1} \cdots f_{3-i_l} v_m,$$

for any $i_1, \dots, i_l \in \{1, 2\}$. The isomorphism χ restricts to isomorphisms of the graded components, $V(k-m, m)_{(p_1, p_2)} \rightarrow V(m, k-m)_{(p_2, p_1)}$.

In Section 3.6 we fixed bases of the homogeneous components $V_{(p_1, p_2)}$ with $p_1 \neq p_2$ of any Verma module V . By Lemma 5.1, under the isomorphism χ the chosen basis of $V(k-m, m)_{(p_1, p_2)}$ is mapped to the chosen basis of $V(m, k-m)_{(p_2, p_1)}$ up to multiplication of the basis vectors by ± 1 . This ± 1 appears due to the formula $\rho\left(\frac{h}{T^i}\right) = -\frac{h}{T^i}$. In particular, we have

$$\chi: \frac{f}{T^i} v_{k-m} \mapsto \frac{e}{T^{i+1}} v_m, \quad \frac{f}{T^i} \frac{f}{T^j} v_{k-m} \mapsto \frac{e}{T^{i+1}} \frac{e}{T^{j+1}} v_m.$$

Let $\sigma_m^*: \widehat{\mathfrak{sl}}_2 \rightarrow \text{End}(V(m, k-m)^*)$ be the contragredient Verma module structure. Let $\sigma_m^* \circ \rho: \widehat{\mathfrak{sl}}_2 \rightarrow \text{End}(V(m, k-m)^*)$ be the twisted module structure. The isomorphism χ induces an isomorphism of modules $\chi^*: (\sigma_m^* \circ \rho, V(m, k-m)^*) \rightarrow (\sigma_{m-k}^*, V(m-k, m)^*)$.

In Section 3.6 we fixed bases in the homogeneous components $V_{(p_1, p_2)}^*$ with $p_1 \neq p_2$ of any contragredient Verma module V^* . Under the isomorphism χ^* , the chosen basis of $V(k-m, m)_{(p_1, p_2)}^*$ is mapped to the chosen basis of $V(m, k-m)_{(p_2, p_1)}^*$ up to multiplication of the basis vectors by ± 1 . In particular, we have

$$\chi^*: \left(\frac{f}{T^i} v_{k-m}\right)^* \mapsto \left(\frac{e}{T^{i+1}} v_m\right)^*, \quad \left(\frac{f}{T^i} \frac{f}{T^j} v_{k-m}\right)^* \mapsto \left(\frac{e}{T^{i+1}} \frac{e}{T^{j+1}} v_m\right)^*.$$

Assume that the relation in formula (3.2) holds in every contragredient Verma module V^* . Then in $V(k-m, m)^*$ it takes the form

$$\begin{aligned} \frac{f}{T^{a-1}} (v_{k-m})^* &= (-m-2+a(k+2)) \left(\frac{f}{T^{a-1}} v_{k-m}\right)^* \\ &+ \sum_{\ell=1}^{a-1} \left[\frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v_{k-m}\right)^* + 2 \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v_{k-m}\right)^* \right]. \end{aligned}$$

The isomorphism χ^* sends this relation to the relation in $V(m, k-m)^*$,

$$\begin{aligned} \frac{e}{T^a} (v_m)^* &= (-m-2+a(k+2)) \left(\frac{e}{T^a} v_m\right)^* \\ &+ \sum_{\ell=1}^{a-1} \left[-\frac{h}{T^\ell} \left(\frac{e}{T^{a-\ell}} v_m\right)^* + 2 \frac{f}{T^{\ell-1}} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{e}{T^{i+1}} \frac{e}{T^{j+1}} v_m\right)^* \right], \end{aligned}$$

which is exactly the relation in formula (3.3). Thus formula (3.2) implies formula (3.3).

5.2 Auxiliary lemma

Let

$$V = V(m, k-m) \quad \text{and} \quad V^* = V(m, k-m)^*.$$

Lemma 5.2. For $x \in V$, $\phi \in V^*$, $k \in \mathbb{Z}_{\geq 0}$, we have

$$\left\langle \frac{f}{T^k} \varphi, x \right\rangle = \langle \varphi, eT^k x \rangle, \quad \left\langle \frac{e}{T^k} \varphi, x \right\rangle = \langle \varphi, fT^k x \rangle, \quad \left\langle \frac{h}{T^k} \varphi, x \right\rangle = \langle \varphi, hT^k x \rangle.$$

Proof. The proof is by induction. We prove the first equality, the others are proved similarly.

We have $[f_2, f_1] = \frac{h}{T}$, hence $[f_1, [f_2, f_1]] = \frac{2f}{T}$. Similarly $[e_1, [e_2, e_1]] = 2eT$. So for $k=1$, we have

$$\left\langle \frac{f}{T} \varphi, x \right\rangle = \left\langle \frac{1}{2} [f_1, [f_2, f_1]] \varphi, x \right\rangle = \left\langle \varphi, \frac{1}{2} [[e_1, e_2], e_1] x \right\rangle = \langle \varphi, eTx \rangle.$$

We have $[f_2, \frac{f}{T^{k-1}}] = \frac{h}{T^k}$, hence $[f_1, [f_2, \frac{f}{T^{k-1}}]] = \frac{2f}{T^k}$. Similarly, $[[eT^{k-1}, fT], e] = 2eT^k$. Then

$$\left\langle \frac{f}{T^k} \varphi, x \right\rangle = \left\langle \frac{1}{2} \left[f_1, \left[f_2, \frac{f}{T^{k-1}} \right] \right] \varphi, x \right\rangle = \langle \varphi, [[eT^{k-1}, e_2], e_1] x \rangle = \langle \varphi, eT^k x \rangle. \quad \blacksquare$$

5.3 The structure of the proof of formula (3.2)

We reformulate formula (3.2) as

$$(m + (a-1)(k+2)) \left(\frac{f}{T^{a-1}} v \right)^* = \frac{f}{T^{a-1}} (v)^* - \sum_{\ell=1}^{a-1} \left[\frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^* + 2 \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^* \right], \quad (5.1)$$

and will prove it in this form.

Each term in (5.1) is an element of the homogeneous component $V_{(a,a-1)}^*$. In Section 3.6 we specified a basis of the dual component $V_{(a,a-1)}$. We will calculate the value of the right-hand side in (5.1) on an arbitrary basis vector and will obtain the value of the left-hand side on that vector.

The basis in $V_{(a,a-1)}$ consists of the vectors

$$\frac{f}{T^{i_1}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v,$$

where

$$0 \leq i_r \leq i_{r-1} \leq \cdots \leq i_1, \quad 1 \leq j_s \leq j_{s-1} \leq \cdots \leq j_1, \quad 1 \leq l_{r-1} \leq l_{r-2} \leq \cdots \leq l_1;$$

$$\sum_{u=1}^r i_u + \sum_{u=1}^s j_u + \sum_{u=1}^{r-1} l_u = a-1.$$

We partition the basis in four groups. Group O consists of the single basis vector $\frac{f}{T^{a-1}} v$. Group I consists of all basis vectors with $r=1$, but different from $\frac{f}{T^{a-1}} v$. Group II consists of all basis vectors with $r=2$. Group III consists of all basis vectors with $r \geq 3$.

Notice that the value of the left-hand side of (5.1) on the basis vector $\frac{f}{T^{a-1}} v$ equals $m + (a-1)(k+2)$. Hence we need to show that the value of the right-hand side on the basis vector $\frac{f}{T^{a-1}} v$ equals $m + (a-1)(k+2)$. Similarly the value of the left-hand side on any basis vector of Groups I–III equals zero. Hence we need to prove that the value of the right-hand side on any basis vector of Groups I–III equals zero. These four statements are the content of Propositions 5.3, 5.4, 5.7, and 5.9 below. These propositions prove Theorem 3.2.

5.4 Group O

Proposition 5.3. *The value of the right-hand side of (5.1) on the basis vector $\frac{f}{T^{a-1}} v$ equals $m + (a-1)(k+2)$.*

Proof. By Lemma 5.2 we have

$$\begin{aligned} \left\langle \frac{f}{T^{a-1}} (v)^*, \frac{f}{T^{a-1}} v \right\rangle &= \left\langle (v)^*, eT^{a-1} \frac{f}{T^{a-1}} v \right\rangle \\ &= \left\langle (v)^*, \left[h + (a-1)c + \frac{f}{T^{a-1}} eT^{a-1} \right] v \right\rangle = m + (a-1)k, \end{aligned}$$

since $eT^{a-1} v$ is of degree $(-a, -a+1)$, hence zero.

By Lemma 5.2, for $\ell \in \{1, \dots, a-1\}$ we have

$$\begin{aligned} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{f}{T^{a-1}} v \right\rangle &= \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, h T^\ell \frac{f}{T^{a-1}} v \right\rangle \\ &= \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, -2 \frac{f}{T^{a-1-\ell}} v \right\rangle = -2. \end{aligned}$$

By Lemma 5.2 for $\ell \in \{1, \dots, a-1\}$ we have

$$\begin{aligned} \left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{a-1}} v \right\rangle &= \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, f T^\ell \frac{f}{T^{a-1}} v \right\rangle \\ &= \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{a-1}} f T^\ell v \right\rangle = 0, \end{aligned}$$

since $f T^\ell v$ is of degree $(-\ell+1, -\ell) \leq (0, -1)$, hence zero. Therefore,

$$\begin{aligned} \left\langle \frac{f}{T^{a-1}} (v)^* - \sum_{\ell=1}^{a-1} \left[\frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^* + 2 \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^* \right], \frac{f}{T^{a-1}} v \right\rangle \\ = m + (a-1)k - (-2)(a-1) = m + (a-1)(k+2). \end{aligned}$$

Proposition 5.3 is proved. ■

5.5 Group I

Proposition 5.4. *The value of the right-hand side of (5.1) on any basis vector of Group I equals zero.*

Proof. Group I consists of basis vectors of the form

$$w = \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v, \quad \text{where } n \in \{1, \dots, a-1\}, \quad j_1 + \cdots + j_s = n, \quad j_i \geq 1.$$

Lemma 5.5. *In the notation above, if $s = 1$, then*

$$\begin{aligned} \left\langle \frac{f}{T^{a-1}} (v)^*, w \right\rangle &= 2nk, \\ \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, w \right\rangle &= \begin{cases} 2nk, & \text{if } \ell = n, \\ -4, & \text{if } \ell > a-1-n, \\ 0, & \text{if } \ell \leq a-1-n, \end{cases} \\ \left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle &= \begin{cases} 2, & \text{if } \ell \leq n, \\ 0, & \text{if } \ell > n. \end{cases} \end{aligned} \tag{5.2}$$

Note that the first line in (5.2) is not mutually exclusive with the second and third lines in (5.2).

Proof. We have $w = \frac{f}{T^{a-1-n}} \frac{h}{T^n} v$. Then

$$\left\langle \frac{f}{T^{a-1}} (v)^*, \frac{f}{T^{a-1-n}} \frac{h}{T^n} v \right\rangle = \left\langle (v)^*, e T^{a-1} \frac{f}{T^{a-1-n}} \frac{h}{T^n} v \right\rangle$$

$$= \left\langle (v)^*, \left[hT^n + \frac{f}{T^{a-1-n}} eT^{a-1} \right] \frac{h}{T^n} v \right\rangle.$$

Note that $eT^{a-1} \frac{h}{T^n} v$ is of degree $(-a+n, -a+1+n) \leq (-1, 0)$, so that $eT^{a-1} \frac{h}{T^n} v = 0$. Hence

$$\left\langle (v)^*, hT^n \frac{h}{T^n} v \right\rangle = \left\langle (v)^*, \left[2nk + \frac{h}{T^n} hT^n \right] v \right\rangle = 2nk.$$

We have

$$\begin{aligned} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{f}{T^{a-1-n}} \frac{h}{T^n} v \right\rangle &= \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, hT^\ell \frac{f}{T^{a-1-n}} \frac{h}{T^n} v \right\rangle, \\ hT^\ell \frac{f}{T^{a-1-n}} \frac{h}{T^n} v &= \left[-2fT^{\ell+n-a+1} + \frac{f}{T^{a-1-n}} hT^\ell \right] \frac{h}{T^n} v \\ &= -2fT^{\ell+n-a+1} \frac{h}{T^n} v + \frac{f}{T^{a-1-n}} hT^\ell \frac{h}{T^n} v. \end{aligned}$$

Note that the second summand is nonzero if and only if $\ell = n$. In that case we have

$$\left\langle \left(\frac{f}{T^{a-1-n}} v \right)^*, \frac{f}{T^{a-1-n}} hT^n \frac{h}{T^n} v \right\rangle = 2nk.$$

For the first summand, if $\ell + n - a + 1 \leq 0$, then $-2fT^{\ell+n-a+1} \frac{h}{T^n} v$ is a basis vector and so pairing with $\left(\frac{f}{T^{a-1-\ell}} v \right)^*$ gives zero. If $\ell + n - a + 1 > 0$, then

$$\begin{aligned} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, -2fT^{\ell+n-a+1} \frac{h}{T^n} v \right\rangle \\ = \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, -2 \left[2 \frac{f}{T^{a-1-\ell}} + \frac{h}{T^n} fT^{\ell+n-a+1} \right] v \right\rangle = -4, \end{aligned}$$

where we used $fT^{\ell+n-a+1} v = 0$.

Finally,

$$\begin{aligned} \left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{a-1-n}} \frac{h}{T^n} v \right\rangle &= \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, fT^\ell \frac{f}{T^{a-1-n}} \frac{h}{T^n} v \right\rangle, \\ fT^\ell \frac{f}{T^{a-1-n}} \frac{h}{T^n} v &= \frac{f}{T^{a-1-n}} fT^\ell \frac{h}{T^n} v = \frac{f}{T^{a-1-n}} \left[2fT^{\ell-n} + \frac{h}{T^n} fT^\ell \right] v = 2 \frac{f}{T^{a-1-n}} \frac{f}{T^{n-\ell}} v, \end{aligned}$$

since $fT^\ell v = 0$. Note that $(a-1-n) + (n-\ell) = a-1-\ell$, hence if $i = a-1-n$ and $j = n-\ell$ (or vice versa depending on what is greater) we have

$$\left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, 2 \frac{f}{T^{a-1-n}} \frac{f}{T^{n-\ell}} v \right\rangle = 2,$$

whenever $n-\ell \geq 0$ and zero otherwise. The lemma is proved. \blacksquare

For $s = 1$ Proposition 5.4 follows from Lemma 5.5:

$$\begin{aligned} \left\langle \frac{f}{T^{a-1}} (v)^* - \sum_{\ell=1}^{a-1} \left[\frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^* + 2 \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^* \right], \frac{f}{T^{a-1-n}} \frac{h}{T^n} v \right\rangle \\ = 2nk - 2nk + 4n - 2 \cdot 2n = 0. \end{aligned}$$

Lemma 5.6. *For $s \geq 2$, we have*

$$\left\langle \frac{f}{T^{a-1}}(v)^*, w \right\rangle = 0, \quad (5.3)$$

$$\sum_{\ell=1}^{a-1} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, w \right\rangle = 0, \quad (5.4)$$

$$\sum_{\ell=1}^{a-1} \left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle = 0. \quad (5.5)$$

Proof. Recall that $w = \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v$ with $j_1 + \cdots + j_s = n$. We have

$$\begin{aligned} \left\langle \frac{f}{T^{a-1}}(v)^*, w \right\rangle &= \langle (v)^*, eT^{a-1}w \rangle, \\ eT^{a-1}w &= eT^{a-1} \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v = \left[hT^n + \frac{f}{T^{a-1-n}} eT^{a-1} \right] \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v. \end{aligned}$$

We have $hT^n \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v = 0$, since hT^n commutes with all $\frac{h}{T^{j_i}}$. Indeed, we have $n > j_i$ since $j_1 + \cdots + j_s = n$, $j_i \geq 1$, and $s \geq 2$.

We also have $\frac{f}{T^{a-1-n}} eT^{a-1} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v = 0$ since $eT^{a-1} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v$ is of degree $(-a+n, -a+1+n) \leq (-1, 0)$, hence zero. This proves (5.3).

We prove (5.4) by induction on s . For $s = 2$ we have

$$\begin{aligned} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, w \right\rangle &= \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, hT^\ell w \right\rangle, \\ hT^\ell w &= hT^\ell \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v = \left[-2fT^{\ell-a+1+n} + \frac{f}{T^{a-1-n}} hT^\ell \right] \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v \\ &= -2fT^{\ell-a+1+n} \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v + \frac{f}{T^{a-1-n}} hT^\ell \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v. \end{aligned}$$

Note that for $\frac{f}{T^{a-1-n}} hT^\ell \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v$ to give a nonzero pairing with $\left(\frac{f}{T^{a-1-\ell}} v \right)^*$ we need $\ell = n$, which implies that hT^ℓ commutes with $\frac{h}{T^{j_1}}$ and $\frac{h}{T^{j_2}}$ ($\ell > j_i$ since $j_1 + j_2 = n = \ell$ and $j_i \geq 1$), so that $\frac{f}{T^{a-1-n}} hT^\ell \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v$ gives zero for all ℓ .

Also note that whenever $\ell \leq a-1-n$, $fT^{\ell-a+1+n} \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v$ is a basis vector and so pairing with $\left(\frac{f}{T^{a-1-\ell}} v \right)^*$ gives zero. If $\ell > a-1-n$, then

$$\begin{aligned} -2fT^{\ell-a+1+n} \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v &= -2 \left[2fT^{\ell-a+1+n-j_1} + \frac{h}{T^{j_1}} fT^{\ell-a+1+n} \right] \frac{h}{T^{j_2}} v \\ &= -4fT^{\ell-a+1+n-j_1} \frac{h}{T^{j_2}} v - 2 \frac{h}{T^{j_1}} fT^{\ell-a+1+n} \frac{h}{T^{j_2}} v. \end{aligned} \quad (5.6)$$

If $\ell \leq a-1-n+j_1$, the first summand gives zero when pairing with $\left(\frac{f}{T^{a-1-\ell}} v \right)^*$, since for such $\ell fT^{\ell-a+1+n-j_1} \frac{h}{T^{j_2}} v$ is a basis vector. For $\ell > a-1-n+j_1$, we have

$$-4fT^{\ell-a+1+n-j_1} \frac{h}{T^{j_2}} v = -4 \left[2 \frac{f}{T^{a-1-\ell}} v + \frac{h}{T^{j_2}} fT^{\ell-a+1+n-j_1} \right] v = -8 \frac{f}{T^{a-1-\ell}} v,$$

since $fT^{\ell-a+1+n-j_1} v$ is of degree $(-\ell+a-1-n+j_1+1, -\ell+a-1-n+j_1) \leq (0, -1)$, hence must be equal to zero. So for $\ell \in \{a-1-n+j_1+1, \dots, a-1\}$ we get

$$\left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, -4fT^{\ell-a+1+n-j_1} \frac{h}{T^{j_2}} v \right\rangle = -8$$

and zero for other values of ℓ . The total number of elements in the set $\{a-1-n+j_1+1, \dots, a-1\}$ equals j_2 .

Whereas, for the second summand in (5.6) we have

$$\begin{aligned} -2 \frac{h}{T^{j_1}} f T^{\ell-a+1+n} \frac{h}{T^{j_2}} v &= -2 \frac{h}{T^{j_1}} \left[2f T^{\ell-a+1+n-j_2} + \frac{h}{T^{j_2}} f T^{\ell-a+1+n} \right] v \\ &= -4 \frac{h}{T^{j_1}} f T^{\ell-a+1+n-j_2} v, \end{aligned}$$

since $f T^{\ell-a+1+n} v$ is of degree $(-\ell + a - 1 - n + 1, -\ell + a - 1 - n) \leq (0, -1)$, hence must be equal to zero.

If $\ell > a - 1 - n + j_2$, then $f T^{\ell-a+1+n-j_2} v$ is of degree $(-\ell + a - 1 - n + j_2 + 1, -\ell + a - 1 - n + j_2) \leq (0, -1)$, hence must be equal to zero. If $\ell \leq a - 1 - n + j_2$, then

$$\begin{aligned} -4 \frac{h}{T^{j_1}} f T^{\ell-a+1+n-j_2} v &= -4 \left[-2 \frac{f}{T^{a-1-\ell}} + f T^{\ell-a+1+n-j_2} \frac{h}{T^{j_1}} \right] v \\ &= 8 \frac{f}{T^{a-1-\ell}} - 4 f T^{\ell-a+1+n-j_2} \frac{h}{T^{j_1}}. \end{aligned}$$

The second summand gives zero when pairing with $(\frac{f}{T^{a-1-\ell}} v)^*$. So for $\ell \in \{a-1-n+1, \dots, a-1-n+j_2\}$ we get

$$\left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, -2 \frac{h}{T^{j_1}} f T^{\ell-a+1+n} \frac{h}{T^{j_2}} v \right\rangle = 8$$

and zero for other values of ℓ . The total number of elements in the set $\{a-1-n+1, \dots, a-1-n+j_2\}$ equals j_2 .

Therefore,

$$\sum_{\ell=1}^{a-1} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v \right\rangle = -8j_2 + 8j_2 = 0$$

and so for $s = 2$ we proved (5.4).

Now suppose that (5.4) holds for all natural numbers up to s . Then

$$h T^\ell \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} v = \left[-2 f T^{\ell-a+1+n} + \frac{f}{T^{a-1-n}} h T^\ell \right] \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} v.$$

Note that for $\frac{f}{T^{a-1-n}} h T^\ell \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} v$ to give a nonzero pairing with $(\frac{f}{T^{a-1-\ell}} v)^*$ we need $\ell = n$. That assumption implies that $h T^\ell$ commutes with $\frac{h}{T^{j_i}}$ for all $i \in \{1, \dots, s+1\}$ since $\ell > j_i$ as $j_1 + \dots + j_{s+1} = n = \ell$ and $j_i \geq 1$. Hence $\frac{f}{T^{a-1-n}} h T^\ell \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} v$ gives zero for all ℓ .

Also note that whenever $\ell \leq a - 1 - n$, the vector $f T^{\ell-a+1+n} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} v$ is a basis vector and so pairing with $(\frac{f}{T^{a-1-\ell}} v)^*$ gives zero.

If $\ell > a - 1 - n$, then

$$\begin{aligned} -2 f T^{\ell-a+1+n} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} v &= -2 \left[2f T^{\ell-a+1+n-j_1} + \frac{h}{T^{j_1}} f T^{\ell-a+1+n} \right] \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \\ &= -4 \frac{f}{T^{a-1-(n-j_1)-\ell}} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v - 2 \frac{h}{T^{j_1}} f T^{-a+1+n+\ell} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v. \end{aligned} \quad (5.7)$$

Note that by induction hypothesis we have

$$0 = \sum_{\ell=1}^{a-1} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{f}{T^{a-1-(n-j_1)}} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle$$

$$= \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, h T^\ell \frac{f}{T^{a-1-(n-j_1)}} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle.$$

So we add this zero term multiplied by -2 to the first summand in (5.7) to get

$$\begin{aligned} & -2 \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, 2 \frac{f}{T^{a-1-(n-j_1)-\ell}} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \\ & \quad - 2 \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, h T^\ell \frac{f}{T^{a-1-(n-j_1)}} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \\ & = -2 \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \left[2 \frac{f}{T^{a-1-(n-j_1)-\ell}} + h T^\ell \frac{f}{T^{a-1-(n-j_1)}} \right] \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \\ & = -2 \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{f}{T^{a-1-(n-j_1)}} h T^\ell \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle, \end{aligned} \tag{5.8}$$

where in the last step we use commutation relations.

Note that for $\frac{f}{T^{a-1-(n-j_1)}} h T^\ell \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v$ to give a nonzero pairing with $\left(\frac{f}{T^{a-1-\ell}} v \right)^*$ we need $\ell = n - j_1$. That assumption implies that $h T^\ell$ commutes with $\frac{h}{T^{j_i}}$ for all $i \in \{2, \dots, s+1\}$ since $\ell > j_i$ as $j_2 + \cdots + j_{s+1} = n - j_1 = \ell$ and $j_i \geq 1$. Hence $\frac{f}{T^{a-1-(n-j_1)}} h T^\ell \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v$ gives zero for all ℓ .

For the second summand in (5.7) we have

$$\begin{aligned} & -2 \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{h}{T^{j_1}} f T^{-a+1+n+\ell} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \\ & = -2 \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{h}{T^{j_1}} \left[2 \frac{f}{T^{a-1-(n-j_2)-\ell}} + \frac{h}{T^{j_2}} f T^{-a+1+n+\ell} \right] \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \\ & = -4 \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{h}{T^{j_1}} \frac{f}{T^{a-1-(n-j_2)-\ell}} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \end{aligned} \tag{5.9}$$

$$- 2 \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} f T^{-a+1+n+\ell} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle. \tag{5.10}$$

In (5.9) we note that

$$\begin{aligned} & \frac{h}{T^{j_1}} \frac{f}{T^{a-1-(n-j_2)-\ell}} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \\ & = \left[-2 \frac{f}{T^{a-1-(n-j_1-j_2)-\ell}} + \frac{f}{T^{a-1-(n-j_2)-\ell}} \frac{h}{T^{j_1}} \right] \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \\ & = -2 \frac{f}{T^{a-1-(n-j_1-j_2)-\ell}} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v + \frac{f}{T^{a-1-(n-j_2)-\ell}} \frac{h}{T^{j_1}} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v. \end{aligned}$$

Note that in both terms the number of h 's is less than or equal to s , so we use the exact same reasoning as in (5.8) to show that

$$\begin{aligned} & \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{f}{T^{a-1-(n-j_1-j_2)-\ell}} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle = 0, \\ & \sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{f}{T^{a-1-(n-j_2)-\ell}} \frac{h}{T^{j_1}} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle = 0, \end{aligned}$$

which implies that the expression in (5.9) equals zero. Similarly, one shows that in (5.10),

$$\sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} f T^{-a+1+n+\ell} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle$$

the factor $f T^{-a+1+n+\ell}$ can be pulled to the right by using the same argument (first commute $f T^{-a+1+n+\ell}$ with $\frac{h}{T^{j_3}}$ and then pull $f T^{-a+1+n-j_3+\ell}$ to the left). Ultimately, we get

$$\sum_{\ell=1}^{a-1} \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} f T^{-a+1+n+\ell} v \right\rangle = 0,$$

since $\ell > a - 1 - n$ and so $f T^{-a+1+n+\ell} v = 0$. Therefore,

$$\sum_{\ell=1}^{a-1} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v \right\rangle = 0,$$

and formula (5.4) is proved.

We prove formula (5.5) by induction on s . For $s = 2$, we have

$$\left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle = \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, f T^\ell w \right\rangle,$$

$$f T^\ell w = f T^\ell \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v = \frac{f}{T^{a-1-n}} f T^\ell \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v.$$

Note that $f T^\ell \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} v$ is of degree $(n - \ell + 1, n - \ell)$, hence nonzero only if $\ell \leq n$. For such ℓ we have

$$\begin{aligned} \frac{f}{T^{a-1-n}} \left[2f T^{\ell-j_1} + \frac{h}{T^{j_1}} f T^\ell \right] \frac{h}{T^{j_2}} v &= 2 \frac{f}{T^{a-1-n}} f T^{\ell-j_1} \frac{h}{T^{j_2}} v + \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} f T^\ell \frac{h}{T^{j_2}} v \\ &= 2 \frac{f}{T^{a-1-n}} f T^{\ell-j_1} \frac{h}{T^{j_2}} v + 2 \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} f T^{\ell-j_2} v. \end{aligned} \quad (5.11)$$

If $\ell \leq j_1$, then the first summand in (5.11) gives zero when pairing with any vector with two f 's. If $\ell > j_1$, then

$$2 \frac{f}{T^{a-1-n}} f T^{\ell-j_1} \frac{h}{T^{j_2}} v = 2 \frac{f}{T^{a-1-n}} \left[2f T^{\ell-j_1-j_2} + \frac{h}{T^{j_2}} f T^{\ell-j_1} \right] v = 4 \frac{f}{T^{a-1-n}} \frac{f}{T^{n-\ell}} v.$$

If $\ell > j_2$, then the second summand in (5.11) is zero simply because $f T^{\ell-j_2} v = 0$. If $\ell \leq j_2$, then

$$2 \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} f T^{\ell-j_2} v = 2 \frac{f}{T^{a-1-n}} \left[-2f T^{\ell-j_1-j_2} + f T^{\ell-j_2} \frac{h}{T^{j_1}} \right] v = -4 \frac{f}{T^{a-1-n}} \frac{f}{T^{n-\ell}} v,$$

since $\frac{f}{T^{a-1-n}} f T^{\ell-j_2} \frac{h}{T^{j_1}} v$ is a basis vector, hence pairing with a vector consisting of two f 's gives zero. Therefore,

$$\sum_{\ell=1}^{a-1} \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, f T^\ell w \right\rangle$$

$$= \sum_{\ell=1}^{a-1} \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, 4 \frac{f}{T^{a-1-n}} \frac{f}{T^{n-\ell}} v \right\rangle \quad (5.12)$$

$$+ \sum_{\ell=1}^{a-1} \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, -4 \frac{f}{T^{a-1-n}} \frac{f}{T^{n-\ell}} v \right\rangle. \quad (5.13)$$

Note that in the expression in (5.12) for each $\ell \in \{j_1 + 1, \dots, n\}$ there exists exactly one pair of indices $(i, j) = (\max\{a-1-n, n-\ell\}, \min\{a-1-n, n-\ell\})$ that gives 4 when pairing. All other pairs (i, j) give zero. Similarly, the expression in (5.13) equals -4 for each $\ell \in \{1, \dots, j_2\}$ and exactly one corresponding pair (i, j) , and zero otherwise. Also note that the number of elements in each set $\{j_1 + 1, \dots, n\}$ and $\{1, \dots, j_2\}$ equals j_2 . Hence we get

$$4j_2 - 4j_2 = 0.$$

Therefore, formula (5.5) is proved for $s = 2$.

Now suppose that formula (5.5) holds for all natural numbers up to s . Then

$$\begin{aligned} fT^\ell \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} v &= \frac{f}{T^{a-1-n}} fT^\ell \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} v \\ &= \frac{f}{T^{a-1-n}} \left[2fT^{\ell-j_1} + \frac{h}{T^{j_1}} fT^\ell \right] \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \\ &= 2 \frac{f}{T^{a-1-n}} fT^{\ell-j_1} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v + \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} fT^\ell \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v. \end{aligned} \quad (5.14)$$

Note that if $\ell \leq j_1$, then the first summand in (5.14) is a basis vector and hence its pairing with a vector consisting of two f 's gives zero. If $\ell > j_1$ we have

$$\begin{aligned} &\sum_{\ell=1}^{a-1} \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{a-1-n}} fT^{\ell-j_1} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \\ &= \sum_{\ell=j_1+1}^{a-1} \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{a-1-n}} fT^{\ell-j_1} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \\ &= \sum_{\ell=j_1+1}^{a-1} \left\langle \frac{e}{T^{\ell-j_1}} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{a-1-n}} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle \\ &= \sum_{k=1}^{a-1-j_1} \left\langle \frac{e}{T^k} \sum_{\substack{i+j=a-1-j_1-k \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{(a-1-j_1)-(n-j_1)}} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle = 0 \end{aligned}$$

by induction hypothesis. For the second summand in (5.14) we have

$$\begin{aligned} \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} fT^\ell \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_{s+1}}} v &= \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \left[2fT^{\ell-j_2} + \frac{h}{T^{j_2}} fT^\ell \right] \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \\ &= 2 \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} fT^{\ell-j_2} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v + \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} fT^\ell \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v. \end{aligned} \quad (5.15)$$

Note that

$$\frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} fT^{\ell-j_2} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v = \frac{f}{T^{a-1-n}} \left[-2fT^{\ell-j_1-j_2} + fT^{\ell-j_2} \frac{h}{T^{j_1}} \right] \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v$$

$$= -2 \frac{f}{T^{a-1-n}} f T^{\ell-j_1-j_2} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v + \frac{f}{T^{a-1-n}} f T^{\ell-j_2} \frac{h}{T^{j_1}} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v,$$

where in each vector the number of h 's is less than or equal to s . Repeating the argument above, we see that by induction hypothesis we get

$$\sum_{\ell=1}^{a-1} \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} f T^{\ell-j_2} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v \right\rangle = 0.$$

Now in the second summand in (5.15),

$$\frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \frac{h}{T^{j_2}} f T^{\ell} \frac{h}{T^{j_3}} \cdots \frac{h}{T^{j_{s+1}}} v,$$

we pull $f T^{\ell}$ to the right and at each step we use induction hypothesis to argue that we keep getting zeros. Ultimately, we get a vector

$$\frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_{s+1}}} f T^{\ell} v,$$

which is zero, since $f T^{\ell}$ has grading $(-\ell + 1, -\ell) \leq (0, -1)$ and so $f T^{\ell} v = 0$. Therefore,

$$\sum_{\ell=1}^{a-1} \left\langle \frac{e}{T^{\ell}} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, \frac{f}{T^{a-1-n}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} v \right\rangle = 0.$$

Formula (5.5) and Lemma 5.6 are proved. ■

Proposition 5.4 is proved. ■

5.6 Group II

Proposition 5.7. *The value on the right-hand side of (5.1) on any basis vector from Group II equals zero.*

Proof. Group II consists of vectors

$$w = \frac{f}{T^{i_1}} \frac{f}{T^{i_2}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{\ell}} v.$$

Lemma 5.8. *We have*

$$\left\langle \frac{f}{T^{a-1}} (v)^*, w \right\rangle = 2^{s+1} (m - lk), \quad (5.16)$$

$$\left\langle \frac{h}{T^{\ell}} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, w \right\rangle = \begin{cases} 2^{s+2} (m - lk), & \text{if } i_1 = i_2 = a - 1 - \ell, \\ 2^{s+1} (m - lk), & \text{if } i_1 \neq i_2 \text{ and } i_1 \text{ or } i_2 = a - 1 - \ell, \\ 0, & \text{otherwise,} \end{cases} \quad (5.17)$$

$$\left\langle \frac{e}{T^{\ell}} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle = \begin{cases} -2^s (m - lk), & \text{if } \ell = a - 1 - i_1 - i_2, \\ 0, & \text{otherwise.} \end{cases} \quad (5.18)$$

Proof. We have

$$\begin{aligned} \left\langle \frac{f}{T^{a-1}}(v)^*, w \right\rangle &= \langle (v)^*, eT^{a-1}w \rangle = \left\langle (v)^*, \left[hT^{a-1-i_1} + \frac{f}{T^{i_1}}eT^{a-1} \right] \frac{f}{T^{i_2}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v \right\rangle \\ &= \left\langle (v)^*, hT^{a-1-i_1} \frac{f}{T^{i_2}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v \right\rangle + \left\langle (v)^*, \frac{f}{T^{i_1}} eT^{a-1} \frac{f}{T^{i_2}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v \right\rangle. \end{aligned} \quad (5.19)$$

Note that $eT^{a-1} \frac{f}{T^{i_2}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l}$ is of degree $(-i_1 - 1, -i_1) \leq (-1, 0)$, hence $eT^{a-1} \frac{f}{T^{i_2}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v = 0$. In the first summand in (5.19) we pull hT^{a-1-i_1} to the right to get

$$\begin{aligned} \left\langle (v)^*, -2fT^{a-1-i_1-i_2} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v \right\rangle &= \cdots = \left\langle (v)^*, -2^{s+1}fT^{a-1-i_1-i_2-j_1-\cdots-a_s} \frac{e}{T^l} v \right\rangle \\ &= \left\langle (v)^*, -2^{s+1}fT^l \frac{e}{T^l} v \right\rangle = \langle (v)^*, 2^{s+1}(h-lc)v \rangle = 2^{s+1}(m-lk), \end{aligned}$$

where at each step we do not write monomials of negative degree, since they give zero when applied to v . This proves formula (5.16).

We have

$$\begin{aligned} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, w \right\rangle &= \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, hT^\ell w \right\rangle, \\ hT^\ell w &= \left[-2fT^{\ell-i_1} + \frac{f}{T^{i_1}}hT^\ell \right] \frac{f}{T^{i_2}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v \\ &= -2fT^{\ell-i_1} \frac{f}{T^{i_2}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v + \frac{f}{T^{i_1}} \left[-2fT^{\ell-i_2} + \frac{f}{T^{i_2}}hT^\ell \right] \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v \\ &= \left[-2\frac{f}{T^{i_2}}fT^{\ell-i_1} - 2\frac{f}{T^{i_1}}fT^{\ell-i_2} + \frac{f}{T^{i_1}}\frac{f}{T^{i_2}}hT^\ell \right] \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v. \end{aligned}$$

Note that the vector $\frac{f}{T^{i_1}} \frac{f}{T^{i_2}} hT^\ell \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v$ after pulling hT^ℓ to the right either becomes a zero vector or a vector with two f 's, which of course gives zero when pairing with a basis vector with one f . Also, note that the only possibility for the vector $\frac{f}{T^{i_2}} fT^{\ell-i_1} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v$ to give a nonzero pairing with $\left(\frac{f}{T^{a-1-\ell}} v \right)^*$ is when $i_2 = a-1-\ell$. Similarly $\frac{f}{T^{i_1}} fT^{\ell-i_2} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v$ gives a nonzero number only if $i_1 = a-1-\ell$. First consider the case $i_1 = i_2 = a-1-\ell$. We have

$$\begin{aligned} -4\frac{f}{T^{a-1-\ell}}fT^{2\ell-a+1} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v \\ = -4\frac{f}{T^{a-1-\ell}} \left[2fT^{2\ell-a+1-j_1} + \frac{h}{T^{j_1}}fT^{2\ell-a+1} \right] \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v. \end{aligned}$$

Note that $fT^{2\ell-a+1} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l}$ is of degree $(-j_1, -j_1) \leq (-1, -1)$, hence

$$fT^{2\ell-a+1} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v = 0.$$

So we get

$$\begin{aligned} -8\frac{f}{T^{a-1-\ell}}fT^{2\ell-a+1-j_1} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^l} v \cdots &= -2^{s+2}\frac{f}{T^{a-1-\ell}}fT^{2\ell-a+1-j_1-\cdots-j_s} \frac{e}{T^l} v \\ &= -2^{s+2}\frac{f}{T^{a-1-\ell}}fT^l \frac{e}{T^l} v = 2^{s+2}\frac{f}{T^{a-1-\ell}}(h-lc)v = 2^{s+2}(m-lk), \end{aligned}$$

where at each step we don't write monomials of negative degree, since they give zero when applied to v .

For $i_1 = a - 1 - \ell \neq i_2$ and $i_2 = a - 1 - \ell \neq i_1$ we have

$$-2 \frac{f}{T^{a-1-\ell}} f T^{2\ell-a+1} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^\ell} v = 2^{s+1}(m - lk),$$

where we performed the exact same computation as above. Therefore, formula (5.17) is proved.

We have

$$\begin{aligned} \left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle &= \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, f T^\ell w \right\rangle, \\ f T^\ell w &= \frac{f}{T^{i_1}} \frac{f}{T^{i_2}} f T^\ell \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^\ell} v = \frac{f}{T^{i_1}} \frac{f}{T^{i_2}} \left[2 f T^{\ell-j_1} + \frac{h}{T^{j_1}} f T^\ell \right] \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^\ell} v. \end{aligned}$$

The only nonzero pairing happens when ℓ is such that $i_1 + i_2 = a - 1 - \ell$. In that case $f T^\ell \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^\ell} v$ has degree $(-j_1, -j_1) \leq (-1, -1)$, hence $f T^\ell \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^\ell} v = 0$. Therefore we have

$$2 \frac{f}{T^{i_1}} \frac{f}{T^{i_2}} f T^{\ell-j_1} \frac{h}{T^{j_2}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^\ell} v = 2^s \frac{f}{T^{i_1}} \frac{f}{T^{i_2}} f T^{\ell-j_1-\cdots-j_s} \frac{e}{T^\ell} v = 2^s \frac{f}{T^{i_1}} \frac{f}{T^{i_2}} f T^\ell \frac{e}{T^\ell} v,$$

where we pulled $f T^{\ell-j_1}$ to the right and did not write monomials of negative degree, since they give zero when applied to v . Hence we get

$$2^s \frac{f}{T^{i_1}} \frac{f}{T^{i_2}} (-h + lc) v.$$

Therefore,

$$\begin{aligned} \left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle \\ = \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left\langle \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, 2^s \frac{f}{T^{i_1}} \frac{f}{T^{i_2}} (-h + lc) v \right\rangle = -2^s(m - lk), \end{aligned}$$

since for $i = i_1, j = i_2$ we get $-2^s(m - lk)$ and zero for other pairs (i, j) . Formula (5.18) and Lemma 5.8 are proved. \blacksquare

By Lemma 5.8, we have

$$\begin{aligned} \left\langle \frac{f}{T^{a-1}} (v)^* - \sum_{\ell=1}^{a-1} \left[\frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^* + 2 \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^* \right], w \right\rangle \\ = 2^{s+1}(m - lk) - 2 \cdot 2^{s+1}(m - lk) + 2 \cdot 2^s(m - lk) = 0. \end{aligned}$$

Note that

$$\left\langle \sum_{\ell=1}^{a-1} \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, w \right\rangle = 2 \cdot 2^{s+1}(m - lk)$$

in both cases $i_1 = i_2$ and $i_1 \neq i_2$. Also note that

$$\left\langle \sum_{\ell=1}^{a-1} \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle \neq 0$$

only if ℓ is such that $i_1 + i_2 = a - 1 - \ell$. Therefore, Proposition 5.7 is proved. \blacksquare

5.7 Group III

Proposition 5.9. *The value of the right-hand side of (5.1) on any basis vector of Group III equals zero.*

Proof. A vector in Group III has the form

$$w = \frac{f}{T^{i_1}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v,$$

where $r \geq 3$.

Lemma 5.10. *For every $\ell \in \{1, \dots, a-1\}$, we have*

$$\left\langle \frac{f}{T^{a-1}}(v)^*, w \right\rangle = 0, \quad (5.20)$$

$$\left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, w \right\rangle = 0, \quad (5.21)$$

$$\left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle = 0. \quad (5.22)$$

Proof. We have

$$\begin{aligned} \left\langle \frac{f}{T^{a-1}}(v)^*, w \right\rangle &= \langle (v)^*, eT^{a-1}w \rangle \\ &= \left\langle (v)^*, \left[hT^{a-1-i_1} + \frac{f}{T^{i_1}} eT^{a-1} \right] \frac{f}{T^{i_2}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v \right\rangle. \end{aligned}$$

Note that $eT^{a-1} \frac{f}{T^{i_2}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v$ is of degree $(-i_1 - 1, -i_1) \leq (-1, 0)$, hence zero. So we have

$$\left\langle (v)^*, \left[-2fT^{a-1-i_1-i_2} + \frac{f}{T^{i_2}} hT^{a-1-i_1} \right] \frac{f}{T^{i_3}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v \right\rangle.$$

As above, note that $hT^{a-1-i_1} \frac{f}{T^{i_3}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v$ is of degree $(-i_2 - 1, -i_2) \leq (-1, 0)$, hence zero. Therefore we obtain

$$\left\langle (v)^*, -2 \frac{f}{T^{i_3}} \cdots \frac{f}{T^{i_r}} fT^{a-1-i_1-i_2} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v \right\rangle = 0,$$

since $fT^{a-1-i_1-i_2} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v$ is of degree $(-i_3 - \cdots - i_r - r + 2, -i_2 - \cdots - i_r) \leq (-1, 0)$ for $r \geq 3$, hence zero. Formula (5.20) is proved.

We have

$$\begin{aligned} \left\langle \frac{h}{T^\ell} \left(\frac{f}{T^{a-1-\ell}} v \right)^*, w \right\rangle &= \left\langle \left(\frac{f}{T^{a-1-\ell}} v \right)^*, hT^\ell w \right\rangle, \\ hT^\ell w &= \left[-2fT^{\ell-i_1} + \frac{f}{T^{i_1}} hT^\ell \right] \frac{f}{T^{i_2}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v \\ &= -2 \frac{f}{T^{i_2}} \cdots \frac{f}{T^{i_r}} fT^{\ell-i_1} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v \\ &\quad + \frac{f}{T^{i_1}} hT^\ell \frac{f}{T^{i_2}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v. \end{aligned} \quad (5.23)$$

$$+ \frac{f}{T^{i_1}} hT^\ell \frac{f}{T^{i_2}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v. \quad (5.24)$$

Observe that for $\ell \leq i_1$ in (5.23) we have a basis vector, hence it gives zero when pairing with $(\frac{f}{T^{a-1-\ell}}v)^*$. If $\ell > i_1$, then we pull $fT^{\ell-i_1}$ to the right and notice that no matter how $fT^{\ell-i_1}$ interacts with h 's and e 's, it does not affect the number of f 's, which is greater or equal than two. Hence, the vector in (5.23) gives zero when pairing with $(\frac{f}{T^{a-1-\ell}}v)^*$.

In (5.24) note that

$$hT^\ell \frac{f}{T^{i_2}} = -2fT^{\ell-i_2} + \frac{f}{T^{i_2}} hT^\ell$$

so that either hT^ℓ is pulled to the right not affecting the number of f 's or it gives $fT^{\ell-i_2}$, for which we apply the same argument as above after pulling it to the right to argue that the pairing of the vector in (5.24) with $(\frac{f}{T^{a-1-\ell}}v)^*$ is zero. Formula (5.21) is proved.

We have

$$\begin{aligned} \left\langle \frac{e}{T^\ell} \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, w \right\rangle &= \left\langle \sum_{\substack{i+j=a-1-\ell \\ i \geq j \geq 0}} \left(\frac{f}{T^i} \frac{f}{T^j} v \right)^*, fT^\ell w \right\rangle, \\ fT^\ell w &= fT^\ell \frac{f}{T^{i_1}} \cdots \frac{f}{T^{i_r}} \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v \\ &= \frac{f}{T^{i_1}} \cdots \frac{f}{T^{i_r}} fT^\ell \frac{h}{T^{j_1}} \cdots \frac{h}{T^{j_s}} \frac{e}{T^{l_1}} \cdots \frac{e}{T^{l_{r-1}}} v. \end{aligned}$$

As in formula (5.21), no matter how fT^ℓ interacts with h 's and e 's, the number of f 's remains unchanged, i.e., we have more than or equal to three f 's, so that pairing with $(\frac{f}{T^i} \frac{f}{T^j} v)^*$ is zero. Formula (5.22) is proved. ■

Proposition 5.9 follows from Lemma 5.10. ■

Theorem 3.2 is proved.

Acknowledgements

The authors thank V. Schechtman for useful discussions. The second author was supported in part by NSF grant DMS-1665239.

References

- [1] Feigin B., Schechtman V., Varchenko A., On algebraic equations satisfied by hypergeometric correlators in WZW models. I, *Comm. Math. Phys.* **163** (1994), 173–184.
- [2] Feigin B., Schechtman V., Varchenko A., On algebraic equations satisfied by hypergeometric correlators in WZW models. II, *Comm. Math. Phys.* **170** (1995), 219–247, [arXiv:hep-th/9407010](#).
- [3] Kac V.G., Kazhdan D.A., Structure of representations with highest weight of infinite-dimensional Lie algebras, *Adv. Math.* **34** (1979), 97–108.
- [4] Khoroshkin S., Schechtman V., Factorizable \mathcal{D} -modules, *Math. Res. Lett.* **4** (1997), 239–257, [arXiv:q-alg/9611018](#).
- [5] Khoroshkin S., Varchenko A., Quiver \mathcal{D} -modules and homology of local systems over an arrangement of hyperplanes, *Int. Math. Res. Pap.* **2006** (2006), 69590, 116 pages, [arXiv:math.QA/0510451](#).
- [6] Malikov F., Feigin B., Fuks D., Singular vectors in Verma modules over Kac–Moody algebras, *Funct. Anal. Appl.* **20** (1986), 103–113.
- [7] Schechtman V., Terao H., Varchenko A., Local systems over complements of hyperplanes and the Kac–Kazhdan conditions for singular vectors, *J. Pure Appl. Algebra* **100** (1995), 93–102, [arXiv:hep-th/9411083](#).
- [8] Schechtman V., Varchenko A., Arrangements of hyperplanes and Lie algebra homology, *Invent. Math.* **106** (1991), 139–194.
- [9] Schechtman V., Varchenko A., Rational differential forms on the line and singular vectors in Verma modules over $\widehat{\mathfrak{sl}}_2$, *Mosc. Math. J.* **17** (2017), 787–802, [arXiv:1511.09014](#).