

# Rigidity of proper holomorphic mappings between generalized Fock-Bargmann-Hartogs domains

Enchao Bi<sup>1\*</sup> & Zhenhan Tu<sup>2</sup>

<sup>1</sup>School of Mathematics and Statistics, Qingdao University, Qingdao, Shandong 266071, P.R. China

<sup>2</sup>School of Mathematics and Statistics, Wuhan University, Wuhan, Hubei 430072, P.R. China

Email: bienchao@whu.edu.cn (E. Bi), zhhtu.math@whu.edu.cn (Z. Tu)

**Abstract** A generalized Fock-Bargmann-Hartogs domain  $D_n^{\mathbf{m}, \mathbf{p}}$  is defined as a domain fibered over  $\mathbb{C}^n$  with the fiber over  $z \in \mathbb{C}^n$  being a generalized complex ellipsoid  $\Sigma_z(\mathbf{m}, \mathbf{p})$ . In general, a generalized Fock-Bargmann-Hartogs domain is an unbounded non-hyperbolic domains without smooth boundary. The main contribution of this paper is as follows. By using the explicit formula of Bergman kernels of the generalized Fock-Bargmann-Hartogs domains, we obtain the rigidity results of proper holomorphic mappings between two equidimensional generalized Fock-Bargmann-Hartogs domains. We therefore exhibit an example of unbounded weakly pseudoconvex domains on which the rigidity results of proper holomorphic mappings can be built.

**Key words:** Automorphism groups, Bergman kernels, Generalized Fock-Bargmann-Hartogs domains, Proper holomorphic mappings

**Mathematics Subject Classification (2010):** 32A07· 32A25· 32H35· 32M05

## 1 Introduction

A holomorphic map  $F : \Omega_1 \rightarrow \Omega_2$  between two domains  $\Omega_1, \Omega_2$  in  $\mathbb{C}^n$  is said to be proper if  $F^{-1}(K)$  is compact in  $\Omega_1$  for every compact subset  $K \subset \Omega_2$ . In particular, an automorphism  $F : \Omega \rightarrow \Omega$  of a domain  $\Omega$  in  $\mathbb{C}^n$  is a proper holomorphic mapping of  $\Omega$  into  $\Omega$ . There are many works about proper holomorphic mappings between various bounded domains with some requirements of the boundary (e.g., Bedford-Bell [3], Diederich-Fornaess [8], Dini-Primicerio [9] and Tu-Wang [24]). However, very little seems to be known about proper holomorphic mapping between the unbounded weakly pseudoconvex domains. There are also some works about automorphism groups of hyperbolic domains (e.g., Isaev [10], Isaev-Krantz [11] and Kim-Verdiani [14]). In this paper, we mainly focus our attention on some unbounded non-hyperbolic weakly pseudoconvex domains.

The Fock-Bargmann-Hartogs domain  $D_{n,m}(\mu)$  is defined by

$$D_{n,m}(\mu) = \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^m : \|w\|^2 < e^{-\mu\|z\|^2}\} \text{ for } \mu > 0,$$

where  $\|\cdot\|$  is the standard Hermitian norm. The Fock-Bargmann-Hartogs domains  $D_{n,m}(\mu)$  are strongly pseudoconvex domains in  $\mathbb{C}^{n+m}$  with smooth real-analytic boundary. We note that each  $D_{n,m}(\mu)$  contains  $\{(z, 0) \in \mathbb{C}^n \times \mathbb{C}^m\} \cong \mathbb{C}^n$ . Thus each  $D_{n,m}(\mu)$  is not hyperbolic in the sense of Kobayashi and  $D_{n,m}(\mu)$  can not be biholomorphic to any bounded domain in  $\mathbb{C}^{n+m}$ . Therefore, each Fock-Bargmann-Hartogs domain  $D_{n,m}(\mu)$  is an unbounded non-hyperbolic domain in  $\mathbb{C}^{n+m}$ .

In 2013, Yamamori [25] gave an explicit formula for the Bergman kernels of the Fock-Bargmann-Hartogs domains in terms of the polylogarithm functions. In 2014, by checking that the Bergman

---

\*Corresponding author.

kernel ensures revised the Cartan's theorem, Kim-Ninh-Yamamori [13] determined the automorphism group of the Fock-Bargmann-Hartogs domains as follows.

**Theorem 1.1** (Kim-Ninh-Yamamori [13]). *The automorphism group  $\text{Aut}(D_{n,m}(\mu))$  is exactly the group generated by all automorphisms of  $D_{n,m}(\mu)$  as follows:*

$$\begin{aligned}\varphi_U &: (z, w) \mapsto (Uz, w), \quad U \in \mathcal{U}(n); \\ \varphi_{U'} &: (z, w) \mapsto (z, U'w), \quad U' \in \mathcal{U}(m); \\ \varphi_v &: (z, w) \mapsto (z + v, e^{-\mu\langle z, v \rangle - \frac{\mu}{2}\|v\|^2} w), \quad (v \in \mathbb{C}^n),\end{aligned}$$

where  $\mathcal{U}(k)$  is the unitary group of degree  $k$ , and  $\langle \cdot, \cdot \rangle$  is the standard Hermitian inner product on  $\mathbb{C}^n$ .

Recently, Tu-Wang [23] has established the rigidity of the proper holomorphic mappings between two equidimensional Fock-Bargmann-Hartogs domains as follows.

**Theorem 1.2** (Tu-Wang [23]). *If  $D_{n,m}(\mu)$  and  $D_{n',m'}(\mu')$  are two equidimensional Fock-Bargmann-Hartogs domains with  $m \geq 2$  and  $f$  is a proper holomorphic mapping of  $D_{n,m}(\mu)$  into  $D_{n',m'}(\mu')$ , then  $f$  is a biholomorphism between  $D_{n,m}(\mu)$  and  $D_{n',m'}(\mu')$ .*

A generalized complex ellipsoid (also called generalized pseudoellipsoid) is a domain of the form

$$\Sigma(\mathbf{n}; \mathbf{p}) = \{(\zeta_1, \dots, \zeta_r) \in \mathbb{C}^{n_1} \times \dots \times \mathbb{C}^{n_r} : \sum_{k=1}^r \|\zeta_k\|^{2p_k} < 1\},$$

where  $\mathbf{n} = (n_1, \dots, n_r) \in \mathbb{N}^r$  and  $\mathbf{p} = (p_1, \dots, p_r) \in (\mathbb{R}_+)^r$ . In the special case where all the  $p_k = 1$ , the generalized complex ellipsoid  $\Sigma(\mathbf{n}; \mathbf{p})$  reduces to the unit ball in  $\mathbb{C}^{n_1 + \dots + n_r}$ . Also, it is known that a generalized complex ellipsoid  $\Sigma(\mathbf{n}; \mathbf{p})$  is homogeneous if and only if  $p_k = 1$  for all  $1 \leq k \leq r$  (cf. Kodama [15]). In general, a generalized complex ellipsoid is not strongly pseudoconvex and its boundary is not smooth. The automorphism group  $\text{Aut}(\Sigma(\mathbf{n}; \mathbf{p}))$  of  $\Sigma(\mathbf{n}; \mathbf{p})$  has been studied by Dini-Primicerio [9], Kodama [15] and Kodama-Krantz-Ma [16].

In 2013, Kodama [15] obtained the result as follows.

**Theorem 1.3** (Kodama [15]). *(i) If 1 does not appear in  $p_1, \dots, p_r$ , then any automorphism  $\varphi \in \text{Aut}(\Sigma(\mathbf{n}; \mathbf{p}))$  is of the form*

$$\varphi(\zeta_1, \dots, \zeta_r) = (\gamma_1(\zeta_{\sigma(1)}), \dots, \gamma_r(\zeta_{\sigma(r)})) \quad (1.1)$$

where  $\sigma \in S_r$  is a permutation of the  $r$  numbers  $\{1, \dots, r\}$  such that  $n_{\sigma(i)} = n_i, p_{\sigma(i)} = p_i, 1 \leq i \leq r$  and  $\gamma_1, \dots, \gamma_r$  are unitary transformation of  $\mathbb{C}^{n_1} (n_{\sigma(1)} = n_1), \dots, \mathbb{C}^{n_r} (n_{\sigma(r)} = n_r)$  respectively.

*(ii) If 1 appears in  $p_1, \dots, p_r$ , we can assume, without loss of generality, that  $p_1 = 1, p_2 \neq 1, \dots, p_r \neq 1$ , then  $\text{Aut}(\Sigma(\mathbf{n}; \mathbf{p}))$  is generated by elements of the form (1.1) and automorphisms of the form*

$$\varphi_a(\zeta_1, \zeta_2, \dots, \zeta_r) = (T_a(\zeta_1), \zeta_2(\psi_a(\zeta_1))^{1/2p_2}, \dots, \zeta_r(\psi_a(\zeta_1))^{1/2p_r}) \quad (1.2)$$

where  $T_a$  is an automorphism of the ball  $\mathbb{B}^{n_1}$  in  $\mathbb{C}^{n_1}$ , which sends a point  $a \in \mathbb{B}^{n_1}$  to the origin and

$$\psi_a(\zeta_1) = \frac{1 - \|a\|^2}{(1 - \langle \zeta_1, a \rangle)^2}.$$

In this paper, we define the generalized Fock-Bargmann-Hartogs domains  $D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$  as follows:

$$D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu) = \{(z, w_{(1)}, \dots, w_{(l)}) \in \mathbb{C}^{n_0} \times \mathbb{C}^{n_1} \times \dots \times \mathbb{C}^{n_l} : \sum_{j=1}^l \|w_{(j)}\|^{2p_j} < e^{-\mu\|z\|^2}\} \quad (\mu > 0),$$

where  $\mathbf{p} = (p_1, \dots, p_l) \in (\mathbb{R}_+)^l$ ,  $\mathbf{n} = (n_1, \dots, n_l)$ ,  $w_{(j)} = (w_{j1}, \dots, w_{jn_j}) \in \mathbb{C}^{n_j}$ , in which  $n_j$  is a positive integer for  $1 \leq j \leq l$ . Here and henceforth, with no loss of generality, we always assume that  $p_i \neq 1$  ( $2 \leq i \leq l$ ) for  $D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$ .

Obviously, each generalized Fock-Bargmann-Hartogs domain  $D_{n_0}^{\mathbf{n}, \mathbf{p}}$  is an unbounded non-hyperbolic domain. In general, a generalized Fock-Bargmann-Hartogs domain is not a strongly pseudoconvex domain and its boundary is not smooth.

In this paper, we prove the following results.

**Theorem 1.4.** *Suppose  $D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$  and  $D_{m_0}^{\mathbf{m}, \mathbf{q}}(\nu)$  are two equidimensional generalized Fock-Bargmann-Hartogs domains. Let  $f : D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu) \rightarrow D_{m_0}^{\mathbf{m}, \mathbf{q}}(\nu)$  be a biholomorphic mapping. Then there exists  $\phi \in \text{Aut}(D_{m_0}^{\mathbf{m}, \mathbf{q}}(\nu))$  such that*

$$\phi \circ f(z, w) = (z, w_{(\sigma(1))}, \dots, w_{(\sigma(l))}) \begin{pmatrix} A & & & & \\ & \Gamma_1 & & & \\ & & \Gamma_2 & & \\ & & & \ddots & \\ & & & & \Gamma_l \end{pmatrix}, \quad (1.3)$$

where  $\sigma \in S_l$  is a permutation such that  $n_{\sigma(j)} = m_j$ ,  $p_{\sigma(j)} = q_j$  ( $1 \leq j \leq l$ ),  $\sqrt{\frac{z}{\mu}}A \in \mathcal{U}(n)$  ( $n := n_0 = m_0$ ), and  $\Gamma_i \in \mathcal{U}(n_i)$  ( $1 \leq i \leq l$ ).

**Corollary 1.5.** *Let  $f : D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu) \rightarrow D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$  be a biholomorphic mapping with  $f(0) = 0$ . Then we have*

$$f(z, w) = (z, w_{(\sigma(1))}, \dots, w_{(\sigma(l))}) \begin{pmatrix} A & & & & \\ & \Gamma_1 & & & \\ & & \Gamma_2 & & \\ & & & \ddots & \\ & & & & \Gamma_l \end{pmatrix},$$

where  $\sigma \in S_l$  is a permutation such that  $n_{\sigma(j)} = n_j$ ,  $p_{\sigma(j)} = p_j$  ( $1 \leq j \leq l$ ),  $A \in \mathcal{U}(n_0)$  and  $\Gamma_i \in \mathcal{U}(n_i)$  ( $1 \leq i \leq l$ ).

As a consequence, it is easy for us to prove the following results.

**Theorem 1.6.** *The automorphism group  $\text{Aut}(D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu))$  is generated by the following mappings:*

$$\varphi_A : (z, w_{(1)}, \dots, w_{(l)}) \mapsto (zA, w_{(1)}, \dots, w_{(l)});$$

$$\varphi_D : (z, w_{(1)}, \dots, w_{(l)}) \mapsto (z, (w_{(\sigma(1))}, \dots, w_{(\sigma(l))})D);$$

$$\varphi_a : (z, w) \mapsto (z + a, w_{(1)}(e^{-2\mu\langle z, a \rangle - \mu\|a\|^2})^{\frac{1}{2p_1}}, \dots, w_{(l)}(e^{-2\mu\langle z, a \rangle - \mu\|a\|^2})^{\frac{1}{2p_l}}),$$

where  $a \in \mathbb{C}^{n_0}$ ,  $A \in \mathcal{U}(n_0)$ ,  $\sigma \in S_l$  is a permutation such that  $n_{\sigma(j)} = n_j$ ,  $p_{\sigma(j)} = p_j$  ( $1 \leq j \leq l$ ), and

$$D = \begin{pmatrix} \Gamma_1 & & & & \\ & \Gamma_2 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \Gamma_l \end{pmatrix},$$

in which  $\Gamma_i \in \mathcal{U}(n_i)$  ( $1 \leq i \leq l$ ).

Now, for  $\mathbf{p}$  and  $\mathbf{q}$ , we introduce notation:

$$\epsilon = \begin{cases} 1, & p_1 = 1 \\ 0, & p_1 \neq 1 \end{cases}, \quad \delta = \begin{cases} 1, & q_1 = 1 \\ 0, & q_1 \neq 1 \end{cases}.$$

**Theorem 1.7.** *Suppose  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  and  $D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu)$  are two equidimensional generalized Fock-Bargmann-Hartogs domains with  $\min\{n_{1+\epsilon}, n_2, \dots, n_l, n_1 + \dots + n_l\} \geq 2$  and  $\min\{m_{1+\delta}, m_2, \dots, m_l, m_1 + \dots + m_l\} \geq 2$ . Then any proper holomorphic mapping between  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  and  $D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu)$  must be a biholomorphism.*

**Remark 1.1.** *The conditions  $\min\{n_{1+\epsilon}, n_2, \dots, n_l\} \geq 2$  can not be removed. For example,  $n_1 = 1$  (i.e.,  $w_1 \in \mathbb{C}$ ),  $p_1 \neq 1$ , and*

$$F(z, w) : (z, w_{(1)}, \dots, w_{(l)}) \rightarrow (z, w_{(1)}^2, w_{(2)}, \dots, w_{(l)}).$$

*Then  $F$  is a proper holomorphic mapping between  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  and  $D_{n_0}^{\mathbf{n}, \mathbf{Q}}(\mu)$  where  $\mathbf{q} = (p_1/2, p_2, \dots, p_l)$ .  $F$  is not a biholomorphism.*

**Corollary 1.8.** *Suppose  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  is a generalized Fock-Bargmann-Hartogs domain with*

$$\min\{n_{1+\epsilon}, n_2, \dots, n_l, n_1 + \dots + n_l\} \geq 2.$$

*Then any proper holomorphic self-mapping of  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  must be an automorphism.*

**Remark 1.2.** *The conditions  $n_1 + \dots + n_l \geq 2$  can not be removed. For instance, with no loss of generality, we can assume  $n_1 = 1$  and  $n_i = 0$  ( $2 \leq i \leq l$ ). Then*

$$F : (z, w_{(1)}) \rightarrow (\sqrt{2}z, w_{(1)}^2)$$

*is a proper holomorphic self-mapping of  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  which is not an automorphism.*

The paper is organized as follows. In Section 2, using the explicit formula for the Bergman kernels of the generalized Fock-Bargmann-Hartogs domains, we prove that a proper holomorphic mapping between two equidimensional generalized Fock-Bargmann-Hartogs domains extends holomorphically to their closures and check that the Cartan's theorem holds also for the generalized Fock-Bargmann-Hartogs domains. In Section 3, we exploit the boundary structure of generalized Fock-Bargmann-Hartogs domains to prove our results in this paper.

## 2 Preliminaries

### 2.1 The Bergman kernel of the domain $D_{n_0}^{\mathbf{n}, \mathbf{P}}$

For a domain  $\Omega$  in  $\mathbb{C}^n$ , let  $A^2(\Omega)$  be the Hilbert space of square integrable holomorphic functions on  $\Omega$  with the inner product:

$$\langle f, g \rangle = \int_{\Omega} f(z) \overline{g(z)} dV(z) \quad (f, g \in \mathcal{O}(\Omega)),$$

where  $dV$  is the Euclidean volume form. The Bergman kernel  $K(z, w)$  of  $A^2(\Omega)$  is defined as the reproducing kernel of the Hilbert space  $A^2(\Omega)$ , that is, for all  $f \in A^2(\Omega)$ , we have

$$f(z) = \int_{\Omega} f(w) K(z, w) dV(w) \quad (z \in \Omega).$$

For a positive continuous function  $p$  on  $\Omega$ , let  $A^2(\Omega, p)$  be the weighted Hilbert space of square integrable holomorphic functions with respect to the weight function  $p$  with the inner product:

$$\langle f, g \rangle = \int_{\Omega} f(z) \overline{g(z)} p(z) dV(z) \quad (f, g \in \mathcal{O}(\Omega)).$$

Similarly, the weighted Bergman kernel  $K_{A^2(\Omega,p)}$  of  $A^2(\Omega,p)$  is defined as the reproducing kernel of the Hilbert space  $A^2(\Omega,p)$ . For a positive integer  $m$ , define the Hartogs domain  $\Omega_{m,p}$  over  $\Omega$  by

$$\Omega_{m,p} = \{(z, w) \in \Omega \times \mathbb{C}^m : \|w\|^2 < p(z)\}.$$

Ligocka [17, 18] showed that the Bergman kernel of  $\Omega_{m,p}$  can be expressed as infinite sum in terms of the weighted Bergman kernel of  $A^2(\Omega, p^k)$  ( $k = 1, 2, \dots$ ) as follows.

**Theorem 2.1** (Ligocka [18]). *Let  $K_m$  be the Bergman kernel of  $\Omega_{m,p}$  and let  $K_{A^2(\Omega,p^k)}$  be the weighted Bergman kernel of  $A^2(\Omega, p^k)$  ( $k = 1, 2, \dots$ ). Then*

$$K_m((z, w), (t, s)) = \frac{m!}{\pi^m} \sum_{k=0}^{\infty} \frac{(m+1)_k}{k!} K_{A^2(\Omega,p^{k+m})}(z, t) \langle w, s \rangle^k,$$

where  $(a)_k$  denotes the Pochhammer symbol  $(a)_k = a(a+1)\cdots(a+k-1)$ .

The Fock-Bargmann space is the weighted Hilbert space  $A^2(\mathbb{C}^n, e^{-\mu\|z\|^2})$  on  $\mathbb{C}^n$  with the Gaussian weight function  $e^{-\mu\|z\|^2}$  ( $\mu > 0$ ). The reproducing kernel of  $A^2(\mathbb{C}^n, e^{-\mu\|z\|^2})$ , called the Fock-Bargmann kernel, is  $\mu^n e^{\mu\langle z, t \rangle} / \pi^n$  (see Bargmann [2]). Thus, the Fock-Bargmann-Hartogs domain  $D_{n,m} = \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^m : \|w\|^2 < e^{-\mu\|z\|^2}\}$  ( $\mu > 0$ ) and the Fock-Bargmann space  $A^2(\mathbb{C}^n, e^{-\mu\|z\|^2})$  are closely related. In 2013, using Theorem 2.1 and the expression of the Fock-Bargmann kernel, Yamamori [25] gave the Bergman kernel of the Fock-Bargmann-Hartogs domain  $D_{n,m}$  as follows.

**Theorem 2.2** (Yamamori [25]). *The Bergman kernel of the Fock-Bargmann-Hartogs domain  $D_{n,m}$  is given by*

$$K_{D_{n,m}}((z, w), (t, s)) = \frac{m! \mu^n}{\pi^{m+n}} \sum_{k=0}^{\infty} \frac{(m+1)_k (k+m)^n}{k!} e^{\mu(k+m)\langle z, t \rangle} \langle w, s \rangle^k,$$

where  $(a)_k$  denotes the Pochhammer symbol  $(a)_k = a(a+1)\cdots(a+k-1)$ .

Following the idea of Theorem 2.1, we compute the Bergman kernel for the generalized Fock-Bargmann-Hartogs domain  $D_{n_0}^{\mathbf{n}, \mathbf{P}}$ . In order to compute the Bergman kernel, we first introduce some notation.

Let

$$\alpha = (\alpha_{(1)}, \dots, \alpha_{(l)}) \in (\mathbb{R}_+)^{n_1} \times \dots \times (\mathbb{R}_+)^{n_l},$$

where  $\alpha_{(i)} = (\alpha_{i1}, \dots, \alpha_{in_i}) \in (\mathbb{R}_+)^{n_i}$  for  $1 \leq i \leq l$ . For  $\alpha \in (\mathbb{R}_+)^n$ , we define

$$\beta(\alpha) = \frac{\prod_{i=1}^l \Gamma(\alpha_i)}{\Gamma(|\alpha|)};$$

see D'Angelo [7]. Here  $\Gamma$  is the usual Euler Gamma function.

**Lemma 2.3** (D'Angelo [7], Lemma 1). *Suppose  $\alpha \in (\mathbb{R}_+)^n$ . Then we have*

$$\int_{B_+^n} r^{2\alpha-1} dV(r) = \frac{\beta(\alpha)}{2^n |\alpha|},$$

$$\int_{S_+^{n-1}} w^{2\alpha-1} d\sigma(w) = \frac{\beta(\alpha)}{2^{n-1}},$$

where  $dV$  is the Euclidean  $n$ -dimensional volume form,  $dS$  is the Euclidean  $(n-1)$ -dimensional volume form, and the subscript “+” denotes that all the variables are positive, that is,  $B_+^n = B^n \cap (\mathbb{R}_+)^n$  and  $S_+^{n-1} = S^{n-1} \cap (\mathbb{R}_+)^n$ , in which  $B^n$  is the unit ball in  $\mathbb{R}^n$  and  $S^{n-1}$  is the unit sphere in  $\mathbb{R}^n$ .

**Theorem 2.4.** Suppose  $\alpha = (\alpha_{(1)}, \dots, \alpha_{(l)}) \in (\mathbb{R}_+)^{n_1} \times \dots \times (\mathbb{R}_+)^{n_l}$ ,  $\alpha_{(i)} = (\alpha_{i1}, \dots, \alpha_{in_i}) \in (\mathbb{R}_+)^{n_i}$ ,  $1 \leq i \leq l$ . Then we have the formula:

$$\int_{\sum_{j=1}^l \|w_{(j)}\|^{2p_j} < t} w^\alpha \bar{w}^\alpha dV(w) = (\pi)^{n_1 + \dots + n_l} \frac{\prod_{i=1}^l \Gamma(\alpha_i + 1) \prod_{i=1}^l \Gamma\left(\frac{|\alpha_{(i)}| + n_i}{p_i}\right)}{\prod_{i=1}^l p_i \prod_{i=1}^l \Gamma(|\alpha_i| + n_i) \Gamma\left(\sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i} + 1\right)} \cdot t^{\sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i}} \quad (2.1)$$

*Proof.* For the integral

$$\int_{\sum_{j=1}^l \|w_{(j)}\|^{2p_j} < t} w^\alpha \bar{w}^\alpha dV(w), \quad (2.2)$$

by applying the polar coordinates  $w = se^{i\theta}$  (namely,  $w_{ij} = s_{ij}e^{i\theta_{ij}}$ ,  $1 \leq j \leq n_i$ ,  $1 \leq i \leq l$ ,  $s = (s_{(1)}, \dots, s_{(l)})$ ), we have

$$(2.2) = (2\pi)^{n_1 + \dots + n_l} \int_{\substack{\sum_{j=1}^l \|s_{(j)}\|^{2p_j} < t \\ s_{ji} > 0, 1 \leq i \leq n_j, 1 \leq j \leq l}} s^{2\alpha+1} dV(s).$$

Using the spherical coordinates in the variables  $s_{(1)}, s_{(2)}, \dots, s_{(l)}$  respectively, we get

$$\begin{aligned} & \int_{\substack{\sum_{j=1}^l \|s_{(j)}\|^{2p_j} < t \\ s_{ji} > 0, 1 \leq i \leq n_j, 1 \leq j \leq l}} s^{2\alpha+1} dV(s) \\ &= \int_{\substack{\sum_{i=1}^l |\rho_i|^{2p_i} < t \\ \rho_i > 0, 1 \leq i \leq l}} \rho_1^{2|\alpha_{(1)}| + 2n_1 - 1} \dots \rho_l^{2|\alpha_{(l)}| + 2n_l - 1} d\rho_1 d\rho_2 \dots d\rho_l \\ & \quad \times \int_{S_+^{n_1-1}} \dots \int_{S_+^{n_l-1}} w_{(1)}^{2\alpha_{(1)}+1} \dots w_{(l)}^{2\alpha_{(l)}+1} d\sigma(w_{(1)}) \dots d\sigma(w_{(l)}). \end{aligned}$$

Let  $\rho_i^{p_i} = r_i$ ,  $1 \leq i \leq l$ . Then we have  $d\rho_i = \frac{1}{p_i} \rho_i^{1-p_i} dr_i = \frac{1}{p_i} r_i^{\frac{1}{p_i}-1} dr_i$ . Therefore, Lemma 2.3 and the above formulas yield

$$(2.2) = (2\pi)^{n_1 + \dots + n_l} \frac{1}{\prod_{i=1}^l p_i} \frac{\beta(\alpha_{(1)} + 1)}{2^{n_1-1}} \dots \frac{\beta(\alpha_{(l)} + 1)}{2^{n_l-1}} \int_{\substack{\sum_{i=1}^l |r_i|^{2p_i} < t \\ r_i > 0, 1 \leq i \leq l}} r_1^{\frac{2|\alpha_{(1)}| + 2n_1 - 1}{p_1}} \dots r_l^{\frac{2|\alpha_{(l)}| + 2n_l - 1}{p_l}} dr_1 \dots dr_l.$$

Let  $r = (r_1, r_2, \dots, r_l) \in (\mathbb{R}_+)^l$  and  $k := t^{-\frac{1}{2}}r$ . Then  $dr = t^{\frac{l}{2}}dk$ . After a straightforward computation, we obtain that (2.2) equals

$$(2\pi)^{n_1 + \dots + n_l} \frac{1}{\prod_{i=1}^l p_i} \frac{\beta(\alpha_{(1)} + 1)}{2^{n_1-1}} \dots \frac{\beta(\alpha_{(l)} + 1)}{2^{n_l-1}} \cdot t^{\sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i}} \int_{B_+^l} k_1^{\frac{2|\alpha_{(1)}| + 2n_1 - 1}{p_1}} \dots k_l^{\frac{2|\alpha_{(l)}| + 2n_l - 1}{p_l}} dk_1 \dots dk_l.$$

Applying Lemma 2.3 to the above formula, we get

$$\begin{aligned} (2.2) &= (\pi)^{n_1 + \dots + n_l} \beta(\alpha_{(1)} + 1) \dots \beta(\alpha_{(l)} + 1) \frac{\beta(\alpha')}{|\alpha'| \prod_{i=1}^l p_i} \cdot t^{\sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i}} \\ &= (\pi)^{n_1 + \dots + n_l} \frac{1}{\prod_{i=1}^l p_i} \frac{\prod_{i=1}^l \Gamma(\alpha_{(i)} + 1) \prod_{i=1}^l \Gamma\left(\frac{|\alpha_{(i)}| + n_i}{p_i}\right)}{\prod_{i=1}^l \Gamma(|\alpha_i| + n_i) \Gamma\left(\sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i} + 1\right)} \cdot t^{\sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i}}, \end{aligned} \quad (2.3)$$

where  $\alpha' = (\frac{|\alpha_{(1)}|+n_1}{p_1}, \dots, \frac{|\alpha_{(l)}|+n_l}{p_l}) \in (\mathbb{R}_+)^l$ .  $\square$

Now we consider the Hilbert space  $A^2(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$  of square-integrable holomorphic functions on  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ .

**Lemma 2.5.** *Let  $f \in A^2(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$ . Then*

$$f(z, w) = \sum_{\alpha} f_{\alpha}(z)w^{\alpha},$$

where the series is uniformly convergent on compact subsets of  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ ,  $f_{\alpha}(z) \in A^2(\mathbb{C}^{n_0}, e^{-\mu\lambda_{\alpha}\|z\|^2})$  for any  $\alpha = (\alpha_{(1)}, \dots, \alpha_{(l)}) \in \mathbb{N}^{n_1} \times \dots \times \mathbb{N}^{n_l}$ ,  $\alpha_{(i)} = (\alpha_{i1}, \dots, \alpha_{in_i}) \in \mathbb{N}^{n_i}$ ,  $1 \leq i \leq l$ ,  $\lambda_{\alpha} = \sum_{i=1}^l \frac{|\alpha_{(i)}|+n_i}{p_i}$ , in which  $A^2(\mathbb{C}^n, e^{-\mu\lambda_{\alpha}\|z\|^2})$  denotes the space of square-integrable holomorphic functions on  $\mathbb{C}^n$  with respect to the measure  $e^{-\mu\lambda_{\alpha}\|z\|^2} dV_{2n}$ .

*Proof.* Since  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  is a complete Reinhardt domain, each holomorphic function on  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  is the sum of a locally uniformly convergent power series. Thus, for  $f \in A^2(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$ , we have

$$f(z, w) = \sum_{\alpha} f_{\alpha}(z)w^{\alpha},$$

where the series is uniformly convergent on compact subsets of  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ . We choose a sequence of compact subsets  $D_k$  ( $1 \leq k < \infty$ )

$$D_k := \{(z, w_{(1)}, \dots, w_{(l)}) \in \mathbb{C}^{n_0} \times \mathbb{C}^{n_1} \times \dots \times \mathbb{C}^{n_l} : \sum_{j=1}^l \|w_{(j)}\|^{2p_j} \leq e^{-\mu\|z\|^2} - \frac{1}{k}\} \cap \overline{B(0, k)},$$

where  $B(0, k)$  is the ball in  $\mathbb{C}^{n_0+n_1+\dots+n_l}$  of the radius  $k$ . Obviously,  $D_k \Subset D_{k+1}$  and  $\bigcup_{k=1}^{\infty} D_k = D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ .

Since  $D_k$  is a circular domain, then

$$f_{\alpha}(z)w^{\alpha} \perp f_{\beta}(z)w^{\beta} \quad (\alpha \neq \beta)$$

in the Hilbert space  $A^2(D_k)$ . Hence we have

$$\|f\|_{L^2(D_k)}^2 = \sum_{|\alpha|=0}^{\infty} \|f_{\alpha}(z)w^{\alpha}\|_{L^2(D_k)}^2.$$

Since  $f(z, w) \in A^2(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$ , we have

$$\|f_{\alpha}(z)w^{\alpha}\|_{L^2(D_k)}^2 \leq \|f\|_{L^2(D_k)}^2 \leq \|f\|_{L^2(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))}^2.$$

Then  $f_{\alpha}(z)w^{\alpha} \in A^2(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$ . Therefore,

$$\begin{aligned} & \int_{D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)} |f_{\alpha}(z)|^2 w^{\alpha} \bar{w}^{\alpha} dV < \infty \\ \implies & \int_{\mathbb{C}^{n_0}} |f_{\alpha}(z)|^2 dV(z) \int_{\sum_{j=1}^l \|w_{(j)}\|^{2p_j} < e^{-\mu\|z\|^2}} w^{\alpha} \bar{w}^{\alpha} dV(w) < \infty. \end{aligned}$$

By (2.1), it follows

$$\int_{\mathbb{C}^{n_0}} |f_{\alpha}(z)|^2 e^{-\mu\lambda_{\alpha}\|z\|^2} dV(z) < \infty.$$

Consequently,  $f_{\alpha}(z) \in A^2(\mathbb{C}^{n_0}, e^{-\mu\lambda_{\alpha}\|z\|^2})$ , where  $\lambda_{\alpha} = \sum_{i=1}^l \frac{|\alpha_{(i)}|+n_i}{p_i}$ .  $\square$

Lemma 2.5 implies that  $f(z)w^\alpha$  where  $f(z) \in A^2(\mathbb{C}^{n_0}, e^{-\mu\lambda_\alpha\|z\|^2})$  form a linearly dense subset of  $A^2(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$ . Now we can express the Bergman kernel of  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  as follows.

**Theorem 2.6.** *The Bergman kernel of  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  can be expressed by the following form*

$$K_{D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)}[(z, w), (s, t)] = \sum_{|\alpha|=0}^{\infty} c_\alpha \frac{\lambda_\alpha^{n_0} \mu^{n_0}}{\pi^{n_0}} e^{\lambda_\alpha \mu \langle z, s \rangle} w^\alpha \bar{t}^\alpha, \quad (2.4)$$

where  $\alpha = (\alpha_{(1)}, \dots, \alpha_{(l)}) \in \mathbb{N}^{n_1} \times \dots \times \mathbb{N}^{n_l}$ ,  $\alpha_{(i)} = (\alpha_{i1}, \dots, \alpha_{in_i}) \in \mathbb{N}^{n_i}$ ,  $1 \leq i \leq l$ , and

$$c_\alpha = \frac{\prod_{i=1}^l p_i \prod_{i=1}^l \Gamma(|\alpha_i| + n_i) \Gamma(\sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i} + 1)}{(\pi)^{n_1 + \dots + n_l} \prod_{i=1}^l \Gamma(\alpha_{(i)} + 1) \prod_{i=1}^l \Gamma(\frac{|\alpha_{(i)}| + n_i}{p_i})}, \quad \lambda_\alpha = \sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i}.$$

*Proof.* Since  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  is a complete Reinhardt domain, it follows

$$K_{D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)}[(z, w), (s, t)] = \sum_{|\beta|=0}^{\infty} c_\beta g_\beta(z, s) w^\beta \bar{t}^\beta,$$

where the sum is locally uniformly convergent, by the invariance of the Bergman kernel  $K_{D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)}$  on  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  under the unitary subgroup action

$$(z_1, \dots, z_{n_0+|\mathbf{n}|}) \rightarrow (e^{\sqrt{-1}\theta_1} z_1, \dots, e^{\sqrt{-1}\theta_{n_0+|\mathbf{n}|}} z_{n_0+|\mathbf{n}|}) \quad (\theta_1, \dots, \theta_{n_0+|\mathbf{n}|} \in \mathbb{R}).$$

For any  $\alpha = (\alpha_{(1)}, \dots, \alpha_{(l)}) \in \mathbb{N}^{n_1} \times \dots \times \mathbb{N}^{n_l}$  with  $\alpha_{(i)} = (\alpha_{i1}, \dots, \alpha_{in_i}) \in \mathbb{N}^{n_i}$  ( $1 \leq i \leq l$ ), any  $f(z) \in A^2(\mathbb{C}^{n_0}, e^{-\mu\lambda_\alpha\|z\|^2})$  ( $\lambda_\alpha = \sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i}$ ), we have  $f(z)w^\alpha \in A^2(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$ . Thus

$$\begin{aligned} f(z)w^\alpha &= \int_{D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)} f(s) t^\alpha K_{D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)}[(z, w), (s, t)] dV \\ &= \int_{\mathbb{C}^{n_0}} f(s) \sum_{\beta=0}^{\infty} c_\beta g_\beta(z, s) w^\beta dV(s) \int_{\sum_{j=1}^l \|t_{(j)}\|^{2p_j} < e^{-\mu\|s\|^2}} t^\alpha \bar{t}^\beta dV(t) \\ &= w^\alpha \int_{\mathbb{C}^{n_0}} f(s) g_\alpha(z, s) [e^{-\mu\|s\|^2}]^{\sum_{i=1}^l \frac{|\alpha_{(i)}| + n_i}{p_i}} dV(s) \quad (\text{by (2.1)}). \end{aligned}$$

By Bargmann [2], we get that the Bergman kernel of  $A^2(\mathbb{C}^{n_0}, e^{-\mu\lambda_\alpha\|z\|^2})$  can be described by the form

$$K_\alpha(z, w) = \frac{\lambda_\alpha^{n_0} \mu^{n_0}}{\pi^{n_0}} e^{\lambda_\alpha \mu \langle z, w \rangle}. \quad (2.5)$$

Thus we obtain

$$g_\alpha(z, s) = \frac{\lambda_\alpha^{n_0} \mu^{n_0}}{\pi^{n_0}} e^{\lambda_\alpha \mu \langle z, s \rangle}.$$

This completes the proof.  $\square$

The transformation rule for Bergman kernels under proper holomorphic mapping (e.g., Th. 1 in Bell [4]) is also valid for unbounded domains (e.g., see Cor. 1 in Trybula [21]). Note that the coordinate functions play a key role in the approach of Bell [4] to extend proper holomorphic mapping, but, in general, are no longer square integrable on unbounded domains. In order to overcome the difficulty, by combining the transformation rule for Bergman kernels under proper holomorphic mapping in Bell [4] and our explicit form (2.4) of the Bergman kernel function for  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ , we prove that a proper holomorphic mapping between two equidimensional generalized Fock-Bargmann-Hartogs domains extends holomorphically to their closures as follows.



**Lemma 2.7.** *Suppose that  $f : D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu) \rightarrow D_{m_0}^{\mathbf{m},\mathbf{q}}(\nu)$  is a proper holomorphic mapping between two equidimensional generalized Fock-Bargmann-Hartogs domains. Then  $f$  extends holomorphically to a neighborhood of the closure  $\overline{D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu)}$ .*

In fact, using the explicit form (2.4) of the Bergman kernel function for  $D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu)$ , we immediately have Lemma 2.7 by a slightly modifying the proof of Th. 2.5 in Tu-Wang [23].

## 2.2 Cartan's Theorem on the $D_{n_0}^{\mathbf{n},\mathbf{p}}$

Suppose  $D$  is a domain in  $\mathbb{C}^N$  and let  $K_D(z, w)$  be its Bergman kernel. From Ishi-Kai [12], we know that if the following conditions are satisfied:

- (a)  $K_D(0, 0) > 0$ ;
- (b)  $T_D(0, 0)$  is positive definite,

where  $T_D$  is an  $N \times N$  matrix

$$T_D(z, w) := \begin{pmatrix} \frac{\partial^2 \log K_D(z, w)}{\partial z_1 \partial \bar{w}_1} & \cdots & \frac{\partial^2 \log K_D(z, w)}{\partial z_1 \partial \bar{w}_N} \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 \log K_D(z, w)}{\partial z_N \partial \bar{w}_1} & \cdots & \frac{\partial^2 \log K_D(z, w)}{\partial z_N \partial \bar{w}_N} \end{pmatrix}.$$

Then the Cartan's theorem can also be applied to the case of unbounded circular domains. The above conditions are obviously satisfied by the bounded domain.

Kim-Ninh-Yamamori [13] proved the following result.

**Lemma 2.8** (Kim-Ninh-Yamamori [13], Th. 4). *Suppose that  $D$  is a circular domain and its Bergman kernel satisfies the above conditions (a) and (b). If  $\varphi \in \text{Aut}(D)$  preserves the origin, then  $\varphi$  is a linear mapping.*

Ishi-Kai [12] proved the generalization of Lemma 2.8 as follows.

**Lemma 2.9** (Ishi-Kai [12], Prop. 2.1). *Let  $D_k$  be a circular domain (not necessarily bounded) in  $\mathbb{C}^N$  with  $0 \in D_k$  ( $k = 1, 2$ ), and let  $\varphi : D_1 \rightarrow D_2$  be a biholomorphism with  $\varphi(0) = 0$ . If  $K_{D_k}(0, 0) > 0$  and  $T_{D_k}(0, 0)$  is positive definite ( $k = 1, 2$ ), then  $\varphi$  is linear.*

Therefore, by using the expressions of Bergman kernels of generalized Fock-Bargmann-Hartogs domains, we have the following result.

**Theorem 2.10.** *Suppose that  $\varphi : D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu) \rightarrow D_{m_0}^{\mathbf{m},\mathbf{q}}(\nu)$  be a biholomorphic mapping between two equidimensional generalized Fock-Bargmann-Hartogs domains with  $\varphi(0) = 0$ . Then  $\varphi$  is linear.*

*Proof.* By using the expressions (2.4) of Bergman kernels of generalized Fock-Bargmann-Hartogs domains and a straightforward computation, we show that the Bergman kernel of every generalized Fock-Bargmann-Hartogs domain satisfies the above conditions (a) and (b). So we get Th. 2.10 by Lemma 2.9.  $\square$

## 3 Proof Of The Main Theorem

To begin, we exploit the boundary structure of  $D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu)$  which is comprised of

$$bD_{n_0}^{\mathbf{n},\mathbf{p}}(\mu) = b_0D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu) \cup b_1D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu) \cup b_2D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu),$$

where

$$b_0D_{n_0}^{\mathbf{n},\mathbf{p}}(\mu) := \{(z, w_{(1)}, \dots, w_{(l)}) \in \mathbb{C}^{n_0} \times \cdots \times \mathbb{C}^{n_l} : \sum_{j=1}^l \|w_{(j)}\|^{2p_j} = e^{-\mu\|z\|^2}, \|w_{(j)}\|^2 \neq 0, 1 + \epsilon \leq j \leq l\};$$

$$b_1 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) := \bigcup_{j=1+\epsilon}^l \{(z, w_{(1)}, \dots, w_{(l)}) \in \mathbb{C}^{n_0} \times \dots \times \mathbb{C}^{n_l} : \sum_{j=1}^l \|w_{(j)}\|^{2p_j} = e^{-\mu\|z\|^2}, \|w_{(j)}\|^2 = 0, p_j > 1\};$$

$$b_2 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) := \bigcup_{j=1+\epsilon}^l \{(z, w_{(1)}, \dots, w_{(l)}) \in \mathbb{C}^{n_0} \times \dots \times \mathbb{C}^{n_l} : \sum_{j=1}^l \|w_{(j)}\|^{2p_j} = e^{-\mu\|z\|^2}, \|w_{(j)}\|^2 = 0, p_j < 1\}.$$

Now we give the following proposition.

**Proposition 3.1.** (1) *The boundary  $b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  is a real analytic hypersurface in  $\mathbb{C}^{n_0+n_1+\dots+n_l}$  and  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  is strongly pseudoconvex at all points of  $b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ .*  
(2)  *$D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  is weakly pseudoconvex but not strongly pseudoconvex at any point of  $b_1 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  and is not smooth at any point of  $b_2 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ .*

*Proof.* Let

$$\rho(z, w_{(1)}, \dots, w_{(l)}) := \sum_{j=1}^l \|w_{(j)}\|^{2p_j} - e^{-\mu\|z\|^2}.$$

Then  $\rho$  is a real analytic definition function of  $b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ . Fix a point  $(z_0, w_{(1)0}, \dots, w_{(l)0}) \in b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  and let  $T = (\zeta, \eta_{(1)}, \dots, \eta_{(l)}) \in T_{(z_0, w_{(1)0}, \dots, w_{(l)0})}^{1,0}(b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$ . Then by definition, we know that

$$w_{(j)0} \neq 0, \quad j = 1 + \epsilon, \dots, l; \quad (3.1)$$

$$\sum_{k=1}^l p_k \|w_{(k)0}\|^{2(p_k-1)} \overline{w_{(k)0}} \cdot \eta_{(k)} + \mu e^{-\mu\|z_0\|^2} \overline{z_0} \cdot \zeta = 0; \quad (3.2)$$

$$\sum_{j=1}^l \|w_{(j)0}\|^{2p_j} - e^{-\mu\|z_0\|^2} = 0. \quad (3.3)$$

Thanks to (3.1), (3.2) and (3.3), the Levi form of  $\rho$  at the point  $(z_0, w_{(1)0}, \dots, w_{(l)0})$  can be computed as follows:

$$\begin{aligned} & L_\rho(T, T) \\ &:= \sum_{i,j=1}^{n_0+n_1+\dots+n_l} \frac{\partial^2 \rho}{\partial T_i \partial \overline{T_j}}(z_0, w_{(1)0}, \dots, w_{(l)0}) T_i \overline{T_j} \\ &= \sum_{k=1}^l p_k(p_k-1) \|w_{(k)0}\|^{2(p_k-2)} |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2 + \sum_{k=1}^l p_k \|w_{(k)0}\|^{2(p_k-1)} \|\eta_{(k)}\|^2 \\ &\quad + \mu e^{-\mu\|z_0\|^2} \|\zeta\|^2 - \mu^2 e^{-\mu\|z_0\|^2} |\overline{z_0} \cdot \zeta|^2 \\ &= \sum_{k=1}^l p_k^2 \|w_{(k)0}\|^{2(p_k-2)} |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2 + \mu e^{-\mu\|z_0\|^2} \|\zeta\|^2 - \mu^2 e^{-\mu\|z_0\|^2} |\overline{z_0} \cdot \zeta|^2 \\ &\quad + \sum_{k=1}^l p_k \|w_{(k)0}\|^{2(p_k-2)} (\|w_{(k)0}\|^2 \|\eta_{(k)}\|^2 - |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2) \end{aligned}$$

$$\begin{aligned}
&= \left( \sum_{k=1}^l \|w_{(k)0}\|^{2p_k} \right)^{-1} \left( \sum_{k=1}^l p_k^2 \|w_{(k)0}\|^{2(p_k-2)} |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2 \right) \left( \sum_{k=1}^l \|w_{(k)0}\|^{2p_k} \right) \\
&\quad - \left( \sum_{k=1}^l \|w_{(k)0}\|^{2p_k} \right)^{-1} \left| \sum_{k=1}^l p_k \|w_{(k)0}\|^{2(p_k-1)} \overline{w_{(k)0}} \cdot \eta_{(k)} \right|^2 \\
&\quad + \sum_{k=1}^l p_k \|w_{(k)0}\|^{2(p_k-2)} \left( \|w_{(k)0}\|^2 \|\eta_{(k)}\|^2 - |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2 \right) + \mu e^{-\mu \|z_0\|^2} \|\zeta\|^2 \\
&= \left( \sum_{k=1}^l \|w_{(k)0}\|^{2p_k} \right)^{-1} \left[ \left( \sum_{k=1}^l p_k^2 \|w_{(k)0}\|^{2(p_k-2)} |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2 \right) \left( \sum_{k=1}^l \|w_{(k)0}\|^{2p_k} \right) \right. \\
&\quad \left. - \left| \sum_{k=1}^l p_k \|w_{(k)0}\|^{2(p_k-1)} \overline{w_{(k)0}} \cdot \eta_{(k)} \right|^2 \right] + \mu e^{-\mu \|z_0\|^2} \|\zeta\|^2 \\
&\quad + \sum_{k=1}^l p_k \|w_{(k)0}\|^{2(p_k-2)} \left( \|w_{(k)0}\|^2 \|\eta_{(k)}\|^2 - |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2 \right) \geq \mu e^{-\mu \|z_0\|^2} \|\zeta\|^2 \geq 0
\end{aligned}$$

by the Cauchy-Schwarz inequality, for all  $T = (\zeta, \eta_{(1)}, \dots, \eta_{(l)}) \in T_{(z_0, w_{(1)0}, \dots, w_{(l)0})}^{1,0}(b_0 D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu))$ . Obviously, if  $\zeta \neq 0$ , then  $L_\rho(T, T) > 0$ .

On the other hand, combining with (3.1), (3.2) and (3.3), we know that the equality holds if and only if

$$\zeta = 0, \quad (3.4)$$

$$\|w_{(k)0}\|^2 \|\eta_{(k)}\|^2 - |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2 = 0, \quad (3.5)$$

$$\begin{aligned}
&\left[ \left( \sum_{k=1}^l p_k^2 \|w_{(k)0}\|^{2(p_k-2)} |\overline{w_{(k)0}} \cdot \eta_{(k)}|^2 \right) \left( \sum_{k=1}^l \|w_{(k)0}\|^{2p_k} \right) \right. \\
&\quad \left. - \left| \sum_{k=1}^l p_k \|w_{(k)0}\|^{2(p_k-1)} \overline{w_{(k)0}} \cdot \eta_{(k)} \right|^2 \right] = 0. \quad (3.6)
\end{aligned}$$

Suppose  $\zeta = 0$ , then  $T = (\zeta, \eta_{(1)}, \dots, \eta_{(l)}) \neq 0$  implies that there exists  $\eta_{i_0} \neq 0$ . If  $L_\rho(T, T) = 0$  for all  $T \neq 0 \in T_{(z_0, w_{(1)0}, \dots, w_{(l)0})}^{1,0}(b_0 D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu))$ , then by (3.1), (3.2), (3.3) and (3.6), we have  $\eta_k = 0$  ( $1 \leq k \leq l$ ). This is a contradiction.

When there exists  $j_0 \geq 1 + \epsilon$  such that  $\|w_{(j_0)0}\|^2 = 0$  and  $p_{j_0} > 1$ , then  $(z_0, w_{(1)0}, \dots, w_{(l)0}) \in b_1 D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$ . Let  $T_0 = (0, \dots, \eta_{(j_0)}, 0, \dots, 0)$ ,  $\|\eta_{(j_0)}\| \neq 0$ . Then  $L_\rho(T_0, T_0) = 0$ . Hence  $D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$  is weakly pseudoconvex but not strongly pseudoconvex on any point of  $b_1 D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$ .

It is obvious that  $D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$  is not smooth at any point of  $b_2 D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$ . The proof is completed.  $\square$

**Lemma 3.1** (Tu-Wang [24]). *Let  $\Sigma(\mathbf{n}; \mathbf{p})$  and  $\Sigma(\mathbf{m}; \mathbf{q})$  be two equidimensional generalized pseudoellipsoids,  $\mathbf{n}, \mathbf{m} \in \mathbb{N}^l$ ,  $\mathbf{p}, \mathbf{q} \in (\mathbb{R}_+)^l$  (where  $p_k, q_k \neq 1$  for  $2 \leq k \leq l$ ). Let  $h : \Sigma(\mathbf{n}; \mathbf{p}) \rightarrow \Sigma(\mathbf{m}; \mathbf{q})$  be a biholomorphic linear isomorphism between  $\Sigma(\mathbf{n}; \mathbf{p})$  and  $\Sigma(\mathbf{m}; \mathbf{q})$ . Then there exists a permutation  $\sigma \in S_r$  such that  $n_{\sigma(i)} = m_i$ ,  $p_{\sigma(i)} = q_i$  and*

$$h(\zeta_1, \dots, \zeta_r) = (\zeta_{\sigma(1)}, \dots, \zeta_{\sigma(r)}) \begin{bmatrix} U_1 & & & \\ & U_2 & & \\ & & \ddots & \\ & & & U_r \end{bmatrix},$$

where  $U_i$  is a unitary transformation of  $\mathbb{C}^{m_i}$  ( $m_i = n_{\sigma(i)}$ ) for  $1 \leq i \leq r$ .

Define

$$V_1 := \{(z, w_{(1)}, \dots, w_{(l)}) \in \mathbb{C}^{n_0} \times \mathbb{C}^{n_1} \times \dots \times \mathbb{C}^{n_l} : w_{(1)} = 0, \dots, w_{(l)} = 0\} (\cong \mathbb{C}^{n_0}),$$

$$V_2 := \{(z, w_{(1)}, \dots, w_{(l)}) \in \mathbb{C}^{m_0} \times \mathbb{C}^{m_1} \times \dots \times \mathbb{C}^{m_l} : w_{(1)} = 0, \dots, w_{(l)} = 0\} (\cong \mathbb{C}^{m_0}).$$

Then we have the following lemma.

**Lemma 3.2.** *Suppose  $D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu)$  and  $D_{m_0}^{\mathbf{m}, \mathbf{q}}(\nu)$  are two equidimensional generalized Fock-Bargmann-Hartogs domains,  $f : D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu) \rightarrow D_{m_0}^{\mathbf{m}, \mathbf{q}}(\nu)$  is a biholomorphic mapping. Then we have  $f(V_1) \subseteq V_2$  and  $f|_{V_1} : V_1 \rightarrow V_2$  is biholomorphic. Consequently  $n_0 = m_0$ .*

*Proof.* Let  $f(z, 0) = (f_1(z), f_2(z))$ , then we get  $\sum_{i=1}^l \|f_{2i}\|^{2q_i} < e^{-\nu\|f_1(z)\|^2} \leq 1$ . Then we obtain that the bounded entire mapping  $f_{2i}(z)$  on  $\mathbb{C}^{n_0}$  is constant ( $1 \leq i \leq l$ ) by Liouville's Theorem. Since  $f(z)$  is biholomorphic,  $f_1(z)$  is an unbounded function. Hence there exist  $\{z_k\}$  such that  $f_1(z_k) \rightarrow \infty$  as  $k \rightarrow \infty$ . It implies  $f_2(z) \equiv 0$ . This proves  $f(V_1) \subseteq V_2$ . Similarly, by making the same argument for  $f^{-1}$ , we have  $f^{-1}(V_2) \subseteq V_1$ . Namely,  $f|_{V_1} : V_1 \rightarrow V_2$  is biholomorphic. Hence  $n_0 = m_0$ .  $\square$

Now we give the proof of Theorem 1.4.

*The proof of Theorem 1.4.* Let  $f(0, 0) = (a, b)$  (thus  $b = 0$  by Lemma 3.2) and define

$$\phi(z, w_{(1)}, \dots, w_{(l)}) := (z - a, w_{(1)}(e^{2\nu\langle z, a \rangle - \nu\|a\|^2})^{\frac{1}{2q_1}}, \dots, w_{(l)}(e^{2\nu\langle z, a \rangle - \nu\|a\|^2})^{\frac{1}{2q_l}}).$$

Obviously,  $\phi \in \text{Aut}(D_{m_0}^{\mathbf{m}, \mathbf{q}}(\nu))$  and  $\phi \circ f(0, 0) = (0, 0)$ . Then  $\phi \circ f$  is linear by Theorem 2.10. We describe  $\phi \circ f$  as follows:

$$\phi \circ f(z, w) = (z, w) \begin{pmatrix} A & B \\ C & D \end{pmatrix} = (zA + wC, zB + wD).$$

According to Lemma 3.2, we have  $f(z, 0) = (f_1(z), 0)$ . Thus  $B = 0$ . Since  $g := \phi \circ f$  is biholomorphic,  $A$  and  $D$  are invertible matrices. We write  $g(z, w)$  as follows:

$$g(z, w) = (z, w) \begin{pmatrix} A & 0 \\ C & D \end{pmatrix} = (z, w_{(1)}, \dots, w_{(l)}) \begin{pmatrix} A & 0 & \dots & 0 \\ C_{11} & D_{11} & \dots & D_{1l} \\ \vdots & \vdots & \ddots & \dots \\ C_{l1} & D_{l1} & \dots & D_{ll} \end{pmatrix},$$

which implies that

$$g^{-1}(z, w) = (z, w) \begin{pmatrix} A^{-1} & 0 \\ -D^{-1}CA^{-1} & D^{-1} \end{pmatrix} = (z, w_{(1)}, \dots, w_{(l)}) \begin{pmatrix} A^{-1} & 0 & \dots & 0 \\ E_{11} & G_{11} & \dots & G_{1l} \\ \vdots & \vdots & \ddots & \vdots \\ E_{l1} & G_{l1} & \dots & G_{ll} \end{pmatrix}.$$

Set  $\Sigma(\mathbf{n}; \mathbf{p}) = \{(w_{(1)}, \dots, w_{(l)}) \in \mathbb{C}^{n_1} \times \dots \times \mathbb{C}^{n_l} : \sum_{j=1}^l \|w_{(j)}\|^{2p_j} < 1\}$ . Then, if  $\sum_{j=1}^l \|w_{(j)}\|^{2p_j} < e^{-\mu\|0\|^2} = 1$ , we obtain

$$\sum_{j=1}^l \|w_{(1)}D_{1j} + \dots + w_{(l)}D_{lj}\|^{2q_j} < e^{-\nu\|wC\|^2} < 1$$

and if  $\sum_{j=1}^l \|w_{(j)}\|^{2q_j} < e^{-\nu\|0\|^2} = 1$ , we have

$$\sum_{j=1}^l \|w_{(1)}G_{1j} + \cdots + w_{(l)}G_{lj}\|^{2p_j} < e^{-\mu\|w(-D^{-1}CA^{-1})\|^2} < 1.$$

Therefore, we conclude that the mapping  $g_2(w) : \Sigma(\mathbf{n}; \mathbf{p}) \rightarrow \Sigma(\mathbf{m}; \mathbf{q})$  given by

$$g_2(w_{(1)}, \dots, w_{(l)}) = wD = (w_{(1)}, \dots, w_{(l)}) \begin{pmatrix} D_{11} & \cdots & D_{1l} \\ \vdots & \ddots & \vdots \\ D_{l1} & \cdots & D_{ll} \end{pmatrix}$$

is a biholomorphic linear mapping. By Lemma 3.1,  $g_2$  can be expressed in the form:

$$g_2(w_{(1)}, \dots, w_{(l)}) = (w_{(\sigma(1))}, \dots, w_{(\sigma(l))}) \begin{pmatrix} \Gamma_1 & & & \\ & \Gamma_2 & & \\ & & \ddots & \\ & & & \Gamma_l \end{pmatrix},$$

where  $\sigma \in S_l$  is a permutation with  $n_{\sigma(j)} = m_j$ ,  $p_{\sigma(j)} = q_j$  ( $j = 1, \dots, l$ ) and  $\Gamma_i \in \mathcal{U}(m_i)$  ( $1 \leq i \leq l$ ). Hence  $g$  can be rewritten as follows:

$$g(z, w) = (z, w) \begin{pmatrix} A & 0 \\ C & D \end{pmatrix} = (z, w_{(\sigma(1))}, \dots, w_{(\sigma(l))}) \begin{pmatrix} A & & & & \\ C_{\sigma(1)1} & \Gamma_1 & & & \\ C_{\sigma(2)1} & & \Gamma_2 & & \\ \vdots & & & \ddots & \\ C_{\sigma(l)1} & & & & \Gamma_l \end{pmatrix}.$$

Next we prove that  $C = 0$ . The linearity of  $g$  yields that  $g(bD_{n_0}^{\mathbf{n}, \mathbf{p}}) = bD_{m_0}^{\mathbf{m}, \mathbf{q}}$ . Let  $(0, w) = (0, 0, \dots, w_{(j)}, 0, \dots, 0) \in bD_{n_0}^{\mathbf{n}, \mathbf{p}}$ , namely,  $\|w_{(j)}\|^2 = (e^{-\mu\|0\|^2})^{\frac{1}{p_j}} = 1$ . As  $\Gamma_j$  ( $1 \leq j \leq l$ ) are unitary matrices, moreover, assuming  $\sigma(i_0) = j$ , we have

$$\|w_{(j)}\|^{2p_j} = \|w_{(\sigma(i_0))}\Gamma_{i_0}\|^{2q_{i_0}} = e^{-\nu\|w_{(\sigma(i_0))}C_{\sigma(i_0)1}\|^2} = 1.$$

This implies  $w_{(j)}C_{j1} = 0$  for all  $\|w_{(j)}\|^2 = 1$ . So  $C_{j1} = 0$  ( $1 \leq j \leq l$ ). Thus we have

$$g(z, w_{(1)}, \dots, w_{(l)}) = (z, w_{(\sigma(1))}, \dots, w_{(\sigma(l))}) \begin{pmatrix} A & & & & \\ & \Gamma_1 & & & \\ & & \Gamma_2 & & \\ & & & \ddots & \\ & & & & \Gamma_l \end{pmatrix}.$$

Lastly, we show  $\sqrt{\frac{\nu}{\mu}}A \in \mathcal{U}(n)$  ( $n := n_0 = m_0$ ). For  $z \in \mathbb{C}^{n_0}$ , take  $(w_{(1)}, \dots, w_{(l)})$  such that  $e^{-\mu\|z\|^2} = \sum_{j=1}^l \|w_{(j)}\|^{2p_j}$ . By  $g(bD_{n_0}^{\mathbf{n}, \mathbf{p}}) = bD_{m_0}^{\mathbf{m}, \mathbf{q}}$ , we have  $\sum_{j=1}^l \|w_{(\sigma(j))}\Gamma_j\|^{2q_j} = e^{-\mu\|zA\|^2}$ . Since  $\Gamma_j$  ( $j = 1, \dots, l$ ) are unitary matrices, we get

$$e^{-\mu\|z\|^2} = \sum_{j=1}^l \|w_{(\sigma(j))}\|^{2p_{\sigma(j)}} = \sum_{j=1}^l \|w_{(\sigma(j))}\Gamma_j\|^{2q_j} = e^{-\mu\|zA\|^2}.$$

Therefore,  $\nu\|zA\|^2 = \mu\|z\|^2$  ( $z \in \mathbb{C}^n$ ). Then we get  $\sqrt{\frac{\nu}{\mu}}A \in \mathcal{U}(n)$ , and the proof is completed.  $\square$

*The proof of Corollary 1.5.* In fact, the significance of the above  $\phi$  is just to ensure  $\phi \circ f(0) = 0$ . Then the proof of Theorem 1.4 implies that Corollary 1.5 is obvious.  $\square$

*The proof of the Theorem 1.6.* Obviously,  $\varphi_A, \varphi_D$  and  $\varphi_a$  are biholomorphic self-mappings of  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ . On the other hand, for  $\varphi \in \text{Aut}(D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu))$ , we assume  $\varphi(0, 0) = (a, b)$  (then  $b = 0$  by Lemma 3.2). Hence  $\varphi_{-a} \circ \varphi$  preserves the origin. Then by Corollary 1.5, we obtain  $\varphi_{-a} \circ \varphi = \varphi_D \circ \varphi_A$  for some  $\varphi_A, \varphi_D$ . Hence  $\varphi = \varphi_a \circ \varphi_D \circ \varphi_A$ , and the proof is complete.  $\square$

*The proof of Theorem 1.7.* Let  $f$  be a proper holomorphic mapping between two equidimensional generalized Fock-Bargmann-Hartogs domains  $D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  and  $D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu)$ . Then by Th. 2.7,  $f$  extends holomorphically to a neighborhood  $\Omega$  of  $\overline{D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)}$  with

$$f(bD_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)) \subset bD_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu).$$

Then by Proposition 3.1 and Lemma 1.3 in Pinčuk [19], we have

$$f(M \cap b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)) \subset b_1 D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu) \cup b_2 D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu) \quad (3.7)$$

where  $M := \{z \in \Omega, \det(\frac{\partial f_i}{\partial z_j}) = 0\}$  is the zero locus of the complex Jacobian of the holomorphic mapping  $f$  on  $\Omega$ .

If  $M \cap bD_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) \neq \emptyset$ , then, from  $\min\{n_{1+\epsilon}, n_2, \dots, n_l\} \geq 2$ , we have  $M \cap b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) \neq \emptyset$ . Take an irreducible component  $M'$  of  $M$  with  $M' \cap b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) \neq \emptyset$ . Then the intersection  $E_{M'}$  of  $M'$  with  $b_0 D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$  is a real analytic submanifold of dimensional  $2(n_0 + n_1 + \dots + n_l) - 3$  on a dense, open subset of  $E_{M'}$ . By (3.7), we have  $f(E_{M'}) \subset b_1 D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu) \cup b_2 D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu)$ . Hence

$$f(M' \cap D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)) \subset \bigcup_{j=1+\delta}^l Pr_i(D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu)), \quad (3.8)$$

where  $Pr_i(D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu)) := \{(z, w_{(1)}, \dots, w_{(l)}) \in D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu), \|w_{(i)}\| = 0\}$  ( $1 + \delta \leq i \leq l$ ), by the uniqueness theorem. Since  $\text{codim} M' = 1$ ,  $\text{codim}[\bigcup_{j=1+\delta}^l Pr_i(D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu))] \geq \min\{m_{1+\delta}, \dots, m_l, m_1 + \dots + m_l\} \geq 2$  and  $f : D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) \rightarrow D_{m_0}^{\mathbf{m}, \mathbf{Q}}(\nu)$  is proper, this is contradiction with (3.8). Thus we have  $M \cap bD_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) = \emptyset$ .

Let  $S := M \cap D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu)$ . Hence we have

$$S \subset D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu), \quad \overline{S} \cap bD_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) = \emptyset.$$

If  $S \neq \emptyset$ , then  $S$  is a complex analytic set in  $\mathbb{C}^{n_0+n_1+\dots+n_l}$  also. For any  $(z, w) \in S$ , we have  $|w_{l_i}|^{2p_i} \leq \sum_{j=1}^l \|w_{(j)}\|^{2p_j} \leq e^{-\mu\|z\|^2} \leq 1$ . Thus

$$|w_{l_i}|^2 \leq 1 \leq 1 + \|(z, w')\|, \quad (3.9)$$

where  $w = (w', w_{l_i})$ . Then  $S$  is an algebraic set of  $\mathbb{C}^{n_0+n_1+\dots+n_l}$  by §7.4 Th. 3 of Chirka [5].

Suppose  $S_1$  is an irreducible component of  $S$ . Let  $\overline{S_1}$  be the closure of  $S_1$  in  $\mathbb{P}^{n_0+n_1+\dots+n_l}$ . Then by §7.2 Prop. 2 of Chirka [5],  $\overline{S_1}$  is a projective algebraic set and  $\dim \overline{S_1} = n_0 + n_1 + \dots + n_l - 1$ . Let  $[\xi, z, w]$  be the homogeneous coordinate in  $\mathbb{P}^{n_0+n_1+\dots+n_l}$ , we embed  $\mathbb{C}^{n_0+n_1+\dots+n_l}$  into  $\mathbb{P}^{n_0+n_1+\dots+n_l}$  as the affine piece  $U_0 = \{[\xi, z, w] \in \mathbb{P}^{n_0+n_1+\dots+n_l}, \xi \neq 0\}$  by  $(z, w) \mapsto [1, z, w]$ . Then we have

$$D_{n_0}^{\mathbf{n}, \mathbf{P}}(\mu) \cap U_0 = \left\{ [\xi, z, w], \xi \neq 0, \sum_{j=1}^l \frac{\|w_{(j)}\|^{2p_j}}{|\xi|^{2p_j}} < e^{-\mu \frac{\|z\|^2}{|\xi|^2}} \right\}.$$

Let  $H = \{\xi = 0\} \subset \mathbb{P}^{n_0+n_1+\dots+n_l}$ . Consider another affine piece  $U_1 = \{[\xi, z, w] \in \mathbb{P}^{n_0+n_1+\dots+n_l}, z_1 \neq 0\}$  with affine coordinate  $(\zeta, t, s) = (\zeta, t_2, \dots, t_{n_0}, s_{(1)}, \dots, s_{(l)})$ . Let  $t' = (1, t_2, \dots, t_{n_0})$ . Since  $\frac{\|w_{(j)}\|^{2p_j}}{|\xi|^{2p_j}} = \frac{\|w_{(j)}\|^{2p_j} |z_1|^{2p_j}}{|z_1|^{2p_j} |\xi|^{2p_j}} = \frac{\|s_{(j)}\|^{2p_j}}{|\zeta|^{2p_j}}$  and  $e^{-\mu \frac{\|z\|^2}{|\xi|^2}} = e^{-\mu \frac{\|z\|^2 |z_1|^2}{|z_1|^2 |\xi|^2}} = e^{-\mu \frac{1+t_2^2+\dots+t_{n_0}^2}{|\zeta|^2}}$ , we obtain

$$\begin{aligned} & D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu) \cap U_0 \cap U_1 \\ &= \left\{ (\zeta, t_2, \dots, t_{n_0}, s_{(1)}, \dots, s_{(l)}) \in \mathbb{C}^{n_0+n_1+\dots+n_l}, \sum_{j=1}^l \frac{\|s_{(j)}\|^{2p_j}}{|\zeta|^{2p_j}} < e^{-\mu \frac{\|t'\|^2}{|\zeta|^2}} \right\}. \end{aligned} \quad (3.10)$$

Let  $S' = \overline{S_1} \cap U_1$  and  $H_1 = H \cap U_1 = \{\zeta = 0\}$  (note  $\xi = \frac{\zeta}{z_1}$ ). For every  $u \in S' \cap H_1$ , there exists a sequence of points  $\{u_k\} \subset \overline{S_1} \cap ((U_0 \cap U_1) \setminus H_1)$  such that  $u_k \rightarrow u$  ( $k \rightarrow \infty$ ). The formula (3.10) implies

$$\|s_{(j)}(u_k)\|^{2p_j} \leq |\zeta(u_k)|^{2p_j} e^{-\mu \frac{\|t'\|^2}{|\zeta(u_k)|^2}}, \quad 1 \leq j \leq l. \quad (3.11)$$

Since  $u \in H_1$ , that means  $\zeta(u) = 0$  and  $\zeta(u_k) \rightarrow 0$  ( $k \rightarrow \infty$ ). Therefore we have  $\|s_{(j)}(u)\|^{2p_j} \leq 0$  ( $1 \leq j \leq l$ ) as  $k \rightarrow \infty$ . Hence

$$S' \cap H_1 \subset \left\{ \zeta = 0, s_{(1)} = \dots = s_{(l)} = 0 \right\}.$$

Then  $\dim(S' \cap H_1) \leq n_0 - 1$ . Shafarevich [20] §6.2 Th. 6 implies

$$n_0 - 1 \geq \dim(S' \cap H_1) \geq \dim S' + \dim H_1 - n_0 - n_1 - \dots - n_l \geq \dim S' - 1. \quad (3.12)$$

This means  $\dim S' \leq n_0$ , and thus  $n_0 + n_1 + \dots + n_l - 1 = \dim S' \leq n_0$ . Therefore, we get  $n_1 + \dots + n_l \leq 1$ , a contradiction with assumption  $\min\{n_{1+\epsilon}, n_2, \dots, n_l, n_1 + \dots + n_l\} \geq 2$ .

Therefore,  $S = \emptyset$  and thus  $f$  is unbranched. Since the generalized Fock-Bargmann-Hartogs domain is simply connected,  $f : D_{n_0}^{\mathbf{n}, \mathbf{p}}(\mu) \rightarrow D_{m_0}^{\mathbf{m}, \mathbf{q}}(\nu)$  is a biholomorphism. The proof is completed.  $\square$

**Acknowledgments** We sincerely thank the referees, who read the paper very carefully and gave many useful suggestions. E. Bi was supported by the Natural Science Foundation of Shandong Province, China (No.ZR2018BA015), and Z. Tu was supported by the National Natural Science Foundation of China (No.11671306).

## References

- [1] Alexander, H.: Proper holomorphic mappings in  $\mathbf{C}^n$ . Indiana Univ. Math. J. **26**, 137-146 (1977)
- [2] Bargmann, V.: On a Hilbert space of analytic functions and an associated integral transform Part II: A family of related function spaces application to distribution theory. Communications on Pure and Applied Mathematics. **20**(1), 1-101 (1967)
- [3] Bedford, E., Bell, S.: Proper self-maps of weakly pseudoconvex domains. Math. Ann. **261**, 47-49 (1982)
- [4] Bell, S.: The Bergman kernel function and proper holomorphic mappings. Trans. Amer. Math. Soc. **270**, 685-691 (1982)
- [5] Chirka, E.M.: Complex analytic sets. Springer Science & Business Media (1989)
- [6] D'Angelo, J.P.: A note on the Bergman kernel. Duke. Math. J. **45**(2), 259-265 (1978)
- [7] D'Angelo, J.P.: An explicit computation of the Bergman kernel function. J. Geom. Anal. **4**(1), 23-34 (1994)
- [8] Diederich, K., Forneaess, J.E.: Proper holomorphic images of strictly pseudoconvex domains. Math. Ann. **259**, 279-286 (1982)
- [9] Dini, G., Primicerio, A.S.: Localization principle of automorphisms on generalized pseudoellipsoids. J. Geom. Anal. **7**(4), 575-584 (1997)
- [10] Isaev, A.V.: Hyperbolic  $n$ -dimensional manifolds with automorphism group of dimension  $n^2$ . Geom. Funct. Anal. **17**(1), 192-219 (2007)

- [11] Isaev, A.V., Krantz, S.G.: On the automorphism groups of hyperbolic manifolds. *J. Reine Angew. Math.* **534**, 187-194 (2001)
- [12] Ishi, H., Kai, C.: The representative domain of a homogeneous bounded domain. *Kyushu J. Math.* **64**(1), 35-47 (2009)
- [13] Kim, H., Ninh, V.T., Yamamori, A.: The automorphism group of a certain unbounded non-hyperbolic domain. *J. Math. Anal. Appl.* **409**(2), 637-642 (2014)
- [14] Kim, K.T., Verdiani, L.: Complex  $n$ -dimensional manifolds with a real  $n^2$ -dimensional automorphism group. *J. Geom. Anal.* **14**(4), 701-713 (2004)
- [15] Kodama, A.: On the holomorphic automorphism group of a generalized complex ellipsoid. *Complex Variables and Elliptic Equations.* **59**(9), 1342-1349 (2014)
- [16] Kodama, A., Krantz, S.G., Ma, D.W.: A characterization of generalized pseudoellipsoids in  $\mathbf{C}^N$  and related results. *Indiana Univ. Math. J.* **41**, 173-195 (1992)
- [17] Ligocka, E.: The regularity of the weighted Bergman projections. *Seminar on Deformations.* Springer, Berlin Heidelberg (1985)
- [18] Ligocka, E.: On the Forelli-Rudin construction and weighted Bergman projections. *Studia Math.* **94**(3), 257-272 (1989)
- [19] Pinčuk, S.I.: On the analytic continuation of biholomorphic mappings. *Math. USSR Sb.* **27**, 375-392 (1975)
- [20] Shafarevich, I.R.: *Basic algebraic geometry.* Springer-Verlag, New York (1977)
- [21] Trybula, M.: Proper holomorphic mappings, Bell's formula, and the Lu Qi-Keng problem on the tetrablock. *Arch. Math.* **101**, 549-558 (2013)
- [22] Tsuboi, T.: Bergman representative domains and minimal domains. *Japanese J. Math.* **29**, 141-148 (1959)
- [23] Tu, Z.H., Wang, L.: Rigidity of proper holomorphic mappings between certain unbounded non-hyperbolic domains. *J. Math. Anal. Appl.* **419**, 703-714 (2014)
- [24] Tu, Z.H., Wang, L.: Rigidity of proper holomorphic mappings between equidimensional Hua domains. *Math. Ann.* **363**, 1-34 (2015)
- [25] Yamamori, A.: The Bergman kernel of the Fock-Bargmann-Hartogs domain and the polylogarithm function. *Complex Var. Elliptic Equ.* **58**(6), 783-793 (2013)