

Ground state heavy tetraquark production in heavy quark fragmentation

S. Mohammad Moosavi Nejad^{a,b*} and Nahid Amiri^a

^(a)*Faculty of Physics, Yazd University, P.O. Box 89195-741, Yazd, Iran and*

^(b)*School of Particles and Accelerators, Institute for Research in Fundamental Sciences (IPM), P.O.Box 19395-5531, Tehran, Iran*

(Dated: January 19, 2022)

During recent years, the study of exotic hadrons including tetraquarks and pentaquarks has attracted a lot of interests and more studies are in progress experimentally and theoretically. It is well-known that at sufficiently large transverse momentum the dominant production mechanism for standard heavy hadrons (mesons/baryons) is actually the fragmentation so that the same mechanism is also proposed for the production of heavy exotic hadrons. This work is the first attempt to study the direct fragmentation of a heavy quark into ground state heavy tetraquarks in leading order of perturbative QCD. In this regard, we will present an analytical expression for the fragmentation production of neutral hidden flavor tetraquarks ($Qq\bar{Q}\bar{q}$) which includes most of the kinematical and dynamical properties of the process.

I. INTRODUCTION

Since the early days of the quark model it has been assumed the possibility of the existence of exotic multi-quark hadrons with the content of the valence quarks/antiquarks different from those in standard mesons and baryons. In this context, the exotic hadrons such as tetraquarks (composed of two quarks and two antiquarks) and pentaquarks (including four quarks and one antiquark) had been anticipated by Murray Gell-Mann and George Zweig in their 1964 papers [1]. The only necessary condition for these new multi-quark states is to be color singlets. However, the absence of convincing experimental evidences for such exotic structures made their investigation of marginal interest for several decades. Nevertheless, the situation dramatically changed in the last two decades when the first explicit experimental evidence of the existence of these exotic hadrons became available, for recent reviews see Refs. [2–4] and references therein. Within the past decade several exotic hadrons, the majority pertaining to the class of heavy tetraquarks with at least two heavy quarks, were clearly confirmed by the BELLE, BESIII and LHCb experimental collaborations. The first experimental evidence for a heavy tetraquark was the $X(3872)$ state [5] observed in 2003 by the Belle collaboration in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays. The decay of this state as well as the state $Z_c(3900)^\pm$ into the J/ψ meson (observed in BESIII) confirmed that these exotic particles are formed of two quarks and two antiquarks. Very recently, a narrow structure around 6.9 GeV, named as $X(6900)$, is observed by the CERN LHCb in the process $pp \rightarrow J/\psi J/\psi X$ [6]. This is the first candidate of fully-heavy tetraquarks observed in experiment which is proposed to be either the first radial (2S) excitation or the second orbital (1D) excitation of the hidden-charm tetraquark $cc\bar{c}\bar{c}$.

Nowadays, convincing candidates for both the exotic tetraquark ($qq\bar{q}\bar{q}$) and pentaquark ($qqqq\bar{q}$) states are found. In the literature, there is no definite consensus about the composition of these states and different interpretations for the tetraquark candidates are proposed, for example: molecules composed from two mesons loosely bound by the meson exchange, compact tetraquarks composed from a diquark and antidiquark bound by strong forces, hadroquarkonium composed of a heavy quarkonium embedded in a light meson, kinematic cusps, etc. The distinction between various models is a very complicated experimental task. Among exotic hadrons (tetraquarks, pentaquarks, hexaquarks, etc), heavy tetraquarks are of particular interest since the presence of a heavy quark increases the binding energy of the bound system and, in conclusion, the possibility that such tetraquarks will have masses below the thresholds for decays to mesons with open heavy flavor. In this case the strong decays, which proceed through the quark and antiquark rearrangements, are kinematically forbidden and the corresponding tetraquarks can decay only weakly or electromagnetically and thus they should have a tiny decay width.

When the strong interactions are concerned, the investigation of production mechanism of heavy standard/exotic hadrons is a powerful tool to understand the dynamics of strong interactions. Basically, two different mechanisms are assumed for the production of heavy hadrons: recombination and fragmentation [7]. In the second mechanism, the fragmentation refers to the process of a parton which carries large transverse momentum and subsequently forms a jet containing the expected hadron [8]. At sufficiently large transverse momentum of heavy hadron production, the

*Electronic address: mmoosavi@yazd.ac.ir

direct leading order production scheme (recombination mechanism) is normally suppressed while the fragmentation mechanism becomes dominant, though it is formally of higher order in the strong coupling constant α_s [8, 9]. In Refs. [10–17], we calculated the perturbative QCD fragmentation function (FF) for a heavy quark to fragment into the S -wave heavy mesons and baryons at leading-order (LO) and next-to-leading order (NLO). In Ref. [18], it has been pointed out that the dominant production mechanism for pentaquarks consisting of a heavy quark is heavy quark fragmentation, similar to heavy-light mesons and baryons consisting of a heavy quark. So, this mechanism is also adopted in our work for the production of heavy tetraquark states and we, for the first time, study the production mechanism of heavy tetraquarks in the fragmentation of a heavy quark at lowest order of perturbative QCD. In our calculation, tetraquarks are assumed as four-quark $Qq\bar{Q}\bar{q}$ states which are tightly bound by the color forces. Our analytical results are presented for the fragmentation function of heavy quark Q into neutral hidden flavor tetraquarks $Qq\bar{Q}\bar{q}$ with $Q = c, b$ and $q = u, d, s$. Having these FFs it would be possible to evaluate the production rate of these heavy states in e^+e^- annihilation and hadron colliders. In fact, the specific importance of FFs is for their model independent predictions of the cross sections at colliders. However, due to a large number of decay modes the results from e^+e^- annihilation seem to be small but sizable rates are expected in energetic hadron colliders where a large number of them is produced. To assess the prospects for the experimental study of heavy tetraquarks in present and future colliders, it is important to know the accurate predictions for their production rates and, in conclusion, their FFs.

This paper is organized as follows. In Sec. II, we present the theoretical formalism for the heavy tetraquark FF in the Suzuki model and present our numerical results, and our discussions and conclusions are given in the last section.

II. HEAVY TETRAQUARK FRAGMENTATION

A motivation for studying the production of multi-quark structures is for better understanding of the dynamics of strong interactions and the confinement mechanism. It is well known that the dominant mechanism to produce standard heavy hadrons at sufficiently large transverse momentum is fragmentation so this mechanism is valid for the production of heavy tetraquarks too. Fragmentation refers to the production of a parton with a large transverse momentum which subsequently decays to form a jet containing an expected hadron. It is hence important to obtain the corresponding fragmentation function in order to properly estimate the production rate of a specific hadronic state. According to the QCD factorization theorem, the cross section for the production of hadron H in the typical scattering process $A + B \rightarrow H + X$, can be expressed as

$$d\sigma = \sum_{a,b,c} \int_0^1 dx_a \int_0^1 dx_b \int_0^1 dz f_{a/A}(x_a, Q) f_{b/B}(x_b, Q) d\hat{\sigma}(a + b \rightarrow c + X) D_c^H(z, Q), \quad (1)$$

where a and b are incident partons in the colliding initial hadrons A and B , respectively, $f_{a/A}$ and $f_{b/B}$ are the nonperturbative parton distribution function (PDFs) at the scale Q^2 of the partonic subprocess $a + b \rightarrow c + X$, c is the fragmenting parton and X stands for any unobserved jet. Here, $D_c^H(z, Q)$ is the FF at the scale Q which can be obtained by evolving from the initial FF $D_c^H(z, \mu_0)$ using the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) renormalization group equations [19]. In electron-positron annihilation, one does not need to deal with the nonperturbative PDFs as in hadron collisions. This process has in general less contributions by background processes compared to hadron collisions and the uncertainties introduced by parton density functions are not appeared. The process of inclusive heavy hadron production in electron-positron annihilation, i.e., $e^+e^- \rightarrow (\gamma, Z) \rightarrow H + X$, has the cross section as [20–22]

$$\frac{d\sigma}{dx}(x, s) = \sum_a \int_x^1 \frac{dy}{y} \frac{d\sigma}{dy}(y, \mu_R, \mu_F) D_a^H\left(\frac{x}{y}, \mu_F\right), \quad (2)$$

where, a stands for the partons $a = g, u, \bar{u}, \dots, b, \bar{b}$ and μ_F and μ_R are the factorization and renormalization scales, respectively. Generally, one can choose two different values for these scales, but a common choice consists of setting $\mu_R = \mu_F = \sqrt{s}$. Here, $x = 2E_H/\sqrt{s}$ measures the energy of hadron in units of the beam energy in the center-of-mass frame. At lowest-order of perturbative QCD, the cross section of relevant partonic subprocesses is given by [23]

$$\frac{d\sigma}{dy}(e^+e^- \rightarrow q\bar{q}) = N_c \sigma_0 (V_{q_i}^2 + A_{q_i}^2) \delta(1 - y). \quad (3)$$

Here, $N_c = 3$ is the number of quark colors, V_{q_i} and A_{q_i} are the effective vector and axial-vector couplings of quark q_i to the photon and Z boson. For small energies ($\sqrt{s} \ll M_Z$), for the summation of squared effective electroweak




Figure 1: The lowest order Feynman diagram contributing to the fragmentation of a heavy quark Q into a heavy tetraquark $X(Qq\bar{Q}\bar{q})$. The four momenta are also labeled.

charges we have $V_{q_i}^2 + A_{q_i}^2 = e_{q_i}^2$ where e_{q_i} is the electric charge of the quark q_i [24]. In Eq. (3), $\sigma_0 = 4\pi\alpha^2/(3s)$ is the leading order total cross section of $e^+e^- \rightarrow \mu^+\mu^-$ for the massless muons, in which α is the Sommerfeld fine-structure constant. Therefore, for the cross section of $e^+e^- \rightarrow H + X$ at LO, one has

$$\frac{d\sigma^{LO}}{dx}(x, s) = N_c\sigma_0 \sum_q e_q^2 [D_q^H(x, \mu) + D_{\bar{q}}^H(x, \mu)] = \frac{8\pi\alpha^2}{s} \sum_q e_q^2 D_q^H(x, \mu), \quad (4)$$

where we assumed $D_q^H = D_{\bar{q}}^H$ for $q = u, d, s, c, b$.

Conventionally, the fragmentation mechanism is described by the function $D_i^H(z, \mu)$ which refers to the probability for a parton i at the fragmentation scale μ to fragment into a hadron H carrying away a fraction z of its momentum. The FFs are related to the low-energy part of the hadroproduction processes and forms the nonperturbative aspect of QCD but, fortunately, it was found that these functions for heavy hadron productions are analytically calculable by virtue of perturbative QCD (pQCD) with limited phenomenological parameters. Historically, the first theoretical effort to describe the procedure of heavy hadron production was made by Bjorken [25] by using a naive quark-parton model and in the following the pQCD scheme was applied by Suzuki [26], Amiri and Ji [27]. Among them, an elaborate model of fragmentation which does contain spin information has been proposed by Suzuki [28]. This model includes most of the kinematical and dynamical properties of the splitting process and gives a detailed insight about the fragmentation process. In this approach, Suzuki calculates the fragmentation function of a heavy quark using the convenient Feynman diagrams for the parton level of the process and also by considering the wave function of heavy bound state which contains the nonperturbative dynamic of hadroproduction process. In fact, the Suzuki model mixes a perturbative picture with nonperturbative dynamics of fragmentation and not only predicts the z -dependence of the FFs, but also their dependence on the transverse momentum of the hadron relative to the jet.

Here, using the Suzuki model we focus on the heavy tetraquark FF for which the only Feynman diagram at leading order in α_s is shown in Fig. 1. Any other possible Feynman diagram is related to higher orders in α_s . Considering this diagram we first set the relevant four-momenta as

$$\begin{aligned} p_\mu &= [p_0, \mathbf{p}_T, p_L] & , & & s_\mu &= [s_0, \mathbf{0}, s_L] \\ s'_\mu &= [s'_0, \mathbf{p}_T, s'_L] & , & & t_\mu &= [t_0, \mathbf{0}, t_L] \\ t'_\mu &= [t'_0, \mathbf{0}, t'_L] & , & & r_\mu &= [r_0, \mathbf{0}, r_L] \end{aligned} \quad (5)$$

where we assumed that the produced tetraquark X moves along the \hat{z} -axes (fragmentation axes). The four-momentum of tetraquark is also written as $\bar{P}_\mu = [\bar{P}_0, \mathbf{0}, \bar{P}_L]$ so that $\bar{P}_L = s_L + r_L + t_L + t'_L$. In the Suzuki model the FF for the

production of unpolarized ground state tetraquark is defined as [28]

$$D_Q^X(z, \mu_0) = \frac{1}{1 + 2s_Q} \sum_{spin} \sum_{color} \int |T_X|^2 \delta^3(\bar{\mathbf{P}} + \mathbf{s}' - \mathbf{p}) d^3 \bar{\mathbf{P}} d^3 \mathbf{s}', \quad (6)$$

where μ_0 is the fragmentation scale, s_Q refers to the spin of the fragmenting quark and the summation is going over the spins and colors. Four momenta are as labeled in Fig. 1. In computing Eq. (6), following Ref. [28], we adopt the infinite momentum frame where the fragmentation variable in the usual light-cone form, i.e. $z = (\bar{P}_0 + \bar{P}_L)/(p_0 + p_L)$, is reduced to a more popular one as $z = \bar{P}_0/p_0 = E_X/E_Q$. From experimental point of view, this definition is more suitable since refers to the energy fraction of the fragmenting heavy quark which is taken away by the produced tetraquark. This scaling variable ranges as $0 \leq z \leq 1$.

In Eq. (6), T_X is the probability amplitude of the tetraquark production which, at the large momentum transfer, is expressed in terms of the hard scattering amplitude T_H and the process-independent distribution amplitude Φ_B as [29, 30]

$$T_X(\bar{P}, s') = \int [dx] T_H(\bar{P}, s', x_i) \Phi_B(x_i, Q^2), \quad (7)$$

where $[dx] = \prod_{i=1}^4 dx_i \delta(1 - \sum_{i=1}^4 x_i)$. Here, x_i 's stand for the tetraquark energy fractions carried by its constituent quarks which are defined as

$$x_1 = \frac{s_0}{\bar{P}_0}, \quad x_2 = \frac{r_0}{\bar{P}_0}, \quad x_3 = \frac{t'_0}{\bar{P}_0}, \quad x_4 = \frac{t_0}{\bar{P}_0}. \quad (8)$$

The energy conservation law requires that $\sum_{i=1}^4 x_i = 1$.

The scheme applied to describe the probability amplitude T_X as the convolution in Eq. (7) is convenient to absorb the soft behavior of the bound state (tetraquark in our example) into the hard scattering amplitude T_H [29]. The short-distance part of the amplitude, i.e. T_H , can be computed perturbatively from quark-gluon subprocesses at leading-order or higher-order QCD approximations. The long-distance distribution amplitude Φ_B which contains the bound state nonperturbative dynamic of the outgoing tetraquark, is the probability amplitude for a $Qq\bar{Q}\bar{q}$ -set to evolve into a particular bound state. The underlying link between hadronic phenomena in QCD at long- and short-distances is the hadronic wave function. In fact, the nonperturbative aspect of the hadroproduction processes is contained in the bound state of the hadron which is described by the wave function Φ_B . Following Ref. [28] and according to the Lepage-Brodsky's approach [31] we neglect the relative motion of the constituent quarks inside the tetraquark therefore we assume, for simplicity, that the constituent quarks are emitted collinearly with each other and move along the \hat{z} -axes. In fact, in the Suzuki model a multi-quark state is replaced by collinear constituents with neglecting their Fermi motion and the nonperturbative aspect of the hadroproduction is included in the wave function of the heavy tetraquark state. By this simplicity, the relative motion of the constituent quarks is effectively nonrelativistic and this allows one to estimate the nonrelativistic wave function of heavy tetraquark as a delta function form, see also [13]. Therefore, the distribution amplitude for a S-wave heavy tetraquark with neglecting the Fermi motion reads [32]

$$\Phi_B \approx f_B \delta(x_i - \frac{m_i}{m_X}). \quad (9)$$

where m_X is the tetraquark mass and f_B refers to the decay constant of hadron. In Ref. [33], we studied the effect of meson wave function on its FF by considering a typical mesonic wave function which is different of the delta function and is the nonrelativistic limit of the solution of Bethe-Salpeter equation with the QCD kernel [29]. There, we showed that the Fermi motion effect can be neglected with a reasonable approximation.

With the delta function approximation for the tetraquark wave function (9), we are assuming that the contribution of each constituent from the tetraquark energy is proportional to its mass, i.e. $x_i = m_i/m_X$ where $m_X = 2(m_q + m_Q)$. Considering the definitions of fragmentation variable ($z = \bar{P}_0/p_0$) and the fractions x_i 's presented in Eq. (8) one also may write the parton energies in terms of the initial heavy quark energy (p_0) as: $s_0 = x_1 z p_0$, $r_0 = x_2 z p_0$, $t'_0 = x_3 z p_0$, $t_0 = x_4 z p_0$ and $s'_0 = (1 - z)p_0$ where $x_1 = x_4 = m_Q/m_X$ and $x_2 = x_3 = m_q/m_X$.

In Eq. (7), the amplitude T_H is, in essence, the partonic cross section to produce a set of heavy quarks with certain quantum numbers that in the QCD perturbation theory considering the Feynman rules is written as

$$T_H = \frac{16\pi^2 \alpha_s(2m_Q) \alpha_s(2m_q) m_X m_Q^2 C_F \Gamma}{2\sqrt{2\bar{P}_0 s'_0 p_0} D_0}, \quad (10)$$

where, $D_0 = \bar{P}_0 + s'_0 - p_0$ is the energy denominator, $\alpha_s(\mu)$ is the strong coupling constant at the renormalization scale μ and C_F is the color factor which is calculated using color line counting rule.

In Eq. (10), Γ represents an appropriate combination of the propagators and the spinorial parts of the amplitude. Considering Fig. 1 for the transition $Q \rightarrow X_{Qq\bar{Q}\bar{q}}$ the transition amplitude at leading order is written as

$$\Gamma = G\bar{u}(t)\gamma^\mu u(p) \left[\bar{u}(t')\gamma_\mu(q_2 + m_q)\gamma_\nu v(r) \right] v(s)\gamma^\nu \bar{u}(s') \quad (11)$$

where G is related to the denominator of propagators as

$$G = \frac{1}{q_1^2(q_2^2 - m_q^2)q_3^2} = \frac{1}{8(m_Q^2 - p \cdot t)^2(m_Q^2 + s \cdot s')^2(m_Q^2 + r \cdot s + r \cdot s' + s \cdot s')^2}. \quad (12)$$

In squaring the amplitude Γ , one needs the following dot products of the relevant four-momenta:

$$\begin{aligned} r \cdot t' &= m_q^2, & s \cdot t &= m_Q^2 \\ r \cdot s &= t \cdot t' = s \cdot t' = m_q m_Q, \\ p \cdot s &= p \cdot t = \frac{m_X m_Q}{2z} + \frac{z m_Q}{2m_X} (m_Q^2 + p_T^2), \\ p \cdot s' &= \frac{1}{2} (m_Q^2 + p_T^2) \left(1 - z + \frac{1}{1-z}\right) - p_T^2, \\ r \cdot s' &= s' \cdot t' = \frac{z m_q}{2m_X (1-z)} (m_Q^2 + p_T^2) + \frac{m_X m_q (1-z)}{2z}, \\ s \cdot s' &= t \cdot s' = \frac{m_Q m_X (1-z)}{2z} + \frac{z m_Q}{2m_X (1-z)} (m_Q^2 + p_T^2), \\ p \cdot t' &= p \cdot r = \frac{m_q m_X}{2z} + \frac{z m_q}{2m_X} (m_Q^2 + p_T^2). \end{aligned} \quad (13)$$

To obtain the FF for an unpolarized S-wave heavy tetraquark $X_{Qq\bar{Q}\bar{q}}$, considering Eqs. (6-10) one has

$$D_c^X(z, \mu_0) = (4\pi^2 f_B \alpha_s (2m_q) \alpha_s (2m_Q) m_X m_Q^2 C_F)^2 \int \frac{d^3 \bar{\mathbf{P}} d^3 \mathbf{s}' \sum_{spin} |\Gamma|^2 \delta^3(\bar{\mathbf{P}} + \mathbf{s}' - \mathbf{p})}{\bar{P}_0 p_0 s'_0 (\bar{P}_0 + s'_0 - p_0)^2}. \quad (14)$$

To perform the phase space integrations we start with the following integral

$$\int \frac{d^3 \bar{\mathbf{P}} \delta^3(\bar{\mathbf{P}} + \mathbf{s}' - \mathbf{p})}{\bar{P}_0 p_0 (\bar{P}_0 + s'_0 - p_0)^2} = \int \frac{\bar{P}_0 d^3 \bar{\mathbf{P}} \delta^3(\bar{\mathbf{P}} + \mathbf{s}' - \mathbf{p})}{p_0 (\bar{P}_0^2 - (p_0 - s'_0)^2)^2} = \frac{z p_0}{p_0 (m_X^2 - (p - s')^2)^2} = \frac{z}{(m_X^2 - 2m_Q^2 + 2p \cdot s')^2}. \quad (15)$$

According to the Suzuki approach, in Eq. (14) instead of integrating over the transverse momentum of outgoing heavy quark Q , we simply replace the integration variable by its average value, i.e.

$$\int \frac{d^3 \mathbf{s}'}{s'_0} g(z, s'_T) = \int g(z, s'_T) \frac{ds'_L d^2 s'_T}{s'_0} = m_Q^2 g(z, \langle p_T^2 \rangle), \quad (16)$$

where the average value $\langle p_T^2 \rangle$ is a free parameter which can be specified phenomenologically.

Putting all in Eq. (14), the FF of heavy tetraquark X reads

$$D_Q^X(z, \mu_0) = N \frac{z \times \sum_{spin} \Gamma \bar{\Gamma}}{(m_X^2 - 2m_Q^2 + 2p \cdot s')^2} = N \frac{z \times \sum_{spin} \Gamma \bar{\Gamma}}{[m_X^2 - (m_Q^2 + \langle p_T^2 \rangle)(1 + z - \frac{1}{1-z})]^2}, \quad (17)$$

where $N = (16\pi^2 f_B \alpha_s (2m_q) \alpha_s (2m_Q) m_q m_Q^4 C_F)^2$ but it is related to the normalization condition: $\int dz D(z, \mu_0) = 1$ [27, 28]. Using the traditional trace technique the squared amplitude ($\sum |\Gamma|^2$) reads

$$\sum_{spin} \Gamma \bar{\Gamma} = \frac{G^2}{z^3 (1-z)} \left(z^6 \langle p_T^2 \rangle^3 + \alpha z^4 \langle p_T^2 \rangle^2 + \beta z^2 \langle p_T^2 \rangle - \gamma \right), \quad (18)$$




Figure 2: Fragmentation of c -quark into heavy tetraquark $X_{cu\bar{c}\bar{u}}$ at lowest-order of perturbative QCD. The uncertainty band due to the variation of charm mass is also shown.

where

$$\begin{aligned}
\alpha &= m_Q^2(11z^2 - 18z + 12) + 6m_q^2(2z^2 - z + 2) + 4m_q m_Q(3z^2 - 4z + 6), \\
\beta &= m_Q^4(19z^4 - 84z^3 + 168z^2 - 144z + 48) + 4m_Q^2 m_q^2(6z^4 - 37z^3 + 102z^2 - 112z + 72) \\
&\quad - 8m_q^4(z^3 - 6z^2 + 6z - 6) - 16m_Q m_q^3(3z^3 - 12z^2 + 14z - 12) + 8m_q m_Q^3(3z^4 - 22z^3 + 54z^2 - 52z + 24), \\
\gamma &= -64m_Q m_q^5(z^3 + 3z^2 - 10z + 6) + 8m_q^4 m_Q^2(z^5 - 6z^4 + 42z^3 - 158z^2 + 236z - 120) + 16m_q^3 m_Q^3(3z^5 - \\
&\quad 24z^4 + 98z^3 - 196z^2 + 192z - 80) - 2m_q^2 m_Q^4(6z^6 - 71z^5 + 422z^4 - 1232z^3 + 1872z^2 - 1424z + 480) - \\
&\quad 4m_q m_Q^5(3z^6 - 40z^5 + 182z^4 - 424z^3 + 544z^2 - 352z + 96) - 32m_q^6(z^3 - 3z + 2) - m_Q^6(z^2 - 2z + 4)(3z^2 - 8z + 4)^2.
\end{aligned} \tag{19}$$

In the Suzuki model, the fragmentation function depends on both the fragmentation variable z and the parameter $\langle p_T^2 \rangle$ (the transverse momentum of produced hadron relative to the parent jet). In Ref. [32], it is shown that the choice of $\langle p_T^2 \rangle = 1 \text{ GeV}^2$ is an extreme value for this quantity and any higher value will produce the peak even at lower- z regions. In this work we adopt this value for the transverse momentum of heavy tetraquark.

The z -dependence of FFs is not yet calculable at each scale, however once they are computed at the initial scale μ_0 their μ evolution can be specified by the DGLAP evolution equations [19]. Here, we set the initial scale of fragmentation to $\mu_0 = m_X + m_Q$ which is the minimum value of the invariant mass of the fragmenting parton. Then, the FF presented in Eq. (17) should be regarded as a model for the heavy quark fragmentation Q into the heavy tetraquark $X(Qq\bar{Q}\bar{q})$ at the initial scale μ_0 . To present our numerical analysis, we adopt the following input parameters [34]:

$$m_c = 1.67 \text{ GeV}, \quad m_b = 4.78 \text{ GeV}, \quad f_B = 0.25 \text{ GeV}, \quad \alpha_s(2m_c) = 0.26, \quad \alpha_s(2m_b) = 0.18. \tag{20}$$

Note that, in Ref. [34] two values are reported for the quark masses. The above masses show the pole masses, i.e. the renormalized quark masses in the on-shell renormalization scheme, which correspond to the values $m_c = 1.27 \pm 0.02 \text{ GeV}$ and $m_b = 4.18 \pm 0.03 \text{ GeV}$ for the \overline{MS} masses (the renormalized quark mass in the modified minimal subtraction scheme [35] which is intimately related to the use of dimensional regularization [36]). The relation between the pole quark mass and the \overline{MS} mass is known to three loops in QCD perturbation theory, details are given in Ref. [37]. In our analysis, to show the effect of variation of heavy quark masses on their corresponding FFs we will consider the charm and bottom quark masses as $1.27 \leq m_c \leq 1.67 \text{ GeV}$ and $4.18 \leq m_b \leq 4.78 \text{ GeV}$.

In Fig. 2, our phenomenological prediction for the FFs of charm into the neutral hidden-charm state $X(cu\bar{c}\bar{u})$ is shown at the starting scale $\mu_0 = m_c + m_X$. The uncertainty band due to the variation of charm quark mass is also




Figure 3: The $b \rightarrow X_{bs\bar{s}}$ FF at leading order perturbative QCD along with the uncertainty band of bottom mass.

plotted. Assuming $m_u \approx m_d$ this result is also valid for the transitions $c \rightarrow X(cd\bar{c}\bar{d}), X(cu\bar{c}\bar{d}), X(cd\bar{c}\bar{u})$. In Fig. 3, the b-quark FF into the neutral hidden-bottom tetraquark including strangeness flavor, i.e. $b \rightarrow X(bs\bar{s})$, is shown considering the mass uncertainty band. Since the b-quark is heavier than the c-quark, as we expected, the peak of b-quark FF shifted toward higher values of z .

Besides the heavy tetraquark FFs, their first moment is also of phenomenological interest and subject to experimental determination. It corresponds to the average fraction of energy that the tetraquark receives from the fragmenting heavy quark and is defined as [24]

$$\langle z \rangle_Q(\mu) = \frac{1}{B_Q(\mu)} \int_0^1 z D_Q^X(z, \mu) dz, \quad (21)$$

where, $B_Q(\mu)$ stands for the branching fraction which is defined as $B_Q(\mu) = \int_0^1 D_Q^X(z, \mu) dz$, where $Q = c, b$. For the tetraquark $X(cu\bar{c}\bar{u})$ the branching fraction and the average energy fraction are $B_c = 2 \times 10^{-3}$ and $\langle z \rangle_c(\mu_0) = 0.635$, respectively, and the corresponding ones for the tetraquark $X(bu\bar{b}\bar{u})$ read $B_b = 1.1 \times 10^{-4}$ and $\langle z \rangle_b = 0.629$. Considering the inclusive differential cross section given in Eq. (4), one can compute the inclusive cross section for the production of heavy tetraquark at LO as

$$\sigma^{LO}(e^+e^- \rightarrow X_{Qq\bar{Q}\bar{q}} + jets) = \frac{8\pi\alpha^2}{s} \sum_q e_q^2 \int_0^1 dz D_q^X(z, \mu), \quad (22)$$

By ignoring the contributions which are appeared at higher orders of perturbation, one has

$$\sigma^{LO}(e^+e^- \rightarrow X_{Qq\bar{Q}\bar{q}} + jets) \approx \frac{8\pi\alpha^2}{s} e_Q^2 B_Q(s). \quad (23)$$

For example, the contributions of $c \rightarrow X_{bu\bar{b}\bar{u}}, b \rightarrow X_{cu\bar{c}\bar{u}}$ or $g \rightarrow X_{bu\bar{b}\bar{u}}/X_{cu\bar{c}\bar{u}}$ are related to very complicated Feynman diagrams which are in higher orders in α_s , because they do not occur directly. Then, their contributions are ignorable in comparison with the one occurs directly, i.e., $b \rightarrow X_{bu\bar{b}\bar{u}}$ and $c \rightarrow X_{cu\bar{c}\bar{u}}$ FFs (17). Finally, having the branching fraction $B_Q(\mu)$ for the transition $B \rightarrow X$ at the center-of-mass energy $\mu = s$ one can compute the corresponding cross section at electron-positron annihilation.

III. CONCLUSION

Besides the standard mesons and baryons there exist hadrons made of more than three quarks/antiquarks or valence gluons. These new types of hadrons consist of exotic states including glueballs, hybrids, tetraquarks, pentaquarks, hexaquarks, etc. In last two decades a large number of these exotic states, especially tetraquarks and pentaquarks, have been observed in the particle factories. However, some of them are now well-established there are also some doubts on existence of some other members. Therefore, exact determinations of the nature, structure and quantum numbers of these exotic states need more experimental efforts. Specifically, there are many exotic states which are proposed in theory but their existence need to be confirmed by the experiments.

It should be noted that, most of heavy multi-quark particles discovered in the experiment have hidden-charm or hidden-bottom quark structure, i.e., they contain $c\bar{c}$ or $b\bar{b}$ in their inner structures. The study of their production mechanism and using it as a probe to the structure of hadrons are among the most active research fields in particle and nuclear physics. In fact, the challenge is to understand the nonperturbative transition from high energy e^+e^- , photon-hadron, and hadron-hadron collisions to physical exotic states.

It is well-known that the dominant mechanism to produce heavy hadrons at high transverse momentum is fragmentation; the production of a high energy parton followed by its fragmentation into the heavy bound states. In this work, using the Suzuki model we, for the first time, studied the fragmentation function of a heavy quark into an unpolarized S-wave heavy tetraquark at leading order perturbative QCD. To be specific, we analytically computed the FF of bottom and charm quark into the neutral hidden flavor tetraquarks ($X|Qq\bar{Q}\bar{q}\rangle$) at the initial scale $\mu_0 = m_X + m_Q$. Using the extracted FFs one can compute the production rates of heavy tetraquarks in hadron colliders.

To have more accurate FFs, one can think of other effects such as the Fermi motion of constituents, the tetraquark mass effect, NLO radiative corrections, etc. Although, including the effects such as the Fermi motion of constituents leads to very complicated integrals which should be solved numerically [33].

-
- [1] M. Gell-Mann, "A Schematic Model of Baryons and Mesons," *Phys. Lett.* **8** (1964) 214.
 - [2] Liu, Y.R.; Chen, H.X.; Chen, W.; Liu, X.; Zhu, S.L. Pentaquark and Tetraquark states. *Prog. Part. Nucl. Phys.* 2019, 107, 237-320; DOI:10.1016/j.ppnp.2019.04.003.
 - [3] Brambilla, N.; Eidelman, S.; Hanhart, C.; Nefediev, A.; Shen, C.P.; Thomas, C.E.; Vairo A.; Yuan, C.Z. The XY Z states: experimental and theoretical status and perspectives. *Phys. Rept.* 2020, 873, 1-154; DOI:10.1016/j.physrep.2020.05.001.
 - [4] Yang, G.; Ping, J.; Segovia, J. Tetra- and penta-quark structures in the constituent quark model. *Symmetry.* 2020, 12, 1869; DOI:10.3390/sym12111869
 - [5] S. K. Choi *et al.* [Belle], "Observation of a narrow charmonium-like state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays," *Phys. Rev. Lett.* **91** (2003), 262001 doi:10.1103/PhysRevLett.91.262001 [arXiv:hep-ex/0309032 [hep-ex]].
 - [6] R. Aaij *et al.* (LHCb Collaboration), Observation of structure in the J/ψ -pair mass spectrum, *Sci. Bull.* 65, 1983 (2020).
 - [7] A. P. Martynenko and V. A. Saleev, "Heavy quark fragmentation functions in the heavy quark effective theory," *Phys. Rev. D* **53** (1996) 6666.
 - [8] E. Braaten and T. C. Yuan, "Gluon fragmentation into heavy quarkonium," *Phys. Rev. Lett.* **71** (1993) 1673.
 - [9] M. Kramer, 1, "Quarkonium production at high-energy colliders," *Prog. Part. Nucl. Phys.* **47** (2001) 141.
 - [10] S. M. Moosavi Nejad, M. Torkian and M. Delpasand, "NLO QCD corrections to heavy baryon fragmentation functions through quark-diquark model," *Phys. Rev. D* **102** (2020) no.3, 034025 doi:10.1103/PhysRevD.102.034025
 - [11] S. M. Moosavi Nejad and P. Sartipi Yarahmadi, "Heavy quark fragmentation functions at next-to-leading perturbative QCD," *Eur. Phys. J. A* **52** (2016) no.10, 315.
 - [12] S. M. Moosavi Nejad and A. Armat, "Study of bottom and charmed baryons in quark-diquark model," *Eur. Phys. J. A* **56** (2020) no.11, 287 doi:10.1140/epja/s10050-020-00291-w
 - [13] S. M. M. Nejad and A. Armat, "Heavy quark perturbative QCD fragmentation functions in the presence of hadron mass," *Eur. Phys. J. Plus* **128** (2013) 121. doi:10.1140/epjp/i2013-13121-2
 - [14] M. Salajegheh, S. M. Moosavi Nejad, H. Khanpour, B. A. Kniehl and M. Soleymaninia, " B -hadron fragmentation functions at next-to-next-to-leading order from a global analysis of e^+e^- annihilation data," *Phys. Rev. D* **99** (2019) no.11, 114001 doi:10.1103/PhysRevD.99.114001 [arXiv:1904.08718 [hep-ph]].
 - [15] S. M. Moosavi Nejad and D. Mahdi, "Gluon fragmentation into S-wave heavy quarkonium," [arXiv:1401.5223 [hep-ph]].
 - [16] S. M. Moosavi Nejad, "The effect of meson wave function on heavy-quark fragmentation function," *Eur. Phys. J. A* **52** (2016) no.5, 127 doi:10.1140/epja/i2016-16127-9
 - [17] M. Salajegheh, S. M. Moosavi Nejad, M. Soleymaninia, H. Khanpour and S. Atashbar Tehrani, "NNLO charmed-meson fragmentation functions and their uncertainties in the presence of meson mass corrections," *Eur. Phys. J. C* **79** (2019) no.12, 999 doi:10.1140/epjc/s10052-019-7521-x [arXiv:1904.09832 [hep-ph]].
 - [18] K. Cheung, "A Note on charmed and bottomed pentaquark production by fragmentation," *Phys. Lett. B* **595** (2004), 283-287 doi:10.1016/j.physletb.2004.06.072 [arXiv:hep-ph/0405281 [hep-ph]].
 - [19] V. N. Gribov and L. N. Lipatov, "Deep inelastic e p scattering in perturbation theory," *Sov. J. Nucl. Phys.* **15** (1972) 438

- [Yad. Fiz. **15** (1972) 781].
- [20] M. Delpasand, S. M. Moosavi Nejad and M. Soleymaninia, “ Λ_c^+ fragmentation functions from pQCD approach and the Suzuki model,” Phys. Rev. D **101** (2020) no.11, 114022 doi:10.1103/PhysRevD.101.114022 [arXiv:2006.07602 [hep-ph]].
- [21] M. Soleymaninia *et al.* [SKM18], “QCD analysis of D^* -meson fragmentation functions and their uncertainties at NNLO,” PoS **Confinement2018** (2019), 108 doi:10.22323/1.336.0108
- [22] M. Soleymaninia, H. Khanpour and S. M. Moosavi Nejad, “First determination of D^{*+} -meson fragmentation functions and their uncertainties at next-to-next-to-leading order,” Phys. Rev. D **97** (2018) no.7, 074014 doi:10.1103/PhysRevD.97.074014 [arXiv:1711.11344 [hep-ph]].
- [23] R. Baier and K. Fey, “FINITE CORRECTIONS TO QUARK FRAGMENTATION FUNCTIONS IN PERTURBATIVE QCD,” Z. Phys. C **2** (1979), 339-349 doi:10.1007/BF01545897
- [24] T. Kneesch, B. A. Kniehl, G. Kramer and I. Schienbein, “Charmed-meson fragmentation functions with finite-mass corrections,” Nucl. Phys. B **799** (2008), 34-59 doi:10.1016/j.nuclphysb.2008.02.015 [arXiv:0712.0481 [hep-ph]].
- [25] J. D. Bjorken, “Properties of Hadron Distributions in Reactions Containing Very Heavy Quarks,” Phys. Rev. D **17** (1978), 171-173 doi:10.1103/PhysRevD.17.171
- [26] M. Suzuki, “Fragmentation of Hadrons from Heavy Quark Partons,” Phys. Lett. B **71** (1977), 139-141 doi:10.1016/0370-2693(77)90761-4
- [27] F. Amiri and C. R. Ji, “Perturbative Quantum Chromodynamic Prediction for the Heavy Quark Fragmentation Function,” Phys. Lett. B **195** (1987), 593-598 doi:10.1016/0370-2693(87)91579-6
- [28] M. Suzuki, “Spin Property of Heavy Hadron in Heavy Quark Fragmentation: A Simple Model,” Phys. Rev. D **33** (1986) 676. doi:10.1103/PhysRevD.33.676
- [29] S. J. Brodsky and C. R. Ji, “Exclusive Production of Higher Generation Hadrons and Form-factor Zeros in Quantum Chromodynamics,” Phys. Rev. Lett. **55** (1985) 2257.
- [30] F. Amiri, B. C. Harms and C. -R. Ji, “Exclusive Heavy Meson Production In Z^0 Decay,” Phys. Rev. D **32** (1985) 2982.
- [31] G. P. Lepage and S. J. Brodsky, “Exclusive Processes in Perturbative Quantum Chromodynamics,” Phys. Rev. D **22** (1980) 2157.
- [32] M. A. Gomshi Nobary, “Heavy quark fragmentation functions,” J. Phys. G **20** (1994) 65.
- [33] S. M. Moosavi Nejad, “The impact of Fermi motion on the heavy quarkonia fragmentation using the light-cone wave function,” Eur. Phys. J. Plus **133** (2018) no.1, 25 doi:10.1140/epjp/i2018-11862-x
- [34] P. A. Zyla *et al.* [Particle Data Group], “Review of Particle Physics,” Prog. Theor. Exp. Phys **2020** (2020) 083C01 and 2021 update.
- [35] G. 't Hooft, “Dimensional regularization and the renormalization group,” Nucl. Phys. B **61** (1973), 455-468 doi:10.1016/0550-3213(73)90376-3
- [36] G. 't Hooft and M. J. G. Veltman, “Regularization and Renormalization of Gauge Fields,” Nucl. Phys. B **44** (1972), 189-213 doi:10.1016/0550-3213(72)90279-9
- [37] K. Melnikov and T. v. Ritbergen, “The Three loop relation between the $\overline{\text{MS}}$ -bar and the pole quark masses,” Phys. Lett. B **482** (2000), 99-108 doi:10.1016/S0370-2693(00)00507-4 [arXiv:hep-ph/9912391 [hep-ph]].