D4-branes wrapped on a spindle

Federico Faedo^{a,b} and Dario Martelli^{a,b,c}

^aDipartimento di Matematica "Giuseppe Peano", Università di Torino, Via Carlo Alberto 10, 10123 Torino, Italy

> ^bINFN, Sezione di Torino & ^cArnold–Regge Center, Via Pietro Giuria 1, 10125 Torino, Italy

Abstract

We construct supersymmetric $\operatorname{AdS}_4 \times \Sigma$ solutions of D = 6 gauged supergravity, where Σ is a two-dimensional orbifold known as a spindle. These uplift to solutions of massive type IIA supergravity using a general prescription, that we describe. We argue that these solutions correspond to the near-horizon limit of a system of N_f D8-branes, together with N D4-branes wrapped on a spindle, embedded as a holomorphic curve inside a Calabi-Yau three-fold. The dual field theories are $d = 3, \mathcal{N} = 2$ SCFTs that arise from a twisted compactification of the d = 5, $\mathcal{N} = 1 USp(2N)$ gauge theory. We show that the holographic free energy associated to these solutions is reproduced by extremizing an off-shell free energy, that we conjecture to arise in the large N limit of the localized partition function of the d = 5theories on $S^3 \times \Sigma$. We formulate a universal proposal for a class of off-shell free energies, whose extremization reproduces all previous results for branes wrapped on spindles, as well as on genus g Riemann surfaces $\Sigma_{\rm g}$. We further illustrate this proposal discussing D4-branes wrapped on $\mathbb{Z} \times \Sigma_{g}$, for which we present a supersymmetric $AdS_2 \times \mathbb{Z} \times \Sigma_g$ solution of D = 6 gauged supergravity along with the associated entropy function.

Contents

1	Introduction	1
2	Uplift of $D = 6$ solutions to massive type IIA2.1 The $D = 6$ gauged supergravity2.2 Improved uplift to massive type IIA2.3 The AdS ₆ solution and its uplift2.4 The AdS ₄ × Σ_g solutions and their uplift	5 6 8 10
3	The AdS ₄ × Σ solutions3.1Local form of the solutions3.2Global analysis I: metric and magnetic fluxes3.3Solution of the regularity conditions3.4Global analysis II: gauge fields and Killing spinors3.5Uplift to massive type IIA and holographic free energy	12 12 14 16 18 21
4	Field theory4.1 $d = 5$ SCFTs dual to the AdS ₆ solution4.2 $d = 3$ SCFTs dual to the AdS ₄ × Σ_g solutions4.3 $d = 3$ SCFTs dual to the AdS ₄ × Σ solutions	 23 23 24 26
5	Gravitational blocks for branes on spindles 5.1 M2-branes	28 31 33 34 34
6	D4-branes wrapped on $\mathbb{Z} \times \Sigma_g$ 6.1 $AdS_2 \times \mathbb{Z} \times \Sigma_g$ solutions6.2 Entropy function	36 37 40
7	Discussion	41
A	More details on the $AdS_4 \times \Sigma_g$ solutions A.1 Relation with the Lagrangian of [1]	44 44 44
в	OSp(2 4) superalgebra from bilinears	46

1 Introduction

A plethora of examples of AdS/CFT dualities have been constructed following the idea of [3] of wrapping branes on supersymmetric cycles. On the field theory side, these constructions realise supersymmetric lower-dimensional theories as "twisted" compactifications of the theories living on the branes. Here the twisting refers to the coupling of the field theory to a background R-symmetry gauge field that gets identified with a

connection on the tangent bundle of the manifolds on which the theory is compactified, so that supersymmetry can be preserved simply taking constant spinors. This is referred to as a topological twist. On the gravity side, one generically expects to find supersymmetric solutions incorporating the backreaction of the large number of branes wrapped, and when the dual theory is a SCFT, the solutions will comprise an AdS factor. Focussing on compactifications on two-dimensional manifolds, these constructions have been realised for M2, D3, D4 and M5-branes wrapping constant curvature Riemann surfaces, which include the round two-sphere as the genus g = 0 case. The references presenting these solutions, along with a discussion of the field theory duals, are summarised in the first row of Table 1.

These solutions usually have been constructed in some $U(1)^{\mathfrak{d}}$ gauged supergravity in D = p + 2 dimensions, where p is the world-volume dimension of the brane, and then lifted to D = 10 or D = 11 supergravities, which is a necessary step in order to compare gravity computations with calculations performed in the dual field theory. An exception to this is the solution corresponding to D4-branes wrapped on $\Sigma_{\rm g}$, that was obtained directly in massive type IIA supergravity [2]. Below we will show that, in fact, that solution can also be obtained in a D = 6, $U(1)^2$ gauged supergravity and uplifted to massive type IIA, provided we take due care of the flux quantization conditions. The case of M2-branes, corresponding to supersymmetric $AdS_2 \times \Sigma_g$ solutions, is particularly interesting, because on general grounds it corresponds to the near-horizon limit of BPS black holes in AdS₄. In this case, one can also add rotation to the AdS₂ \times S² solutions. The black holes are interpreted as "flows" across dimensions, with the AdS_4 conformal boundary representing the parent three-dimensional SCFT in the UV and the AdS₂ nearhorizon region corresponding to the one-dimensional IR theory. Such flows have also been constructed for higher-dimensional AdS solutions, although usually they are known only numerically.

	M2	D3	D4	M5
$\Sigma_{\rm g}$	[4]	[5]	[2]	[6]
\mathbb{Z}	[7]	[8]	here	[9]

Table 1: In the first row, the references discussing supersymmetric $AdS \times \Sigma_g$ solutions for different branes, where Σ_g is a Riemann surface of genus g, equipped with a constant curvature metric. In the second row, the references discussing the "simplest" supersymmetric $AdS \times \Sigma$ solutions, where Σ is the spindle.

The solution presented in [8] opened up a new, unexpected, direction of exploration in the landscape of AdS/CFT constructions. This comprises a supersymmetric $\operatorname{AdS}_3 \times \Sigma$ background of minimal D = 5 gauged supergravity, where $\Sigma = \mathbb{WCP}_{[n_-,n_+]}^1$ is a weighted projective space, also known as a spindle. This uplifts to an $\operatorname{AdS}_3 \times M_7$ solution of type IIB supergravity and it has been argued to be dual to a class of d = 4, $\mathcal{N} = 1$ SCFTs compactified on the spindle with a novel type of twist, different from the topological twist, that was later dubbed "anti-twist". A similar construction, for $\operatorname{AdS}_2 \times \Sigma$ solutions of minimal D = 4 gauged supergravity, was presented in [7]. These have been later extended to spindle solutions of STU gauged supergravities in D = 5 [10, 11] and D = 4 [12, 13], respectively. A supersymmetric $\operatorname{AdS}_5 \times \Sigma$ solution corresponding to M5-branes wrapped on the spindle was constructed in [9] and, differently from the previous constructions, it realises supersymmetry by means of a "topologically topological twist". Namely, the background R-symmetry gauge field is identified with a connection on the tangent bundle of the spindle, as for the topological twist, but the corresponding local curvatures are not equal. It turns out that these local solutions comprising spindles contain, as interesting degenerate limits, solutions corresponding to branes wrapped on disks or Riemann surfaces with non-constant curvature [13–18].

In this paper we will construct an $\operatorname{AdS}_4 \times \Sigma$ solution, corresponding to D4-branes wrapped on the spindle, thus filling the outstanding entry in Table 1. We will show that our construction realises the topologically topological twist, as for the M5-brane solution in [9], with which it shares some similarities. We will first present the solution in a D = 6 gauged supergravity model and then we will discuss how to uplift this to a globally consistent solution in massive type IIA supergravity. We will elucidate the global structure of the Killing spinors, identifying the precise bundles of which they are sections and showing how they differ from the Killing spinors of the previous constructions for M2 [7] and D3-branes [8]. Our solution completes the panorama of the "basic" branes wrapped on spindles.

While for the SCFTs compactified on Riemann surfaces with the standard topological twist various supersymmetric partition functions have been computed and studied in the large N limit, for compactifications on spindles similar results are not yet available. For theories in d = 4 and d = 6 this lack of knowledge can be bypassed employing the recipe of [8] for extracting the trial central charge of the (d - 2)-dimensional theories from the anomaly polynomials of the parent theories. In d = 3 an entropy function was obtained in [19], from the on-shell gravitational action of the suitably regularised black hole solutions, employing the method of [20]. Extremizing this reproduces the entropy associated to the $AdS_2 \times \Sigma$ solution of [7]. An extension of this entropy function was conjectured in [12] and shown to reproduce correctly the entropy of multi-charge spindle solutions. Taking inspiration from that, in this paper we will propose a conjectural offshell free energy, whose extremization will, remarkably, reproduce the gravitational free energy associated to our solutions.

In the last part of the paper we will propose a universal class of off-shell free energies for various branes wrapped on spindles, analogous to the entropy functions, to which these reduce in d = 3. Specifically, we conjecture that for a large class of SCFTs in dimensions d = 3, 4, 5, 6, possessing large N gravity duals, when these are compactified on a spindle Σ , the exact superconformal R-symmetry of the SCFTs in dimension d - 2is determined extremizing the following off-shell free energies

$$F^{\pm}(\varphi_i,\epsilon;\mathfrak{n}_i,n_+,n_-,\sigma) = \frac{1}{\epsilon} \left(\mathcal{F}_d(\varphi_i+\mathfrak{n}_i\epsilon) \pm \mathcal{F}_d(\varphi_i-\mathfrak{n}_i\epsilon) \right),$$
(1.1)

where the variables φ_i, ϵ and the magnetic fluxes \mathbf{n}_i satisfy the constraints

$$\sum_{i=1}^{\mathfrak{d}} \varphi_i - \frac{n_+ - \sigma n_-}{n_+ n_-} \epsilon = 2, \qquad \sum_{i=1}^{\mathfrak{d}} \mathfrak{n}_i = \frac{n_+ + \sigma n_-}{n_+ n_-}.$$
(1.2)

The form (1.1) is suggested by the idea of gluing universal contributions called gravitational blocks, advocated in [21] for the entropy functions of SCFTs compactified on different manifolds. The "building blocks" are the functions \mathcal{F}_d above, which have different interpretations in the different dimensions d, being proportional to either the central charge or the sphere partition function of the SCFTs. They are also related to the prepotentials of the various gauged supergravities in dimension D = d + 1. Their precise form will be given later, see Table 2. The sign $\sigma = \pm 1$ labels the different twists that may occur on spindles. The sign $\sigma = +1$ corresponds to the topologically topological twist, which includes the standard topological twist as a special case, while the sign $\sigma = -1$ corresponds to the anti-twist, realised by M2 [7] and D3-branes [8]. The sign \pm depends on the gluing, in the language of [21], and we shall comment below on its relation to the sign of σ . For example, in d = 3, taking $n_+ = n_- = 1$ in the above formulas leads to the entropy functions for the supersymmetric black holes with $\operatorname{AdS}_2 \times S^2$ near-horizon geometry [21]. In this case, for $\sigma = +1$ we must take F^- and this reduces to the entropy function [22] of the supersymmetric AdS₄ black holes with a topological twist [4]. On the contrary, for $\sigma = -1$ we must take F^+ and this reduces to the entropy function [23] for the supersymmetric rotating Kerr-Newmann AdS₄ black holes [24].

More generally, we will provide evidence that in D = 4, 6 the gluing sign \pm coincides with $-\sigma$, while in D = 5, 7 they appear to be independent. In D = 4, the fact that $-\sigma$ coincides with the sign \pm may be understood as follows. The AdS/CFT correspondence implies that, in the large N limit, the free energies F^{\pm} should be identified with the appropriately regularised gravitational on-shell action of the dual supergravity solutions. In [25] it has been proved, in the context of minimal gauged supergravity, that the onshell action of any (Euclidean) supersymmetric solution takes the form of a sum over contributions from fixed points of the canonical Killing vector field, defined as a bilinear in the Killing spinors of the solution. The relative sign of these contributions is determined by the chirality of the Killing spinors at the fixed points, and in all the known supergravity solutions comprising spindle (including S^2 as a special case) we have that the chiralities at the north and south poles of the spindles are the same for the topologically topological twist and opposite for the anti-twist. A general proof of this fact is given in [26].

Our proposal reproduces all the previously known results for $\operatorname{AdS} \times \mathbb{Z}$ solutions, including the $\operatorname{AdS} \times S^2$ solutions as special cases¹. Moreover, in d = 5, taking $\sigma = +1$, corresponding to the topologically topological twist, we will show that the extremization of the function $F^-(\varphi_i, \epsilon; \mathfrak{n}_i, n_+, n_-, +1)$ in (1.1) precisely reproduces the gravitational S^3 free energy of our solution (see eq. (3.71)). We then conjecture that this should arise in the large N limit of the localized partition function on $S^3 \times \mathbb{Z}$, with the topologically topological twist. To add weight to our proposal, we will also discuss D4-branes wrapped on the four-dimensional orbifold $\mathbb{Z} \times \Sigma_g$. The effective field theory obtained from the twisted compactification of the d = 5 SCFT is expected to be superconformal, at least in some ranges of the magnetic fluxes. We will discuss the corresponding supersymmetric $\operatorname{AdS}_2 \times \mathbb{Z} \times \Sigma_g$ solutions of D = 6 gauged supergravity and show that the entropy function constructed from the "spindly" gravitational blocks correctly reproduces the geometric entropy.

The rest of the paper is organised as follows. In section 2 we discuss the uplift of solutions of a D = 6, $U(1)^2$ gauged supergravity model to massive type IIA. As a warm-up, we illustrate this obtaining the (global) solutions of [27] and [2] from known solutions

¹As we shall discuss, for $\sigma = +1$, by formally replacing $\frac{n_+ - n_-}{n_+ n_-} \mapsto 0$ and $\frac{n_+ + n_-}{n_+ n_-} \mapsto \chi(\Sigma_g)$, our formulas cover also the AdS× Σ_g solutions.

in D = 6. In section 3 we construct new supersymmetric $\operatorname{AdS}_4 \times \Sigma$ solutions and discuss global properties of these both in D = 6 and D = 10. In section 4 we discuss aspects of the field theory duals of these solutions. In particular, we conjecture a field-theoretic large N off-shell free energy and show that extremizing this reproduces the holographic free energy associated to our solutions. In section 5 we discuss how our proposal fits in a general scheme of off-shell free energies for field theories compactified on spindles (as well as on genus g Riemann surfaces), comprising and extending entropy functions and trial central charges, previously discussed in the literature. In section 6 we begin investigating D4-branes wrapped on four-dimensional orbifolds, focussing on a class of supersymmetric $\operatorname{AdS}_2 \times \Sigma \times \Sigma_g$ solutions. We conclude with a discussion in section 7. Appendix A contains technical details useful for comparing known solutions, in different conventions. In appendix B we demonstrate how the Killing spinors of the $\operatorname{AdS}_4 \times \Sigma$ solutions encapsulate the OSp(2|4) superalgebra of the dual SCFTs.

2 Uplift of D = 6 solutions to massive type IIA

In this paper we discuss solutions of a D = 6 gauged supergravity with gauge group $U(1)^2$, comprising two gauge fields A_1, A_2 , a two-form B and two real scalar fields $\vec{\varphi} = (\varphi_1, \varphi_2)$. These can be uplifted locally to solutions of massive type IIA supergravity by means of the consistent truncation formulas presented in [28]. However, we will see that globally the solutions uplifted through this ansatz are incompatible with quantization of the fluxes and need to be supplemented by an additional parameter that arises in D = 10.

2.1 The D = 6 gauged supergravity

The D = 6 supergravity model of interest can also be obtained as a sub-sector of an extension of Romans F(4) gauged supergravity [29], coupled to three vector multiplets [30]. The bosonic part of the action reads²

$$S_{6D} = \frac{1}{16\pi G_{(6)}} \int d^6 x \sqrt{-g} \left(R - V - \frac{1}{2} |d\vec{\varphi}|^2 - \frac{1}{2} \sum_{i=1}^2 X_i^{-2} |F_i|^2 \right),$$
(2.1)

where $F_i = dA_i$ and the scalar fields $\vec{\varphi}$ are parameterised as

$$X_i = e^{-\frac{1}{2}\vec{a}_i \cdot \vec{\varphi}}$$
 with $\vec{a}_1 = (2^{1/2}, 2^{-1/2}), \quad \vec{a}_2 = (-2^{1/2}, 2^{-1/2}).$ (2.2)

The scalar potential is

$$V = m^2 X_0^2 - 4g^2 X_1 X_2 - 4mg X_0 (X_1 + X_2), \qquad (2.3)$$

with g the gauge coupling and m the mass parameter, and where for later convenience we defined $X_0 = (X_1X_2)^{-3/2}$. Here we consistently set B = 0 because in the first part of the paper we will restrict to configurations with $F_1 \wedge F_2 = 0$. We will restore the two-form B in section 6.1. It is worth mentioning that locally the ratio m/g can be set to any non-zero

²Here and in what follows we define, for any *p*-form ω , $|\omega|^2 = \frac{1}{p!} \omega_{\mu_1 \dots \mu_p} \omega^{\mu_1 \dots \mu_p}$.

value rescaling the scalar fields X_i and the field strengths F_i . In particular, m can be absorbed in the coupling constant transforming

$$X_i \mapsto \left(\frac{m}{lg}\right)^{1/4} X_i, \qquad F_i \mapsto \left(\frac{m}{lg}\right)^{1/4} F_i, \qquad (2.4)$$

and defining the new gauge coupling as $\tilde{g} = \left(\frac{mg^3}{l}\right)^{1/4}$, with l a positive constant. The action (2.1) keeps the same form, with the scalar potential becoming

$$V = \tilde{g}^2 \left[l^2 X_0^2 - 4X_1 X_2 - 4l X_0 (X_1 + X_2) \right].$$
(2.5)

However, for the time being we will keep both parameters m and g.

A solution to the equations of motion of the model is supersymmetric if and only if it satisfies also the following set of Killing spinor equations [30]:

$$\mathcal{D}_{\mu}\epsilon^{A} + \frac{1}{8} \left[g(X_{1} + X_{2}) + mX_{0} \right] \Gamma_{\mu}\epsilon^{A} + \frac{i}{32} \left(X_{1}^{-1}F_{1} + X_{2}^{-1}F_{2} \right)_{\nu\lambda} \left(\Gamma_{\mu}^{\ \nu\lambda} - 6\delta_{\mu}^{\nu}\Gamma^{\lambda} \right) (\sigma^{3})^{A}{}_{B}\epsilon^{B} = 0 , \qquad (2.6)$$

$$\frac{1}{4} (X_1^{-1} \partial_\mu X_1 + X_2^{-1} \partial_\mu X_2) \Gamma^\mu \epsilon^A - \frac{1}{8} [g(X_1 + X_2) - 3mX_0] \epsilon^A - \frac{i}{32} (X_1^{-1} F_1 + X_2^{-1} F_2)_{\mu\nu} \Gamma^{\mu\nu} (\sigma^3)^A_{\ B} \epsilon^B = 0, \qquad (2.7)$$

$$\frac{1}{2} (X_1^{-1} \partial_\mu X_1 - X_2^{-1} \partial_\mu X_2) \Gamma^\mu (\sigma^3)_{AB} \epsilon^B - g(X_1 - X_2) (\sigma^3)_{AB} \epsilon^B - \frac{i}{4} (X_1^{-1} F_1 - X_2^{-1} F_2)_{\mu\nu} \Gamma^{\mu\nu} \epsilon_A = 0, \qquad (2.8)$$

where

$$\mathcal{D}_{\mu}\epsilon^{A} \equiv \partial_{\mu}\epsilon^{A} + \frac{1}{4}\omega_{\mu}^{\ ab}\Gamma_{ab}\epsilon^{A} - \frac{\mathrm{i}}{2}g(A_{1} + A_{2})_{\mu}(\sigma^{3})^{A}_{\ B}\epsilon^{B}.$$
(2.9)

These follow from setting to zero the supersymmetry variations of the fermionic fields of the theory with three vector multiplets [30], that do not vanish automatically in the sub-truncation that we are considering. Here $(\sigma^3)^A_B$ is the usual third Pauli matrix and $\{\Gamma_a, \Gamma_b\} = 2\eta_{ab}$. The SU(2) indices A, B are raised and lowered as $\epsilon^A = \varepsilon^{AB}\epsilon_B$ and $\epsilon_A = \epsilon^B \varepsilon_{BA}$, where $\varepsilon_{AB} = -\varepsilon_{BA}$ and its inverse matrix ε^{AB} is defined such that $\varepsilon^{AB}\varepsilon_{AC} = \delta^B_C$. The supersymmetry parameter ϵ^A is an eight-component symplectic-Majorana spinor, hence it satisfies the condition

$$\varepsilon^{AB} \epsilon_B^* = \mathcal{B}_6 \epsilon_A \,, \tag{2.10}$$

where \mathcal{B}_6 is related to the six-dimensional charge conjugation matrix \mathcal{C}_6 by $\mathcal{B}_6 = i \mathcal{C}_6 \Gamma^0$.

2.2 Improved uplift to massive type IIA

Any solution to the equations of motion of this theory can be embedded in massive type IIA supergravity by means of the dimensional reduction ansatz presented in $[28]^3$,

³The original ansatz was written for a four-scalar system, but the model with two scalars can be easily obtained setting to zero two of the four scalars in [28]. The consistency of this sub-truncation was conjectured in [31].

provided the gauge coupling and mass parameter are related as m = 2g/3. The metric in the string frame and the dilaton are given by

$$ds_{\text{s.f.}}^{2} = \mu_{0}^{-1/3} (X_{1}X_{2})^{-1/4} \left\{ \Delta^{1/2} ds_{6}^{2} + g^{-2} \Delta^{-1/2} \left[X_{0}^{-1} d\mu_{0}^{2} + X_{1}^{-1} \left(d\mu_{1}^{2} + \mu_{1}^{2} \sigma_{1}^{2} \right) + X_{2}^{-1} \left(d\mu_{2}^{2} + \mu_{2}^{2} \sigma_{2}^{2} \right) \right] \right\},$$
(2.11)

$$e^{\Phi} = \mu_0^{-5/6} \Delta^{1/4} (X_1 X_2)^{-5/8}, \qquad (2.12)$$

where ds_6^2 is the six-dimensional metric and we defined the one-forms $\sigma_i \equiv d\phi_i - gA_i$. The angular coordinates ϕ_1 , ϕ_2 have canonical 2π periodicities, and the warp factor is

$$\Delta = \sum_{a=0}^{2} X_a \mu_a^2 \,. \tag{2.13}$$

The coordinates μ_a , with a = 0, 1, 2, satisfy the constraint $\sum \mu_a^2 = 1$, which can be solved for example defining

$$\mu_0 = \sin\xi, \qquad \mu_1 = \cos\xi \sin\eta, \qquad \mu_2 = \cos\xi \cos\eta, \qquad (2.14)$$

and taking $\eta \in [0, \pi/2]$, $\xi \in (0, \pi/2]$, where the range of ξ arises from the necessity of having $\mu_0 > 0$. At any point in the six-dimensional space-time, the metric inside the square brackets in (2.11) parameterises a four-dimensional hemisphere, that we will denote by \mathbb{S}^4 . This metric is in general squashed and it reduces to the metric on "half the round four-sphere" when $X_1 = X_2 = 1$. The only non-vanishing fields of the RR sector are the ten-dimensional Romans mass

$$F_{(0)} = \frac{2g}{3} \tag{2.15}$$

and the four-form flux. This is conveniently written in terms of its Hodge dual as

$$\star_{10} F_{(4)} = gU \operatorname{vol}(M_6) - \frac{1}{g^2} \sum_i X_i^{-2} \mu_i(\star_6 F_i) \wedge \mathrm{d}\mu_i \wedge \sigma_i + \frac{1}{g} \sum_a X_a^{-1} \mu_a(\star_6 \mathrm{d}X_a) \wedge \mathrm{d}\mu_a \,, \quad (2.16)$$

where $vol(M_6)$ is the volume form of the six-dimensional space and

$$U = 2\sum_{a=0}^{2} X_{a}^{2} \mu_{a}^{2} - \left[\frac{4}{3}X_{0} + 2(X_{1} + X_{2})\right] \Delta.$$
 (2.17)

The Hodge star operator \star_{10} in (2.16) is computed using the string frame metric (2.11), while \star_6 is defined using the six-dimensional metric ds_6^2 .

Provided the equations of motion of the six-dimensional supergravity hold, the above field configuration solves the equations of motion of massive type IIA supergravity, whose action in the string frame reads⁴

$$S_{\rm mIIA} = \frac{1}{16\pi G_{(10)}} \left\{ \int d^{10}x \sqrt{-g} \left[e^{-2\Phi} \left(R + 4 |d\Phi|^2 - \frac{1}{2} |H_{(3)}|^2 \right) - \frac{1}{2} \left(F_{(0)}^2 + |F_{(2)}|^2 + |F_{(4)}|^2 \right) \right] - \frac{1}{2} \int \left(B_{(2)} \wedge dC_{(3)} \wedge dC_{(3)} + 2F_{(0)} B_{(2)}^3 \wedge dC_{(3)} + 6F_{(0)}^2 B_{(2)}^5 \right) \right\}, \qquad (2.18)$$

⁴Here $B_{(2)}^n$ denotes the wedge product of $B_{(2)}$ with itself n times, divided by n!.

where the field strengths are defined from the NS two-form $B_{(2)}$ and the RR potentials $C_{(1)}$ and $C_{(3)}$ as

$$H_{(3)} = dB_{(2)}, \quad F_{(2)} = dC_{(1)} + F_{(0)}B_{(2)}, \quad F_{(4)} = dC_{(3)} - H_{(3)} \wedge C_{(1)} + \frac{1}{2}F_{(0)}B_{(2)} \wedge B_{(2)}.$$
(2.19)

It will be important to notice that the equations of motion of massive type IIA are invariant if the fields are transformed as

$$d\hat{s}_{s.f.}^{2} = \lambda^{2} ds_{s.f.}^{2}, \qquad e^{\hat{\Phi}} = \lambda^{2} e^{\Phi}, \qquad \hat{B}_{(2)} = \lambda^{2} B_{(2)}, \hat{F}_{(0)} = \lambda^{-3} F_{(0)}, \qquad \hat{C}_{(n-1)} = \lambda^{n-3} C_{(n-1)}, \qquad (2.20)$$

with n = 2, 4, where λ is any strictly positive constant. However, this scaling symmetry holds only at the classical level in supergravity and it is broken upon imposing the Dirac quantization conditions on the fluxes. As we will discuss momentarily, this additional parameter will be crucial for ensuring that six-dimensional solutions yield globally regular solutions in D = 10, in particular that the fluxes are correctly quantized.

Notice that the reduction ansatz of [28] applies only after setting m = 2g/3 in the six-dimensional theory and it implies that the Romans mass of the ten-dimensional theory is fixed in terms of the gauge coupling constant g as in (2.15). It is natural to suspect that there may exist a more general truncation ansatz that relates the six-dimensional mass parameter m to the ten-dimensional parameter λ , so that the Romans mass $F_{(0)}$ is an independent parameter. This would be a mechanism analogous to the ten-dimensional origin of dyonic four-dimensional supergravity discussed in [32]. We leave this interesting question for the future and proceed to discuss different globally regular solutions of massive type IIA originating in D = 6.

In summary, our strategy will be as follows. We construct the solutions in D = 6 and after setting m = 2g/3 we uplift these to local solutions in D = 10, using the formulas in [28]. Then we introduce the parameter λ and proceed to quantize the fluxes, finding globally consistent solutions of massive type IIA supergravity.

2.3 The AdS_6 solution and its uplift

The equations of motion following from the action (2.1) admit the well-known supersymmetric vacuum with constant scalars $X_1 = X_2 = \left(\frac{3m}{2g}\right)^{1/4}$, vanishing gauge fields $A_1 = A_2 = 0$ and metric

$$ds_6^2 = \frac{9}{2 \left(6mg^3\right)^{1/2}} ds_{AdS_6}^2, \qquad (2.21)$$

where $ds_{AdS_6}^2$ is the metric on AdS₆ with unit radius. We now set m = 2g/3 and uplift this solution to massive type IIA using the formulas in [28]. After introducing the parameter λ using the local scaling symmetry (2.20), we obtain

$$ds_{s.f.}^{2} = \lambda^{2} (\sin \xi)^{-1/3} L_{AdS_{6}}^{2} \left[ds_{AdS_{6}}^{2} + \frac{4}{9} (d\xi^{2} + \cos^{2}\xi \, ds_{S^{3}}^{2}) \right],$$

$$e^{\Phi} = \lambda^{2} (\sin \xi)^{-5/6},$$

$$F_{(0)} = \frac{1}{\lambda^{3} L_{AdS_{6}}}, \qquad F_{(4)} = -\lambda \frac{10 \cos^{3}\xi \sin^{1/3}\xi}{3g^{3}} \, d\xi \wedge \operatorname{vol}(S^{3}),$$

(2.22)

where $ds_{S^3}^2$ denotes the metric on a unit radius round three-sphere

$$ds_{S^3}^2 = d\eta^2 + \sin^2\eta \, d\phi_1^2 + \cos^2\eta \, d\phi_2^2 \,, \qquad (2.23)$$

and $vol(S^3)$ its associated volume form. The quantization conditions of the (non-zero) fluxes in massive type IIA read

$$(2\pi\ell_s)F_{(0)} = n_0 \in \mathbb{Z}$$
 and $\frac{1}{(2\pi\ell_s)^3} \int_{\mathbb{S}^4} F_{(4)} = N \in \mathbb{Z}$, (2.24)

where ℓ_s is the string length. In the solution (2.22) these imply that⁵

$$g^{8} = \frac{1}{(2\pi\ell_{s})^{8}} \frac{18\pi^{6}}{N^{3}n_{0}}, \qquad \lambda^{8} = \frac{8\pi^{2}}{9Nn_{0}^{3}}.$$
(2.25)

It is clear from the second equation that setting $\lambda = 1$ leads to an inconsistent relation between the integers N and n_0 . This problem arises because without introducing λ there is only one free dimensionless parameter $(g\ell_s)$ and two conditions to impose. Thus the scaling symmetry (2.20) plays a crucial role in making the uplifted solution globally consistent. After imposing (2.25) the uplifted solution (2.22) can be matched with the solution of [27] identifying

$$L_{\text{AdS}_6}^2 = \frac{9}{4} \ell_s^2 \left(\frac{3(8-N_f)}{4\pi C}\right)^{1/2} Q_4^{3/4}, \qquad \lambda^2 = \left(\frac{3(8-N_f)}{4\pi}\right)^{-5/6} C^{1/6} Q_4^{-1/4}, \qquad (2.26)$$

with $n_0 = 8 - N_f$, where N_f is the number of D8-branes, and Q_4 is related to the number of D4-branes N by

$$N = \frac{3Q_4}{8\pi} \left(\frac{3(8-N_f)}{4\pi C^2}\right)^{1/3}.$$
 (2.27)

Note that the constant C is a trivial parameter which can be set to any non-zero value redefining $Q_4 \mapsto C^{2/3}Q_4$. Although both the six-dimensional vacuum and the tendimensional solution of [27] are well-known, clarifying their relationship will allow us to discuss the six-dimensional origin of more interesting solutions in the following.

As discussed in [33], the effective six-dimensional Newton constant that should be proportional to the large N limit of the S^5 free energy of the dual field theory is divergent, due to the singularity of the ten-dimensional solution on the boundary of the hemisphere \mathbb{S}^4 , *i.e.* for $\xi \to 0$. This problem was circumvented in [33] by calculating the holographic entanglement entropy across a three-sphere and then extracting from this the free energy F_{S^5} . We refer the reader to [33] for the details and here, for completeness, we only quote the result

$$F_{S^5} = -\frac{3^5 \pi^6 \lambda^4}{5(2\pi \ell_s g)^8} = -\frac{9\sqrt{2\pi}}{5} \frac{N^{5/2}}{\sqrt{8 - N_f}}.$$
(2.28)

⁵For consistency, we need to identify the flux of $F_{(4)}$ with -N, where N is the number of D4-branes.

2.4 The $AdS_4 \times \Sigma_g$ solutions and their uplift

In this section we will discuss a class of supersymmetric solutions of the D = 6, $U(1)^2$ supergravity comprising an AdS₄ factor, that were constructed in [1] (see also [31]). We will use these to illustrate our procedure for uplifting to solutions of massive type IIA supergravity, showing that the uplifted solutions coincide with the solutions constructed in [2], by directly solving the supersymmetry conditions of [34] in ten dimensions. The solutions have the form of a product $AdS_4 \times \Sigma_g$, where Σ_g is a genus g Riemann surface equipped with a constant curvature metric. As usual, we can distinguish three cases for the curvature $\kappa = \pm 1, 0$. When $\kappa = +1$ we have the round two-sphere with g = 0, while for $\kappa = -1$ locally we have the metric on the two-dimensional hyperbolic space H^2 , which can be quotiented to obtain a constant curvature Riemann surface with genus g > 1. In order to encompass both non-zero curvature cases⁶ we denote by $ds_{\Sigma_g}^2$ the metric on Σ_g and define the one-form ω_g such that $d\omega_g = vol(\Sigma_g)$. Explicitly, we can take

$$ds_{\Sigma_{g}}^{2} = d\theta^{2} + \sin^{2}\theta \, d\phi^{2} , \qquad \omega_{g} = -\cos\theta \, d\phi \qquad (\kappa = +1) , ds_{\Sigma_{g}}^{2} = d\theta^{2} + \sinh^{2}\theta \, d\phi^{2} , \qquad \omega_{g} = \cosh\theta \, d\phi \qquad (\kappa = -1) .$$
(2.29)

In our conventions, the metric, scalars and gauge potentials take the form

$$ds_{6}^{2} = L_{AdS_{4}}^{2} ds_{AdS_{4}}^{2} + e^{2G} ds_{\Sigma_{g}}^{2},$$

$$X_{1} = k_{8}^{1/8} k_{2}^{1/2}, \qquad X_{2} = k_{8}^{1/8} k_{2}^{-1/2},$$

$$A_{1} = \frac{p_{1}}{2g} \omega_{g}, \qquad A_{2} = \frac{p_{2}}{2g} \omega_{g},$$

(2.30)

where

$$L_{\text{AdS}_4}^2 = \frac{k_8^{3/4}}{m^2}, \qquad e^{2G} = \left[\frac{p_2^2 k_2^{3/2} - p_1^2 k_2^{-1/2}}{16mg^3(k_2 - 1)}\right]^{1/2}, \qquad (2.31)$$
$$k_8 = \frac{4m^2 k_2}{g^2(1 + k_2)^2}, \qquad k_2 = -\frac{3(p_1 - p_2) + \sqrt{9p_1^2 - 14p_1p_2 + 9p_2^2}}{2p_2},$$

with p_1 , p_2 two constant parameters obeying the supersymmetry constraint⁷

$$p_1 + p_2 = 2\kappa \,. \tag{2.32}$$

The fluxes of the gauge fields through $\Sigma_{\rm g}$ are given by⁸

$$P_i = \frac{g}{2\pi} \int_{\Sigma_g} F_i = p_i \kappa (1 - g) \in \mathbb{Z}, \qquad (2.33)$$

where the quantization condition above arises from the requirement that gA_i be welldefined connection one-forms on U(1) bundles over Σ_g . The six-dimensional solution is therefore specified by the genus g and one integer, say $p_1(1-g)$.

⁶The case $\kappa = 0$ is straightforward to include, but we will not discuss this further.

⁷A sign ambiguity in the above formulas [1] has been fixed by noticing that the solution corresponding to the opposite sign is obtained by changing both signs of p_1 and p_2 .

⁸Recall that, for $\kappa \neq 0$, the integrated volume of the Riemann surface Σ_g is $Vol(\Sigma_g) = 4\pi\kappa(1-g)$.

We now set m = 2g/3 and, after uplifting to massive type IIA using the formulas in [28], we introduce the parameter λ . The ten-dimensional metric and dilaton are

$$ds_{s.f.}^{2} = \lambda^{2} k_{8}^{3/4} g^{-2} \mu_{0}^{-1/3} \left\{ \tilde{\Delta}^{1/2} \left[\frac{9}{4} ds_{AdS_{4}}^{2} + e^{2G} k_{8}^{-3/4} g^{2} ds_{\Sigma_{g}}^{2} \right] + \tilde{\Delta}^{-1/2} k_{8}^{-1} \left[k_{8}^{1/2} d\mu_{0}^{2} + k_{2}^{-1/2} (d\mu_{1}^{2} + \mu_{1}^{2}\sigma_{1}^{2}) + k_{2}^{1/2} (d\mu_{2}^{2} + \mu_{2}^{2}\sigma_{2}^{2}) \right] \right\}.$$
(2.34)

$$e^{\Phi} = \lambda^2 \mu_0^{-5/6} \tilde{\Delta}^{1/4} k_8^{-1/8} , \qquad (2.35)$$

while the Romans mass and four-form flux read

$$F_{(0)} = \frac{2g}{3\lambda^3},$$
(2.36)

$$F_{(4)} = \frac{\lambda \mu_0^{1/3}}{g^3 k_8^{1/2} \tilde{\Delta}} \left\{ \frac{\tilde{U}}{\tilde{\Delta}} \frac{\mu_1 \mu_2}{\mu_0} \, \mathrm{d}\mu_1 \wedge \mathrm{d}\mu_2 \wedge \sigma_1 \wedge \sigma_2 \right.$$
(2.37)

$$-g\left[F_{1}\wedge \mathrm{d}\phi_{2}\wedge\left(\mu_{0}\mu_{2}\mathrm{d}\mu_{2}-\mu_{2}^{2}\frac{k_{8}^{1/2}}{k_{2}^{1/2}}\mathrm{d}\mu_{0}\right)+F_{2}\wedge \mathrm{d}\phi_{1}\wedge\left(\mu_{0}\mu_{1}\mathrm{d}\mu_{1}-\mu_{1}^{2}k_{8}^{1/2}k_{2}^{1/2}\mathrm{d}\mu_{0}\right)\right]\right\}.$$

We have $\sigma_i = \mathrm{d}\phi_i - \frac{p_i}{2}\omega_{\mathrm{g}}$ and we defined

$$\tilde{\Delta} = k_8^{-1/2} \mu_0^2 + k_2^{1/2} \mu_1^2 + k_2^{-1/2} \mu_2^2,$$

$$\tilde{U} = -\frac{2}{3} \left[k_8^{-1/2} \mu_0^2 + 3k_8^{1/2} \left(1 - \mu_0^2 \right) + 2\tilde{\Delta} \right].$$
(2.38)

We have checked that the above configuration satisfies the ten-dimensional equations of motion and that $dF_{(4)} = 0$. Imposing flux quantization (2.24) we obtain the relations (2.25), exactly as for the vacuum solution, showing that $\lambda = 1$ would be again inconsistent. We have therefore obtained a globally regular (modulo the ever-present singularity at $\xi = 0$) ten-dimensional solution with an AdS₄ factor, parameterised by the genus g and the integers $n_0, N, p_1(1 - g)$. This is precisely the solution presented in [2], as we show in detail in appendix A.2.

Writing the ten-dimensional metric (2.34) in the form

$$ds_{s.f.}^2 = e^{2A} \left(ds_{AdS_4}^2 + ds_{M_6}^2 \right), \qquad (2.39)$$

we have that $ds_{M_6}^2$ is the metric on the internal space M_6 , that is the total space of an \mathbb{S}^4 bundle over Σ_g , namely

$$\mathbb{S}^4 \hookrightarrow M_6 \to \Sigma_{\rm g} \,. \tag{2.40}$$

Recall that \mathbb{S}^4 is a hemisphere of the four-sphere due to the fact that the warp factor is singular at $\mu_0 \to 0$, corresponding to the location of the O8-plane [27]. However, we can also think of the internal geometry as an S^4 bundle over Σ_g , before the inclusion of the O8-plane. Either way, there is a $U(1) \times U(1) \subset SU(2)_R \times SU(2)_F$ symmetry acting on \mathbb{S}^4 and the gauge fields gA_i are connections on the associated circle bundles, twisting these over Σ_g , with Chern numbers P_i . The solution can be interpreted as follows [2]: one starts with a geometry of the type $\mathbb{R}^{1,2} \times \mathbb{R} \times Y_6$, with Y_6 a local Calabi-Yau three-fold of the form

$$\mathcal{O}(-p_1) \oplus \mathcal{O}(-p_2) \hookrightarrow Y_6 \to \Sigma_g,$$
 (2.41)

where $-gA_i$ are Hermitian connections on the line bundles $\mathcal{O}(-p_i)$. Then (2.32) implies that the total space of Y_6 has vanishing first Chern class and hence is a Calabi-Yau threefold. At the origin of \mathbb{R} there are an O8-plane and $N_f = 8 - n_0$ coincident D8-branes, and in addition one wraps N D4-branes over the zero section Σ_g of Y_6 .

At low energies the effective theory on the D4/D8-system will be a d = 3, $\mathcal{N} = 2$ field theory, obtained from the compactification on $\Sigma_{\rm g}$ of the d = 5, $\mathcal{N} = 1$ SCFT with gauge group USp(2N) [27], with the standard topological twist. The supergravity solution above strongly suggests that in the large N limit this is a SCFT and its S^3 free energy can be computed holographically, from the full ten-dimensional solution. Specifically, this can be obtained using the formula presented in [32] adapted to the string frame, and it reads [2]

$$F_{S^3 \times \Sigma_{\rm g}} = \frac{16\pi^3}{(2\pi\ell_s)^8} \int e^{8A-2\Phi} \operatorname{vol}(M_6) = \frac{16\pi\kappa(1-{\rm g})N^{5/2} \left({\rm z}^2-\kappa^2\right)^{3/2} \left(\sqrt{\kappa^2+8{\rm z}^2}-\kappa\right)}{5\sqrt{8-N_f} \left(\kappa\sqrt{\kappa^2+8{\rm z}^2}-\kappa^2+4{\rm z}^2\right)^{3/2}},$$
(2.42)

where the parameter z is related to our parameters as $p_1 = \kappa + z$, $p_2 = \kappa - z$. This expression has been reproduced exactly by a direct field theory computation, using the large N expansion of the localized partition function on $S^3 \times \Sigma_g$ [35, 36].

3 The $AdS_4 \times \mathbb{Z}$ solutions

3.1 Local form of the solutions

Our starting point are the following local solutions to the equations of the D = 6 gauged supergravity model of section 2.1

$$ds^{2} = (y^{2}h_{1}h_{2})^{1/4} \left(ds^{2}_{AdS_{4}} + \frac{y^{2}}{F} dy^{2} + \frac{F}{h_{1}h_{2}} dz^{2} \right) ,$$

$$A_{i} = \left(\alpha_{i} - \frac{y^{3}}{h_{i}} \right) dz , \qquad X_{i} = \left(y^{2}h_{1}h_{2} \right)^{3/8}h_{i}^{-1} , \qquad (3.1)$$

$$F(y) = m^{2}h_{1}h_{2} - y^{4} , \qquad h_{i}(y) = \frac{2g}{3m}y^{3} + q_{i} ,$$

where $ds^2_{AdS_4}$ denotes the unit radius metric on AdS₄ and q_1 , q_2 are two real parameters. The real constants α_i are pure gauge and we have included them as they will play a crucial role for understanding the global properties of the solution. These backgrounds can be obtained by doing an analytic continuation [37] of a class of six-dimensional BPS black holes [38]. A curvature singularity lies at y = 0, hence without loss of generality, in the subsequent analysis we will make sure that the globally regular solutions will be restricted to y > 0.

Before turning attention to the global structure of the solutions, we will demonstrate that they are supersymmetric by constructing the local form of the Killing spinors solving the equations (2.6) - (2.8). We employ the following orthonormal frame

$$e^{\hat{a}} = y^{1/4} (h_1 h_2)^{1/8} \hat{e}^{\hat{a}}, \qquad e^4 = \frac{y^{5/4} (h_1 h_2)^{1/8}}{F^{1/2}} \,\mathrm{d}y, \qquad e^5 = \frac{y^{1/4} F^{1/2}}{(h_1 h_2)^{3/8}} \,\mathrm{d}z, \qquad (3.2)$$

where $\hat{e}^{\hat{a}}$, $\hat{a} = 0, ..., 3$, is the vierbein on AdS₄, whose coordinates are denoted as $x^{\hat{\mu}}$. Equation (2.6) then splits in the following equations

$$\left(\partial_{\hat{\mu}} + \frac{1}{4}\omega_{\hat{\mu}}^{\ \hat{a}\hat{b}}\Gamma_{\hat{a}\hat{b}}\right)\epsilon^{A} - \frac{i}{2}\,\hat{e}_{\hat{\mu}}^{\hat{a}}\,\Gamma_{\hat{a}}^{\ 45}(\sigma^{3})^{A}_{\ B}\epsilon^{B} = 0\,,$$

$$\partial_{y}\epsilon^{A} - \frac{1}{16y}\left[(2+y\tilde{h}')\epsilon^{A} - i\frac{4y^{2}}{F^{1/2}}(4-y\tilde{h}')\Gamma^{5}(\sigma^{3})^{A}_{\ B}\epsilon^{B}\right] = 0\,,$$

$$\partial_{z}\epsilon^{A} - i\frac{g}{2}\left(\alpha_{1} + \alpha_{2} - \frac{2m}{g}\right)(\sigma^{3})^{A}_{\ B}\epsilon^{B} = 0\,,$$
(3.3)

where we defined $\tilde{h} \equiv \log(h_1h_2)$. Equations (2.7) and (2.8) yield the same constraint

$$F^{1/2}\Gamma^{4}\epsilon^{A} + m(h_{1}h_{2})^{1/2}\epsilon^{A} + i y^{2}\Gamma^{45}(\sigma^{3})^{A}{}_{B}\epsilon^{B} = 0.$$
(3.4)

From the third equation in (3.3) we see immediately that setting

$$\alpha_1 + \alpha_2 = \frac{2m}{g} \tag{3.5}$$

leads to Killing spinors independent of z and we will adopt this choice in the reminder of this section. We consider the specific decomposition of the gamma matrices

$$\Gamma^{\hat{a}} = \gamma^{\hat{a}} \otimes I_2 , \qquad \Gamma^{\hat{i}+3} = \gamma_5 \otimes \rho^{\hat{i}} , \qquad (3.6)$$

where $\gamma^{\hat{a}}$ are the (Lorentzian) gamma matrices in D = 4, $\gamma_5 = i \gamma^0 \gamma^1 \gamma^2 \gamma^3$ is the related chiral matrix and $\rho^{\hat{i}}$, $\hat{i} = 1, 2$, are the (Euclidean) gamma matrices in D = 2. For $\rho^{\hat{i}}$ we choose the following representation

$$\rho^{\hat{i}} = \sigma^{\hat{i}}, \qquad \rho_* = -i\,\rho^1\rho^2 = \sigma^3,$$
(3.7)

and we take $\mathcal{B}_2 = -\sigma^2$ so that $\mathcal{B}_6 = \mathcal{B}_4 \otimes \mathcal{B}_2$. The ansatz for the symplectic-Majorana Killing spinors ϵ^A is

$$\epsilon^A = \vartheta_+ \otimes \eta^A_+ + \vartheta_- \otimes \eta^A_-, \qquad (3.8)$$

where $\vartheta = \vartheta(x^{\hat{\mu}})$ is a Majorana Killing spinor on AdS_4 and ϑ_{\pm} are its chiral components, *i.e.* $\gamma_5 \vartheta_{\pm} = \pm \vartheta_{\pm}$. Thus we have $\vartheta_{\pm}^* = \mathcal{B}_4 \vartheta_{\mp}$ and $\hat{\nabla}_{\hat{\mu}} \vartheta_{\pm} = \frac{1}{2} \gamma_{\hat{\mu}} \vartheta_{\mp}$. The spinors $\eta_{\pm}^A = \eta_{\pm}^A(y)$ are two-component Dirac spinors defined on the spindle⁹.

Adopting the decomposition (3.6), the symplectic-Majorana condition (2.10) and the Killing spinor equations (3.3), (3.4) can be solved to give

$$\eta_{+}^{1} = \xi y^{1/8} (h_{1}h_{2})^{-3/16} \begin{pmatrix} f_{1}^{1/2} \\ -f_{2}^{1/2} \end{pmatrix}, \qquad \eta_{-}^{1} = -\xi y^{1/8} (h_{1}h_{2})^{-3/16} \begin{pmatrix} f_{1}^{1/2} \\ f_{2}^{1/2} \end{pmatrix}, \qquad (3.9)$$
$$\eta_{+}^{2} = i \xi^{*} y^{1/8} (h_{1}h_{2})^{-3/16} \begin{pmatrix} f_{2}^{1/2} \\ -f_{1}^{1/2} \end{pmatrix}, \qquad \eta_{-}^{2} = i \xi^{*} y^{1/8} (h_{1}h_{2})^{-3/16} \begin{pmatrix} f_{2}^{1/2} \\ f_{1}^{1/2} \end{pmatrix},$$

where ξ is a complex constant and we have defined

$$f_1(y) \equiv m(h_1h_2)^{1/2} + y^2, \qquad f_2(y) \equiv m(h_1h_2)^{1/2} - y^2,$$
 (3.10)

⁹Notice that η_{\pm}^{A} are *not* chiral spinors, despite the notation suggests otherwise. The index A = 1, 2 is an internal index and we will see below that these four Dirac spinors are actually not all independent.

which satisfy $F(y) = f_1(y)f_2(y)$. Notice that the four two-dimensional spinors above can be expressed in terms of just one of them, say η^1_+ , by means of the relations

$$\eta_{-}^{1} = -\sigma^{3}\eta_{+}^{1}, \qquad \eta_{+}^{2} = -\mathrm{i}\,\sigma^{1}(\eta_{+}^{1})^{*}, \qquad \eta_{-}^{2} = \sigma^{2}(\eta_{+}^{1})^{*}. \tag{3.11}$$

Notice also that all the spinors never vanish, as it can be seen from their norm, given by

$$(\eta_{+}^{1})^{\dagger} \eta_{+}^{1} = 2|\xi|^{2} m \, y^{1/4} (h_{1}h_{2})^{1/8} \,. \tag{3.12}$$

We can now count the number of supersymmetries preserved by our $\operatorname{AdS}_4 \times \Sigma$ solution. ϑ is a Majorana spinor, hence it has four real degrees of freedom, while the spinors η_{\pm}^A are fully determined by the complex constant ξ . Therefore, there are eight real independent Killing spinors, that is half the number of supersymmetries of the six-dimensional $\mathcal{N} =$ (1,1) theory, hence the solution is 1/2-BPS. The eight Killing spinors correspond to four Poincaré supercharges Q and four superconformal supercharges S in the d = 3, $\mathcal{N} = 2$ SCFTs. The precise identification of the spinors with the supercharges is discussed in appendix **B**. In particular, we show that ∂_z is part of the superconformal R-symmetry, namely the U(1) generator in the OSp(2|4) superalgebra.

3.2 Global analysis I: metric and magnetic fluxes

From now on we set m = 2g/3 without loss of generality. In order to have a well-defined metric on the spindle \mathbb{Z} ,

$$ds_{\Sigma}^{2} = \frac{y^{2}}{F} dy^{2} + \frac{F}{h_{1}h_{2}} dz^{2}, \qquad (3.13)$$

and positive scalars X_i we need to take F > 0, $h_1 > 0$, $h_2 > 0$ in a closed interval not containing the curvature singularity in y = 0, thus without loss of generality we restrict to y > 0. Taking a look at the explicit form of F and its first derivative

$$F(y) = \frac{4g^2}{9} \left[y^6 - \frac{9}{4g^2} y^4 + (q_1 + q_2) y^3 + q_1 q_2 \right],$$

$$F'(y) = \frac{8g^2}{3} y^2 \left[y^3 - \frac{3}{2g^2} y + \frac{q_1 + q_2}{2} \right],$$
(3.14)

from the expression of F' we see that there exist at most three turning points, hence at most four distinct real roots of F. Since the coefficient of y^6 in F(y) = 0 is positive and we restricted to y > 0, we need at least three positive roots and, to this end, Descartes' rule of signs implies the necessary conditions

$$q_1 + q_2 > 0$$
 and $q_1 q_2 < 0$. (3.15)

Without loss of generality we take $q_1 > 0$, $q_2 < 0$. Recalling that

$$F = \frac{4g^2}{9}h_1h_2 - y^4, \qquad (3.16)$$

we have $h_1h_2 > 0$ when F > 0. Moreover, for positive y and q_1 , $h_1 > 0$ and, accordingly, $h_2 > 0$ as well.

The conditions for \mathbb{Z} to be a spindle are obtained studying $ds_{\mathbb{Z}}^2$ in the neighbourhood of the zeros of F. Denoting $[y_N, y_S]$ the range of the coordinate y, as the latter approaches one of the end-points of this interval, say y_{α} , the metric becomes

$$\mathrm{d}s_{\mathbb{Z}}^2 \simeq \mathrm{d}\varrho^2 + \varrho^2 \, \frac{g^2 F'(y_\alpha)^2}{9y_\alpha^6} \, \mathrm{d}z^2 \,, \tag{3.17}$$

where we defined $\rho^2 = |y - y_{\alpha}|$. As a consequence, ds_{Σ}^2 is a smooth orbifold metric on the spindle if the following conditions hold

$$\frac{gF'(y_N)}{3y_N^3}\Delta z = \frac{2\pi}{n_-}, \qquad \frac{gF'(y_S)}{3y_S^3}\Delta z = -\frac{2\pi}{n_+}, \qquad (3.18)$$

where the minus sign in the second relation is due to the fact that $F'(y_S) < 0$. Here n_{\pm} are two co-prime integers and Δz is the periodicity of the z coordinate. The Euler characteristic of metric (3.13) can be computed noticing that

$$\sqrt{g_{\Sigma}} R_{\Sigma} = \frac{\mathrm{d}}{\mathrm{d}y} \left(\frac{F \,\partial_y(h_1 h_2) - (h_1 h_2) \,\partial_y F}{y \,(h_1 h_2)^{3/2}} \right) \,. \tag{3.19}$$

We then find

$$\chi(\mathbb{\Sigma}) = \frac{1}{4\pi} \int_{\mathbb{\Sigma}} R_{\mathbb{\Sigma}} \operatorname{vol}(\mathbb{\Sigma}) = \frac{n_+ + n_-}{n_+ n_-}, \qquad (3.20)$$

where we employed $F(y_{\alpha}) = 0$ and the following identity:

$$\frac{gF'(y_{\alpha})}{3y_{\alpha}^3} = \frac{4g^3}{9} \left[\frac{3}{2g^2} - \frac{(q_1 + q_2)y_{\alpha}^3 + 2q_1q_2}{y_{\alpha}^4} \right].$$
(3.21)

We now proceed to the quantization conditions for the magnetic fluxes of our $AdS_4 \times \mathbb{Z}$ solution across the spindle. The integrated fluxes of F_i are given by

$$P_{i} = \frac{g}{2\pi} \int_{\Sigma} F_{i} = -g \, q_{i} \, \frac{y_{S}^{3} - y_{N}^{3}}{h_{i}(y_{N})h_{i}(y_{S})} \, \frac{\Delta z}{2\pi} = \frac{p_{i}}{n_{+}n_{-}} \,, \qquad p_{i} \in \mathbb{Z} \,, \tag{3.22}$$

where the quantization of the p_i arises from the requirement that gA_i be well-defined connection one-forms on U(1) bundles over \mathbb{Z} (*c.f.* appendix A of [7]). In particular, the total flux reads

$$P_{1} + P_{2} = \frac{4g^{3}}{9} \left[\frac{(q_{1} + q_{2})y_{S}^{3} + 2q_{1}q_{2}}{y_{S}^{4}} - \frac{(q_{1} + q_{2})y_{N}^{3} + 2q_{1}q_{2}}{y_{N}^{4}} \right] \frac{\Delta z}{2\pi}$$

$$= \frac{n_{+} + n_{-}}{n_{+}n_{-}} = \chi(\mathbb{\Sigma}), \qquad (3.23)$$

where we made used the relation (3.21) and $F(y_{\alpha}) = 0$. From (3.23) it follows that

$$p_1 + p_2 = n_+ + n_-, \qquad (3.24)$$

and therefore the two integers p_i can be conveniently parameterised as

$$p_1 = \frac{n_+ + n_-}{2} (1 + \mathbf{z}), \qquad p_2 = \frac{n_+ + n_-}{2} (1 - \mathbf{z}), \qquad (3.25)$$

where z is an appropriate rational number¹⁰. The situation is analogous to that obtained for M5-branes wrapped on the spindle [9] as here we also see that the "topologically topological twist" is realised by the solution, in contrast to the "anti-twist" that was first encountered for D3-branes and M2-branes wrapped on spindles. In [26] it is shown that very generally on the spindle only these two types of twists are possible. As explained in this reference, the occurrence of the twist case in our solution is correlated with the behavior of the Killing spinors at the north and south poles of the spindle, which we will discuss in section 3.4.

3.3 Solution of the regularity conditions

In this subsection we elaborate on the conditions worked out in the global analysis above. Specifically, we will aim to derive expressions for y_N , y_S and Δz in terms of the spindle parameters n_{\pm} and the flux parameter p_1 (or z). Imposing $F(y_{\alpha}) = 0$ we obtain the sum and product of q_1 and q_2 in terms of the roots y_N and y_S as

$$q_{1} + q_{2} = -(y_{S}^{3} + y_{N}^{3}) + \frac{9}{4g^{2}} \frac{y_{S}^{4} - y_{N}^{4}}{y_{S}^{3} - y_{N}^{3}},$$

$$q_{1}q_{2} = y_{S}^{3} y_{N}^{3} \left(1 - \frac{9}{4g^{2}} \frac{y_{S} - y_{N}}{y_{S}^{3} - y_{N}^{3}}\right),$$
(3.26)

while conditions (3.18) yield

$$\frac{n_{-}}{y_{N}}\left(y_{N}^{3} - \frac{3}{2g^{2}}y_{N} + \frac{q_{1} + q_{2}}{2}\right) = -\frac{n_{+}}{y_{S}}\left(y_{S}^{3} - \frac{3}{2g^{2}}y_{S} + \frac{q_{1} + q_{2}}{2}\right).$$
(3.27)

Plugging the first of the equations (3.26) into (3.27) and changing variables as

$$y_N \equiv w(1-x), \qquad y_S \equiv w(1+x),$$
 (3.28)

we obtain the equation

$$2w^{2}g^{2}(x^{2}+3)^{2}(x-\mu) - 3x(x^{2}+5) + 9\mu(x^{2}+1) = 0, \qquad (3.29)$$

where we defined $\mu_{\pm} \equiv n_{+} \pm n_{-}$ and $\mu \equiv \mu_{-}/\mu_{+}$. From $0 < y_{N} < y_{S}$ it immediately follows that w > 0 and 0 < x < 1. This quadratic equation for w can be easily solved,

$$w = \frac{1}{g(x^2+3)} \sqrt{\frac{9\mu(x^2+1) - 3x(x^2+5)}{2(\mu-x)}},$$
(3.30)

giving the following expressions for the two roots in terms of the parameter μ and the new variable x

$$y_N = \frac{1-x}{g(x^2+3)} \sqrt{\frac{9\mu(x^2+1) - 3x(x^2+5)}{2(\mu-x)}},$$

$$y_S = \frac{1+x}{g(x^2+3)} \sqrt{\frac{9\mu(x^2+1) - 3x(x^2+5)}{2(\mu-x)}}.$$
(3.31)

¹⁰We must take z < -1 to ensure that $p_1 < 0$, $p_2 > 0$, as follows from (3.22) and the signs of q_i . Moreover z must be chosen such that p_1 and p_2 are integers

Notice that in order for y_N, y_S to be real, there are two alternative set of conditions, namely

$$3\mu(x^2+1) - x(x^2+5) > 0$$
 and $\mu - x > 0$, (3.32)

$$3\mu(x^2+1) - x(x^2+5) < 0$$
 and $\mu - x < 0$. (3.33)

Below we will turn attention to the variable x and we will check which of these two conditions hold. From (3.22) and (3.18) we obtain the following useful relations

$$\frac{p_1 p_2}{(n_+ n_-)^2} = \frac{16g^6}{81} \frac{q_1 q_2 (y_S^3 - y_N^3)^2}{y_S^4 y_N^4} \left(\frac{\Delta z}{2\pi}\right)^2 , \qquad (3.34)$$

$$\frac{y_N}{n_-} + \frac{y_S}{n_+} = -\frac{8g^3}{9} \left[(y_S^3 - y_N^3) - \frac{3}{2g^2} (y_S - y_N) \right] \frac{\Delta z}{2\pi} \,. \tag{3.35}$$

Extracting Δz from (3.34) and inserting it into (3.35), along with the expression for the roots y_N, y_S from (3.31), we obtain

$$\mu_{+}\sqrt{-x^{4} + (9\mu^{2} - 5)x^{2} - 12\mu x + 9\mu^{2} - 4\eta\sqrt{-2p_{1}p_{2}}x} = 0, \qquad (3.36)$$

where $\eta = \text{sign}(\mu - x)$. Recalling that x > 0, in order for this equation to have a real solution we need $\eta = +1$, hence any solution will have to satisfy the conditions (3.32). In particular, being x positive by construction, we need $\mu > 0$ as well, hence $\mu_{-} > 0$ and $n_{-} < n_{+}$. Bearing these existence conditions in mind and recalling that $p_{1,2} = \mu_{+}(1\pm z)/2$, from (3.36) we find that x must be a root of the following quartic equation

$$P(x) \equiv x^{4} + (8z^{2} - 3 - 9\mu^{2})x^{2} + 12\mu x - 9\mu^{2} = 0.$$
(3.37)

From (3.35) we can then obtain the periodicity of z, namely

$$\Delta z = \chi \frac{3\pi \left(x^2 + 3\right) \left(\mu - x\right)}{8gx^2}, \qquad (3.38)$$

and from (3.26) we can express q_1 and q_2 in terms of x as

$$q_{1,2} = w \frac{3(1-x^2)}{g^2(x^2+3)^2(\mu-x)} \left[3\mu(1+x^2) - 2x \mp x(x^2+3)z \right].$$
(3.39)

All the relevant quantities are now given in terms of a solution x of equation (3.37), with 0 < x < 1 and meeting the constraints (3.32). We are left with the analysis of the existence and uniqueness of such solution. Focussing on the second condition of (3.32), namely $x < \mu$, we know that, by definition, $\mu < 1$, thus this constraint reduces the range of existence of x to $(0, \mu)$. Recalling that $z^2 > 1$ and $\mu^2 < 1$, we have

$$P(0) = -9\mu^{2} < 0,$$

$$P(\mu) = 8(z^{2} - \mu^{2})\mu^{2} > 0.$$
(3.40)

Since P(x) is continuous, there must exist at least one zero in the interval $(0, \mu)$. Let us now prove by contradiction that the first condition of (3.32) holds. Assuming that

$$3\mu(x^2+1) - x(x^2+5) \le 0, \qquad (3.41)$$

multiplying this by the positive quantity x, and using equation (3.37) to get rid of the quartic term, we obtain

$$3\mu [x(x^{2}+5) - 3\mu (x^{2}+1)] + 8(z^{2}-1)x^{2} \le 0, \qquad (3.42)$$

hence

$$3\mu(x^{2}+1) - x(x^{2}+5) \ge \frac{8(z^{2}-1)x^{2}}{3\mu} > 0, \qquad (3.43)$$

in contradiction with the starting hypothesis, which must be false. In this way, we showed that there exists at least one real solution to equation (3.37), lying inside the range 0 < x < 1.

We shall now prove that this root is unique inside the interval $(0, \mu)$. The first derivative of P(x) reads

$$P'(x) = 4x^3 + 2(8z^2 - 3 - 9\mu^2)x + 12\mu.$$
(3.44)

When $8z^2 - 3 - 9\mu^2 \ge 0$, P'(x) > 0 for $x \in (0, \mu)$, hence P(x) is strictly increasing and can thus have only one root inside the considered range. In the other case, we can focus on the zeros of the polynomial, multiply the condition P'(x) > 0 by x and remember that, at these points, P(x) = 0, ending up with

$$-\left(8z^{2}-3-9\mu^{2}\right)x^{2}+18\mu(\mu-x)>0.$$
(3.45)

This inequality is true since $8z^2 - 3 - 9\mu^2 < 0$ for hypothesis and $\mu - x > 0$. Hence, in every zero of P(x) in the range $(0, \mu)$ the polynomial must be increasing, but being P(x) continuous the root must be unique.

3.4 Global analysis II: gauge fields and Killing spinors

We shall now discuss global properties of the Killing spinors and the gauge fields A_i , following closely the exposition in [7]. Recall that the Killing spinors that we wrote in section 3.1 were obtained in the frame (3.2) and with the gauge fields in the gauge given by the expressions (3.1), subject to

$$\alpha_1 + \alpha_2 = \frac{4}{3}, \qquad (3.46)$$

which was motivated by the fact in this gauge they are independent of z. However, both the frame and the gauge fields are singular at the north and south poles of the spindle, where the azimuthal coordinate z is ill-defined. In order to shed light on the global properties of the spinors and of the gauge fields, we shall therefore cover the spindle by two patches U_{\pm} as usual, and check that spinors and gauge fields can be correctly glued across the equator, where they overlap, identifying the correct bundles of which they are sections and on which they are connections, respectively.

We begin introducing the angular coordinate φ defined as

$$\varphi = \frac{2\pi}{\Delta z} z \,, \tag{3.47}$$

with canonical 2π periodicity, so that the two gauge potentials in (3.1) read

$$A_i = \left(\alpha_i - \frac{y^3}{h_i}\right) \frac{\Delta z}{2\pi} \,\mathrm{d}\varphi\,,\tag{3.48}$$

and we define the *R*-symmetry gauge field $A_R \equiv g(A_1 + A_2)$. The open sets U_{\pm} cover the two hemispheres containing the south and north poles, respectively. Specifically, we have $y_S \in U_+ \simeq \mathbb{R}/\mathbb{Z}_{n_+}$ and $y_N \in U_- \simeq \mathbb{R}^2/\mathbb{Z}_{n_-}$. In these two patches independently we can perform the gauge transformations

$$U_{\pm}: \quad A_i^{\pm} = A_i + \Lambda_i^{\pm} \,\mathrm{d}\varphi\,, \qquad (3.49)$$

so that the transformed gauge fields A_i^{\pm} are non-singular in their respective patches provided that

$$\Lambda_{i}^{-} = \frac{1}{4g} \left(s_{i} \chi z \, \frac{1+x}{x} + \frac{2}{n_{-}} \right) - \left(\alpha_{i} - \frac{2}{3} \right) \frac{\Delta z}{2\pi} \,, \Lambda_{i}^{+} = \frac{1}{4g} \left(s_{i} \chi z \, \frac{1-x}{x} - \frac{2}{n_{+}} \right) - \left(\alpha_{i} - \frac{2}{3} \right) \frac{\Delta z}{2\pi} \,,$$
(3.50)

where $s_1 = +1$, $s_2 = -1$. We then have that $A_i^-|_{y=y_N} = 0$ and $A_i^+|_{y=y_S} = 0$, implying that both gauge fields are non-singular at the poles, as required. The corresponding gauge transformations for the *R*-symmetry gauge field A_R are given by

$$\Lambda_R^- = g(\Lambda_1^- + \Lambda_2^-) = \frac{1}{n_-}, \qquad \Lambda_R^+ = g(\Lambda_1^+ + \Lambda_2^+) = -\frac{1}{n_+}, \qquad (3.51)$$

so that on the overlap $U_{-} \cap U_{+}$ the gauge fields transform as

$$A_i^- = A_i^+ + \left(\Lambda_i^- - \Lambda_i^+\right) d\varphi = A_i^+ + \frac{p_i}{gn_+n_-} d\varphi,$$

$$A_R^- = A_R^+ + \left(\Lambda_R^- - \Lambda_R^+\right) d\varphi = A_R^+ + \frac{n_+ + n_-}{n_+n_-} d\varphi,$$
(3.52)

meaning that gA_i are connections on $O(p_i)$ bundles and A_R is a connection on the $O(n_+ + n_-)$ bundle (which is the tangent bundle) on the spindle, respectively. From the covariant derivative (2.9) we see that the *R*-symmetry charge of ϵ^1 is 1/2, while that of ϵ^2 is -1/2. This implies that a gauge transformation $A_R \mapsto A_R + \Lambda_R d\varphi$ acts on the Killing spinors as $\epsilon^1 \mapsto e^{i\varphi\Lambda_R/2}\epsilon^1$ and $\epsilon^2 \mapsto e^{-i\varphi\Lambda_R/2}\epsilon^2$, with the spinors η^1_{\pm} and η^2_{\pm} behaving accordingly.

We now move to the analysis of the global structure of the Killing spinors. Since the frame spanning the spindle $\{e^4, e^5\}$ in (3.2) is again singular at the poles, we consider two distinct local frames in each of the two patches. We define ρ_{\pm} the geodesic distance between y and each root contained in U_{\pm} , y_S and y_N respectively. We can thus write (cf. (3.17))

$$U_{-}: e^{4} \sim \mathrm{d}\varrho_{-}, \qquad e^{5} \sim \varrho_{-} \frac{\mathrm{d}\varphi}{n_{-}},$$

$$U_{+}: e^{4} \sim -\mathrm{d}\varrho_{+}, \qquad e^{5} \sim \varrho_{+} \frac{\mathrm{d}\varphi}{n_{+}},$$
(3.53)

where the sign of e^4 in U_+ is due to the fact that approaching y_S the coordinate y is increasing, while ϱ_+ is decreasing. In the patch U_- we introduce the complex coordinate $z_- = \varrho_- e^{i\varphi/n_-} = x_- + i y_-$, which is non-singular in y_N and, thus, defines a smooth oneform dz_- on the orbifold. This one-form, in turn, determines a non-singular frame, that can be obtained rotating the initial frame as follows:

$$\begin{pmatrix} e^4\\ e^5 \end{pmatrix} \mapsto \begin{pmatrix} \cos\frac{\varphi}{n_-} & -\sin\frac{\varphi}{n_-}\\ \sin\frac{\varphi}{n_-} & \cos\frac{\varphi}{n_-} \end{pmatrix} \begin{pmatrix} e^4\\ e^5 \end{pmatrix} \sim \begin{pmatrix} \mathrm{d}x_-\\ \mathrm{d}y_- \end{pmatrix} \,. \tag{3.54}$$

This is an $SO(2) \cong U(1)$ rotation of the frame in the patch U_{-} , which induces a transformation of the spinors given by the action of the exponential of the spinor representation of the infinitesimal version of the SO(2) frame rotation. Explicitly, this is a U(1) rotation of the components of the spinors by means of the matrix

$$\exp\left(-\frac{\varphi}{2n_{-}}\sigma^{1}\sigma^{2}\right) = \begin{pmatrix} e^{-i\varphi/2n_{-}} & 0\\ 0 & e^{i\varphi/2n_{-}} \end{pmatrix}.$$
 (3.55)

Performing in U_{-} the frame rotation (3.54) and the gauge transformation (3.49) with gauge parameter Λ_{R}^{-} (3.51), the spinors undergo an *R*-symmetry rotation plus a U(1) rotation. The total action on, *e.g.*, η_{+}^{1} is

$$U_{-}: \quad \eta_{+}^{1} \simeq \begin{pmatrix} f_{1}^{1/2} \\ -f_{2}^{1/2} \end{pmatrix} \mapsto e^{i\varphi/2n_{-}} \begin{pmatrix} e^{-i\varphi/2n_{-}} & 0 \\ 0 & e^{i\varphi/2n_{-}} \end{pmatrix} \begin{pmatrix} f_{1}^{1/2} \\ -f_{2}^{1/2} \end{pmatrix} = \begin{pmatrix} f_{1}^{1/2} \\ -e^{i\varphi/n_{-}} f_{2}^{1/2} \end{pmatrix}.$$
(3.56)

The φ coordinate is not well-defined in y_N , however since $f_2(y_N) = 0$, the transformed spinor is smooth and well-defined at this pole of the spindle and, thus, in the whole patch U_- . Of course the same is true for the spinors η_-^1 and η_{\pm}^2 . A similar analysis can be performed in the patch U_+ defining the non-singular coordinate $z_+ = -\varrho_+ e^{-i\varphi/n_+}$, related to the initial frame by the rotation

$$\begin{pmatrix} e^4\\ e^5 \end{pmatrix} \mapsto \begin{pmatrix} \cos\frac{\varphi}{n_+} & \sin\frac{\varphi}{n_+}\\ -\sin\frac{\varphi}{n_+} & \cos\frac{\varphi}{n_+} \end{pmatrix} \begin{pmatrix} e^4\\ e^5 \end{pmatrix} .$$
(3.57)

This transformation is analogous to the frame rotation in U_{-} , but is performed in the opposite direction, and the same happens to the spinors. The corresponding spinor rotation and gauge transformation combine so that the spinor η_{+}^{1} transforms as

$$U_{+}: \quad \eta_{+}^{1} \simeq \begin{pmatrix} f_{1}^{1/2} \\ -f_{2}^{1/2} \end{pmatrix} \mapsto e^{-i\varphi/2n_{+}} \begin{pmatrix} e^{i\varphi/2n_{+}} & 0 \\ 0 & e^{-i\varphi/2n_{+}} \end{pmatrix} \begin{pmatrix} f_{1}^{1/2} \\ -f_{2}^{1/2} \end{pmatrix} = \begin{pmatrix} f_{1}^{1/2} \\ -e^{-i\varphi/n_{+}} f_{2}^{1/2} \end{pmatrix},$$
(3.58)

giving, again, a well-defined spinor in U_+ .

In conclusion, we have shown that the Killing spinors on the spindle are smooth and well-defined, in the appropriate orbifold sense, in line with all the previous constructions of supersymmetric solutions involving spindles. The spinor transition function in going from the path U_+ to the patch U_- reads

$$\begin{pmatrix} e^{-i\varphi/2n_{-}} & 0\\ 0 & e^{i\varphi/2n_{-}} \end{pmatrix} \cdot \begin{pmatrix} e^{-i\varphi/2n_{+}} & 0\\ 0 & e^{i\varphi/2n_{+}} \end{pmatrix} = \begin{pmatrix} e^{-i\varphi\frac{n_{+}+n_{-}}{2n_{+}n_{-}}} & 0\\ 0 & e^{i\varphi\frac{n_{+}+n_{-}}{2n_{+}n_{-}}} \end{pmatrix}, \quad (3.59)$$

where the sign of the rotation in U_+ is reversed because we started with a non-singular spinor in the patch U_+ . This identifies the positive and negative chirality spin bundles $S^{(\pm)}$ on the spindle as the bundles $O(\mp \frac{1}{2}(n_+ + n_-))$, respectively. Recalling that our spinors are also charged under A_R and therefore they are sections of the bundles $O(\pm (n_+ + n_-))^{1/2} = O(\pm \frac{1}{2}(n_+ + n_-))$, we conclude that, for example

$$\frac{1}{2}(1+\rho_*)\eta_+^1 \quad \text{is a section of} \quad O(-\frac{1}{2}(n_++n_-)) \otimes O(\frac{1}{2}(n_++n_-)) = O(0) , \\ \frac{1}{2}(1-\rho_*)\eta_+^1 \quad \text{is a section of} \quad O(\frac{1}{2}(n_++n_-)) \otimes O(\frac{1}{2}(n_++n_-)) = O(n_++n_-) ,$$
(3.60)

as it was indeed obvious from the explicit transition functions obtained from passing from the expression in (3.56) to that in (3.58).

Notice that all the Killing spinors have definite chirality at the north and south poles of the spindle, and this is the same at both poles. For example, at the poles the spinor η^1_+ reads

$$\eta_{+}^{1} = \xi \sqrt{2} (m y_{\alpha})^{3/8} \begin{pmatrix} 1\\ 0 \end{pmatrix} \qquad \text{for} \qquad \alpha = N, S , \qquad (3.61)$$

which have both positive chirality. The other spinors behave similarly. As discussed in [26], this behaviour is indeed consistent with having a global topological twist.

3.5 Uplift to massive type IIA and holographic free energy

By means of the reduction ansatz of [28] we can uplift our six-dimensional $AdS_4 \times \Sigma$ background (3.1) to massive type IIA and subsequently introduce λ as in (2.20). The metric and dilaton read

$$ds_{s.f.}^{2} = \lambda^{2} \mu_{0}^{-1/3} \left\{ y^{-1} \Delta_{h}^{1/2} \left(ds_{AdS_{4}}^{2} + ds_{\Sigma}^{2} \right) + g^{-2} \Delta_{h}^{-1/2} \left[y^{3} d\mu_{0}^{2} + h_{1} \left(d\mu_{1}^{2} + \mu_{1}^{2} \sigma_{1}^{2} \right) + h_{2} \left(d\mu_{2}^{2} + \mu_{2}^{2} \sigma_{2}^{2} \right) \right] \right\},$$
(3.62)

$$e^{\Phi} = \lambda^2 \mu_0^{-5/6} y^{-3/2} \Delta_h^{1/4} , \qquad (3.63)$$

the Romans mass is

$$F_{(0)} = \frac{2g}{3\lambda^3}, \qquad (3.64)$$

while the four-form flux takes the form

$$F_{(4)} = \frac{\lambda \mu_0^{1/3} h_1 h_2}{g^3 \Delta_h} \left\{ \frac{U_h}{\Delta_h} \frac{\mu_1 \mu_2}{\mu_0} \, \mathrm{d}\mu_1 \wedge \mathrm{d}\mu_2 \wedge \sigma_1 \wedge \sigma_2 - g \sum_{i \neq j} F_i \wedge \mathrm{d}\phi_j \wedge \left(\mu_0 \mu_j \, \mathrm{d}\mu_j - y^3 h_j^{-1} \mu_j^2 \, \mathrm{d}\mu_0\right) + \frac{y^3}{\Delta_h} \sum_{i \neq j} \frac{h_j (h'_i - 3y^{-1} h_i)}{h_i} \mu_i^2 \, \mathrm{d}y \wedge \sigma_i \wedge \sigma_j \wedge \left(\mu_0 \mu_j \, \mathrm{d}\mu_j - y^3 h_j^{-1} \mu_j^2 \, \mathrm{d}\mu_0\right) \right\}.$$
(3.65)

For convenience, we defined the functions

$$\Delta_{h} = h_{1}h_{2}\,\mu_{0}^{2} + y^{3}h_{2}\,\mu_{1}^{2} + y^{3}h_{1}\,\mu_{2}^{2},$$

$$U_{h} = 2\left[(y^{3} - h_{1})(y^{3} - h_{2})\mu_{0}^{2} - y^{6}\right] - \frac{4}{3}\Delta_{h}.$$
(3.66)

The quantization of the fluxes proceeds as in (2.24) and it yields again the relations (2.25), which fix the parameters g and λ in terms of the integers N and n_0 . Therefore our ten-dimensional solution is characterised by five integers: the pair n_0 and N, determining the dual five-dimensional theory, n_{\pm} defining the spindle, and z related to the magnetic charges threading this.

The ten-dimensional geometry has a form analogous to that of the solutions in section 2.4. The internal six-dimensional space M_6 has a fibration structure

$$\mathbb{S}^4 \hookrightarrow M_6 \to \mathbb{Z}, \qquad (3.67)$$

with the twisting of the bundle specified by the connection one-forms gA_i with Chern numbers

$$P_i = \frac{g}{2\pi} \int_{\mathbb{Z}} \mathrm{d}A_i = \frac{p_i}{n_+ n_-}, \qquad p_i \in \mathbb{Z}, \qquad (3.68)$$

subject to

$$P_1 + P_2 = \frac{n_+ + n_-}{n_+ n_-} = \chi(\mathbb{Z}).$$
(3.69)

The solution can then be interpreted as follows: one starts with a geometry of the type $\mathbb{R}^{1,2} \times \mathbb{R} \times Y_6$, where Y_6 is the total space of the vector bundle

$$\mathcal{O}(-p_1) \oplus \mathcal{O}(-p_2) \hookrightarrow Y_6 \to \mathbb{Z},$$
 (3.70)

and the condition (3.69) guarantees that the first Chern class of this bundle vanishes, thus Y_6 is a local Calabi-Yau three-fold. At the origin of \mathbb{R} there are an O8-plane and $N_f = 8 - n_0$ coincident D8-branes, and in addition one wraps N D4-branes over the zero section Σ of Y_6 .

At low energies the effective theory on the D4/D8-system will be a d = 3, $\mathcal{N} = 2$ field theory, obtained from the compactification on Σ of the d = 5, $\mathcal{N} = 1$ SCFT with gauge group USp(2N) [27], with the "topologically topological twist". The supergravity solution above strongly suggests that in the large N limit this is a SCFT and its S^3 free energy can be computed holographically as before [32]. Specifically, we have

$$F_{S^3 \times \Sigma} = \frac{16\pi^3}{(2\pi\ell_s)^8} \frac{3\pi^2\lambda^4}{10g^4} \left(y_S^3 - y_N^3\right) \Delta z$$

= $\chi \frac{\sqrt{3}\pi N^{5/2}}{5\sqrt{8 - N_f}} \frac{[3\mu(x^2 + 1) - x(x^2 + 5)]^{3/2}}{x(x^2 + 3)(\mu - x)^{1/2}}.$ (3.71)

Notice that the dependence of x on the parameter z and μ could be made explicit by writing out the solution to the quartic (3.37), however this is extremely cumbersome and we will refrain from doing so. Alternatively, one could think of x and μ as the two independent parameters, with z given in terms of these two by solving (3.37), which is a simple quadratic equation. In any case, in the next section we will reproduce the expression (3.71) analytically, starting from a conjectural large N free energy of the dual field theories.

Noticing that μ is a free "small" parameter¹¹, it is useful to expand in series of μ near to $\mu \to 0$ (holding χ fixed), which formally corresponds to reducing to a spindle with equal conical deficits, and in particular it includes the two-sphere for $n_+ = n_- = 1$. The root of the quartic equation (3.37) meeting the required constraints then has the following expansion

$$x = \frac{3}{2+t}\mu + \frac{27(1+t)(3+t)}{2t(2+t)^4}\mu^3 + \mathcal{O}(\mu^5), \qquad (3.72)$$

where we have defined $t \equiv \sqrt{8z^2 + 1}$, with t > 3. Inserting this in the free energy (3.71) we obtain the expansion

$$F_{S^3 \times \mathbb{Z}} = \chi \frac{\pi N^{5/2}}{5\sqrt{8 - N_f}} \left[\frac{(t-3)^{3/2}}{(t-1)^{1/2}} + \frac{6(t-3)^{3/2}}{(2+t)(t-1)^{3/2}} \mu^2 \right] + \mathcal{O}(\mu^4), \qquad (3.73)$$

¹¹Although $\mu \in \mathbb{Q}$, we can treat it formally as a real variable taking values in the interval (0, 1).

which, after setting $\chi = 2$, at leading order in μ agrees with the free energy (2.42) for g = 0 and $\kappa = 1$ [2]. This suggests that it may be possible to recover the $AdS_4 \times S^2$ solution in [2] by performing a suitable scaling limit of our solutions, but we will not attempt to do so here. For future reference, let us also record the expansion for Δz , that reads

$$\Delta z = \chi \frac{\pi}{8g} \left[\frac{(t-1)(2+t)}{\mu} - \frac{3(9+34t+25t^2+4t^3)}{2t(2+t)^2} \mu \right] + \mathcal{O}(\mu^3).$$
(3.74)

4 Field theory

We conjecture that the solutions we have constructed in section 3 are holographically dual to three-dimensional SCFTs obtained by compactifying on a spindle Σ the five-dimensional SCFTs dual to the solution of [27]. In the reminder of this section we will provide evidence for this by proposing an off-shell free energy whose extremization reproduces exactly the holographic free energy (3.71). This is an extension of the entropy functions that have been shown to provide an efficient method for reproducing the entropy of supersymmetric AdS black holes in various dimensions. A priori, this function should be derived from first principles, by computing (minus the logarithm of) the localized partition function of the d = 5 SCFT, placed on the background of $S^3 \times \Sigma$, and then taking the large N limit. This strategy has been implemented in [35,36] for the background $S^3 \times \Sigma_g$ and, indeed, it led to reproducing the holographic free energy (2.42) previously obtained in [2]. Instead, we will follow a short-cut inspired by the "gravitational blocks" advocated in [21]. We will infer from the supergravity description the main ingredients involved in the fieldtheoretic construction and we will propose a large N off-shell free energy on $S^3 \times \Sigma$ obtained by suitably gluing the S^5 free energy of the d = 5 theories. We will then show that extremizing it will reproduce exactly the holographic free energy (3.71).

4.1 d = 5 SCFTs dual to the AdS₆ solution

Let us begin by recalling the salient features of the five-dimensional theory that is holographically dual to the $\operatorname{AdS}_6 \times \mathbb{S}^4$ background of massive type IIA, arising in the nearhorizon limit of N D4-branes and N_f D8-branes, that we reviewed in section 2.3. This is an $\mathcal{N} = 1$ gauge theory with gauge group USp(2N), coupled to N_f massless hypermultiplets in the fundamental representation and one hypermultiplet in the antisymmetric representation of USp(2N) [39]. At low energies, this theory flows to an interacting SCFT, with global symmetry $SU(2)_R \times SU(2)_F \times E_{N_f+1}$, where the first two factors are realised as symmetries of the $\operatorname{AdS}_6 \times \mathbb{S}^4$ solution [40]. Placing this theory on a rigid S^5 background, one can compute the exact localized partition function Z_{S^5} and consider the associated free energy, namely

$$F_{S^5} \equiv -\log Z_{S^5} \,, \tag{4.1}$$

as a good measure of the degrees of freedom of the theory. This was computed in [33], that also showed that in the large N limit it becomes

$$F_{S^5} = -\frac{9\sqrt{2}\pi}{5} \frac{N^{5/2}}{\sqrt{8 - N_f}} \tag{4.2}$$

and is reproduced by a holographic calculation in the solution of [27]. The S^5 free energy may also be "refined", promoting it to an off-shell free energy, regarded as a function of the fugacities for the $U(1) \times U(1)$ Cartan subgroup of $SU(2)_R \times SU(2)_F$, which we will denote as Δ_i , with i = 1, 2. A priori, the *R*-symmetry can mix with any flavour symmetry and the Δ_i parameterise this mixing. For the present theory this is actually not necessary, as the *R*-symmetry is non-Abelian, nevertheless this will be useful in the sequel. We can then write

$$F_{S^5}(\Delta_i) = -\frac{9\sqrt{2}\pi}{5} \frac{N^{5/2}}{\sqrt{8 - N_f}} (\Delta_1 \Delta_2)^{3/2} \equiv \frac{27}{4} \mathcal{F}(\Delta_i) , \qquad (4.3)$$

with the fugacities obeying, in a canonical normalization, the R-symmetry constraint

$$\Delta_1 + \Delta_2 = 2. \tag{4.4}$$

For later convenience we have defined the auxiliary function $\mathcal{F}(\Delta_i)$. Extremizing $F_{S^5}(\Delta_i)$ gives $\Delta_1^* = \Delta_2^* = 1$ and inserting these values back one reproduces the initial free energy

$$F_{S^5} \equiv F_{S^5}(\Delta_i^*) = -\frac{9\sqrt{2}\pi}{5} \frac{N^{5/2}}{\sqrt{8 - N_f}}.$$
(4.5)

Let us now move to discussing compactifications of this theory to d = 3 dimensions and the corresponding off-shell free energies.

4.2 d = 3 SCFTs dual to the AdS₄ × Σ_g solutions

We can obtain three-dimensional $\mathcal{N} = 2$ theories by compactifying the above d = 5SCFT on a Riemann surface $\Sigma_{\rm g}$ of arbitrary genus g, performing the standard topological twist [2]. Specifically, we place the theory on $\Sigma_{\rm g}$ and couple it to two background gauge fields A_i for the $U(1) \times U(1)$ Cartan subgroup of $SU(2)_R \times SU(2)_F$, with appropriately quantized magnetic fluxes¹²

$$\mathbf{n}_i = \frac{1}{2\pi} \int_{\Sigma_{\mathbf{g}}} F_i = p_i \kappa (1 - \mathbf{g}) \in \mathbb{Z}.$$
(4.6)

The topological twist implies that the *R*-symmetry gauge field $A_R = A_1 + A_2$ is identified with a connection on the tangent bundle, thus

$$\frac{1}{2\pi} \int_{\Sigma_{g}} dA_{R} = \chi(\Sigma_{g}) = 2(1 - g) = \mathfrak{n}_{1} + \mathfrak{n}_{2}, \qquad (4.7)$$

and the Killing spinors become just constant. It is then convenient to parameterise the magnetic fluxes as

$$\mathbf{n}_1 = (1 - g)(1 + \kappa z), \qquad \mathbf{n}_2 = (1 - g)(1 - \kappa z).$$
 (4.8)

¹²Here and in the following we shall rename the background gauge fields as $gA_i \mapsto A_i$, which is more natural from the field theory point of view. The magnetic fluxes \mathbf{n}_i correspond precisely to the fluxes P_i defined in (2.33). However, we denote these with different symbols to emphasise the fact that the P_i were defined as integrals of supergravity fields, living in D = 6, while the \mathbf{n}_i are defined as integrals of background gauge fields, living in d = 5.

The exact *R*-symmetry of the d = 3 theory will be determined by extremizing the off-shell S^3 free energy [41], viewed as a function of the fugacities Δ_i . Equivalently, this quantity may be thought of as the off-shell free energy of the d = 5 theory on $S^3 \times \Sigma_g$. The latter quantity was computed in [35,36] using localization, and in the large *N* limit it was shown to reproduce the holographic free energy (2.42). Below we will show that it can also be reproduced by a formula obtained by gluing two gravitational blocks. We begin defining the following conjectural large *N* off-shell free energy¹³

$$F(\Delta_i, \epsilon; \mathfrak{n}_i) \equiv \frac{1}{\epsilon} \left(\mathcal{F}(\Delta_i^+) - \mathcal{F}(\Delta_i^-) \right) , \qquad (4.9)$$

where $\mathcal{F}(\Delta_i)$ is defined in (4.3) and

$$\Delta_i^+ \equiv \Delta_i + \mathfrak{n}_i \epsilon \,, \qquad \Delta_i^- \equiv \Delta_i - \mathfrak{n}_i \epsilon \,, \tag{4.10}$$

with Δ_i satisfying (4.4).

Notice that in addition to the fugacities Δ_i of the parent d = 5 theory and the magnetic fluxes \mathbf{n}_i , (4.9) depends a priori also on the parameter ϵ , although we shall see below that the extremization equations automatically set $\epsilon = 0$. At least in the case g = 0, ϵ may be interpreted as the fugacity associated to the $U(1)_J \subset SU(2)_J$ rotational symmetry of the two-sphere. Extremizing (4.9) with respect to ϵ and Δ_i , subject to (4.4), we easily find the critical values

$$\Delta_1^* = 1 + \frac{\kappa + \sqrt{8z^2 + \kappa^2}}{4z}, \qquad \epsilon^* = 0.$$
(4.11)

For the two-sphere, the fact that ϵ^* vanishes means that the *R*-symmetry of the compactified theory does not have a component along $U(1)_J$, as expected. In any case, inserting the critical values in (4.9) we get

$$F(\Delta_i^*, 0; \mathbf{z}) = \frac{16\pi (1 - g)\kappa N^{5/2} \left(\mathbf{z}^2 - \kappa^2\right)^{3/2} \left(\sqrt{\kappa^2 + 8\mathbf{z}^2} - \kappa\right)}{5\sqrt{8 - N_f} \left(\kappa\sqrt{\kappa^2 + 8\mathbf{z}^2} - \kappa^2 + 4\mathbf{z}^2\right)^{3/2}},$$
(4.12)

which agrees with (2.42). Since $\epsilon^* = 0$, we could have started setting $\epsilon = 0$ in (4.9), thus reducing to the known ϵ -independent off-shell free energy

$$\lim_{\epsilon \to 0} F(\Delta_i, \epsilon; \mathfrak{n}_i) = 2 \sum_{j=1}^2 \mathfrak{n}_j \frac{\partial \mathcal{F}(\Delta_i)}{\partial \Delta_j} = -\frac{4\sqrt{2\pi}}{5} \frac{N^{5/2}}{\sqrt{8 - N_f}} (\Delta_1 \Delta_2)^{1/2} (\mathfrak{n}_2 \Delta_1 + \mathfrak{n}_1 \Delta_2) \,, \quad (4.13)$$

corresponding to the standard topological twist [31, 35, 36, 43]. In particular, extremizing (4.13) reproduces (4.12) with Δ_1^* given in (4.11).

We notice that Δ_1^* , Δ_2^* match with the scaling dimensions of two particular 1/2-BPS operators, corresponding to D2-branes wrapped on calibrated surfaces, embedded in the internal six-dimensional geometries, and sitting at the center of AdS₄, described in [2]. For g > 1 we can set z = 0 so that $\Delta_1^* = \Delta_2^* = 1$ and the free energy (4.12) reduces to the universal relation [44]

$$F(\Delta_i^*, 0; \mathfrak{n}_1 = \mathfrak{n}_2 = 1 - g) = \frac{16\pi(g-1)N^{5/2}}{5\sqrt{2}\sqrt{8 - N_f}} = -\frac{8}{9}(g-1)F_{S^5}.$$
(4.14)

¹³For g = 0, the form (4.9) has been recently proved in [42] for $S_b^3 \times S_\epsilon^2$.

4.3 d = 3 SCFTs dual to the AdS₄ × \mathbb{Z} solutions

We now consider the compactification of the d = 5, $\mathcal{N} = 1$ SCFT on a spindle, that we expect to be dual to the solutions we constructed in section 3. In particular, we perform a global topological twist, which means that we place the theory on Σ and couple it to two background gauge fields A_i for the $U(1) \times U(1)$ Cartan subgroup of $SU(2)_R \times SU(2)_F$, with appropriately quantized magnetic fluxes

$$\mathfrak{n}_i = \frac{1}{2\pi} \int_{\mathbb{Z}} F_i = \frac{p_i}{n_+ n_-}, \qquad p_i \in \mathbb{Z}.$$
(4.15)

As for the standard topological twist, the *R*-symmetry gauge field $A_R = A_1 + A_2$ becomes a connection on the tangent bundle, thus

$$\frac{1}{2\pi} \int_{\Sigma} dA_R = \chi(\Sigma) = \frac{n_+ + n_-}{n_+ n_-} = \mathfrak{n}_1 + \mathfrak{n}_2, \qquad (4.16)$$

but crucially, this does not imply that the metric on the spindle has constant curvature, nor that the spinors are chiral and constant. Specifically, the rigid Killing spinors are expected to behave precisely as the spinors η_{\pm}^{A} arising in the supergravity solution. As before in the paper, we will continue to parameterise the magnetic fluxes as

$$\mathfrak{n}_1 = \frac{\chi}{2}(1+\mathbf{z}), \qquad \mathfrak{n}_2 = \frac{\chi}{2}(1-\mathbf{z}).$$
(4.17)

Taking inspiration from the entropy function proposed in [12], we conjecture that the large N off-shell free energy for these theories is given by

$$F(\Delta_i, \epsilon; \mathfrak{n}_i, n_+, n_-) \equiv \frac{1}{\epsilon} \left(\mathcal{F}(\Delta_i^+) - \mathcal{F}(\Delta_i^-) \right) , \qquad (4.18)$$

where $\mathcal{F}(\Delta_i)$ is defined in (4.3) and

$$\Delta_i^+ \equiv \Delta_i + \epsilon \left(\mathfrak{n}_i + \frac{1}{2} \frac{n_+ - n_-}{n_+ n_-} \right) , \qquad \Delta_i^- \equiv \Delta_i - \epsilon \left(\mathfrak{n}_i - \frac{1}{2} \frac{n_+ - n_-}{n_+ n_-} \right) . \tag{4.19}$$

The Δ_i are the fugacities parameterising the *R*-symmetry within the $U(1) \times U(1) \subset$ $SU(2)_R \times SU(2)_F$ global symmetries of the d = 5 theory, and are therefore still subject to the constraint (4.4), while ϵ is an equivariant parameter for the spindle, that is a fugacity for the $U(1)_J$ rotational symmetry. In general, we expect that this will parameterise a non-trivial mixing of the *R*-symmetry of the parent theory, with the $U(1)_J$ of the spindle. The off-shell free energy (4.18) bares a close resemblance to the off-shell central charge for D3-branes wrapped on spindle in [10] and we shall elaborate on this in the next section.

Employing the parametrisation (4.17) we see that upon redefining $\chi \epsilon = \hat{\epsilon}$, the off-shell free energy (4.18) becomes $F = \chi \cdot f(\Delta_i, \hat{\epsilon}; \mu, \mathbf{z})$, implying that the free energy at the critical point must be a function of μ and \mathbf{z} only, with an overall factor of χ . Notice that setting $n_+ = n_- = 1$ the present setup reduces to the case of twisted compactification on S^2 , that is the case g = 0 discussed in the previous subsection.

In order to implement the constraint (4.4) it is useful to introduce a Lagrange multiplier and consider the extremization of the following function

$$\mathcal{S}(\Delta_i, \epsilon, \Lambda; \mathfrak{n}_i, n_+, n_-) = F(\Delta_i, \epsilon; \mathfrak{n}_i, n_+, n_-) + \Lambda(\Delta_1 + \Delta_2 - 2), \qquad (4.20)$$

that is analogous to the entropy functions studied in the literature. The corresponding extremality equations read

$$\Lambda + \frac{1}{\epsilon} \left[\frac{\partial \mathcal{F}(\Delta_i^+)}{\partial \Delta_j^+} - \frac{\partial \mathcal{F}(\Delta_i^-)}{\partial \Delta_j^-} \right] = 0,$$

$$\frac{F}{\epsilon} - \frac{1}{\epsilon} \sum_{j=1}^2 \left[\left(\mathfrak{n}_j + \frac{n_+ - n_-}{2n_+ n_-} \right) \frac{\partial \mathcal{F}(\Delta_i^+)}{\partial \Delta_j^+} + \left(\mathfrak{n}_j - \frac{n_+ - n_-}{2n_+ n_-} \right) \frac{\partial \mathcal{F}(\Delta_i^-)}{\partial \Delta_j^-} \right] = 0, \qquad (4.21)$$

$$\Delta_1 + \Delta_2 - 2 = 0.$$

These are four equations for the variables Δ_1 , Δ_2 , ϵ , Λ and, in order to solve these, it is convenient to process them further. Noticing that (4.18) is homogeneous of degree two in Δ_i and ϵ , by Euler's theorem we have

$$\sum_{j=1}^{2} \Delta_{j} \frac{\partial F}{\partial \Delta_{j}} + \epsilon \frac{\partial F}{\partial \epsilon} = 2F.$$
(4.22)

The extremization equations for \mathcal{S} written as

$$\frac{\partial F}{\partial \Delta_i} + \Lambda = 0, \qquad \qquad \frac{\partial F}{\partial \epsilon} = 0, \qquad (4.23)$$

then immediately imply $\Lambda = -F$. We can therefore eliminate Λ from the system and write the remaining two independent equations as

$$\frac{\partial \mathcal{F}(\Delta_i^+)}{\partial \Delta_1^+} - \frac{\partial \mathcal{F}(\Delta_i^-)}{\partial \Delta_1^-} = \frac{\partial \mathcal{F}(\Delta_i^+)}{\partial \Delta_2^+} - \frac{\partial \mathcal{F}(\Delta_i^-)}{\partial \Delta_2^-}, \qquad (4.24)$$
$$\left(1 + \frac{2\epsilon}{n_+}\right) F = 2\sum_{j=1}^2 \mathfrak{n}_j \frac{\partial \mathcal{F}(\Delta_i^+)}{\partial \Delta_j^+},$$

where one has to use also the constraint $\Delta_1 + \Delta_2 = 2$. Notice that taking $\epsilon \to 0$ the second equation reduces to the first equality in (4.13).

After some work, we determined the critical values

$$\Delta_1^* = 1 + \frac{2\mathbf{z}x}{(x^2 + 3)(\mu - x)}, \qquad \epsilon^* = \frac{1}{\chi} \frac{4x^2}{(x^2 + 3)(\mu - x)}, \qquad (4.25)$$

in terms of the parameter x, that is the unique root in the interval (0, 1) of the quartic (3.37) that we introduced in the discussion of the gravitational solution. Inserting these values back into (4.18) we obtain

$$F(\Delta_i^*, \epsilon^*; \mathbf{z}, \chi, \mu) = \chi \frac{\sqrt{3\pi}N^{5/2}}{5\sqrt{8-N_f}} \frac{[3\mu(x^2+1) - x(x^2+5)]^{3/2}}{x(x^2+3)(\mu-x)^{1/2}},$$
(4.26)

which, remarkably, agrees exactly with the gravitational free energy (3.71)!

To arrive at the solution (4.25), we first solved the extremality equations perturbatively in μ around $\mu = 0$, obtaining agreement with the expansion (3.73), up to high powers of μ . We then noticed that the result for ϵ^* could be rewritten as

$$\epsilon^* = \frac{3}{4g} \frac{2\pi}{\Delta z} \,, \tag{4.27}$$

which is a universal relation holding in all previous spindle solutions [7–13]. Using this, we then obtained the result for Δ_1^* in (4.25).

As discussed in section 3, in the limit $n_{+} = n_{-} = 1$ (4.26) reproduces the free energy (4.12) for g = 0, corresponding to the compactification of the d = 5 theory on the two-sphere, with the standard topological twist. Moreover, expanding Δ_{1}^{*} in (4.25) in series of μ around $\mu = 0$ we find that

$$\Delta_1^* = 1 + \frac{1 + \sqrt{8z^2 + 1}}{4z} + \mathcal{O}(\mu^2), \qquad (4.28)$$

again in agreement with the two-sphere value given in (4.11). It would be interesting to reproduce Δ_1^* , Δ_2^* by computing the scaling dimensions of some supersymmetric probe D2-branes wrapped on calibrated two-cycles in the ten-dimensional geometry (3.62).

5 Gravitational blocks for branes on spindles

The off-shell free energy that we discussed in the previous section may be regarded as a particular instance of a general class of off-shell free energies F^{\pm} , for twisted compactifications of d-dimensional theories on the spindle Σ . Below we will state our conjecture and we will then illustrate how it encapsulates and generalises various extremal functions discussed in the literature. From the constructions of M2, D3 and M5-branes wrapped on a spindle and the results we discussed so far in this paper, it has emerged that supersymmetry on a spindle can be preserved in two different ways, that can be referred to as *twist* and *anti-twist*. These are characterised by two types of background *R*-symmetry gauge field A_R , with fluxes given by

$$\frac{1}{2\pi} \int_{\Sigma} \mathrm{d}A_R = \frac{n_+ + \sigma n_-}{n_+ n_-} \,, \tag{5.1}$$

where $\sigma = +1$ for the twist and $\sigma = -1$ for the anti-twist. The sign $\sigma = +1$ corresponds to the choice we made in section 4.3, and that was also realised for M5-branes wrapped on the spindle in [9]. The sign $\sigma = -1$ has been realised by the supergravity solutions for D3branes [8,10,11] and M2-branes [7,12,13]. Below we will treat both cases simultaneously, with the understanding that not all cases may have a counterpart as gravity solutions. In [26] it is proved that these are the only two possible twists preserving supersymmetry on the spindle.

A large class of SCFTs in different dimensions are expected to be characterised by supersymmetric partition functions $Z_{\mathcal{M}}$, where \mathcal{M} are rigid geometries comprising a metric on an appropriate curved space, the background gauge fields for the *R*-symmetry and possibly other flavour symmetries. The associated free energy is defined by

$$F_{\mathcal{M}} \equiv -\log Z_{\mathcal{M}} \tag{5.2}$$

and it is regarded as a function of the fugacities Δ_i associated to the Cartan subgroup of the continuous global symmetry group of the theory. For theories with an Abelian *R*symmetry, that we are interested in, the fugacities obey a constraint that we can always normalise to be

$$\sum_{i=1}^{\mathfrak{d}} \Delta_i = 2 \,, \tag{5.3}$$

where \mathfrak{d} is the rank of the global symmetry group, of which the Abelian *R*-symmetry is part. In general, these are complicated matrix models, which however simplify drastically in special limits, such as the large *N* limit or Cardy-like limits, reducing to simple local functions of the fugacities Δ_i . For example, in four dimensions the partition function corresponding to $\mathcal{M} = S^1 \times S^3$ is the (refined) superconformal index and in either limits its logarithm is related to the trial central charge $a_4(\Delta_i)$ of the theory. In all SCFTs possessing an Abelian *R*-symmetry, it has been either proved or conjectured that the exact superconformal *R*-symmetry is determined by extremizing these quantities.

We conjecture that for a general class of d-dimensional SCFTs compactified on the spindle, with either the twist or the anti-twist, the exact R-symmetry is determined by extremizing the following off-shell free energies

$$F^{\pm}(\Delta_i, \epsilon; \mathfrak{n}_i, n_+, n_-, \sigma) = \frac{1}{\epsilon} \left(\mathcal{F}_d(\Delta_i^+) \pm \mathcal{F}_d(\Delta_i^-) \right) , \qquad (5.4)$$

where the variables Δ_i^+ , Δ_i^- are defined as

$$\Delta_i^+ \equiv \Delta_i + \epsilon \left(\mathfrak{n}_i + \frac{r_i}{2} \frac{n_+ - \sigma n_-}{n_+ n_-} \right), \qquad \Delta_i^- \equiv \Delta_i - \epsilon \left(\mathfrak{n}_i - \frac{r_i}{2} \frac{n_+ - \sigma n_-}{n_+ n_-} \right), \tag{5.5}$$

and the magnetic fluxes through the spindle, n_i , satisfy the constraint

$$\sum_{i=1}^{\mathfrak{d}} \mathfrak{n}_i = \frac{n_+ + \sigma n_-}{n_+ n_-} \,. \tag{5.6}$$

The building blocks are the functions $\mathcal{F}_d(\Delta_i)$ summarised in Table 2. They are proportional to: the S^3 off-shell free energy of the ABJM theory, the trial central charge of the $\mathcal{N} = 4$ SYM theory, the S^5 off-shell free energy of the d = 5, $\mathcal{N} = 1$ SCFT and the trial central charge of the d = 6, (2,0) SCFT, respectively. In the first two cases, it is straightforward to replace these with the corresponding quantities for more general d = 3, $\mathcal{N} = 2$ theories and d = 4, $\mathcal{N} = 1$ theories, but in this paper we will not pursue this.

	d = 3	d = 4	d = 5	d = 6
\mathcal{F}_d	$b_3(\Delta_1\Delta_2\Delta_3\Delta_4)^{1/2}$	$b_4(\Delta_1\Delta_2\Delta_3)$	$b_5(\Delta_1\Delta_2)^{3/2}$	$b_6(\Delta_1\Delta_2)^2$
b_d	$-\frac{\sqrt{2\pi}}{3}N^{3/2}$	$-\frac{3}{2}N^{2}$	$-\frac{2^{5/2}\pi}{15}\frac{N^{5/2}}{\sqrt{8-N_f}}$	$-\frac{9}{256}N^3$
	$-F_{S^{3}}$	$-6a_{4}$	$\frac{4}{27}F_{S^5}$	$-\frac{63}{4096}a_6$
±	$-\sigma$		$-\sigma$	—

Table 2: Various $\mathcal{F}_d(\Delta_i)$ in different spacetime dimension d. The constants b_d are given in terms of the free energy on S^d (d = 3, 5) or the central charge a_d (d = 4, 6). In d = 6and d = 5 the rank of the global symmetry group is $\mathfrak{d} = 2$, in d = 4 it is $\mathfrak{d} = 3$, while in d = 3 it is $\mathfrak{d} = 4$. In the last row we summarised the relations between the gluing sign \pm and the sign σ , characterising the type of twist.

We expect that the form (5.4) arises, in the large N limit, from a fixed point formula, with the "blocks" $\mathcal{F}_d(\Delta_i)$ evaluated at the north and south poles of the spindle. The superscripts in F^{\pm} refer to the relative choice of sign \pm in (5.4), corresponding to the type of gluing of the contributions of the two hemispheres of the spindle [21]. We will give circumstantial evidence that in D = 4, 6 the type of gluing is correlated with the type of twist, specifically that the gluing sign is $-\sigma$. On the other hand, in D = 5, 7 the results of the explicit supergravity solutions are all reproduced by the minus gluing sign. In the examples that we discuss below, we will explain which choice of twist and gluing is relevant, but a more systematic understanding of these choices is clearly desirable.

The variable ϵ is a fugacity associated to the $U(1)_J$ rotational symmetry of the spindle and the significance of the fact that at the critical point of (5.4) this takes a non-zero value is that the *R*-symmetry of the parent *d*-dimensional theory mixes with $U(1)_J$ to give the exact superconformal *R*-symmetry of the (d-2)-dimensional theory arising in the IR, when this flows to an SCFT. The constants r_i are arbitrary, but subject to the constraint

$$\sum_{i=1}^{\mathfrak{d}} r_i = 2 \,, \tag{5.7}$$

and parameterise the ambiguities of defining flavour symmetries [10]. In the previous section we picked the most symmetric choice, corresponding to $r_i = \frac{\delta}{2} = 1$. However, it is simple to show that the functions (5.4) evaluated at the critical point are independent of the choice of r_i . Introducing a new set of variables defined as

$$\varphi_i \equiv \Delta_i + \frac{r_i}{2} \frac{n_+ - \sigma n_-}{n_+ n_-} \epsilon \,, \tag{5.8}$$

the off-shell free energies simply read

$$F^{\pm}(\varphi_i,\epsilon;\mathfrak{n}_i) = \frac{1}{\epsilon} \left(\mathcal{F}_d(\varphi_i + \mathfrak{n}_i\epsilon) \pm \mathcal{F}_d(\varphi_i - \mathfrak{n}_i\epsilon) \right), \qquad (5.9)$$

where, from now on, we will omit n_+ , n_- , σ from the arguments of the function, in order not to clutter the subsequent formulas. The variables φ_i , ϵ satisfy the constraint

$$\sum_{i=1}^{\mathfrak{d}} \varphi_i - \frac{n_+ - \sigma n_-}{n_+ n_-} \epsilon = 2, \qquad (5.10)$$

inherited from (5.3) and (5.7). Since (5.9) does not depend on the constants r_i , it follows that the critical values φ_i^* , ϵ^* and $F^{\pm}(\varphi_i^*, \epsilon^*; \mathbf{n}_i)$ do not depend on the r_i either.

Our proposal unifies previous proposals concerning entropy functions [12, 21, 45, 46]and central charges [10], extending these to compactifications of *d*-dimensional theories on spindles with both twist and anti-twist. Below we shall illustrate this, recovering known results and discussing some generalisations. The constrained extremization problem can be carried out introducing a Lagrange multiplier. Defining

$$\mathcal{S}^{\pm}(\varphi_i,\epsilon,\Lambda;\mathfrak{n}_i,n_+,n_-) = F^{\pm}(\varphi_i,\epsilon;\mathfrak{n}_i,n_+,n_-) + \Lambda\left(\sum_{j=1}^{\mathfrak{d}}\varphi_j - \frac{n_+ - \sigma n_-}{n_+ n_-}\epsilon - 2\right), \quad (5.11)$$

and using the fact that (5.4) is homogeneous of degree¹⁴ h in φ_i and ϵ , by means of Euler's theorem we have that $\Lambda = -\frac{h}{2}F^{\pm}$, and we can therefore eliminate Λ from the system.

 $^{^{14}}h = 1$ for d = 3, h = 2 for d = 4, 5 and h = 3 for d = 6.

The resulting extremization equations can be written as

$$\frac{h\epsilon}{2}F^{\pm} = \frac{\partial \mathcal{F}_d(\varphi_i^+)}{\partial \varphi_j^+} \pm \frac{\partial \mathcal{F}_d(\varphi_i^-)}{\partial \varphi_j^-}, \qquad (j = 1, \dots, \mathfrak{d})$$

$$\left(1 + \frac{\sigma h\epsilon}{n_+}\right)F^{\pm} = 2\sum_{j=1}^{\mathfrak{d}} \mathfrak{n}_j \frac{\partial \mathcal{F}_d(\varphi_i^+)}{\partial \varphi_j^+}, \qquad (5.12)$$

where $\varphi_i^{\pm} \equiv \varphi_i \pm \mathfrak{n}_i \epsilon$. Notice that the last equation can be replaced by

$$\left(1 - \frac{h\epsilon}{n_{-}}\right)F^{\pm} = \mp 2\sum_{j=1}^{\mathfrak{d}}\mathfrak{n}_{j}\frac{\partial\mathcal{F}_{d}(\varphi_{i}^{-})}{\partial\varphi_{j}^{-}}.$$
(5.13)

5.1 M2-branes

Supergravity solutions describing M2-branes wrapped on the spindle were first constructed in [7] in minimal D = 4 gauged supergravity and generalised to $U(1)^2$ gauged supergravity in [12, 13]. They realise the anti-twist, $\sigma = -1$. The corresponding dual field theory is the ABJM model compactified on the spindle, with two background gauge fields with magnetic fluxes

$$\mathbf{n}_1 + \mathbf{n}_2 = \frac{n_+ + \sigma n_-}{2n_+ n_-} \,, \tag{5.14}$$

for $\sigma = -1$. However, it is straightforward to carry out the extremization leaving the twist unspecified. Picking the plus gluing sign in (5.9) gives

$$F^{+}(\varphi_{i},\epsilon;\mathfrak{n}_{i}) = -\frac{2\sqrt{2}\pi N^{3/2}}{3} \left(\mathfrak{n}_{1}\mathfrak{n}_{2}\epsilon + \frac{\varphi_{1}\varphi_{2}}{\epsilon}\right), \qquad (5.15)$$

with φ_i , ϵ satisfying the constraint

$$\varphi_1 + \varphi_2 - \frac{n_+ - \sigma n_-}{2n_+ n_-} \epsilon = 1.$$
(5.16)

For $\sigma = -1$ this is exactly the entropy function proposed in [12], in the case of vanishing electric charges and angular momentum. Performing the extremization of (5.15), subject to (5.16), we get

$$\varphi_1^* = \frac{-n_+ + \sigma n_- + \sqrt{16n_-^2 n_+^2 \mathfrak{n}_1 \mathfrak{n}_2 + (n_+ - n_- \sigma)^2}}{2\sqrt{16n_-^2 n_+^2 \mathfrak{n}_1 \mathfrak{n}_2 + (n_+ - n_- \sigma)^2}},$$

$$\epsilon^* = -\frac{2n_- n_+}{\sqrt{16n_-^2 n_+^2 \mathfrak{n}_1 \mathfrak{n}_2 + (n_+ - \sigma n_-)^2}},$$
(5.17)

and inserting these values back in (5.15) we find

$$F^{+}(\varphi_{i}^{*}, \epsilon^{*}; \mathfrak{n}_{i}) = \frac{\sqrt{2\pi}N^{3/2}}{3} \frac{-n_{+} + \sigma n_{-} + \sqrt{16n_{+}^{2}n_{-}^{2}\mathfrak{n}_{1}\mathfrak{n}_{2} + (n_{+} - \sigma n_{-})^{2}}}{2n_{+}n_{-}}, \qquad (5.18)$$

which coincides with the entropy in [12] upon setting $\sigma = -1$. In minimal D = 4 gauged supergravity the entropy function (5.15), for $\varphi_1 = \varphi_2$ and $\mathfrak{n}_1 = \mathfrak{n}_2$, was derived in [19],

and its plus gluing sign is consistent with the fact that it indeed arises as the sum of contributions from the north and south poles of the spindle, which are the fixed points of the canonical Killing vector field [25].

The precise map between the variables here and those used in [12] is as follows

$$P_i^{\text{there}} = 2\mathfrak{n}_i, \qquad Q^{\text{there}} = 0, \qquad \varphi_i^{\text{there}} = \pm \frac{\pi i}{2} \varphi_i^{\text{here}}, \qquad \omega^{\text{there}} = \pm 2\pi i \epsilon, \qquad (5.19)$$

and using this, the constraint (5.16) becomes

$$2(\varphi_1^{\text{there}} + \varphi_2^{\text{there}}) - \frac{\chi}{4}\omega^{\text{there}} = \pm i\pi.$$
(5.20)

Note that with vanishing angular momentum and electric charges the entropy function is purely real (or purely imaginary) and therefore the critical points are purely real.

For the other type of gluing we obtain (for either choice of σ) the function

$$F^{-}(\varphi_i,\epsilon;\mathfrak{n}_i) = -\frac{2\sqrt{2}\pi N^{3/2}}{3}\left(\mathfrak{n}_2\varphi_1 + \mathfrak{n}_1\varphi_2\right), \qquad (5.21)$$

whose extremization, however, leads to a degenerate result.

The entropy function for the general four-charge model, for either type of gluing and either type of twist, reads

$$F^{\pm}(\varphi_{i},\epsilon;\mathfrak{n}_{i}) = -\frac{\sqrt{2\pi}N^{3/2}}{3\epsilon} \left(\sqrt{(\varphi_{1}+\mathfrak{n}_{1}\epsilon)(\varphi_{2}+\mathfrak{n}_{2}\epsilon)(\varphi_{3}+\mathfrak{n}_{3}\epsilon)(\varphi_{4}+\mathfrak{n}_{4}\epsilon)} \pm \sqrt{(\varphi_{1}-\mathfrak{n}_{1}\epsilon)(\varphi_{2}-\mathfrak{n}_{2}\epsilon)(\varphi_{3}-\mathfrak{n}_{3}\epsilon)(\varphi_{4}-\mathfrak{n}_{4}\epsilon)}\right).$$
(5.22)

The magnetic fluxes obey

$$\mathbf{n}_1 + \mathbf{n}_2 + \mathbf{n}_3 + \mathbf{n}_4 = \frac{n_+ + \sigma n_-}{n_+ n_-}, \qquad (5.23)$$

and the variables φ_i, ϵ satisfy the constraint

$$\varphi_1 + \varphi_2 + \varphi_3 + \varphi_4 - \frac{n_+ - \sigma n_-}{n_+ n_-} \epsilon = 2.$$
 (5.24)

For $n_+ = n_- = 1$ these reduce to the entropy functions proposed in [21]. In particular, for $\sigma = +1$, F^- reduces to the entropy function of the supersymmetric AdS₄ black holes with a topological twist [4], whereas for $\sigma = -1$, F^+ reduces to the entropy function for the supersymmetric rotating Kerr-Newmann AdS₄ black holes [24]. More generally, it is natural to expect that the correct gluing for either type of twists is $-\sigma$, as reported in Table 2. It would be nice to corroborate this proposal showing that extremizing the entropy functions $F^{-\sigma}$ in (5.22) reproduces the entropy of the AdS₂ × Σ solutions of STU gauged supergravity.

5.2 D3-branes

Supergravity solutions describing D3-branes wrapped on the spindle were first constructed in [8] in minimal D = 5 gauged supergravity and generalised to $U(1)^3$ gauged supergravity in [10, 11]. They realise the anti-twist, $\sigma = -1$. The corresponding dual field theory is $\mathcal{N} = 4$ SYM compactified on the spindle, with three background gauge fields with magnetic fluxes satisfying

$$\mathfrak{n}_1 + \mathfrak{n}_2 + \mathfrak{n}_3 = \frac{n_+ + \sigma n_-}{n_+ n_-}, \qquad (5.25)$$

for $\sigma = -1$. However, it is straightforward to carry out the extremization leaving the twist unspecified. In this case we find that, for *both* types of twist, the correct gluing sign that reproduces all known results is the lower sign in (5.9), which gives

$$F^{-}(\varphi_{i},\epsilon;\mathfrak{n}_{i}) = -3N^{2}\left(\mathfrak{n}_{1}\varphi_{2}\varphi_{3} + \varphi_{1}\mathfrak{n}_{2}\varphi_{3} + \varphi_{1}\varphi_{2}\mathfrak{n}_{3} + \mathfrak{n}_{1}\mathfrak{n}_{2}\mathfrak{n}_{3}\epsilon^{2}\right), \qquad (5.26)$$

with φ_i , ϵ satisfying the constraint

$$\varphi_1 + \varphi_2 + \varphi_3 - \frac{n_+ - \sigma n_-}{n_+ n_-} \epsilon = 2.$$
 (5.27)

Extremizing (5.26) subject to the constraint (5.27) we find

$$\varphi_{1}^{*} = \frac{\mathfrak{n}_{1} (\mathfrak{n}_{1} - \mathfrak{n}_{2} - \mathfrak{n}_{3})}{2\left(\frac{\sigma}{n_{+}n_{-}} - (\mathfrak{n}_{1}\mathfrak{n}_{2} + \mathfrak{n}_{1}\mathfrak{n}_{3} + \mathfrak{n}_{2}\mathfrak{n}_{3})\right)},$$

$$\varphi_{2}^{*} = \frac{\mathfrak{n}_{2} (\mathfrak{n}_{2} - \mathfrak{n}_{3} - \mathfrak{n}_{1})}{2\left(\frac{\sigma}{n_{+}n_{-}} - (\mathfrak{n}_{1}\mathfrak{n}_{2} + \mathfrak{n}_{1}\mathfrak{n}_{3} + \mathfrak{n}_{2}\mathfrak{n}_{3})\right)},$$

$$\epsilon^{*} = \frac{\frac{n_{+} - \sigma n_{-}}{n_{+}n_{-}}}{2\left(\frac{\sigma}{n_{+}n_{-}} - (\mathfrak{n}_{1}\mathfrak{n}_{2} + \mathfrak{n}_{1}\mathfrak{n}_{3} + \mathfrak{n}_{2}\mathfrak{n}_{3})\right)},$$

(5.28)

and inserting these back in (5.26) we get

$$F^{-}(\varphi_{i}^{*}, \epsilon^{*}; \mathfrak{n}_{i}) = \frac{3N^{2}\mathfrak{n}_{1}\mathfrak{n}_{2}\mathfrak{n}_{3}}{\frac{\sigma}{n_{+}n_{-}} - (\mathfrak{n}_{1}\mathfrak{n}_{2} + \mathfrak{n}_{1}\mathfrak{n}_{3} + \mathfrak{n}_{2}\mathfrak{n}_{3})}.$$
(5.29)

Setting $\sigma = -1$ this reduces to the a_2 central charge obtained in [10, 11]. On the other hand, setting $\sigma = +1$ we reproduce also the central charge for D3-branes wrapped on the spindle with the twist, for which supergravity solutions were recently presented in [26]. Further setting $n_+ = n_- = 1$, (5.29) reduces to the result for D3-branes wrapped on S^2 with the standard topological twist [5]. Equivalently, we can extremize over the variables Δ_i , ϵ , with $\Delta_1 + \Delta_2 + \Delta_3 = 2$ obtaining the same result as in (5.29) for the critical central charge. However, the variables Δ_i are affected by the ambiguity related to the choice of the constants r_i .

Notice that for the other type of gluing we obtain (for either choice of σ) the function

$$F^{+}(\varphi_{i},\epsilon;\mathfrak{n}_{i}) = -3N^{2}\left[\left(\mathfrak{n}_{1}\varphi_{2}\mathfrak{n}_{3}+\varphi_{1}\mathfrak{n}_{2}\mathfrak{n}_{3}+\mathfrak{n}_{1}\mathfrak{n}_{2}\varphi_{3}\right)\epsilon + \frac{\varphi_{1}\varphi_{2}\varphi_{3}}{\epsilon}\right].$$
 (5.30)

It may be possible that this corresponds to a different type of twisted compactification of D3-branes on the spindle with a corresponding class of supergravity constructions.

5.3 D4-branes

Let us return to the D4-branes and write down the main ingredients involved in considering simultaneously the case of twist and anti-twist. The magnetic fluxes satisfy

$$\mathbf{n}_1 + \mathbf{n}_2 = \frac{n_+ + \sigma n_-}{n_+ n_-} \,. \tag{5.31}$$

In terms of the variables φ_i, ϵ satisfying the constraint

$$\varphi_1 + \varphi_2 - \frac{n_+ - \sigma n_-}{n_+ n_-} \epsilon = 2, \qquad (5.32)$$

the off-shell free energies read

$$F^{\pm}(\varphi_i,\epsilon;\mathfrak{n}_i) = -\frac{4\sqrt{2}\pi}{15\epsilon} \frac{N^{5/2}}{\sqrt{8-N_f}} \Big[\big((\varphi_1+\mathfrak{n}_1\epsilon)(\varphi_2+\mathfrak{n}_2\epsilon)\big)^{3/2} \pm \big((\varphi_1-\mathfrak{n}_1\epsilon)(\varphi_2-\mathfrak{n}_2\epsilon)\big)^{3/2} \Big] \,.$$
(5.33)

As for the M2-branes, we expect that the correct gluing for either type of twists is $-\sigma$, as reported in Table 2. Extremizing $F^{-}(\varphi_i, \epsilon; \mathfrak{n}_i)$, subject to the constraint (5.32), we find (leaving σ unspecified)

$$\varphi_1^* = 1 + \frac{2x[(n_+ - \sigma n_-)x + n_+ n_-(\mathfrak{n}_1 - \mathfrak{n}_2)]}{(x^2 + 3)[(n_+ - \sigma n_-) - (n_+ + \sigma n_-)x]},$$

$$\epsilon^* = \frac{4n_+ n_- x^2}{(x^2 + 3)[(n_+ - \sigma n_-) - (n_+ + \sigma n_-)x]},$$
(5.34)

where x is the only solution in the interval (0, 1) of the quartic equation

$$(n_{+} + \sigma n_{-})^{2} x^{4} + 4 [2n_{+}^{2}n_{-}^{2}(\mathfrak{n}_{1} - \mathfrak{n}_{2})^{2} - 3(n_{+}^{2} + n_{-}^{2} - \sigma n_{+}n_{-})]x^{2} + 12(n_{+}^{2} - n_{-}^{2})x - 9(n_{+} - \sigma n_{-})^{2} = 0.$$

$$(5.35)$$

Inserting the values of φ_i^* and ϵ^* back in (5.33) we get

$$F^{-}(\varphi_{i}^{*}, \epsilon^{*}; \mathfrak{n}_{i}) = \frac{\sqrt{3}\pi}{5} \frac{N^{5/2}}{\sqrt{8 - N_{f}}} \frac{[3(n_{+} - \sigma n_{-})(x^{2} + 1) - (n_{+} + \sigma n_{-})x(x^{2} + 5)]^{3/2}}{n_{+}n_{-}x(x^{2} + 3)[(n_{+} - \sigma n_{-}) - (n_{+} + \sigma n_{-})x]^{1/2}},$$
(5.36)

which reduces to the results of the previous section for $\sigma = +1$.

We expect the choice of minus gluing sign to be correlated to the fact that in our solutions the Killing spinors on the spindle have the same chirality at the north and south poles. It would be interesting to investigate the extremization of F^+ and to find out whether for $\sigma = -1$ it has a physical critical point, corresponding to $AdS_4 \times \Sigma$ solutions with the anti-twist, yet to be constructed.

5.4 M5-branes

Supergravity solutions describing M5-branes wrapped on the spindle were constructed in [9] in a D = 7, $U(1)^2$ gauged supergravity model. They realise the twist, $\sigma = +1$. The corresponding dual field theory is the d = 6, (2,0) SCFT compactified on the spindle, with two background gauge fields with magnetic fluxes

$$\mathfrak{n}_1 + \mathfrak{n}_2 = \frac{n_+ + \sigma n_-}{n_+ n_-} \,, \tag{5.37}$$

for $\sigma = +1$. However, it is straightforward to carry out the extremization for either type of twist. In this case we find that the correct gluing sign that reproduces the results of [9] is the lower sign in (5.9), which gives

$$F^{-}(\varphi_{i},\epsilon;\mathfrak{n}_{i}) = -\frac{9}{64}N^{3}\left(\mathfrak{n}_{2}\varphi_{1}+\mathfrak{n}_{1}\varphi_{2}\right)\left(\mathfrak{n}_{1}\mathfrak{n}_{2}\epsilon^{2}+\varphi_{1}\varphi_{2}\right),\qquad(5.38)$$

with φ_i, ϵ satisfying the constraint

$$\varphi_1 + \varphi_2 - \frac{n_+ - \sigma n_-}{n_+ n_-} \epsilon = 2.$$
(5.39)

Performing the extremization of (5.38), subject to the constraint (5.39), we find

$$\varphi_{1}^{*} = 1 + \frac{(\mathbf{s} + \mathbf{n}_{1} + \mathbf{n}_{2})[2n_{+}^{2}n_{-}^{2}(\mathbf{n}_{1}^{2} - \mathbf{n}_{2}^{2}) + 3(n_{+} - \sigma n_{-})^{2} + n_{+}^{2}n_{-}^{2}(\mathbf{n}_{1} - \mathbf{n}_{2})\mathbf{s}]}{12n_{+}n_{-}(\sigma - n_{+}n_{-}\mathbf{n}_{1}\mathbf{n}_{2})[\mathbf{s} + 2(\mathbf{n}_{1} + \mathbf{n}_{2})]},$$

$$\epsilon^{*} = \frac{(n_{+} - \sigma n_{-})(\mathbf{s} + \mathbf{n}_{1} + \mathbf{n}_{2})}{2(\sigma - n_{+}n_{-}\mathbf{n}_{1}\mathbf{n}_{2})[\mathbf{s} + 2(\mathbf{n}_{1} + \mathbf{n}_{2})]},$$
(5.40)

where

$$\mathbf{s} \equiv \sqrt{7(\mathbf{n}_1^2 + \mathbf{n}_2^2) + 2\mathbf{n}_1\mathbf{n}_2 - 6\frac{n_+^2 + n_-^2}{n_+^2n_-^2}},$$
(5.41)

and inserting these values back in (5.38) we get

$$F^{-}(\varphi_{i}^{*}, \epsilon^{*}; \mathfrak{n}_{i}) = \frac{3N^{3}}{8} \frac{\mathfrak{n}_{1}^{2}\mathfrak{n}_{2}^{2}(\mathfrak{s} + \mathfrak{n}_{1} + \mathfrak{n}_{2})}{(\frac{\sigma}{n_{+}n_{-}} - \mathfrak{n}_{1}\mathfrak{n}_{2})[\mathfrak{s} + 2(\mathfrak{n}_{1} + \mathfrak{n}_{2})]^{2}}.$$
(5.42)

Setting $\sigma = +1$ this reduces to the a_4 central charge obtained in [9] integrating the M5brane anomaly polynomial on the spindle. However, in [9] the extremization was carried out over the variables Δ_i , subject to $\Delta_1 + \Delta_2 = 2$, and it was found that the critical values are given by $\Delta_1^* = \Delta_2^* = 1$, while the critical ϵ^* coincides with the one given in (5.40), up to a convention-dependent factor, specifically, $\epsilon_{\text{here}}^* = -\frac{1}{2}\epsilon_{\text{there}}^*$. One can check that the variables Δ_i utilised in [9] correspond (for $\sigma = +1$) to the choice of constants $r_1 = 2 - r_2$ given by

$$r_1 = 1 + \frac{n_+^2 n_-^2 (\mathfrak{n}_1 - \mathfrak{n}_2) [\mathbf{s} + 2(\mathfrak{n}_1 + \mathfrak{n}_2)]}{3(n_+ - \sigma n_-)^2} \,. \tag{5.43}$$

It would be interesting to find out whether the critical points for $\sigma = -1$ correspond to $AdS_5 \times \Sigma$ solutions with the anti-twist, yet to be constructed.

We note that, as for the case of $AdS_4 \times \Sigma_g$ solutions, the extremization of the offshell free energy (5.38) reproduces also the central charge¹⁵ of d = 4 SCFTs dual to the

¹⁵For g = 0, (5.38) can also be viewed as the off-shell free energy on $S^3 \times S_{\epsilon}^2$ of $\mathcal{N} = 2$ SYM in d = 5. Using this, the form (5.38) has been recently proved in [42] for $S_b^3 \times S_{\epsilon}^2$.

 $AdS_5 \times \Sigma_g$ solutions [6]. As anticipated in footnote 1, in this case the variables φ_i and the magnetic fluxes are subject to the constraints

$$\varphi_1 + \varphi_2 = 2, \qquad \mathbf{n}_1 + \mathbf{n}_2 = 2(1 - g), \qquad (5.44)$$

and as usual the latter may be parameterised as

$$\mathbf{n}_1 = (1 - g) (1 + \mathbf{z} \kappa) , \qquad \mathbf{n}_2 = (1 - g) (1 - \mathbf{z} \kappa) .$$
 (5.45)

The free energy (5.38) is extremized by

$$\varphi_1^* = 1 + \frac{\kappa + \sqrt{\kappa^2 + 3z^2}}{3z}, \qquad \epsilon^* = 0,$$
(5.46)

to which corresponds the critical value [6]

$$F^{-}(\varphi_{i}^{*}, 0; \mathfrak{n}_{i}) = (1 - g)N^{3} \frac{\kappa^{2} - 9z^{2} + \kappa(\kappa^{2} + 3z^{2})^{3/2}}{48z^{2}}.$$
 (5.47)

Since $\epsilon^* = 0$, we could have started setting $\epsilon = 0$ in (5.38), thus reducing to the known ϵ -independent off-shell free energy

$$\lim_{\epsilon \to 0} F^{-}(\varphi_{i}, \epsilon; \mathfrak{n}_{i}) = 2 \sum_{j=1}^{2} \mathfrak{n}_{j} \frac{\partial \mathcal{F}_{6}(\varphi_{i})}{\partial \varphi_{j}} = -\frac{9N^{3}}{64} \varphi_{1} \varphi_{2} \left(\mathfrak{n}_{2} \varphi_{1} + \mathfrak{n}_{1} \varphi_{2}\right) , \qquad (5.48)$$

corresponding to the standard topological twist. In particular, extremizing (5.48) reproduces (5.47). For g > 1 we can set z = 0 so that $\varphi_1^* = \varphi_2^* = 1$ and the free energy (5.48) reduces to the universal relation [44]

$$F^{-}(\varphi_{i}^{*}, 0; \mathfrak{n}_{1} = \mathfrak{n}_{2} = 1 - g) = \frac{9N^{3}}{32}(g - 1) = \frac{63}{512}(g - 1)a_{6}.$$
 (5.49)

Notice that for the other type of gluing we obtain (for either choice of σ) the function

$$F^{+}(\varphi_{i},\epsilon;\mathfrak{n}_{i}) = -\frac{9}{128}N^{3}\left(\mathfrak{n}_{1}^{2}\mathfrak{n}_{2}^{2}\epsilon^{3} + \left(\mathfrak{n}_{1}^{2}\varphi_{2}^{2} + \mathfrak{n}_{2}^{2}\varphi_{1}^{2} + 4\mathfrak{n}_{1}\mathfrak{n}_{2}\varphi_{1}\varphi_{2}\right)\epsilon + \frac{\varphi_{1}^{2}\varphi_{2}^{2}}{\epsilon}\right).$$
(5.50)

It may be possible that this corresponds to a different type of twisted compactification of M5-branes on the spindle with a corresponding class of supergravity constructions.

6 D4-branes wrapped on $\mathbb{Z} \times \Sigma_{g}$

In this section we will begin investigating constructions corresponding to D4-branes wrapped on orbifolds of dimension higher than two, focussing on a class of explicit solutions that can be easily obtained uplifting to massive type IIA supergravity the AdS₂ solutions of four-dimensional minimal gauged supergravity of [7]. Generically, for fourdimensional orbifolds \mathbb{M}_4 , the field theories are d = 1 SCQMs obtained from twisted compactifications of the d = 5, $\mathcal{N} = 1 USp(2N)$ gauge theory. The off-shell free energies are in this case entropy functions, whose extremization determines the entropy of supersymmetric AdS_6 black holes with $AdS_2 \times M_4$ near-horizon geometry. As we have discussed, we expect that these will take the form of a sum of gravitational blocks over the set of fixed points of the canonical Killing vector, which includes also the non-orbifold geometries as special cases. In particular, the entropy function for the product of two constant-curvature Riemann surfaces $\Sigma_{g_1} \times \Sigma_{g_2}$ with the standard topological twist may be recovered considering¹⁶

$$S = -\frac{1}{4\epsilon_1\epsilon_2} \Big[\mathcal{F}_5(\varphi_i + \mathfrak{n}_i\epsilon_1 + \mathfrak{s}_i\epsilon_2) - \mathcal{F}_5(\varphi_i - \mathfrak{n}_i\epsilon_1 + \mathfrak{s}_i\epsilon_2) \\ - \mathcal{F}_5(\varphi_i + \mathfrak{n}_i\epsilon_1 - \mathfrak{s}_i\epsilon_2) + \mathcal{F}_5(\varphi_i - \mathfrak{n}_i\epsilon_1 - \mathfrak{s}_i\epsilon_2) \Big],$$
(6.1)

subject to the constraints

$$\mathfrak{n}_1 + \mathfrak{n}_2 = 2(1 - g_1), \qquad \mathfrak{s}_1 + \mathfrak{s}_2 = 2(1 - g_2), \qquad \varphi_1 + \varphi_2 = 2.$$
(6.2)

Extremizing (6.1) with respect to ϵ_1 and ϵ_2 sets $\epsilon_1 = \epsilon_2 = 0$ so that the entropy function reduces to [43]

$$\lim_{\epsilon_1,\epsilon_2\to 0} S(\varphi_i,\epsilon_1,\epsilon_2;\mathfrak{n}_i,\mathfrak{s}_i) = -\sum_{j,k=1}^2 \mathfrak{n}_j \mathfrak{s}_k \frac{\partial^2 \mathcal{F}_5(\varphi_i)}{\partial \varphi_j \partial \varphi_k} = \frac{4\sqrt{2\pi}}{15} \frac{N^{5/2}}{\sqrt{8-N_f}} \sum_{j,k=1}^2 \mathfrak{n}_j \mathfrak{s}_k \frac{\partial^2 (\varphi_1 \varphi_2)^{3/2}}{\partial \varphi_j \partial \varphi_k}.$$
(6.3)

Extremizing this with respect to φ_i reproduces the entropy of an associated class of $\operatorname{AdS}_2 \times \Sigma_{g_1} \times \Sigma_{g_2}$ supersymmetric solutions [31]. Below we will discuss the case of $\mathbb{M}_4 = \mathbb{Z} \times \Sigma_g$, leaving $\mathbb{M}_4 = \mathbb{Z}_1 \times \mathbb{Z}_2$ and more general orbifolds for future work. Analogous solutions, corresponding to M5-branes wrapped on $\mathbb{Z} \times \Sigma_g$, were presented in [11].

6.1 $\operatorname{AdS}_2 \times \Sigma \times \Sigma_g$ solutions

A class of supersymmetric $\operatorname{AdS}_2 \times \Sigma \times \Sigma_g$ backgrounds may be easily obtained by lifting the $\operatorname{AdS}_2 \times \Sigma$ solutions of four-dimensional minimal gauged supergravity constructed in [7] to D = 6 matter-coupled gauged supergravity, using the consistent truncation presented in [46]. In our conventions, the bosonic part of the action reads

$$S_{6D} = \frac{1}{16\pi G_{(6)}} \int d^6 x \sqrt{-g} \left[R - V - \frac{1}{2} |d\vec{\varphi}|^2 - \frac{1}{2} \sum_{i=1}^2 X_i^{-2} |F_i|^2 - \frac{1}{8} (X_1 X_2)^2 |H|^2 - \frac{m^2}{4} (X_1 X_2)^{-1} |B|^2 - \frac{1}{16} \frac{\varepsilon^{\mu\nu\rho\sigma\tau\lambda}}{\sqrt{-g}} B_{\mu\nu} \left(F_{1\,\rho\sigma} F_{2\,\tau\lambda} + \frac{m^2}{12} B_{\rho\sigma} B_{\tau\lambda} \right) \right],$$
(6.4)

where $F_i = dA_i$, H = dB and the scalar fields X_i with the scalar potential V are given by (2.2) and (2.3), respectively. This model is the complete version of the truncation

¹⁶The overall factor of $-\frac{1}{4}$ may be fixed by splitting the compactification on $\Sigma_{g_1} \times \Sigma_{g_2}$ in two steps, first reducing from d = 5 to d = 3 and then from d = 3 to d = 1. This factor is then consistent with the rules summarised in Table 2.

presented in section 2.1, with non-vanishing two-form B. Below for simplicity we will restrict our attention to the static solution, which in the notation of [7] corresponds to setting j = 0. The six-dimensional solution then reads

$$ds_{6}^{2} = e^{-2C} L_{AdS_{4}}^{2} \left[\frac{y^{2}}{4} ds_{AdS_{2}}^{2} + \frac{y^{2}}{q(y)} dy^{2} + \frac{q(y)}{4y^{2}} dz^{2} \right] + e^{2C} ds_{\Sigma_{g}}^{2} ,$$

$$X_{1} = k_{8}^{1/8} k_{2}^{1/2} , \qquad X_{2} = k_{8}^{1/8} k_{2}^{-1/2} ,$$

$$B = \mathbf{a} \frac{9k_{8}^{1/2}}{8g^{2}} \operatorname{vol}(AdS_{2}) , \qquad (6.5)$$

$$F_{1} = \frac{\mathbf{a}}{y^{2}} \frac{3k_{8}^{1/2} k_{2}^{1/2}}{4g} dy \wedge dz + \frac{\kappa + \mathbf{z}}{2g} \operatorname{vol}(\Sigma_{g}) ,$$

$$F_{2} = \frac{\mathbf{a}}{y^{2}} \frac{3k_{8}^{1/2} k_{2}^{-1/2}}{4g} dy \wedge dz + \frac{\kappa - \mathbf{z}}{2g} \operatorname{vol}(\Sigma_{g}) ,$$

where the function q(y) and the constant **a** are given by¹⁷

$$q(y) = y^4 - (2y - a)^2$$
, $a = \frac{n_+^2 - n_-^2}{n_+^2 + n_-^2}$. (6.6)

The constants k_2 and k_8 are those appearing in (2.31), namely

$$k_2 = \frac{3\mathbf{z} + \sqrt{\kappa^2 + 8\mathbf{z}^2}}{\mathbf{z} - \kappa}, \qquad k_8 = \frac{16k_2}{9(1+k_2)^2}, \tag{6.7}$$

while C reads

$$e^{-2C} = m^2 k_8^{1/4} k_4$$
 with $k_4 = \frac{18}{-3\kappa + \sqrt{\kappa^2 + 8z^2}}$. (6.8)

As usual, the parameter g is fixed in terms of m, while L_{AdS_4} is related to m by eq. (4.24) of [46],

$$g = \frac{3m}{2}, \qquad L_{\text{AdS}_4} = \frac{k_8^{1/4} k_4^{-1/2}}{m^2}.$$
 (6.9)

Notice that formally taking $\mathbf{a} = 0$ the solution (6.5) reduces precisely to the $AdS_4 \times \Sigma_g$ solution discussed in section 2.4.

Let us now consider the quantization of the fluxes. For the fluxes through the Riemann surface we have

$$\mathfrak{s}_{1} = \frac{g}{2\pi} \int_{\Sigma_{g}} F_{1} = (1 + \mathbf{z} \kappa) (1 - g) \in \mathbb{Z},$$

$$\mathfrak{s}_{2} = \frac{g}{2\pi} \int_{\Sigma_{g}} F_{2} = (1 - \mathbf{z} \kappa) (1 - g) \in \mathbb{Z},$$
(6.10)

with $\mathfrak{s}_1 + \mathfrak{s}_2 = 2(1 - g)$. Recalling that [7]

$$\left(\frac{1}{y_1} - \frac{1}{y_2}\right)\frac{\Delta z}{2\pi} = \frac{n_+^2 + n_-^2}{n_+ n_- (n_+ + n_-)},\tag{6.11}$$

¹⁷In order to be consistent with our notation we have exchanged n_+ and n_- with respect to [7].

where y_1, y_2 are the two relevant roots of q(y), the fluxes through the spindle are given by

$$\mathfrak{n}_{1} = \frac{g}{2\pi} \int_{\Sigma} F_{1} = \frac{3}{4} k_{8}^{1/2} k_{2}^{1/2} \frac{n_{+} - n_{-}}{n_{+} n_{-}} = \frac{p_{1}}{n_{+} n_{-}},$$

$$\mathfrak{n}_{2} = \frac{g}{2\pi} \int_{\Sigma} F_{2} = \frac{3}{4} k_{8}^{1/2} k_{2}^{-1/2} \frac{n_{+} - n_{-}}{n_{+} n_{-}} = \frac{p_{2}}{n_{+} n_{-}},$$
(6.12)

where $p_i \in \mathbb{Z}$. We then note that

$$\frac{3}{2}k_8^{1/2}k_2^{1/2} = \Delta_1^*, \qquad \frac{3}{2}k_8^{1/2}k_2^{-1/2} = 2 - \Delta_1^* \equiv \Delta_2^*, \tag{6.13}$$

where Δ_1^* was given in (4.11), thus we have

$$\mathfrak{n}_{i} = \Delta_{i}^{*} \frac{n_{+} - n_{-}}{2n_{+}n_{-}} \equiv n \,\Delta_{i}^{*} \,, \tag{6.14}$$

that will be important in the field theory extremization. The constraint $\Delta_1^* + \Delta_2^* = 2$ implies

$$\mathfrak{n}_1 + \mathfrak{n}_2 = \frac{n_+ - n_-}{n_+ n_-}, \qquad (6.15)$$

showing that there is an anti-twist over the spindle. Note that from the relations (6.14) we obtain

$$\left(1 + \frac{\kappa + \sqrt{8z^2 + \kappa^2}}{4z}\right)(n_+ - n_-) = 2p_1, \qquad (6.16)$$

which is a non-trivial Diophantine equation. One example of solution is given by the set of values $\kappa = -1$, $\mathbf{z} = 6$, $n_+ - n_- = 6$, $p_1 = 5$ and we have checked that other combinations exist. Moreover, when $\kappa = -1$ it is possible to smoothly take the limit $\mathbf{z} \to 0$, in which case the equation reduces to $n_+ - n_- = 2p_1$.

We can now uplift the solution (6.5) to massive type IIA using the recipe described in section 2.2. For simplicity, we will only present the relevant ingredients for the computation of the entropy, namely the metric and the dilaton, that read

$$ds_{\text{s.f.}}^{2} = \lambda^{2} \mu_{0}^{-1/3} \left\{ \tilde{\Delta}^{1/2} \left[e^{-2C} L_{\text{AdS}_{4}}^{2} \left(\frac{y^{2}}{4} ds_{\text{AdS}_{2}}^{2} + \frac{y^{2}}{q(y)} dy^{2} + \frac{q(y)}{4y^{2}} dz^{2} \right) + e^{2C} ds_{\Sigma_{g}}^{2} \right] + g^{-2} \tilde{\Delta}^{-1/2} k_{8}^{-1/4} \left[k_{8}^{1/2} d\mu_{0}^{2} + k_{2}^{-1/2} \left(d\mu_{1}^{2} + \mu_{1}^{2} \sigma_{1}^{2} \right) + k_{2}^{1/2} \left(d\mu_{2}^{2} + \mu_{2}^{2} \sigma_{2}^{2} \right) \right] \right\},$$

$$e^{\Phi} = \lambda^{2} \mu_{0}^{-5/6} \tilde{\Delta}^{1/4} k_{8}^{-1/8}, \qquad (6.18)$$

with $\sigma_i = \mathrm{d}\phi_i - gA_i$ and $\tilde{\Delta}$ given in (2.38). Expressing the metric in string frame as

$$ds_{s.f.}^2 = e^{2A} \left(ds_{AdS_2}^2 + ds_{M_8}^2 \right), \qquad (6.19)$$

the effective two-dimensional Newton constant is given by

$$\frac{1}{G_{(2)}} = \frac{32\pi^2}{(2\pi\ell_s)^8} \int e^{8A-2\Phi} \operatorname{vol}(M_8), \qquad (6.20)$$

and therefore the entropy reads

$$S = \frac{1}{4G_{(2)}} = \frac{1}{(2\pi\ell)^8} \frac{9(3\pi\lambda)^4 k_8^{1/2}}{20g^8 k_4} 4\pi\kappa(1-g) A_h , \qquad (6.21)$$

where A_h is the area of the horizon of the four-dimensional black hole with $L_{AdS_4} = 1$, specifically [7]

$$A_h = \frac{1}{2}(y_2 - y_1)\Delta z = \pi \frac{-(n_+ + n_-) + \sqrt{2}\sqrt{n_+^2 + n_-^2}}{n_+ n_-}.$$
 (6.22)

Both the Romans mass $F_{(0)}$ and the part of the four-form flux $F_{(4)}$ along the fourhemisphere \mathbb{S}^4 remain unaltered with respect to (2.36) and (2.37), therefore the quantization of the fluxes in ten dimensions is unchanged, giving the relations (2.25). The final expression of the entropy is

$$S = \frac{8\kappa(1-g)N^{5/2}(z^2-\kappa^2)^{3/2}(\sqrt{\kappa^2+8z^2}-\kappa)}{5\sqrt{8-N_f}(\kappa\sqrt{\kappa^2+8z^2}-\kappa^2+4z^2)^{3/2}}A_h.$$
 (6.23)

6.2 Entropy function

As before, we can deduce the main ingredients of the field theory construction from the supergravity solution. In particular, the magnetic fluxes \mathfrak{s}_i show that there is a (standard) topological twist on Σ_g , while from the form of the \mathfrak{n}_i we see that there is an anti-twist on Σ . Then applying our conjecture twice, we obtain the following entropy function

$$S(\varphi_i, \epsilon_1, \epsilon_2; \mathbf{n}_i, \mathbf{s}_i) = -\frac{1}{4\epsilon_1\epsilon_2} \Big[\mathcal{F}_5(\varphi_i + \mathbf{n}_i\epsilon_1 + \mathbf{s}_i\epsilon_2) + \mathcal{F}_5(\varphi_i - \mathbf{n}_i\epsilon_1 + \mathbf{s}_i\epsilon_2) - \mathcal{F}_5(\varphi_i - \mathbf{n}_i\epsilon_1 - \mathbf{s}_i\epsilon_2) - \mathcal{F}_5(\varphi_i - \mathbf{n}_i\epsilon_1 - \mathbf{s}_i\epsilon_2) \Big],$$

$$(6.24)$$

subject to the constraints

$$\mathfrak{n}_1 + \mathfrak{n}_2 = \frac{n_+ - n_-}{n_+ n_-}, \qquad \mathfrak{s}_1 + \mathfrak{s}_2 = 2(1 - g), \qquad \varphi_1 + \varphi_2 - \frac{n_+ + n_-}{n_+ n_-} \epsilon_1 = 2.$$
(6.25)

Extremizing this with respect to ϵ_2 sets $\epsilon_2 = 0$ and after renaming $\epsilon_1 \mapsto \epsilon$ we obtain

$$S(\varphi_i,\epsilon;\mathfrak{n}_i,\mathfrak{s}_i) = \frac{c}{\epsilon} \left[\sqrt{(\varphi_1 + \mathfrak{n}_1\epsilon)(\varphi_2 + \mathfrak{n}_2\epsilon)} (\mathfrak{s}_1(\varphi_2 + \mathfrak{n}_2\epsilon) + \mathfrak{s}_2(\varphi_1 + \mathfrak{n}_1\epsilon)) + (\epsilon \to -\epsilon) \right],$$
(6.26)

where

$$c \equiv \frac{\sqrt{2\pi}}{5} \frac{N^{5/2}}{\sqrt{8 - N_f}},$$
(6.27)

which has to be extremized subject to the constraints (6.25).

So far we have not used the additional input (6.14) given by the supergravity solution discussed above. This strongly suggests that the extremization of (6.26), without imposing

(6.14), will give the entropy of a more general supergravity solution¹⁸. However, below we will proceed enforcing (6.14). Doing so, it is convenient to introduce the rescaled variables

$$\varphi_1 \equiv \Delta_1^* \lambda_1 , \qquad \varphi_2 \equiv \Delta_2^* \lambda_2 , \qquad (6.28)$$

subject to the constraint

$$\Delta_1^* \lambda_1 + \Delta_2^* \lambda_2 - \frac{n_+ + n_-}{n_+ n_-} \epsilon = 2, \qquad (6.29)$$

in terms of which the entropy function reads

$$S(\lambda_i, \epsilon; \mathbf{z}) = \frac{c}{\epsilon} \left[\sqrt{\Delta_1^* \Delta_2^* (\lambda_1 + n\epsilon) (\lambda_2 + n\epsilon)} (\mathfrak{s}_1 \Delta_1^* (\lambda_2 + n\epsilon) + \mathfrak{s}_2 \Delta_2^* (\lambda_1 + n\epsilon)) + (\epsilon \to -\epsilon) \right].$$
(6.30)

Extremizing this we find the critical values

$$\epsilon^* = \eta \frac{2n_+ n_-}{\sqrt{2n_+^2 + 2n_-^2}}, \qquad \lambda_1^* = \lambda_2^* = 1 + \eta \frac{n_+ + n_-}{\sqrt{2n_+^2 + 2n_-^2}}, \tag{6.31}$$

where the sign ambiguity $\eta = \pm 1$ arises by solving the equations over the complex numbers¹⁹. Inserting these values back into the entropy function, we compute

$$S(\lambda_i^*, \epsilon^*; \mathbf{z}) = 2 \frac{\sqrt{2\pi}}{5} \frac{N^{5/2}}{\sqrt{8 - N_f}} \sqrt{\Delta_1^* \Delta_2^*} (\mathfrak{s}_1 \Delta_2^* + \mathfrak{s}_2 \Delta_1^*) \frac{-n_+ - n_- - \eta \sqrt{2n_-^2 + 2n_+^2}}{n_+ n_-} = \frac{8\pi\kappa (1 - \mathbf{g}) N^{5/2} (\mathbf{z}^2 - \kappa^2)^{3/2} (\sqrt{\kappa^2 + 8\mathbf{z}^2} - \kappa)}{5\sqrt{8 - N_f} (\kappa\sqrt{\kappa^2 + 8\mathbf{z}^2} - \kappa^2 + 4\mathbf{z}^2)^{3/2}} \frac{-n_+ - n_- - \eta \sqrt{2n_+^2 + 2n_+^2}}{n_+ n_-},$$
(6.32)

where we need to pick $\eta = -1$ in order to get a positive entropy that agrees with the one computed from the ten-dimensional supergravity solution.

It should be straightforward to incorporate electric charge and rotation (along the spindle), by promoting the extremization of (6.30) to the Legendre transform. The entropy obtained in this way should match that of the supergravity solution obtained by uplifting to massive type IIA the rotating $AdS_2 \times \Sigma$ solution of [7]²⁰.

7 Discussion

The work of [8] opened the way to a novel class of examples of the AdS/CFT correspondence, by constructing a supergravity solution with an AdS₃ factor, comprising the two-dimensional orbifold Σ , known as the spindle. This was interpreted as the nearhorizon limit of D3-branes wrapped on the spindle with the corresponding field theory

¹⁸The recent paper [47] discusses $AdS_2 \times \Sigma \times \Sigma_g$ supergravity solutions and it would be interesting to understand the relationship to our work.

 $^{^{19}}$ Generically, in the presence of rotation, we are forced to work with the complex numbers [19, 20] and we therefore continue to do so also in the static case.

²⁰In the rotating solution, setting $n_{+} = n_{-} = 1$ will reduce to a rotating $AdS_2 \times S_{\epsilon}^2 \times \Sigma_g$ solution in D = 6, whose entropy function has been recently discussed in [42].

duals being a class of d = 4, $\mathcal{N} = 1$ SCFTs compactified on the spindle, with a new type of supersymmetry-preserving twist, that was dubbed anti-twist. Following this, over the past year various extensions, including analogous constructions for M2 and M5-branes, have appeared, realising either the anti-twist [8], or a global generalization [9] of the standard topological twist. These constructions have left out the notable class of solutions describing D4-branes wrapped on the spindle, that we constructed in this paper. We found these $AdS_4 \times \Sigma$ solutions in D = 6 gauged supergravity and then uplifted them to massive type IIA supergravity. The type of twist realised in our solutions is the one previously found in [9] for M5-branes, which was referred to as a global topological twist. Differently from the standard topological twist, the *R*-symmetry gauge field does not cancel the spin connection, despite the fact that its integrated flux is equal to the Euler characteristic of the spindle. As a consequence the Killing spinors are then non-trivial sections (in particular, they are not simply constant) of the same bundles that occur in standard topological twist.

There are several aspects of our solution that may be interesting to investigate in the future. For example, it may be instructive to cast it in the form of the classification of supersymmetric AdS_4 solutions of massive type IIA supergravity [34], or to study supersymmetric probe D-branes in our background in order to extract further information about the dual field theories. Another question that arises from our work is whether there exists a more general consistent truncation of massive type IIA supergravity to D = 6, analogous to the dyonic consistent truncation to D = 4 supergravity found in [32]. It is also intriguing to investigate whether there exist supergravity solutions corresponding to D4-branes wrapped on the spindle with the anti-twist. Finally, as an important step towards improving control on the dual field theories, it would be worthwhile computing an appropriately regularised on-shell action, that should prove the validity of our conjectural off-shell free energy (4.18). To do this, one may need to know the full "black two-brane" solution, interpolating between AdS_6 asymptotically and $AdS_4 \times \Sigma$ in the near-horizon, similarly to [19]. However, it may be possible to employ the strategy of [25] to prove that the supergravity (Euclidean) on-shell action localises at the poles of the spindle, which are the fixed points for the canonical Killing vector field associated to any (Euclidean) supersymmetric solution of the theory [48].

There is also a number of assorted interesting questions in the field theory side. The most direct one is to study the five-dimensional SCFTs in the background of $S^3 \times \Sigma$, computing the localized partition function from first principles, and then showing that in the large N limit the associated free energy reduces to (4.18). This is indeed an open problem for SCFTs compactified on spindles in different dimension.

It is compelling that the five-dimensional off-shell free energy that we conjectured here and the entropy function that was conjectured in [12] fit into a broader conjecture for the free energies (*i.e.* minus the logarithm of partition functions) of SCFTs in various dimensions, namely (5.9). These extend the idea of gravitational blocks put forward in [21] in two directions. Firstly, from compactifications on smooth manifolds, to the realm of compactifications on orbifolds. It is remarkable that compactifications on spindles can be incorporated by a simple modification of the constraint obeyed by the fugacities (see eq. (5.10)). This depends on the type of twist performed and it includes the standard topological twist and the no-twist as special cases. Secondly, we have pointed out that the form (5.9) should hold also for observables of higher-dimensional theories, beyond the entropy, associated to AdS_2 solutions dual d = 1 SCFTs.

Using the AdS/CFT correspondence, the free energies (5.9) are generically expected to arise as gravitational (Euclidean) on-shell actions, and their structure is suggested by the fact that for supersymmetric solutions, this form should arise from summing contributions at fixed points of the canonical Killing vector field, defined as a bilinear in the Killing spinors of the solution [25, 49]. An immediate issue that would be nice to clarify is a better justification of the \pm signs in (5.9) and their relationship with the type of twist. For example, while for field theories in odd dimensions we have proposed that the gluing sign should be $-\sigma$, for field theories in even dimensions, the explicit examples indicate that one should always pick the minus sign. It would interesting to find out whether the functions that have not been considered so far, may have critical points that correspond to new gravitational objects.

It is quite clear that the structure of (5.9) will extend to compactifications of SCFTs on higher-dimensional²¹ orbifolds and in general we expect that the functions to extremize will take the form of a sum of contributions from fixed points of the canonical Killing vector on the compactification orbifold \mathbb{M} . Assuming this is toric, for concreteness, there will be as many equivariant parameters ϵ_I as the rank of the torus acting on M. One of the simplest examples of this type of construction is given by twisted compactifications of the d = 5 SCFT on $\mathbb{M} = \mathbb{Z} \times \Sigma_{g}$, that we discussed in section 6. We have shown that the corresponding entropy function can be obtained iterating our conjecture for the off-shell free energies twice and extremizing this reproduces the entropy of a corresponding class of $AdS_2 \times \Sigma \times \Sigma_g$ supergravity solutions. More generally, we expect that there should exist solutions of the type $\operatorname{AdS}_2 \times \mathbb{Z}_1 \times \mathbb{Z}_2$ in D = 6 supergravity and of the type $\operatorname{AdS}_3 \times \mathbb{Z}_1 \times \mathbb{Z}_2$ in D = 7 supergravity, for which the corresponding entropy function and trial central charge can be obtained following the rules we proposed in this paper. Interestingly, the former case would be identified with the near-horizon limit of a novel class of supersymmetric black holes in AdS_6 , that may be accelerating. Work along these lines is underway and we hope to report in the near future.

As we continue to navigate the landscape of supergravity solutions corresponding to branes wrapping orbifolds, it will be revealing to employ the approach developed in [50–53], adapting it to situations with orbifold singularities. While this is already well developed for solutions arising from M2 and D3-branes, it is tantalising to think that an analogous geometric approach may be concocted for studying AdS_5 solutions of D = 11supergravity or other backgrounds with an AdS factor.

Acknowledgments

The research of FF is supported by the project "Nonlinear Differential Equations" of Università degli Studi di Torino. DM would like to thank J. Gauntlett and J. Sparks for insightful comments and enjoyable collaborations on related topics.

²¹The recent paper [42] discusses related setups, for compactifications of d = 5 and d = 6 SCFTs on various smooth manifolds, including $S_{\epsilon}^2 \times \Sigma_{g}$ and $S_{\epsilon_1}^2 \times S_{\epsilon_2}^2$.

A More details on the $AdS_4 \times \Sigma_g$ solutions

A.1 Relation with the Lagrangian of [1]

Here we make contact between the D = 6 Lagrangian (2.1), that we use in the paper, and the Lagrangian used in [1], given explicitly in [54]. In order to minimise confusion we have relabelled some of the quantities in [54] as follows

$$\varphi_{1,2} \mapsto \hat{\varphi}_{1,2}, \qquad F^{3,6} \mapsto \frac{1}{2} \mathcal{F}^{3,6}, \qquad m \mapsto m_1, \qquad (A.1)$$

where the 1/2 factor is due to the unusual definition of the field strengths as, e.g., $\mathcal{F}_{\mu\nu}^3 = \frac{1}{2}(\partial_{\mu}\mathcal{A}_{\nu}^3 - \partial_{\nu}\mathcal{A}_{\mu}^3)$. Consistency between the equations of motion and the supersymmetry equations requires $\hat{\varphi}_1 = 0$, and in this case the D = 6 Lagrangian in [54] reads

$$e^{-1}\mathcal{L} = \frac{1}{4}R - \hat{V} - |d\sigma|^2 - \frac{1}{4}|d\hat{\varphi}_2|^2 - \frac{1}{16}e^{-2\sigma}\cosh(2\hat{\varphi}_2)|\mathcal{F}^3|^2 - \frac{1}{16}e^{-2\sigma}\cosh(2\hat{\varphi}_2)|\mathcal{F}^6|^2 + \frac{1}{16}e^{-2\sigma}\sinh(2\hat{\varphi}_2)\mathcal{F}^3_{\mu\nu}\mathcal{F}^{6\mu\nu},$$
(A.2)

where the scalar potential is

$$\hat{V} = -g_1^2 e^{2\sigma} - 4g_1 m_1 e^{-2\sigma} \cosh \hat{\varphi}_2 + m_1^2 e^{-6\sigma} .$$
(A.3)

Defining the quantities

$$\vec{\varphi} = -\sqrt{2} \left(\hat{\varphi}_2, 2\sigma \right) \iff X_1 = e^{\sigma + \hat{\varphi}_2}, \quad X_2 = e^{\sigma - \hat{\varphi}_2},$$

$$F_1 = \frac{1}{2} (\mathcal{F}^3 + \mathcal{F}^6), \qquad F_2 = \frac{1}{2} (\mathcal{F}^3 - \mathcal{F}^6), \qquad (A.4)$$

$$g = g_1, \qquad m = 2m_1,$$

we obtain the Lagrangian given in (2.1), divided by 4.

A.2 Equivalence with the solutions of [2]

Below we show that the ten-dimensional background (2.34) - (2.37), obtained from the uplift of the D = 6 solution of [1], is equivalent to the solution to massive type IIA supergravity constructed in [2]. Let us start with the metric (4.14) of [2]

$$ds_{s.f.}^2 = \frac{L_{AdS_4}^2}{(Hy)^{1/2}\hat{F}_0} \left(ds_{AdS_4}^2 + e^{2\nu} ds_{\Sigma_g}^2 + \frac{H}{4} ds_{\mathbb{S}^4}^2 \right),$$
(A.5)

where we restored the dimension-full AdS_4 radius L_{AdS_4} that was set to one in [2] and we renamed the constant F_0 in [2] as \hat{F}_0 . The metric on the squashed hemisphere reads

$$ds_{\mathbb{S}^4}^2 = \frac{8}{3} d\mu_0^2 + 2 \left(d\hat{\mu}_1^2 + \hat{\mu}_1^2 \eta_+^2 + d\hat{\mu}_2^2 + \hat{\mu}_2^2 \eta_-^2 \right),$$
(A.6)

where the coordinates $\mu_0, \hat{\mu}_1, \hat{\mu}_2$ satisfy the constraint $\mu_0^2 + a_+ \hat{\mu}_1^2 + a_- \hat{\mu}_2^2 = 1$, and the one-forms η_{\pm} are $\eta_{\pm} = d\phi_{\pm} - m_{\pm}\omega_g$. The functions H and y are given by

$$H = \frac{2}{3\mu_0^2 + 4\left(a_+^2\hat{\mu}_1^2 + a_-^2\hat{\mu}_2^2\right)}, \qquad y^3\hat{F}_0^2 = \frac{3}{2}\mu_0^2, \qquad (A.7)$$

while $e^{2\nu}$ is a constant and reads

$$e^{2\nu} = \frac{-\kappa + \sqrt{\kappa^2 + 8z^2}}{4}.$$
 (A.8)

The constants a_{\pm} , m_{\pm} can be expressed in terms of κ and a constant parameter z as

$$a_{\pm} = \frac{1 \pm \epsilon}{2}, \qquad \epsilon = \frac{\kappa \pm \sqrt{\kappa^2 + 8\mathbf{z}^2}}{4\mathbf{z}}, \qquad m_{\pm} = \frac{\kappa \pm \mathbf{z}}{2}.$$
 (A.9)

In order to compare the two solutions we take

$$k_2 = \frac{a_+}{a_-}, \qquad k_8 = \frac{16}{9}a_+a_-, \qquad (A.10)$$

which are consistent with the constraint $a_+ + a_- = 1$ required by the definition of a_{\pm} , and relate our coordinates μ_1 , μ_2 to those in [2] as

$$\mu_1 = \sqrt{a_+}\hat{\mu}_1, \qquad \mu_2 = \sqrt{a_-}\hat{\mu}_2.$$
 (A.11)

We then identify $\phi_1 = \phi_+, \ \phi_2 = \phi_-$ and

$$p_1 = 2m_+ = \kappa + z, \qquad p_2 = 2m_- = \kappa - z, \qquad (A.12)$$

thus having $\sigma_1 = \eta_+$, $\sigma_2 = \eta_-$. As consistency checks we have that $2(m_+ + m_-) = p_1 + p_2 = 2\kappa$ and that the definition of k_2 in (2.31) agrees with (A.10). Lastly, comparing the overall factors in the metric and the dilaton of the two solutions, we find

$$g^{2} = \left(\frac{3}{2}\right)^{3/2} k_{8}^{3/4} L_{\text{AdS}_{4}}^{-2}, \qquad \lambda^{2} = \left(\frac{3}{2}\right)^{-1/6} k_{8}^{1/4} \hat{F}_{0}^{-2/3}.$$
(A.13)

The Romans mass (2.36) then reads $F_{(0)} = \frac{\hat{F}_0}{L_{\text{AdS}_4}}$, reducing to the expression in [2] for $L_{\text{AdS}_4} = 1$. Writing our four-form flux (2.37) in the variables used in [2], we find

$$F_{(4)} = \frac{H}{8} \left(\frac{2\mu_0}{3\hat{F}_0} \right)^{1/3} L^3_{\text{AdS}_4} \left\{ W \frac{\hat{\mu}_1 \hat{\mu}_2}{\mu_0} \, \mathrm{d}\hat{\mu}_1 \wedge \mathrm{d}\hat{\mu}_2 \wedge \eta_+ \wedge \eta_- \right. \\ \left. - \operatorname{vol}(\Sigma_{\mathrm{g}}) \wedge \left[\frac{m_+}{a_+} \, \mathrm{d}\phi_- \wedge \left(3\mu_0 \hat{\mu}_2 \, \mathrm{d}\hat{\mu}_2 - 4a_- \hat{\mu}_2^2 \, \mathrm{d}\mu_0 \right) \right. \\ \left. + \frac{m_-}{a_-} \, \mathrm{d}\phi_+ \wedge \left(3\mu_0 \hat{\mu}_1 \, \mathrm{d}\hat{\mu}_1 - 4a_+ \hat{\mu}_1^2 \, \mathrm{d}\mu_0 \right) \right] \right\},$$
(A.14)

where we defined

$$W = 9H\mu_0^2 + 16H(a_+^3\hat{\mu}_1^2 + a_-^3\hat{\mu}_2^2) - 12.$$
 (A.15)

This expression agrees with the $F_{(4)}$ given in [2] only partially²². A couple of useful identities that we used for comparing the two solutions are

$$\frac{m_{+}}{a_{+}} = -2e^{2\nu}[1 - 2(a_{+} - a_{-})], \qquad \frac{m_{-}}{a_{-}} = -2e^{2\nu}[1 - 2(a_{-} - a_{+})].$$
(A.16)

²²While the above $F_{(4)}$ satisfies the equations of motion and the Bianchi identity, we found that the four-form flux given in [2] is not closed.

B OSp(2|4) superalgebra from bilinears

Below we show that the OSp(2|4) superalgebra can be reconstructed from the Killing spinors of our $AdS_4 \times \mathbb{Z}$ solution, as expected. The fact that the Killing vectors of AdS can be constructed as bilinears in the Killing spinors is well known. Here we demonstrate that the D = 6 solution captures part of the *R*-symmetry of the dual d = 3, $\mathcal{N} = 2$ SCFTs. Specifically, it includes the component of the *R*-symmetry along the spindle isometry. The full *R*-symmetry would arise from working with the Killing spinors of the uplifted ten-dimensional solution.

Let us recall the relevant ingredients from section 3.1. The Killing spinors of the D = 6 solution are a pair of symplectic-Majorana spinors, taking the form

$$\epsilon^A = \vartheta_+ \otimes \eta_+^A + \vartheta_- \otimes \eta_-^A, \tag{B.1}$$

where A = 1, 2 is a symplectic index. The spinors ϑ_{\pm} are the chiral components of a Majorana Killing spinor $\vartheta = \vartheta(x^{\hat{\mu}})$ in AdS₄, obeying

$$\vartheta^* = \mathcal{B}_4 \vartheta, \qquad \hat{\nabla}_{\hat{\mu}} \vartheta = \frac{1}{2} \gamma_{\hat{\mu}} \vartheta,$$
(B.2)

thus we have $\vartheta = \vartheta_+ + \vartheta_-$ with $\gamma_5 \vartheta_{\pm} = \pm \vartheta_{\pm}$. The spinors $\eta_{\pm}^A = \eta_{\pm}^A(y)$ are Dirac spinors defined on the spindle. Defining the following vector bilinear

$$\mathbf{K}^{\mu} = \mathbf{i}\,\overline{\epsilon'^{1}}\Gamma^{\mu}\epsilon^{1}\,,\tag{B.3}$$

where ϵ^1 and ϵ'^1 are two Killing spinors, straightforward manipulations lead to

$$\mathbf{K}^{\hat{\mu}} = 2\mathrm{i}\,m(\xi')^*\xi\,\bar{\vartheta}'\gamma^{\hat{\mu}}\vartheta\,,\qquad \mathbf{K}^y = 0\,,\qquad \mathbf{K}^z = 2(\xi')^*\xi\,\bar{\vartheta}'\vartheta\,.\tag{B.4}$$

From (B.2) it immediately follows that

$$\nabla_{(\hat{\mu}}(\bar{\vartheta}'\gamma_{\hat{\nu})}\vartheta) = 0 = \nabla_{\hat{\mu}}(\bar{\vartheta}'\vartheta), \qquad (B.5)$$

hence

$$\mathsf{K}^{\mu}\partial_{\mu} = 2(\xi')^{*}\xi \left(\mathrm{i}\,m\bar{\vartheta}'\gamma^{\hat{\mu}}\vartheta\,\partial_{\hat{\mu}} + \bar{\vartheta}'\vartheta\,\partial_{z}\right) \tag{B.6}$$

are Killing vectors for the D = 6 solution. One can check that $\overline{\epsilon'^2}\Gamma^{\mu}\epsilon^2 = -(\overline{\epsilon'^1}\Gamma^{\mu}\epsilon^1)^*$, while $\overline{\epsilon'^1}\Gamma^{\mu}\epsilon^2$, $\overline{\epsilon'^2}\Gamma^{\mu}\epsilon^1$ and $\overline{\epsilon'^A}\Gamma^7\Gamma^{\mu}\epsilon^B$ are not Killing vectors, so there are no further vector bilinears to consider.

To make contact with the OSp(2|4) superalgebra we need to look in more detail into the structure of the Killing spinors in AdS₄. Writing the unit radius AdS₄ metric as

$$ds_{AdS_4}^2 = u^2 \eta_{mn} dx^m dx^n + \frac{du^2}{u^2},$$
 (B.7)

and defining the coordinates $x^{\hat{\mu}} = (t, x^1, x^2, u) \equiv (x^m, u)$, the Killing spinors on AdS₄ can be written as [55]

$$\epsilon_{\text{AdS}_4} = e^{\frac{1}{2}(\ln u)\,\gamma_3} \left[1 + \frac{1}{2u} x^m \gamma_m (1 - \gamma_3) \right] \psi \,, \tag{B.8}$$

with ψ a constant spinor and $\{\gamma_m, \gamma_n\} = u^2 \eta_{mn}$. This can be conveniently split into two independent Killing spinors as $\epsilon_{AdS_4} = \epsilon_+ + \epsilon_-$, where

$$\epsilon_{+} = u^{1/2}\psi_{+}, \qquad \epsilon_{-} = u^{-1/2} \left(1 + x^{m}\gamma_{m}\right)\psi_{-}, \qquad (B.9)$$

with

$$\psi_{\pm} \equiv \frac{I_4 \pm \gamma_3}{2} \psi \qquad \Longrightarrow \qquad \gamma_3 \psi_{\pm} = \pm \psi_{\pm} \,. \tag{B.10}$$

Note that ϵ_{\pm} and ψ_{\pm} are not chiral spinors, since γ_3 is not the chiral matrix. A Majorana condition can be consistently imposed on ϵ_{AdS_4} , which implies that ψ and, in turn, ψ_{\pm} are Majorana spinors. We can then identify ϑ with ϵ_{AdS_4} and write the Killing vectors in (B.4) as

$$\mathbf{K}^{\hat{\mu}} = 2\mathbf{i} \, m(\xi')^* \xi \left(\vec{\epsilon}'_+ \gamma^{\hat{\mu}} \epsilon_+ + \vec{\epsilon}'_+ \gamma^{\hat{\mu}} \epsilon_- + \vec{\epsilon}'_- \gamma^{\hat{\mu}} \epsilon_+ + \vec{\epsilon}'_- \gamma^{\hat{\mu}} \epsilon_- \right), \\
\mathbf{K}^z = 2(\xi')^* \xi \left(\vec{\epsilon}'_+ \epsilon_+ + \vec{\epsilon}'_+ \epsilon_- + \vec{\epsilon}'_- \epsilon_+ + \vec{\epsilon}'_- \epsilon_- \right).$$
(B.11)

As standard, the two independent ϵ_+ spinors are identified with the real Poincaré supercharges Q and the ϵ_- with the real superconformal supercharges S. For this reason we shall refer to the different contributions in (B.11) as $K^{\hat{\mu}}_{QQ}$, $K^{\hat{\mu}}_{QS}$, $K^{\hat{\mu}}_{SQ}$ and $K^{\hat{\mu}}_{SS}$, respectively, and likewise for K^z . Specifically, the components along the spindle are

$$\begin{aligned}
\mathbf{K}_{QQ}^{z} &= 0, & \mathbf{K}_{QS}^{z} &= 2(\xi')^{*} \xi \, \psi'_{+} \psi_{-}, \\
\mathbf{K}_{SS}^{z} &= 0, & \mathbf{K}_{SQ}^{z} &= 2(\xi')^{*} \xi \, \bar{\psi}'_{-} \psi_{+},
\end{aligned} \tag{B.12}$$

and the components along AdS_4 are

$$\begin{split} \mathbf{K}_{QQ}^{m} &= 2\mathrm{i}\,m(\xi')^{*}\xi\,u\,(\bar{\psi}'_{+}\gamma^{m}\psi_{+})\,, & \mathbf{K}_{QQ}^{u} &= 0\,, \\ \mathbf{K}_{QS}^{m} &= 2\mathrm{i}\,m(\xi')^{*}\xi\,\left[x^{m}(\bar{\psi}'_{+}\psi_{-}) + x^{n}(\bar{\psi}'_{+}\gamma^{m}_{\ n}\psi_{-})\right]\,, & \mathbf{K}_{QS}^{u} &= -2\mathrm{i}\,m(\xi')^{*}\xi\,u\,\bar{\psi}'_{+}\psi_{-}\,, \\ \mathbf{K}_{SQ}^{m} &= 2\mathrm{i}\,m(\xi')^{*}\xi\,\left[-x^{m}(\bar{\psi}'_{-}\psi_{+}) + x^{n}(\bar{\psi}'_{-}\gamma^{m}_{\ n}\psi_{+})\right]\,, & \mathbf{K}_{SQ}^{u} &= 2\mathrm{i}\,m(\xi')^{*}\xi\,u\,\bar{\psi}'_{-}\psi_{+}\,, \\ \mathbf{K}_{SS}^{m} &= (x\cdot x + u^{-2})\,b^{m} - 2x^{m}(x\cdot b)\,, & \mathbf{K}_{SS}^{u} &= 2u\,(x\cdot b)\,, \end{split}$$
(B.13)

with $b^m \equiv 2i m(\xi')^* \xi u (\bar{\psi}'_- \gamma^m \psi_-)$ and $x \cdot b = \eta_{mn} x^m b^n$. We now define the additional parameters

$$a^{m} \equiv 2i m(\xi')^{*} \xi u (\bar{\psi}'_{+} \gamma^{m} \psi_{+}),$$

$$\Lambda^{m}_{\ n} \equiv 2i m(\xi')^{*} \xi (\bar{\psi}'_{+} \gamma^{m}_{\ n} \psi_{-} + \bar{\psi}'_{-} \gamma^{m}_{\ n} \psi_{+}),$$

$$\lambda \equiv 2i m(\xi')^{*} \xi (\bar{\psi}'_{+} \psi_{-} - \bar{\psi}'_{-} \psi_{+}),$$

$$\mathbf{r} \equiv 2(\xi')^{*} \xi (\bar{\psi}'_{+} \psi_{-} + \bar{\psi}'_{-} \psi_{+}).$$
(B.14)

It can be proven that in D = 4, for any pair of generic Majorana spinors Ψ_1 and Ψ_2 , we have

$$(\bar{\Psi}_1 \gamma^{\hat{\mu}_1} \dots \gamma^{\hat{\mu}_n} \Psi_2)^* = -\bar{\Psi}_1 \gamma^{\hat{\mu}_1} \dots \gamma^{\hat{\mu}_n} \Psi_2 \qquad (n \ge 0), \qquad (B.15)$$

namely any bilinear constructed with Majorana spinors is pure imaginary. It follows that the bilinears inside the parentheses in all the parameters are pure imaginary. This is consistent with the $\mathcal{N} = 1$ AdS₄ superalgebra, which is obtained by formally setting $2m(\xi')^*\xi = 1$ in a^m , b^m , $\Lambda^m{}_n$ and λ and $\mathbf{r} = 0$. However, because we have $\mathcal{N} = 2$, the parameters are complexified by ξ and ξ' . The Killing vector K can then be expressed as

$$\mathbf{K} = \mathbf{K}_{QQ} + \mathbf{K}_{QS} + \mathbf{K}_{SQ} + \mathbf{K}_{SS} \,, \tag{B.16}$$

with

$$\begin{aligned} \mathsf{K}_{QQ} &= a^m \partial_m \equiv P \,, \\ \mathsf{K}_{SS} &= (x \cdot x + u^{-2}) \, b^m \partial_m - 2(x \cdot b)(x^m \partial_m - u \, \partial_u) \equiv K \,, \\ \mathsf{K}_{QS} &+ \mathsf{K}_{SQ} &= \lambda (x^m \partial_m - u \, \partial_u) + \Lambda^m_{\ n} \, x^n \partial_m + \mathbf{r} \, \partial_z \equiv D + M + R \,, \end{aligned} \tag{B.17}$$

which have the formal structure of the anticommutators of the OSp(2|4) superalgebra. Specifically, the left hand side of the equations above may be identified with the anticommutators $\{Q, Q\}$, $\{S, S\}$, and $\{Q, S\}$, respectively. On the right hand side we find the bosonic generators of the SO(3, 2) isometry of AdS_4 , namely translations P, Lorentz transformations M, dilatations D, and special conformal transformations K. Moreover, we find that $R \equiv \mathbf{r} \partial_z$ is the generator of the R-symmetry. The superalgebra may be completed by computing the spinorial Lie derivatives $\mathcal{L}_V \epsilon_+$, $\mathcal{L}_V \epsilon_-$ along the above Killing vectors.

References

- [1] P. Karndumri, "Twisted compactification of N = 2 5D SCFTs to three and two dimensions from F(4) gauged supergravity", *JHEP* **09** (2015) 034, arXiv:1507.01515 [hep-th].
- [2] I. Bah, A. Passias, and P. Weck, "Holographic duals of five-dimensional SCFTs on a Riemann surface", JHEP 01 (2019) 058, arXiv:1807.06031 [hep-th].
- [3] J. M. Maldacena and C. Nunez, "Supergravity description of field theories on curved manifolds and a no go theorem", Int. J. Mod. Phys. A 16 (2001) 822–855, arXiv:hep-th/0007018.
- [4] S. L. Cacciatori and D. Klemm, "Supersymmetric AdS(4) black holes and attractors", JHEP 01 (2010) 085, arXiv:0911.4926 [hep-th].
- [5] F. Benini and N. Bobev, "Two-dimensional SCFTs from wrapped branes and c-extremization", JHEP 06 (2013) 005, arXiv:1302.4451 [hep-th].
- [6] I. Bah, C. Beem, N. Bobev, and B. Wecht, "Four-Dimensional SCFTs from M5-Branes", JHEP 06 (2012) 005, arXiv:1203.0303 [hep-th].
- [7] P. Ferrero, J. P. Gauntlett, J. M. P. Ipiña, D. Martelli, and J. Sparks, "Accelerating black holes and spinning spindles", *Phys. Rev. D* 104 no. 4, (2021) 046007, arXiv:2012.08530 [hep-th].
- [8] P. Ferrero, J. P. Gauntlett, J. M. Pérez Ipiña, D. Martelli, and J. Sparks, "D3-Branes Wrapped on a Spindle", *Phys. Rev. Lett.* **126** no. 11, (2021) 111601, arXiv:2011.10579 [hep-th].
- [9] P. Ferrero, J. P. Gauntlett, D. Martelli, and J. Sparks, "M5-branes wrapped on a spindle", JHEP 11 (2021) 002, arXiv:2105.13344 [hep-th].
- [10] S. M. Hosseini, K. Hristov, and A. Zaffaroni, "Rotating multi-charge spindles and their microstates", JHEP 07 (2021) 182, arXiv:2104.11249 [hep-th].

- [11] A. Boido, J. M. P. Ipiña, and J. Sparks, "Twisted D3-brane and M5-brane compactifications from multi-charge spindles", *JHEP* 07 (2021) 222, arXiv:2104.13287 [hep-th].
- [12] P. Ferrero, M. Inglese, D. Martelli, and J. Sparks, "Multi-charge accelerating black holes and spinning spindles", arXiv:2109.14625 [hep-th].
- [13] C. Couzens, K. Stemerdink, and D. van de Heisteeg, "M2-branes on Discs and Multi-Charged Spindles", arXiv:2110.00571 [hep-th].
- [14] I. Bah, F. Bonetti, R. Minasian, and E. Nardoni, "M5-brane sources, holography, and Argyres-Douglas theories", *JHEP* 11 (2021) 140, arXiv:2106.01322 [hep-th].
- [15] C. Couzens, N. T. Macpherson, and A. Passias, " $\mathcal{N} = (2, 2) \text{ AdS}_3$ from D3-branes wrapped on Riemann surfaces", arXiv:2107.13562 [hep-th].
- [16] M. Suh, "D3-branes and M5-branes wrapped on a topological disc", arXiv:2108.01105 [hep-th].
- [17] M. Suh, "D4-D8-branes wrapped on a manifold with non-constant curvature", arXiv:2108.08326 [hep-th].
- [18] M. Suh, "M2-branes wrapped on a topological disc", arXiv:2109.13278 [hep-th].
- [19] D. Cassani, J. P. Gauntlett, D. Martelli, and J. Sparks, "Thermodynamics of accelerating and supersymmetric AdS4 black holes", *Phys. Rev. D* 104 no. 8, (2021) 086005, arXiv:2106.05571 [hep-th].
- [20] A. Cabo-Bizet, D. Cassani, D. Martelli, and S. Murthy, "Microscopic origin of the Bekenstein-Hawking entropy of supersymmetric AdS₅ black holes", *JHEP* 10 (2019) 062, arXiv:1810.11442 [hep-th].
- [21] S. M. Hosseini, K. Hristov, and A. Zaffaroni, "Gluing gravitational blocks for AdS black holes", JHEP 12 (2019) 168, arXiv:1909.10550 [hep-th].
- [22] F. Benini, K. Hristov, and A. Zaffaroni, "Black hole microstates in AdS₄ from supersymmetric localization", JHEP 05 (2016) 054, arXiv:1511.04085 [hep-th].
- [23] J. Nian and L. A. Pando Zayas, "Microscopic entropy of rotating electrically charged AdS₄ black holes from field theory localization", *JHEP* 03 (2020) 081, arXiv:1909.07943 [hep-th].
- [24] K. Hristov, S. Katmadas, and C. Toldo, "Matter-coupled supersymmetric Kerr-Newman-AdS₄ black holes", *Phys. Rev. D* 100 no. 6, (2019) 066016, arXiv:1907.05192 [hep-th].
- [25] P. Benetti Genolini, J. M. Perez Ipiña, and J. Sparks, "Localization of the action in AdS/CFT", JHEP 10 (2019) 252, arXiv:1906.11249 [hep-th].
- [26] P. Ferrero, J. P. Gauntlett, and J. Sparks, "Supersymmetric spindles", arXiv:2112.01543 [hep-th].

- [27] A. Brandhuber and Y. Oz, "The D-4 D-8 brane system and five-dimensional fixed points", Phys. Lett. B 460 (1999) 307-312, arXiv:hep-th/9905148.
- [28] M. Cvetic, S. S. Gubser, H. Lu, and C. N. Pope, "Symmetric potentials of gauged supergravities in diverse dimensions and Coulomb branch of gauge theories", *Phys. Rev. D* 62 (2000) 086003, arXiv:hep-th/9909121.
- [29] L. J. Romans, "The F(4) Gauged Supergravity in Six-dimensions", Nucl. Phys. B 269 (1986) 691.
- [30] R. D'Auria, S. Ferrara, and S. Vaula, "Matter coupled F(4) supergravity and the AdS(6) / CFT(5) correspondence", JHEP 10 (2000) 013, arXiv:hep-th/0006107.
- [31] S. M. Hosseini, K. Hristov, A. Passias, and A. Zaffaroni, "6D attractors and black hole microstates", JHEP 12 (2018) 001, arXiv:1809.10685 [hep-th].
- [32] A. Guarino, D. L. Jafferis, and O. Varela, "String Theory Origin of Dyonic N=8 Supergravity and Its Chern-Simons Duals", *Phys. Rev. Lett.* **115** no. 9, (2015) 091601, arXiv:1504.08009 [hep-th].
- [33] D. L. Jafferis and S. S. Pufu, "Exact results for five-dimensional superconformal field theories with gravity duals", JHEP 05 (2014) 032, arXiv:1207.4359 [hep-th].
- [34] A. Passias, D. Prins, and A. Tomasiello, "A massive class of $\mathcal{N} = 2 \text{ AdS}_4 \text{ IIA}$ solutions", *JHEP* **10** (2018) 071, arXiv:1805.03661 [hep-th].
- [35] P. M. Crichigno and D. Jain, "The 5d Superconformal Index at Large N and Black Holes", JHEP 09 (2020) 124, arXiv:2005.00550 [hep-th].
- [36] P. M. Crichigno, D. Jain, and B. Willett, "5d Partition Functions with A Twist", JHEP 11 (2018) 058, arXiv:1808.06744 [hep-th].
- [37] H. Lu, C. N. Pope, and J. F. Vazquez-Poritz, "From AdS black holes to supersymmetric flux branes", Nucl. Phys. B 709 (2005) 47–68, arXiv:hep-th/0307001.
- [38] M. Cvetic, H. Lu, and C. N. Pope, "Gauged six-dimensional supergravity from massive type IIA", *Phys. Rev. Lett.* 83 (1999) 5226-5229, arXiv:hep-th/9906221.
- [39] N. Seiberg, "Five-dimensional SUSY field theories, nontrivial fixed points and string dynamics", Phys. Lett. B 388 (1996) 753-760, arXiv:hep-th/9608111.
- [40] A. Passias, "A note on supersymmetric AdS₆ solutions of massive type IIA supergravity", JHEP 01 (2013) 113, arXiv:1209.3267 [hep-th].
- [41] D. L. Jafferis, "The Exact Superconformal R-Symmetry Extremizes Z", JHEP 05 (2012) 159, arXiv:1012.3210 [hep-th].
- [42] S. M. Hosseini, I. Yaakov, and A. Zaffaroni, "The joy of factorization at large N: five-dimensional indices and AdS black holes", arXiv:2111.03069 [hep-th].

- [43] S. M. Hosseini, I. Yaakov, and A. Zaffaroni, "Topologically twisted indices in five dimensions and holography", JHEP 11 (2018) 119, arXiv:1808.06626 [hep-th].
- [44] N. Bobev and P. M. Crichigno, "Universal RG Flows Across Dimensions and Holography", JHEP 12 (2017) 065, arXiv:1708.05052 [hep-th].
- [45] S. M. Hosseini and A. Zaffaroni, "Universal AdS Black Holes in Theories with 16 Supercharges and Their Microstates", *Phys. Rev. Lett.* **126** no. 17, (2021) 171604, arXiv:2011.01249 [hep-th].
- [46] S. M. Hosseini and K. Hristov, "4d F(4) gauged supergravity and black holes of class *F*", *JHEP* 02 (2021) 177, arXiv:2011.01943 [hep-th].
- [47] S. Giri, "Black holes with spindles at the horizon", arXiv:2112.04431 [hep-th].
- [48] L. F. Alday, M. Fluder, C. M. Gregory, P. Richmond, and J. Sparks,
 "Supersymmetric solutions to Euclidean Romans supergravity", *JHEP* 02 (2016) 100, arXiv:1505.04641 [hep-th].
- [49] P. B. Genolini and P. Richmond, "Supersymmetry of higher-derivative supergravity in AdS4 holography", *Phys. Rev. D* 104 no. 6, (2021) L061902, arXiv:2107.04590 [hep-th].
- [50] C. Couzens, J. P. Gauntlett, D. Martelli, and J. Sparks, "A geometric dual of c-extremization", JHEP 01 (2019) 212, arXiv:1810.11026 [hep-th].
- [51] S. M. Hosseini and A. Zaffaroni, "Geometry of *I*-extremization and black holes microstates", JHEP 07 (2019) 174, arXiv:1904.04269 [hep-th].
- [52] J. P. Gauntlett, D. Martelli, and J. Sparks, "Toric geometry and the dual of *I*-extremization", JHEP 06 (2019) 140, arXiv:1904.04282 [hep-th].
- [53] J. P. Gauntlett, D. Martelli, and J. Sparks, "Fibred GK geometry and supersymmetric AdS solutions", JHEP 11 (2019) 176, arXiv:1910.08078 [hep-th].
- [54] M. Suh, "Supersymmetric AdS_6 black holes from matter coupled F(4) gauged supergravity", *JHEP* **02** (2019) 108, arXiv:1810.00675 [hep-th].
- [55] H. Lu, C. N. Pope, and J. Rahmfeld, "A Construction of Killing spinors on S**n", J. Math. Phys. 40 (1999) 4518-4526, arXiv:hep-th/9805151.