

DISPERSIVE ESTIMATES FOR THE WAVE EQUATION OUTSIDE A CYLINDER IN \mathbb{R}^3

FELICE IANDOLI, OANA IVANOVICI

ABSTRACT. We consider the wave equation with Dirichlet boundary conditions in the exterior of a cylinder in \mathbb{R}^3 and we construct a global in time parametrix to derive sharp dispersion estimates for all frequencies (low and high) and, as a corollary, Strichartz estimates, all matching the \mathbb{R}^3 case.

1. GENERAL SETTING

We consider the linear wave equation on an exterior domain $\Omega \subset \mathbb{R}^3$ with smooth boundary; let Δ_D be the Laplacian with constant coefficients and Dirichlet boundary conditions,

$$\begin{cases} (\partial_t^2 - \Delta_D)u = 0, & \text{in } \Omega, \\ u|_{t=0} = u_0, \partial_t u|_{t=0} = u_1, & u|_{x=0} = 0. \end{cases} \quad (1.1)$$

A basic homogeneous (local) estimate says that on any smooth Riemannian manifold (Ω, g) *without* boundary, a solution u to the wave equation satisfies (for $T < \infty$)

$$\|u\|_{L^q(0,T)L^r(\Omega)} \leq C_T (\|u_0\|_{\dot{H}^\beta(\Omega)} + \|u_1\|_{\dot{H}^{\beta-1}(\Omega)}), \quad (1.2)$$

where $\beta = d(\frac{1}{2} - \frac{1}{r}) - \frac{1}{q}$ is dictated by scaling and the pair (q, r) is wave-admissible, i.e such that $\frac{2}{q} + \frac{d-1}{r} \leq \frac{d-1}{2}$ and $(q, r, d) \neq (2, \infty, 3)$. Here $\dot{H}^\beta(\Omega)$ denotes the homogeneous L^2 Sobolev space over Ω . If (1.2) holds for $T = \infty$, Strichartz estimates are said to be global. Such inequalities were established long ago for Minkowski space (flat metrics) and can be generalized to any smooth Riemannian manifold (Ω, g) because of their local character (finite propagation speed). They are sharp on every Riemannian manifold (Ω, g) with $\partial\Omega = \emptyset$.

The aforementioned results for \mathbb{R}^d and manifolds without boundary are now well understood. Euclidean results go back to R.Strichartz's pioneering work [17], where he proved the particular case $q = r$ for the wave and Schrödinger equations. This was later generalized to mixed $L_t^q L_x^r$ norms by J.Ginibre and G.Velo [3] for Schrödinger equations, where (q, r) is sharp admissible and $q > 2$; wave estimates were obtained by J.Ginibre and G.Velo [4, 5], H.Lindblad and C.Sogge [8], as well as L.Kapitanski for a smooth variable coefficients metric,[9]. Endpoint cases for both equations were finally settled by M.Keel and T.Tao [10]. On manifolds without boundary, by finite speed of propagation it suffices to work in coordinate charts and to establish estimates for variable coefficients operators in \mathbb{R}^d . For operators with $C^{1,1}$ coefficients, Strichartz estimates were shown by H.Smith [14] (see also D.Tataru [18] for metrics with C^α coefficients).

The canonical path leading to such Strichartz estimates is to obtain a stronger, fixed time, dispersion estimate, which is then combined with energy conservation, interpolation and a duality argument to obtain (1.2). If $e^{\pm it\sqrt{-\Delta_{\mathbb{R}^d}}}$ are the half-wave propagators in $(\mathbb{R}^d, (\delta_{i,j}))$, $\chi \in C_0^\infty(]0, \infty[)$ then the following holds:

$$\|\chi(hD_t)e^{\pm it\sqrt{-\Delta_{\mathbb{R}^d}}}\|_{L^1(\mathbb{R}^d) \rightarrow L^\infty(\mathbb{R}^d)} \leq C(d)h^{-d} \min\{1, (h/|t|)^{\frac{d-1}{2}}\}. \quad (1.3)$$

Our aim in the present paper is to prove dispersion for (1.1) when $\partial\Omega$ is a cylinder in \mathbb{R}^3 : a parametrix near diffractive points may be explicitly obtained in a similar way as in [7] (where the case of the wave and Schrödinger equations outside a ball of \mathbb{R}^3 was dealt with by the second author and G.Lebeau) and the diffractive effects in the shadow region are much weaker; however, dealing with the case when both the source and the observation points are located very close to the boundary at a long distance is a real hurdle. In fact, this situation corresponds to rays that remain close to the boundary for a large time interval and propagate near points where the curvature vanishes : to our knowledge, a parametrix near such points, allowing for

The authors were supported by ERC grant ANADEL 757 996.
Key words : Dispersive estimates, wave equation, Dirichlet boundary condition. *AMS subject classification* : 35Rxx, 58Jxx.

sharp amplitude estimates, was only constructed in [11] inside a cylindrical domain of \mathbb{R}^3 . However, while in [11] the time is bounded (as at the time we did not know to handle the reflections in very large time in the interior case), the parametrix we construct here is global in time, depending on the angle of the initial directions of propagation and on the initial distance of the data to the boundary: different values of these parameters completely modify its construction; dealing with points where the curvature vanishes requires handling separately different situations (involving Hankel and Bessel functions). We expect that in order to deal with general boundaries with no convexity or concavity assumption, and allowing for possibly vanishing curvatures along lower dimensional submanifolds, we need to understand a variety of simple models and the exterior of the cylinder is the first of them after the exterior of a sphere.

Let us provide some details : introducing cylindrical coordinates in \mathbb{R}^3 , our domain becomes $\Omega = \{(r, \theta, z), r \geq 1, \theta \in [0, 2\pi), z \in \mathbb{R}\}$ and $\Delta_D = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} + \frac{\partial^2}{\partial z^2}$. With h a small parameter and $\tau = h\partial_t/i$, $\eta = h\partial_y/i$, $\xi = h\partial_x/i$, $\vartheta = h\partial_z/i$, the characteristic set of $\partial_t^2 - \Delta_D$ is $\tau^2 = \xi^2 + \frac{1}{r^2}\eta^2 + \vartheta^2$ and the boundary is $\{r = 1\}$. In [7], G.Lebeau and the second author constructed a global in time parametrix for the wave equation outside a ball in \mathbb{R}^3 , which allowed them to obtain sharp dispersion bounds. In the particular case of [7] the model domain was $\{(r, \theta, \omega), r \geq 1, \theta \in [0, \pi), \omega \in [0, 2\pi)\}$ and the Laplace operator was given by $\Delta_F := \frac{\partial^2}{\partial r^2} + \frac{d-1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} (\frac{\partial^2}{\partial \theta^2} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \omega^2})$. The main difficulty came from rays that hit the boundary without being deviated (corresponding to $\xi = 0$, $\eta = 1$ and r near 1; in fact, due to the rotational symmetry, in the exterior of the ball the characteristic equation is $\xi^2 + \frac{1}{r^2}\eta^2 = \tau^2$) : for this regime, the most efficient tool is the Melrose-Taylor parametrix (see [20]), as it provides us with the form of the solution to (1.1) near diffractive points $\xi = 0$, $r = 1$ (recall that this parametrix was first used by H.Smith and Ch.Sogge in [16] to obtain, in a direct way, local in time sharp Strichartz bounds for waves). In the case of the exterior of a cylinder, the ‘‘diffractive regime’’ would correspond to $(\eta/\tau)^2 + (\vartheta/\tau)^2 = 1$, $\xi = 0$, $r = 1$ (instead of $(\eta/\tau)^2 = 1$, $\xi = 0$, $r = 1$ of [7]) : it turns out that when ϑ/τ is very close to 1 the Melrose-Taylor parametrix fails to apply (essentially because one cannot perform any kind of stationary phase arguments anymore in the oscillatory integrals that allow to obtain the form of the solution near the boundary in terms of Airy functions). In particular, the situation $\vartheta/\tau = 1$ correspond to rays that (start and) remain close to the boundary for all time and at our knowledge has been encountered only in [11] where the author studied dispersive bounds for (1.1) in the interior of a cylindrical domain $\{(r, \theta, z), r \leq 1, \theta \in [0, 2\pi), z \in \mathbb{R}\} \subset \mathbb{R}^3$ with Dirichlet Laplacian $\Delta_D = \partial_r^2 + (2-r)\partial_\theta^2 + \partial_z^2$ (and obtained a ‘‘sharp loss’’ of 1/4 due to swallowtail type singularities in the wave front set) ; notice however that in [11] the time is bounded so when ϑ/τ is close to 1 the estimates follow easy by Sobolev embedding (and a parametrix is naturally obtained in terms of a spectral sum). In the exterior of a cylinder, our aim is to construct the parametrix globally in time, which makes this situation more difficult (and the case $1 - \vartheta/\tau \sim 2^{-j}$ already very delicate when compared to the exterior of a ball).

Throughout the rest of the paper $A \lesssim B$ means that there exists a constant C such that $A \leq CB$, such a constant may change from line to line and it is independent of all parameters, and $A \sim B$ means that $B \lesssim A \lesssim B$. We may now state our main results.

Theorem 1.1. *Let $\Theta \subset \mathbb{R}^3$ be the cylinder in \mathbb{R}^3 and set $\Omega = \mathbb{R}^3 \setminus \Theta$. Let Δ_D denote the Laplace operator in Ω with Dirichlet boundary condition and let $\chi \in C_0^\infty(0, \infty)$. The following estimate holds for all $t > 0$*

$$\|\chi(hD_t)e^{\pm it\sqrt{-\Delta_D}}\|_{L^1(\Omega) \rightarrow L^\infty(\Omega)} \lesssim h^{-3} \min\{1, \frac{h}{t}\}. \quad (1.4)$$

Moreover, let $\chi_0 \in C_0^\infty(-2, 2)$, equal to 1 on $[0, 3/2]$. Then $\|\chi_0(D_t)e^{\pm it\sqrt{-\Delta_D}}\|_{L^1(\Omega) \rightarrow L^\infty(\Omega)} \lesssim 1/(1+t)$.

Theorem 1.2. *Under the assumptions of Theorem 1.1, Strichartz estimates for the wave flow outside a cylinder in \mathbb{R}^3 hold as in the flat case, globally in time.*

Theorem 1.2 follows from (1.4) using the usual TT^* argument and the conservation of energy. In the remaining of this work we focus on the proof of Theorem 1.1, first in the high-frequency situation which is by far the most difficult one. The small frequency case will be sketched in the last part.

We recall a classical notion of asymptotic expansion: a function $f(w)$ admits an asymptotic expansion for $w \rightarrow 0$ when there exists a (unique) sequence $(c_n)_n$ such that, for any n , $\lim_{w \rightarrow 0} w^{-(n+1)}(f(w) - \sum_0^n c_j w^j) = c_{n+1}$. We denote $f(w) \sim_w \sum_n c_n w^n$.

1.0.1. *The incoming wave.* Let \mathbb{D} denote the unit disk in \mathbb{R}^2 and let $\Theta := \mathbb{D} \times \mathbb{R} \subset \mathbb{R}^3$. We set $\Omega := \mathbb{R}^3 \setminus \Theta$, then $\partial\Omega = \mathbb{S}^1 \times \mathbb{R}$ is the infinite cylinder. We introduce cylindrical coordinates as follows: a point of Q of Ω with coordinates $(x_1, x_2, x_3) \in \mathbb{R}^3$ is defined by (r, θ, z) where $r > 1$, $\theta \in [0, 2\pi)$ and $z \in \mathbb{R}$ and where $x_1 = r \cos(\theta)$, $x_2 = r \sin(\theta)$, $x_3 = z$. We also set $r = 1 + x$, $x \geq 0$, $y := \pi/2 - \theta$, $\theta \in [0, 2\pi)$, $z \in \mathbb{R}$. In these coordinates, the Laplacian becomes

$$\Delta = \frac{\partial^2}{\partial x^2} + \frac{1}{(1+x)} \frac{\partial}{\partial x} + \frac{1}{(1+x)^2} \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}. \quad (1.5)$$

In the new coordinate system, $x \rightarrow (x, y, z)$ is the ray orthogonal to $\partial\Omega$ at $(0, y, z) \in \partial\Omega$. Any point in $Q \in \Omega$ can be written under the form $Q = (0, y, z) + x\vec{\nu}_{(y,z)}$, where (y, z) is the orthogonal projection of Q on $\partial\Omega$ and $\vec{\nu}_{(y,z)}$ the outward unit normal to $\partial\Omega$ pointing towards Ω . The dual variable to (x, y, z) is denoted (ξ, η, ϑ) . The principal symbol of $\partial_t^2 - \Delta$ associated to (1.5) is $p(x, \xi, \eta, \vartheta, \tau) = -\tau^2 + \xi^2 + (1+x)^{-2}\eta^2 + \vartheta^2$. The time variable and its dual are t and τ . We let $\mathcal{Q} = \{(x, y, z, t, \xi, \eta, \vartheta, \tau), x = 0\}$, $\mathcal{P} = \{(x, y, z, t, \xi, \eta, \vartheta, \tau), p = 0\}$. The cotangent bundle of $\partial\Omega \times \mathbb{R}$ is the quotient of \mathcal{Q} by the action of translation in ξ , and we take as coordinates $(y, z, t, \eta, \vartheta, \tau)$. A point $(y, z, t, \eta, \vartheta, \tau) \in T^*(\partial\Omega \times \mathbb{R})$ is classified as one of three distinct types: it is said to be *hyperbolic* if there are two distinct nonzero real solutions ξ to $p|_{x=0} = 0$. These two solutions yield two distinct bicharacteristics, one of which enters Ω as t increases (the *incoming ray*) and one which exits Ω as t increases (the *outgoing ray*). The point is *elliptic* if there are no real solutions ξ to $p|_{x=0} = 0$. In the remaining case $\tau^2 = \eta^2 + \vartheta^2$, there is a unique solution $\xi = 0$ to $p|_{x=0} = 0$ which yields a glancing ray, and the point is said to be a *glancing point*. A glancing ray has exactly second order contact with the boundary if we have in addition $\eta^2 \frac{d}{dx}(1+x)^{-2}|_{x=0} = -\eta^2/2 < 0$, which means if $\eta \neq 0$. We set $\alpha = \eta/\tau$, $\gamma = \vartheta/\tau$: the glancing condition becomes $\alpha^2 + \gamma^2 = 1$, while the hyperbolic (or elliptic) regime satisfy $1 - \alpha^2 - \gamma^2 > 0$ (or $1 - \alpha^2 - \gamma^2 < 0$). A point in $T^*(\partial\Omega \times \mathbb{R})$ such that $1 \geq \alpha^2 > 0$ may be a glancing point of order exactly two. When $\alpha = 0$, it is a glancing point of order ∞ (as, in this case, $H_p^j x = 0$ for all $j \geq 1$).

Remark 1.3. *When $1 - \gamma^2 - \alpha^2 \geq 1/16$, then on the boundary $\xi^2/\tau^2 = (1 - \gamma^2 - \alpha^2) \geq 1/16$ in which case the corresponding point in the cotangent bundle is hyperbolic. The proof of Theorem 1.1 for such points follows as in the case of the half-space, so we will focus on the situation $1 - \gamma^2 - \alpha^2 \leq 1/16$, when $|\xi/\tau| \lesssim 1/4$.*

Let Δ be the Laplacian in \mathbb{R}^3 , then the solution $u_{free}(Q, Q_0, t)$ to the free wave equation $(\partial_t^2 - \Delta)u_{free} = 0$ in \mathbb{R}^3 with $u_{free}|_{t=0} = \delta_{Q_0}$, $\partial_t u_{free}|_{t=0} = 0$, where δ_{Q_0} is the Dirac distribution at $Q_0 \in \mathbb{R}^3$, is given by :

$$u_{free}(Q, Q_0, t) := \frac{1}{(2\pi)^3} \int e^{i(Q-Q_0)\xi} \cos(t|\xi|) d\xi. \quad (1.6)$$

If $w_{in}(Q, Q_0, \tau) := \widehat{1_{t>0} u_{free}}(Q, Q_0, \tau)$ denotes its Fourier transform in time, then the following holds :

$$w_{in}(Q, Q_0, \tau) = \frac{i\tau e^{-i\tau|Q-Q_0|}}{4\pi |Q-Q_0|}. \quad (1.7)$$

Consider the equation (1.1) with initial data $(\delta_{Q_0}, 0)$, where $Q_0 \in \Omega$ is an arbitrary point

$$\begin{cases} (\partial_t^2 - \Delta_D)u = 0 & \text{in } \Omega \times \mathbb{R}, \\ u|_{t=0} = \delta_{Q_0}, \quad \partial_t u|_{t=0} = 0, \quad u|_{\partial\Omega} = 0. \end{cases} \quad (1.8)$$

Let $u(Q, Q_0, t) = \cos(t\sqrt{-\Delta_D})(\delta_{Q_0})(Q)$ denote the solution to (1.8) : in order to prove Theorem 1.1 we construct u for all t and then deduce global in time dispersive bounds. We may assume, without loss of generality, that $\text{dist}(Q_0, \partial\Omega) \geq \text{dist}(Q, \partial\Omega)$: indeed, when this is not the case we can use the symmetry of the Green function to change Q_0 and Q . We may assume that, in the coordinates (r, θ, z) , the source point is of the form $Q_0 = (s, 0, 0)$, where $s - 1 > 0$ represents the distance from Q_0 to the boundary. Let Q be an arbitrary point of Ω , then $Q := (r \cos \theta, r \sin \theta, z)$. We introduce the distance between Q and Q_0 as follows

$$\tilde{\phi}(r, \theta, z, s) := |Q - Q_0| = \sqrt{r^2 - 2sr \cos \theta + s^2 + z^2}. \quad (1.9)$$

In the normal coordinates (x, y, z) we have $Q_0 = (s - 1, \frac{\pi}{2}, 0)$ and $Q = ((1+x) \sin y, (1+x) \cos y, z)$; letting $\phi(x, y, z, s) := \tilde{\phi}(1+x, \frac{\pi}{2} - y, z, s)$, we have $\phi(x, y, z, s) = \sqrt{(1+x)^2 - 2s(1+x) \sin y + s^2 + z^2}$. The coordinates (x, y, z) will be particularly useful when working near a glancing point; near hyperbolic (or elliptic) points we keep the cylindrical coordinates (r, θ, z) . We will switch them when necessary.

Let u_{free} be given in (1.6). By finite speed of propagation, for any sufficiently small time $0 < t < d(Q_0, \partial\Omega)$, the solution to (1.8) in Ω is just $1_{t>0}u_{free}$, whose Fourier transform equals w_{in} . In the following, we decompose w_{in} according to the initial directions of propagation as follows : let $\psi_0(\beta)$ be a smooth function supported near 1, equal to 1 for $1 \geq \beta \geq 1/36$, equal to 0 for $\beta \leq 1/64$ and such that $0 \leq \psi_0 \leq 1$. Let also $\psi \in C_0^\infty(1/4, 4)$ equal to 1 near 1 such that $1 - \psi_0(\beta) = \sum_{j \geq 1} \psi(2^{2j}\beta)$. Write $w_{in} = w_0 + \sum_{j \geq 1} w_j$, with

$$w_0(Q, Q_0, \tau) = \frac{\tau^2}{(2\pi)^2} \frac{i\tau}{4\pi} \int \frac{\psi_0(1-\gamma^2)}{\phi(x, \tilde{y}, \tilde{z}, s)} e^{i\tau((y-\tilde{y})\alpha + (z-\tilde{z})\gamma)} e^{-i\tau\phi(x, \tilde{y}, \tilde{z}, s)} d\alpha d\gamma d\tilde{y} d\tilde{z}, \quad (1.10)$$

$$w_j(Q, Q_0, \tau) = \frac{\tau^2}{(2\pi)^2} \frac{i\tau}{4\pi} \int \frac{\psi(2^{2j}(1-\gamma^2))}{\phi(x, \tilde{y}, \tilde{z}, s)} e^{i\tau((y-\tilde{y})\alpha + (z-\tilde{z})\gamma)} e^{-i\tau\phi(x, \tilde{y}, \tilde{z}, s)} d\alpha d\gamma d\tilde{y} d\tilde{z}. \quad (1.11)$$

Let $\psi_j(\beta) := \psi(2^{2j}\beta)$. We set $u_{free}^+ := 1_{t>0}u_{free} = \int e^{i\tau t} w_{in}(Q, Q_0, \tau) d\tau$. Using (1.10) and (1.11), we decompose as follows $u_{free} = u_{free,0} + \sum_{j \geq 1} u_{free,j}$ and set $u_{free,j}^+ := 1_{t>0}u_{free,j}$, where $\mathcal{F}(u_{free,j}^+) = w_j$.

The paper is organized as follows : in Section 2 we consider $h \in (0, h_0)$ for some small $h_0 \in (0, 1)$ and $s \geq \sqrt{2}$ and we show that, for all $u_{free,j}^+$ with $0 \leq j \leq \frac{1}{3} \log_2(s/h)$, we may construct the outgoing wave in a similar way to that used in [7] in the exterior of a ball as each w_j hits the obstacle at hyperbolic or glancing points of order exactly 2. The assumptions on s and j are necessary to construct the reflected waves near glancing points and to make sure that stationary phase methods do apply. In Section 3 we obtain dispersive bounds first for each $j \leq \frac{1}{3} \log_2(s/h)$ and show that the sum over j is still bounded as expected. Both Sections 2 and 3 deal separately with the glancing and hyperbolic regimes, and also with the cases $\text{dist}(Q, \partial\Omega) \geq \sqrt{2} - 1$ or $\text{dist}(Q, \partial\Omega) \leq \sqrt{2} - 1$ as each case needs to be handled in a different way. In Section 4 we consider $h \in (0, h_0)$ and either $s \leq \sqrt{2}$ or $s \geq \sqrt{2}$ and $j \gtrsim \frac{1}{3} \log_2(s/h)$: in these cases we cannot construct the reflected waves as before, either because the data is too close to the boundary or because the phase functions of w_j don't oscillate anymore. We obtain an explicit parametrix in terms of Bessel and Hankel functions and proceed with the dispersive bounds. In the last Section we explain why the last parametrix still allows to obtain dispersion in the case of small frequencies.

2. PARAMETRIX FOR (1.1) WHEN $s \geq \sqrt{2}$, $h \in (0, h_0)$ AND $2^{-3j}s/h \gtrsim 1$

We consider the source point to be of the form $Q_0 = (s, 0, 0)$, where $s - 1$ represents the distance from Q_0 to the $\partial\Omega$. In this section we consider $s \geq \sqrt{2}$. Let $h_0 \in (0, 1)$ be small and $h \in (0, h_0)$.

Lemma 2.1. *Let $Q_0 = (s, 0, 0)$ with $s \geq \sqrt{2}$ and j such that $2^{-3j}s/h \gtrsim 1$. Then $u_{free,j,h}(\cdot, Q_0, t)$ solves the free wave equation and $u_{free,j,h}(\cdot, Q_0, 0)|_{\partial\Omega} = O(h^\infty)$. Moreover, $u_{free,j,h}(P, Q_0, t)|_{P \in \partial\Omega} = O((h/t)^\infty)$ for $P = (0, \cdot, z)$ with $|z| \geq 4t$.*

Proof. The first statement follows from the fact that Δ commutes with D_z ; for j as above, the second statement follows using non-stationary phase arguments for the phase $\tau(t + (z - \tilde{z})\gamma + (y - \tilde{y})\alpha - \phi(0, \tilde{y}, \tilde{z}, s))$ of $u_{free,j,h}$. If $|z| \geq 4t$, the phase is also non-stationary with respect to τ which allows to conclude. \square

Our goal in this section is to construct, for each $0 \leq j \leq \frac{1}{3} \log_2(s/(hM))$, the solution u_j to the Dirichlet wave equation on Ω whose incoming part (before reflection) equals $u_{free,j}$. To do that, we first set

$$\underline{u}_j(Q, Q_0, t) := \begin{cases} u_j(Q, Q_0, t), & \text{if } Q \in \Omega, \\ 0, & \text{if } Q \in \overline{\Omega}. \end{cases} \quad (2.1)$$

Then, using Duhamel formula and with $u_j^+ := 1_{t>0}u_j$, \underline{u}_j reads as follows

$$\underline{u}_j|_{t>0} = u_{free,j}^+ - u_j^\#, \quad u_j^\#(Q, Q_0, t) := \int_{\partial\Omega} \frac{\partial_\nu u_j^+(P, Q_0, t - |Q - P|)}{4\pi|Q - P|} d\sigma(P). \quad (2.2)$$

Let $h_0 \in (0, 1)$ small enough and $h \in (0, h_0)$. Let $\chi \in C_0^\infty([\frac{1}{2}, 2])$ be a smooth cutoff equal to 1 on $[\frac{3}{4}, \frac{3}{2}]$ and such that $0 \leq \chi \leq 1$. As we are interested in evaluating $\chi(hD_t)\underline{u}_j(Q, Q_0, t)$, let

$$u_{free,j,h}^+ := \chi(hD_t)u_{free,j}^+, \quad u_{j,h}^\# := \chi(hD_t)u_j^\#(Q, Q_0, t). \quad (2.3)$$

As the free wave flow $u_{free,j,h}$ satisfies the usual dispersive estimates, we are reduced to evaluating the sum over $j \leq \frac{1}{3} \log_2(s/(M))$ of $u_{j,h}^\#(Q, Q_0, t)$ (or, when possible, of $\chi(hD_t)u_j^+ := u_{j,h}^+$). Using (2.2) we have

$$u_{j,h}^\#(Q, Q_0, t) = \int e^{it\tau} \chi(h\tau) \int_{P \in \partial\Omega} \mathcal{F}(\partial_\nu u_j^+|_{\partial\Omega})(P, Q_0, \tau) \frac{1}{4\pi|Q-P|} e^{-i\tau|Q-P|} d\sigma(P) d\tau, \quad (2.4)$$

where $\mathcal{F}(\partial_\nu u_j^+|_{\partial\Omega})(P, Q_0, \tau)$ denotes the Fourier transform in time of $\partial_\nu u_j^+|_{\partial\Omega}(P, Q_0, t)$.

Definition 2.2. For a source point Q_0 as above, we define its apparent contour \mathcal{C}_{Q_0} as the set of points $P \in \partial\Omega$ such that the ray Q_0P is tangent to $\partial\Omega$: in other words, for $\tilde{\phi}$ defined in (1.9), we have

$$\mathcal{C}_{Q_0} := \{P \in \partial\Omega \text{ with coordinates } (1, \theta, z) \text{ such that } \partial_r \tilde{\phi}(1, \theta, z, s) = 0\}.$$

As $\partial_r \tilde{\phi} = (r - s \cos \theta) / \tilde{\phi}$ cancels at $r = 1$ when $\cos \theta = \frac{1}{s}$, we find $\mathcal{C}_{Q_0} := \{P = (1, \arccos(1/s), z), z \in \mathbb{R}\}$. In the coordinates (x, y, z) we have $\mathcal{C}_{Q_0} = \{P = (0, y, z), y = \arcsin(1/s)\}$. In the following we set $\theta_* := \arccos(1/s) = \frac{\pi}{2} - \arcsin(1/s)$ and $y_* := \arcsin(1/s)$.

Definition 2.3. Let $h \in (0, h_0)$. We define $j(s, h) := \sup\{j, 2^{-3j}s/h \geq 1\}$ so that $2^{-3j(s,h)}s/h \sim 1$.

On the support of $\psi_j(1 - \gamma^2)$ we have $\sqrt{1 - \gamma^2} \sim 2^{-2j}$ and in this section we consider only $0 \leq j \leq j(s, h)$. In the following we deal separately with the case $\frac{\alpha}{\sqrt{1 - \gamma^2}}$ near 1, when the possible glancing points have exactly second order contact with the boundary and the case $\frac{\alpha}{\sqrt{1 - \gamma^2}}$ outside a small neighborhood of 1.

Let $\chi_0 \in C_0^\infty([-2, 2])$ and equal to 1 on $[-\frac{3}{2}, \frac{3}{2}]$, fix $\varepsilon > 0$ small enough and set $\chi_\varepsilon(\cdot) := \chi_0((\cdot - 1)/\varepsilon)$. We let $w_{j,gl}$ be defined by (1.10), (1.11) with additional cutoff $\chi_\varepsilon(\frac{\alpha}{\sqrt{1 - \gamma^2}})$ supported for $|\frac{\alpha}{\sqrt{1 - \gamma^2}} - 1| \leq 2\varepsilon$. Define also $w_{j,he}$ as in (1.10), (1.11) with additional cutoff $1 - \chi_{\varepsilon_1}(\frac{\alpha}{\sqrt{1 - \gamma^2}})$. Then $u_{free,j}^+ = u_{free,j,gl}^+ + u_{free,j,he}^+$,

$$u_{free,j,gl}^+ := \int e^{it\tau} w_{j,gl} d\tau, \quad u_{free,j,he}^+ := \int e^{it\tau} w_{j,he} d\tau, \quad u_{free,j,h}^+ = \chi(hD_t) u_{free,j}^+.$$

2.1. The glancing part of u_j^+ for $0 \leq j \leq j(s, h)$. We construct $u_{j,gl}^+$, then $\partial_\nu u_{j,gl}^+$, in order to obtain the "glancing part" of $u_j^\#$ from formula (2.2). For $j = 0$, the following result due to Melrose and Taylor holds:

Proposition 2.4. Microlocally near a glancing point of exactly second order contact with the boundary there exist smooth phase functions $\iota(x, y, z, \alpha, \gamma)$ and $\zeta(x, y, z, \alpha, \gamma)$ such that $\phi_\pm = \iota \pm (-\zeta)^{3/2}$ satisfy the eikonal equation and there exist symbols a, b satisfying the transport equation such that, for any parameters α, γ in a conic neighborhood of a glancing direction and for $\tau > 1$ large enough,

$$G_\tau(x, y, z, \alpha, \gamma) := e^{i\tau\iota(x, y, z, \alpha, \gamma)} \left(aA_+(\tau^{2/3}\zeta) + b\tau^{-1/3}A'_+(\tau^{2/3}\zeta) \right) A_+^{-1}(\tau^{2/3}\zeta_0) \quad (2.5)$$

satisfies $(\tau^2 + \Delta)G_\tau = e^{i\tau\iota(x, y, z, \alpha, \gamma)} \left(a_\infty A_+(\tau^{2/3}\zeta) + b_\infty \tau^{-1/3} A'_+(\tau^{2/3}\zeta) \right) A_+^{-1}(\tau^{2/3}\zeta_0)$, where the symbols verify $a_\infty, b_\infty \in O(\tau^{-\infty})$ and where we set $\zeta_0 = \zeta|_{x=0}$. Moreover, the following properties hold

- ι and ζ are homogeneous of degree 0 and $-1/3$ and satisfy $\langle d\iota, d\iota \rangle - \zeta \langle d\zeta, d\zeta \rangle = 1$, $\langle d\iota, d\zeta \rangle = 0$, where $\langle \cdot, \cdot \rangle$ is the polarization of p ; the phase ζ_0 is independent of y, z so that $\zeta_0(\alpha, \gamma)$ vanishes at a glancing direction; the diffractive condition means that $\partial_x \zeta|_{x=0} < 0$ near a glancing point;
- the symbols $a(x, y, z, \alpha, \gamma)$ and $b(x, y, z, \alpha, \gamma)$ belong to the class $\mathcal{S}_{(1,0)}^0$ and satisfy the appropriate transport equations. Moreover $a|_{x=0}$ is elliptic at the glancing point with essential support included in a small, conic neighborhood of it, while $b|_{x=0} = 0$.

The functions ι and ζ of the Melrose-Taylor parametrix solve the system of equations

$$\begin{cases} (\partial_x \iota)^2 + \frac{(\partial_y \iota)^2}{(1+x)^2} + (\partial_z \iota)^2 - \zeta \left((\partial_x \zeta)^2 + \frac{(\partial_y \zeta)^2}{(1+x)^2} + (\partial_z \zeta)^2 \right) = 1, \\ \partial_x \iota \partial_x \zeta + \frac{\partial_y \iota \partial_y \zeta}{(1+x)^2} + \partial_z \iota \partial_z \zeta = 0. \end{cases} \quad (2.6)$$

The system (2.6) admits the pair of solutions $\iota(y, z, \alpha, \gamma) = y\alpha + z\gamma$, $\zeta(x, \alpha, \gamma) = \alpha^{2/3} \tilde{\zeta}((1+x)\sqrt{1-\gamma^2}/\alpha)$, where for $\rho := (1+x)\frac{\sqrt{1-\gamma^2}}{\alpha}$, $\tilde{\zeta}$ is the (unique) solution to $\frac{1}{\rho^2} - \tilde{\zeta}(\rho)[\tilde{\zeta}'(\rho)]^2 = 1$, $\tilde{\zeta}(1) = 0$.

Lemma 2.5. *The equation $-\tilde{\zeta}(\partial_\rho \tilde{\zeta})^2 + 1/\rho^2 = 1$, $\tilde{\zeta}(1) = 0$ has a unique solution of the form*

$$\frac{2}{3}(-\tilde{\zeta}(\rho))^{3/2} = \int_1^\rho \frac{\sqrt{w^2 - 1}}{w} dw = \sqrt{\rho^2 - 1} - \arccos\left(\frac{1}{\rho}\right), \quad (2.7)$$

if $\rho > 1$, while for $\rho < 1$ we have

$$\frac{2}{3}\tilde{\zeta}(\rho)^{3/2} = \int_\rho^1 \frac{\sqrt{1 - w^2}}{w} dw = \log[(1 + \sqrt{1 - \rho^2})/\rho] - \sqrt{1 - \rho^2}. \quad (2.8)$$

We note that at $\rho = 1$ we have $\tilde{\zeta} = 0$ and $\lim_{\rho \rightarrow 1} \frac{(-\tilde{\zeta})(\rho)}{\rho - 1} = 2^{1/3}$.

Corollary 2.6. *Let $\tilde{\psi}_0(\beta) \in C^\infty[\frac{1}{81}, 2]$ be a smooth function supported near 1 and such that $\tilde{\psi}_0 = 1$ on the support of ψ_0 . Consider the operator $M_\tau : \mathcal{E}'(\mathbb{R}^2) \rightarrow \mathcal{D}(\mathbb{R}^3)$, where $\mathcal{E}'(\mathbb{R}^2)$ is the dual space of $C^\infty(\mathbb{R}^2)$,*

$$M_\tau(f)(x, y, z) := \left(\frac{\tau}{2\pi}\right)^2 \int G_\tau(x, y, z, \alpha, \gamma) \tilde{\psi}_0(1 - \gamma^2) \hat{f}(\tau\alpha, \tau\gamma) d\alpha d\gamma.$$

Near the glancing region $(\tau^2 + \Delta)M_\tau(f) \in O(\tau^{-\infty})$ (up to the boundary) for all $f \in \mathcal{E}'(\mathbb{R}^2)$. Moreover, the restriction to the boundary $M_\tau(f)|_{\partial\Omega} =: J_\tau(f)$ defined by

$$J_\tau(f)(y, z) = \left(\frac{\tau}{2\pi}\right)^2 \int e^{i\tau(\iota(y, z, \alpha, \gamma) - \tilde{y}\alpha - \tilde{z}\gamma)} a(0, y, z, \alpha, \gamma, \tau) \tilde{\psi}_0(1 - \gamma^2) f(\tilde{y}, \tilde{z}) d\alpha d\gamma d\tilde{y} d\tilde{z},$$

has a microlocal inverse J_τ^{-1} as $a(x, y, z, \alpha, \gamma, \tau)$ is the elliptic symbol of Proposition 2.4.

We define the following operator $T_\tau : \mathcal{E}'(\mathbb{R}^2) \rightarrow \mathcal{D}(\mathbb{R}^3)$ for $F \in \mathcal{E}'(\mathbb{R}^2)$

$$T_\tau(F)(x, y, z) = \left(\frac{\tau}{2\pi}\right)^2 \tau^{1/3} \int e^{i\tau(y\alpha + z\gamma + \frac{\alpha^3}{3} + \sigma\zeta(x, \alpha, \gamma))} (a + b\frac{\sigma}{1}) \tilde{\psi}_0(1 - \gamma^2) \hat{F}(\tau\alpha, \tau\gamma) d\alpha d\gamma. \quad (2.9)$$

According to [15, Lemma A.2], T_τ is an elliptic FIO near a glancing point and $(\tau^2 + \Delta)T_\tau(F) \in O_{C^\infty}(\tau^{-\infty})$.

Lemma 2.7. *Let $Q_0 = (s, 0, 0)$ with $s \geq \sqrt{2}$, $y_* = \arcsin(1/s)$ and assume $\tau > 1$ is large enough. Then there exists a unique function F_τ satisfying $w_{0,gl}(x, y, z, \tau) = T_\tau(F_\tau)(x, y, z)$ for (x, y) in a neighborhood of $(0, y_*)$. Moreover, F_τ is explicit and has the following form*

$$\hat{F}(\tau\alpha, \tau\gamma) = \tau^{\frac{1}{6}} e^{-i\tau\sqrt{1-\gamma^2}\Gamma_0(\frac{\alpha}{\sqrt{1-\gamma^2}}, s)} f(\alpha, \gamma, \tau) \frac{\chi_\varepsilon(\frac{\alpha}{\sqrt{1-\gamma^2}})\psi_0(1-\gamma^2)}{(1-\gamma^2)^{5/12}(s^2-1)^{1/4}}, \quad (2.10)$$

where $f(\alpha, \gamma, \tau)$ is an elliptic symbol of order 0 $\psi_0(1-\tau^2)$ is the smooth cutoff from (1.10) and χ_ε is the smooth cut-off introduced to define $w_{0,gl}$. For $|\tilde{\alpha} - 1| \leq 2\varepsilon$, $\Gamma_0(\tilde{\alpha}, s) = y_*\tilde{\alpha} + \sqrt{s^2 - 1} + \frac{(1-\tilde{\alpha})^2}{2\sqrt{s^2-1}}(1 + O(1-\tilde{\alpha}))$.

The proof of Lemma 2.7 follows exactly as in [7] as $\sqrt{1-\gamma^2} \geq 1/8$ on the support of ψ_0 . Our goal is to describe, microlocally near the glancing regime, $u_{j,gl,h}^+ := \chi(hD_t)u_{j,gl}^+(\cdot, t)$ for all $0 \leq j \leq j(s, h)$. For $j = 0$:

Proposition 2.8. *For $Q = (x, y, z)$ near the glancing region we have*

$$u_{0,gl}^+(Q, Q_0, t) = \frac{1}{(2\pi)^2} \int e^{it\tau} (w_{0,gl}(x, y, z, \tau) - M_\tau(J_\tau^{-1}(w_{0,gl}|_{\partial\Omega})(x, y, z))) d\tau,$$

where, for F_τ provided by Lemma 2.7 satisfying $w_{0,gl}(\cdot, \tau) = T_\tau(F_\tau)$, $M_\tau(J_\tau^{-1}(w_{0,gl}|_{\partial\Omega}))$ reads as

$$\left(\frac{\tau}{2\pi}\right)^2 \int e^{i\tau(y\alpha + z\gamma)} \left(aA_+(\tau^{2/3}\zeta) + b\tau^{-1/3}A'_+(\tau^{2/3}\zeta)\right) \frac{A(\tau^{2/3}\zeta_0)}{A_+(\tau^{2/3}\zeta_0)} \tilde{\psi}_0(1 - \gamma^2) \hat{F}_\tau(\tau\alpha, \tau\gamma) d\alpha d\gamma. \quad (2.11)$$

Corollary 2.9. *For $P = (0, y, z) \in \partial\Omega$ near \mathcal{C}_{Q_0} we have*

$$\begin{aligned} \mathcal{F}(\partial_x u_{0,gl}^+)(P, Q_0, \tau) &= \left(\frac{\tau}{2\pi}\right)^2 \tau^{2/3+1/6} \int e^{i\tau(y\alpha + z\gamma - \sqrt{1-\gamma^2}\Gamma_0(\frac{\alpha}{\sqrt{1-\gamma^2}}, s))} \\ &\quad \times b_\partial f \frac{\chi_\varepsilon(\frac{\alpha}{\sqrt{1-\gamma^2}})\psi_0(1-\gamma^2)}{(s^2-1)^{1/4}} \frac{(1-\gamma^2)^{-5/12+1/3}}{A_+(\tau^{2/3}\zeta_0(\alpha, \gamma))} d\alpha d\gamma, \end{aligned} \quad (2.12)$$

where $\zeta_0(\alpha, \gamma) = \alpha^{2/3}\tilde{\zeta}(\sqrt{1-\gamma^2}/\alpha)$ with $\tilde{\zeta}$ defined in Lemma 2.5. For $\tilde{\alpha}$ near 1, $b_\partial(y, z, \tilde{\alpha}, \tau)$ is elliptic of order 0 in τ with main contribution $a_0\tilde{\alpha}^{-1/3}(\partial_\rho \tilde{\zeta})(\rho)|_{\rho=\frac{1}{\tilde{\alpha}}}$, $a_0 = a|_{x=0}$.

Proof. Using (2.11), we can compute the normal derivatives of each of the two contributions of $u_{0,gl}^+$ and then take the difference. As such, for $P = (0, y, z)$ near the glancing region, we obtain the following

$$\mathcal{F}(\partial_x u_{0,gl}^+)(P, Q_0, \tau) = \left(\frac{\tau}{2\pi}\right)^2 \int e^{i\tau(y\alpha + z\gamma)} \tau^{2/3} (1 - \gamma^2)^{1/3} b_\partial \left(A' - A'_+ \frac{A}{A_+} \right) (\tau^{2/3} \zeta_0) \widehat{F}_\tau(\tau\alpha, \tau\gamma) d\alpha d\gamma, \quad (2.13)$$

where $(1 - \gamma^2)^{1/3} b_\partial(y, z, \alpha, \gamma, \tau) = a(0, y, z, \alpha, \gamma, \tau) (\partial_x \zeta)(0, \alpha, \gamma) + \tau^{-1} \partial_x b(0, y, z, \alpha, \gamma, \tau)$. As $a_0 := a|_{x=0}$ is elliptic and as $\partial_x \zeta|_{x=0} = (1 - \gamma^2)^{1/3} \frac{1}{\tilde{\alpha}^{1/3}} (\partial_\rho \tilde{\zeta})(\rho)|_{\rho=\frac{1}{\tilde{\alpha}}}$ for $\tilde{\alpha} = \frac{\alpha}{\sqrt{1-\gamma^2}}$ on the support of χ_ε then $b_\partial(y, z, \tilde{\alpha}, \tau) := \frac{a_0}{\tilde{\alpha}^{1/3}} (\partial_\rho \tilde{\zeta})(\rho)|_{\rho=\frac{1}{\tilde{\alpha}}}$ is elliptic, close to 1 on the support of the symbol. Replacing \widehat{F}_τ by (2.10) and using the Wronskian relation $A'(z)A_+(z) - A'_+(z)A(z) = ie^{-i\pi/3}$ allows to conclude. \square

In the remaining of this section we show that, if $j \leq j(s, h)$, similar integral formulas hold for each $w_{j,gl}$; moreover we explicitly compute the corresponding functions $F_{j,\tau}$ to determine $\mathcal{F}(\partial_x u_{j,gl}^+)(P, Q_0, \tau)$, $P \in \partial\Omega$. For $j = 0$ we may follow closely the approach in [7, Section 3.1.1] (as the glancing order contact is exactly 2) to provide a detailed proof. Let $1 \leq j \leq j(s, h)$: we use the explicit form of $w_{j,gl}$ and write it as an oscillatory integral involving the Airy function $A(\tau^{2/3} \zeta)$ and its derivative. After the changes of variables $\alpha = \sqrt{1 - \gamma^2} \tilde{\alpha}$ and $\tilde{z} = \phi(x, \tilde{y}, 0, s) z_1$ in (1.11), $w_{j,gl}(Q, Q_0, \tau)$ becomes

$$\frac{\tau^2}{(2\pi)^2} \frac{i\tau}{4\pi} \int \frac{\psi_j(1 - \gamma^2)}{\sqrt{1 + z_1^2}} \chi_\varepsilon(\tilde{\alpha}) \sqrt{1 - \gamma^2} e^{i\tau(z\gamma + \sqrt{1-\gamma^2}(y-\tilde{y})\tilde{\alpha} - \phi(x, \tilde{y}, 0, s)(z_1\gamma + \sqrt{1+z_1^2}))} d\tilde{\alpha} d\gamma d\tilde{y} dz_1.$$

The critical point w.r.t. z_1 satisfies $\gamma + \frac{z_1}{\sqrt{1+z_1^2}} = 0$ hence $1 + z_1^2 = \frac{1}{1-\gamma^2}$. Set $z_1 = -\sqrt{\frac{w^2}{1-\gamma^2} - 1}$, then the phase is stationary w.r.t. w at $w = 1$ and at this point, the second order derivative of the phase equals $\tau \phi(x, \tilde{y}, 0, s) \sqrt{1 - \gamma^2} \sim 2^{-j} s/h$. As $j \leq j(s, h)$, then $2^{-j} \geq 2^{-j(s, h)} \sim (s/h)^{-1/3}$, hence $2^{-j} s/h \gtrsim (s/h)^{2/3}$ with $s \geq \sqrt{2}$. For w near 1 the stationary phase applies and the critical value of the phase depending of z_1 becomes $-\phi(x, \tilde{y}, 0, s) \sqrt{1 - \gamma^2}$. For $w \notin [1/\sqrt{2}, \sqrt{2}]$ we perform integrations by parts with large parameter $\sim 2^{-j} s/h > (s/h)^{2/3}$. We obtain, modulo $O((h/2^{-j} s)^\infty)$ contributions,

$$w_{j,gl}(Q, Q_0, \tau) = C \tau^{2+1/2} \int \frac{\psi_j(1 - \gamma^2) \chi_\varepsilon(\tilde{\alpha})}{\phi^{1/2}(x, \tilde{y}, 0, s)} (1 - \gamma^2)^{\frac{1}{2} + \frac{1}{2} - \frac{1}{2} - \frac{1}{4}} e^{i\tau(z\gamma + \sqrt{1-\gamma^2}((y-\tilde{y})\tilde{\alpha} - \phi(x, \tilde{y}, 0, s)))} d\tilde{\alpha} d\gamma d\tilde{y}. \quad (2.14)$$

For $\tilde{\alpha}$ near 1, we can perform a suitable change of variable w.r.t. \tilde{y} such that the phase $\tilde{y}\tilde{\alpha} + \phi(x, \tilde{y}, 0, s)$ transforms into an Airy type phase function of the form $\sigma^3/3 + \sigma\tilde{\zeta}(\frac{1+x}{\tilde{\alpha}}) + \Gamma_0(\tilde{\alpha}, s)$, where $\tilde{\zeta}$ is the function defined in Lemma 2.5. Let $\varphi(x, \tilde{y}, \tilde{\alpha}, s) := \tilde{y}\tilde{\alpha} + \phi(x, \tilde{y}, 0, s)$. As $\partial_{\tilde{y}} \phi(x, \tilde{y}, 0, s) = -\frac{s(1+x)\cos \tilde{y}}{\phi}$, $\partial_{\tilde{y}}^2 \phi(x, \tilde{y}, 0, s) = \frac{s(1+x)\sin \tilde{y} - (\partial_{\tilde{y}} \phi)^2}{\phi^2}$, then $\partial_{\tilde{y}}^2 \varphi(x, \tilde{y}, \tilde{\alpha}, s) = 0$ when $\tilde{y} = y_*(x) := \arcsin(\frac{1+x}{s})$ and there $\partial_x \phi(x, \tilde{y}, 0, s)|_{y_*(x)} = 0$ and $\partial_{\tilde{y}} \phi(x, \tilde{y}, 0, s)|_{y_*(x)} = -(1+x)$. For \tilde{y} near $y_*(x)$ there are two critical points $y_\pm = y_\pm(x, \tilde{\alpha})$ satisfying

$$s(1+x)\sin(y_\pm) = \tilde{\alpha}^2 \pm \sqrt{s^2 - \tilde{\alpha}^2} \sqrt{(1+x)^2 - \tilde{\alpha}^2}, \phi(x, y_\pm, 0, s) = \sqrt{s^2 - \tilde{\alpha}^2} \mp \sqrt{(1+x)^2 - \tilde{\alpha}^2}. \quad (2.15)$$

Lemma 2.10. *Let $\tilde{y} = y_*(x) + Y$. There exists a unique change of variables $Y \mapsto \sigma$ which is smooth and satisfying $\frac{dY}{d\sigma} \notin \{0, \infty\}$ such that, for $\tilde{\zeta}$ given by Lemma 2.5, we have*

$$\varphi(x, y_*(x) + Y, \tilde{\alpha}, s) = \frac{\sigma^3}{3} + \sigma \tilde{\alpha}^{2/3} \tilde{\zeta}\left(\frac{1+x}{\tilde{\alpha}}\right) + \Gamma_0(\tilde{\alpha}, s), \quad (2.16)$$

where $\Gamma_0(\tilde{\alpha}, s) := \sqrt{s^2 - 1} + \arcsin(\frac{1}{s})\tilde{\alpha} + \frac{(1-\tilde{\alpha})^2}{2\sqrt{s^2-1}}(1 + O(1-\tilde{\alpha}))$ for $\tilde{\alpha}$ near 1.

Proof. As the phase φ has degenerate critical points of order exactly two, it follows from [2] that there exists a unique change of variables $Y \mapsto \sigma$ which is smooth and satisfying $\frac{dY}{d\sigma} \notin \{0, \infty\}$ and that there exist smooth functions $\zeta^\#(x, \tilde{\alpha}, s)$ and $\Gamma(x, \tilde{\alpha}, s)$ such that

$$\varphi(x, y_*(x) + Y, \tilde{\alpha}, s) = \frac{\sigma^3}{3} + \sigma \zeta^\#(x, \tilde{\alpha}, s) + \Gamma(x, \tilde{\alpha}, s). \quad (2.17)$$

As the change of coordinates is regular the critical points $Y_\pm := y_\pm(x, \tilde{\alpha}) - y_*(x)$ of φ must correspond to $\sigma_\pm = \pm \sqrt{-\zeta^\#(x, \tilde{\alpha}, s)}$. Write $\zeta^\#(x, \tilde{\alpha}, s) := \tilde{\alpha}^{2/3} \tilde{\zeta}^\#(\frac{1+x}{\tilde{\alpha}}, \tilde{\alpha}, s)$. We will show that $\tilde{\zeta}^\#$ satisfies the same

equation as $\tilde{\zeta}$ in (2.5). As the critical values of the two functions in (2.17) must coincide, we have

$$\varphi(x, y_*(x) + Y_{\pm}, \tilde{\alpha}, s) = \mp \frac{2}{3} (-\tilde{\zeta}^{\#})^{\frac{3}{2}}(x, \tilde{\alpha}, s) + \Gamma(x, \tilde{\alpha}, s), \quad (2.18)$$

from which we deduce $\frac{4}{3}\tilde{\alpha}(-\tilde{\zeta}^{\#})^{\frac{3}{2}}(\frac{1+x}{\tilde{\alpha}}, x, s) = \varphi(x, y_-, \tilde{\alpha}, s) - \varphi(x, y_+, \tilde{\alpha}, s)$. Taking the derivative with respect to x in the last equation yields (with $y_{\pm} = y_*(x) + Y_{\pm}$)

$$\begin{aligned} 2(-\partial_x \tilde{\zeta}^{\#})(-\tilde{\zeta}^{\#})^{\frac{1}{2}} &= \partial_x \phi(x, y_*(x) + Y_-, 0, s) - \partial_x \phi(x, y_*(x) + Y_+, 0, s) \\ &\quad - \partial_x y_+ \partial_y \varphi(x, y_+, \tilde{\alpha}, s) + \partial_x y_- \partial_y \varphi(x, y_-, \tilde{\alpha}, s). \end{aligned} \quad (2.19)$$

The last two terms in the second line of (2.19) vanish as y_{\pm} are the critical points of the function φ with respect to y ; for the same reason we have that $\partial_y \phi(x, y_{\pm}(x, \tilde{\alpha}), 0, s) = -\tilde{\alpha}$. As $\phi(x, y, 0, s)$ satisfies the eikonal equation $(\partial_x \phi)^2(x, y, 0, s) + \frac{1}{(1+x)^2} (\partial_y \phi)^2(x, y, 0, s) = 1$, then $(\partial_x \phi(x, y_{\pm}(x), 0, s))^2 = 1 - \frac{\tilde{\alpha}^2}{(1+x)^2}$. Moreover, $\partial_x \phi|_{y_{\pm}} = \frac{s}{\phi(x, y_{\pm}, 0, s)} (\tilde{\rho} - \sin(y_{\pm}))$ (with $\tilde{\rho} = \frac{1+x}{s}$) which is non positive in the “ y_+ case” and positive in the “ y_- case”. Eventually we obtain, using (2.19) and the right signs of $\partial_x \phi$, $-\tilde{\zeta}^{\#} [-\partial_x \tilde{\zeta}^{\#}]^2 = 1 - \frac{\tilde{\alpha}^2}{(1+x)^2}$, which is the same equation as in Lemma 2.5 with $\rho = \frac{1+x}{\tilde{\alpha}} = (1+x) \frac{\sqrt{1-\gamma^2}}{\alpha}$. As the degenerate critical point occurs at $\sigma = 0$, hence at $\zeta^{\#} = 0$, we deduce by uniqueness of the solution that $\tilde{\zeta}^{\#} = \tilde{\zeta} = \tilde{\zeta}(\frac{1+x}{\tilde{\alpha}})$.

Next, we compute the explicit form of the function $\Gamma(x, \tilde{\alpha}, s)$. Taking the sum in (2.18) gives $\Gamma(x, \tilde{\alpha}, s) = \frac{1}{2}(\varphi(x, y_+(x), \tilde{\alpha}, s) + \varphi(x, y_-(x), \tilde{\alpha}, s))$; taking the derivative w.r.t. x yields $\partial_x \Gamma(x, \tilde{\alpha}, s) = 0$. As such, Γ is independent of x and we define $\Gamma_0(\tilde{\alpha}, s) := \Gamma(0, \tilde{\alpha}, s)$, then

$$\Gamma_0(\tilde{\alpha}, s) = \frac{1}{2}((y_+ + y_-)\tilde{\alpha} + \phi(0, y_+, 0, s) + \phi(0, y_-, 0, s)),$$

where $y_{\pm} = y_{\pm}(0)$. For small $x \geq 0$ and for y in a neighborhood of $y_* = y_*(0)$, y remains sufficiently close to $y_*(x)$: shrinking the support if necessary, we may assume $|y - y_*(x)| < 1/2$. For $|y_{\pm} - y_*| < 1/2$ we may compute, using (2.15) with $x = 0$, the first approximation of y_{\pm} : we have

$$y_{\pm} = \arcsin\left(\frac{\tilde{\alpha}^2}{s} \pm \sqrt{1 - \tilde{\alpha}^2} \sqrt{1 - \frac{\tilde{\alpha}^2}{s^2}}\right), \quad y_* = \arcsin\left(\frac{1}{s}\right). \quad (2.20)$$

As $\Gamma_0(\tilde{\alpha}, s) = \frac{1}{2}(\varphi(0, y_+, \tilde{\alpha}, s) + \varphi(0, y_-, \tilde{\alpha}, s))$ and $\partial_y \varphi|_{y_{\pm}} = 0$ then $\partial_{\tilde{\alpha}} \Gamma_0 = \frac{1}{2}(y_+ + y_-) + \frac{1}{2} \sum_{\pm} \partial_{\tilde{\alpha}} y_{\pm} \partial_y \varphi|_{y_{\pm}} = \frac{1}{2}(y_+ + y_-)$. This yields $\Gamma_0(1, s) = \sqrt{s^2 - 1} + \arcsin \frac{1}{s}$ and $\partial_{\tilde{\alpha}} \Gamma_0(1, s) = \arcsin(1/s)$. We need the higher order derivatives: using (2.20), it follows that $(y_+ + y_-)$ reads as an asymptotic expansion of even powers of $\sqrt{1 - \tilde{\alpha}^2}$ and with main term $\arcsin(\frac{\tilde{\alpha}^2}{s})$. We find, with $Z_{\pm} = \frac{\tilde{\alpha}^2}{s} \pm \sqrt{1 - \tilde{\alpha}^2} \sqrt{1 - \frac{\tilde{\alpha}^2}{s^2}}$, $Z_{\pm}|_{\tilde{\alpha}=1} = \frac{1}{s}$,

$$\frac{1}{2} \partial_{\tilde{\alpha}} (y_+ + y_-) = \frac{\tilde{\alpha}}{s} \left(\frac{1}{\sqrt{1 - Z_+^2}} + \frac{1}{\sqrt{1 - Z_-^2}} \right) - \frac{\tilde{\alpha}(s^2 + 1 - 2\tilde{\alpha}^2)}{2s^2 \sqrt{1 - \tilde{\alpha}^2} \sqrt{1 - \frac{\tilde{\alpha}^2}{s^2}}} \left(\frac{1}{\sqrt{1 - Z_+^2}} - \frac{1}{\sqrt{1 - Z_-^2}} \right).$$

As $\left(\frac{1}{\sqrt{1 - Z_+^2}} - \frac{1}{\sqrt{1 - Z_-^2}} \right) = \frac{Z_+^2 - Z_-^2}{\sqrt{1 - Z_+^2} \sqrt{1 - Z_-^2} (\sqrt{1 - Z_+^2} + \sqrt{1 - Z_-^2})}$ and $Z_+^2 - Z_-^2 = 4 \frac{\tilde{\alpha}^2}{s} \sqrt{1 - \tilde{\alpha}^2} \sqrt{1 - \frac{\tilde{\alpha}^2}{s^2}}$,

$$\frac{1}{2} \partial_{\tilde{\alpha}} (y_+ + y_-) = \frac{\tilde{\alpha}}{s} \left(\frac{1}{\sqrt{1 - Z_+^2}} + \frac{1}{\sqrt{1 - Z_-^2}} \right) - \frac{2\tilde{\alpha}^3 (s^2 + 1 - 2\tilde{\alpha}^2)}{s^3 \sqrt{1 - Z_+^2} \sqrt{1 - Z_-^2} (\sqrt{1 - Z_+^2} + \sqrt{1 - Z_-^2})}.$$

At $\tilde{\alpha} = 1$ we obtain $\partial_{\tilde{\alpha}}^2 \Gamma_0(1, s) = \frac{1}{2} \partial_{\tilde{\alpha}} (y_+ + y_-)|_{\tilde{\alpha}=1} = \frac{1}{\sqrt{s^2 - 1}}$. In the same way we notice that all the higher order derivatives of Γ_0 come with a factor $\frac{1}{\sqrt{s^2 - 1}}$. The proof is achieved. \square

After the changes of coordinates $\tilde{y} = y_*(x) + Y$, $Y \rightarrow \sigma$, $\sigma = (\tau \sqrt{1 - \gamma^2})^{-1/3} \tilde{\sigma}$ we obtain $w_{j,gl}(Q, Q_0, \tau)$ as follows (with $Y = Y(\sigma) = Y((\tau \sqrt{1 - \gamma^2})^{-1/3} \tilde{\sigma})$)

$$\tau^{2+\frac{1}{2}-\frac{1}{3}} \int \frac{\psi_j(1 - \gamma^2)(1 - \gamma^2)^{\frac{1}{4}-\frac{1}{6}} \chi_{\varepsilon_1}(\tilde{\alpha}) dY}{\phi^{1/2}(x, y_*(x) + Y, 0, s)} e^{i\tau(z\gamma + \sqrt{1 - \gamma^2}(y\tilde{\alpha} - \Gamma_0(\tilde{\alpha}, s)))} e^{-i(\frac{\tilde{\sigma}^3}{3} + \tilde{\sigma}(\tau \sqrt{1 - \gamma^2})^{2/3} \tilde{\zeta}(\frac{1+x}{\tilde{\alpha}}))} d\tilde{\sigma} d\tilde{\alpha} d\gamma.$$

At this point we let again $\alpha = \sqrt{1 - \gamma^2} \tilde{\alpha}$. Following [2], we integrate by parts in $\tilde{\sigma}$ and apply the Malgrange theorem to write $w_{j,gl}$ under the form $w_{j,gl} = T_{j,\tau}(F_{j,\tau})$, where the operator $T_{j,\tau}$ has the same phase as T_τ and symbols a_j, b_j which are asymptotic expansions with small parameter $h/2^{-j}s$ and where the function $\widehat{F}_{j,\tau}$ has phase $-\tau\sqrt{1 - \gamma^2}\Gamma_0(\frac{\alpha}{\sqrt{1 - \gamma^2}}, s)$ and symbol $\frac{\tau^{\frac{1}{2} - \frac{3}{4}}}{(s^2 - 1)^{1/4}}(1 - \gamma^2)^{\frac{1}{2} - \frac{1}{2}}\psi_j(1 - \gamma^2)f_j$, where f_j is an asymptotic expansion with parameter $h/2^{-j}s$. Notice that, if for $j = 0$ the powers of $(1 - \gamma^2)$ in play no role in (2.10) or in (2.12) as $\psi_0(1 - \gamma^2)$ is supported in $[\frac{1}{64}, 2]$, for $1 \leq j \leq j(s, h)$ it is essential to keep track of them.

2.2. The "non-glancing" parts of u_j^+ , $0 \leq j \leq j(s, h)$. In this section we describe the form of $u_{j,he,h}^+$ whose incoming part equals $u_{free,j,he,h}^+ := \int e^{i\tau\chi(h\tau)w_{j,he}}d\tau$. We obtain as before $w_{j,he}(Q, Q_0, \tau)$ under the form (2.14) but where $\chi_\varepsilon(\tilde{\alpha})$ is now replaced by $(1 - \chi_\varepsilon(\tilde{\alpha}))$. The phase $\tau(z\gamma + \sqrt{1 - \gamma^2}((y - \tilde{y})\tilde{\alpha} - \phi(x, \tilde{y}, 0, s)))$ has two critical points $y_\pm(x, \tilde{\alpha})$ satisfying (2.15) such that $|y_+(x, \tilde{\alpha}) - y_-(x, \tilde{\alpha})| \gtrsim \varepsilon$, as $\tilde{\alpha}$ stays away from a fixed neighborhood of 1 on the support of $1 - \chi_\varepsilon$ and it is stationary with respect to $\tilde{\alpha}$ when $\tilde{y} = y$. The stationary phase applies with large parameter $\tau\sqrt{1 - \gamma^2} \sim 2^{-j}/h$ and gives, modulo $O((h2^j)^\infty)$ terms,

$$w_{j,he}(Q, Q_0, \tau) = \tau^{1+1/2} \int \frac{\psi_j(1 - \gamma^2)(1 - \chi_\varepsilon(\partial_y \phi(x, y, 0, s)))}{\sqrt{\phi(x, y, 0, s)}} (1 - \gamma^2)^{\frac{1}{4} - \frac{1}{2}} \tilde{\sigma}_{free,j,he}^\pm e^{i\tau(z\gamma - \sqrt{1 - \gamma^2}\phi(x, y, 0, s))} d\gamma. \quad (2.21)$$

Recall from (2.15) that $\phi(x, y_\pm(x, \tilde{\alpha}), 0, s) = \sqrt{s^2 - \tilde{\alpha}^2} \mp \sqrt{(1+x)^2 - \tilde{\alpha}^2}$. Here $\tilde{\sigma}_{free,j,he}^\pm$ are classical symbols that read as asymptotic expansion with small parameter $h2^j$. Let now $1 - \gamma^2 = 2^{-2j}\varphi^2$, then $\varphi \sim 1$ on the support of $\psi_j(2^{-2j}\varphi^2) = \psi(\varphi^2)$ and $d\gamma/d\varphi \sim 2^{-2j}$. The phase $\tau(z\sqrt{1 - 2^{-2j}\varphi^2} - 2^{-j}\varphi\phi(x, y, 0, s))$ is stationary when $2^{-j}(-z) \sim \phi(x, y, 0, s)$ and its second order derivative equals $\tau(-z)2^{-2j}/\sqrt{1 - 2^{-2j}\varphi^2}$. At the critical points $\tau(-z)2^{-2j} \sim 2^{-j}s/h \gtrsim (s/h)^{2/3}$, so the stationary phase yields, modulo $O((h/s)^\infty)$,

$$w_{j,he}(Q, Q_0, \tau) = \tau \tilde{\psi} \left(\frac{\phi(x, y, 0, s)}{2^{-j}(-z)} \right) \frac{(1 - \chi_\varepsilon(\partial_y \phi(x, y, 0, s)))}{\phi(x, y, 0, s)} 2^{-2j+j/2+j/2} \sigma_{free,j,he}^\pm e^{-i\tau\phi(x, y, z, s)}, \quad (2.22)$$

where $\tilde{\psi}$ is a smooth cutoff supported near 1, equal to 0 near 0 and such that $\tilde{\psi} = 1$ on the support of ψ . The symbols $\sigma_{free,j,he}^\pm$ are asymptotic expansions with main contribution $\tilde{\sigma}_{free,j,he}^\pm$ and small parameter $h2^j/s$. If we denote $\Sigma_{free,j}$ the factor of $e^{-i\tau\phi(x, y, z, s)}$ in (2.22), then $u_{free,j,he,h}^+ = \int e^{i\tau(t - \phi(x, y, z, s))} \chi(h\tau) \Sigma_{free,j} d\tau$. After the reflection on the boundary, the solution to the wave equation with Dirichlet boundary condition reads as $\int e^{i\tau(t - \phi_R(x, y, z, s))} \chi(h\tau) \Sigma_{R,j} d\tau$, where ϕ_R satisfies the eikonal equation (2.6) and the boundary condition $\phi_R|_{x=0} = \phi|_{x=0}$ and $\partial_x \phi_R|_{x=0} = -\partial_x \phi|_{x=0}$. The symbol $\Sigma_{R,j}$ is an asymptotic expansion with small parameter $(\tau 2^{-j}s)^{-1}$ that reads as $\Sigma_{R,j}(\cdot, \tau) = \sum_k \tau^{-k} \Sigma_{R,k}$, where $\Sigma_{R,k}$ solve a system of the transport equations and $\Sigma_{R,j}|_{x=0} = \Sigma_{free,j}|_{x=0}$. We obtain $\partial_x u_{j,he,h}^+|_{x=0} = \int e^{i\tau(t - \phi(0, y, z, s))} \chi(h\tau) (-i)\tau \Sigma_j(y, z, s, \tau) d\tau$, where Σ_j is a classical symbol that reads as an asymptotic expansion with small parameters $\tau^{-1}, (\tau 2^j s)^{-1}$ and whose main contribution equals $2i\partial_x \phi(0, y, z, s) \Sigma_j|_{x=0}$.

Remark 2.11. *On the support of $1 - \chi_\varepsilon$ we have $1 - \partial_y \phi|_{x=0} \gtrsim \varepsilon$: from the eikonal equation, we obtain the following lower bound: $(\partial_x \phi)^2|_{x=0} = (1 - (1+x)^{-2}(\partial_y \phi)^2 - (\partial_z \phi)^2)|_{x=0} \geq c(\varepsilon)$, where $c(\varepsilon) > 0$ depends only on ε . As $\partial_x \phi|_{x=0} = \frac{1-s \sin y}{\phi(0, \theta, z, s)}$, this implies $s|\sin y_* - \sin y| \geq c(\varepsilon)\phi(0, y, z, s)$, where $y_* = \arcsin(1/s)$.*

For all $0 \leq j \leq j(s, h)$ we eventually find, for all $P = (0, y, z) \in \partial\Omega$,

$$\partial_x u_{j,he,h}^+(P, Q_0, t) = \int e^{i\tau(t - \phi(0, y, z, s))} \chi(h\tau) \frac{\tau^2}{\psi(0, y, z, s)} 2^{-j} \sigma_{j,he}(y, z, s, \tau) d\tau, \quad (2.23)$$

where $\sigma_{2^{-2j},he}$ is an asymptotic expansion with small parameters $\tau^{-1}, (\tau 2^{-3j}s)^{-1}$ supported for $s|\sin y_* - \sin y| \geq c(\varepsilon)\phi(0, y, z, s)$ and $2^{-j}(-z) \sim \phi(0, y, 0, s)$.

3. HIGH-FREQUENCY CASE. DISPERSIVE ESTIMATES WHEN $d(Q_0, \partial\Omega) \geq \sqrt{2} - 1$

3.1. Dispersion for the glancing part when $d(Q, \partial\Omega) \geq \sqrt{2} - 1$. Let $Q_0 = (s, 0, 0)$, $Q = ((1 + x_Q) \sin y_Q, (1 + x_Q) \cos y_Q, z_Q)$ in Ω , and assume $s \geq r := 1 + x_Q \geq \sqrt{2}$. We prove the following :

Proposition 3.1. *There exists $C > 0$ such that for all $t > h$, the following holds uniformly with respect to Q, Q_0 such that $s \geq r \geq \sqrt{2}$ where $s = 1 + x_{Q_0}$, $r = 1 + x_Q$: $\sum_{0 \leq j \leq j(s,h)} |u_{j,gl,h}^\#(Q, Q_0, t)| \leq \frac{C}{h^{2t}}$.*

Proof. We write the details of the proof for $j = 0$ while keeping track of the factors $\sqrt{1 - \gamma^2}$. The proof of dispersive bounds for $1 \leq j \leq j(s, h)$ will follow exactly in the same way as all stationary arguments follow for such values of j and we will be able to sum up all the contributions as these bounds have additional non-positive powers of 2^j . Let $j = 0$ and set $I_{0,gl}(Q, Q_0, \tau) := \int_{P \in \partial\Omega} \frac{\mathcal{F}(\partial_x u_{j,gl}^+(P, Q_0, \tau))}{4\pi\tau|P-Q|} e^{-i\tau|P-Q|} d\sigma(P)$. Then

$$u_{0,gl,h}^\#(Q, Q_0, t) = \frac{1}{4\pi} \int \chi(h\tau) e^{it\tau} \tau I_{0,gl}(Q, Q_0, \tau) d\tau. \quad (3.1)$$

Writing $|P - Q| = \phi(x_Q, y - y_Q, z - z_Q, 1)$ for a point $P = (\sin y, \cos y, z)$ on the boundary $\partial\Omega$ and replacing (2.12) in (2.4) we find, after the change of coordinates $\alpha = \sqrt{1 - \gamma^2} \tilde{\alpha}$,

$$\begin{aligned} I_{0,gl}(Q, Q_0, \tau) &= \int \tau^{-1+2+\frac{5}{8}} e^{i\tau(z\gamma + \sqrt{1-\gamma^2}(y\tilde{\alpha} - \Gamma_0(\tilde{\alpha}, s)) - \phi(x_Q, y - y_Q, z - z_Q, 1))} \\ &\times \frac{f(\alpha, \gamma, \tau) b_\partial(y, z, \tilde{\alpha}, \tau)}{\phi(x_Q, y - y_Q, z - z_Q, 1)} \frac{(1-\gamma^2)^{-\frac{5}{12} + \frac{1}{3} + \frac{1}{2}} \psi_0(1-\gamma^2) \chi_\varepsilon(\tilde{\alpha})}{(s^2-1)^{1/4} A_+(\tau^{\frac{2}{3}} \zeta_0(\alpha, \gamma))} d\tilde{\alpha} d\gamma dy dz. \end{aligned} \quad (3.2)$$

Lemma 3.2. *There exists a constant $C > 0$ such that $|I_{0,gl}(Q, Q_0, \tau)| \leq C/t$ uniformly with respect to Q, Q_0 and t such that $\sqrt{s^2 - 1 + z_Q^2} \sim t$. Moreover, for $\frac{t}{\sqrt{s^2 - 1 + z_Q^2}} \notin [1/4, 4]$, we have $|I_{0,gl}(Q, Q_0, \tau)| \leq \frac{C}{\sqrt{s^2 - 1}}$.*

If $\frac{t}{\sqrt{s^2 - 1 + z_Q^2}} \in [1/4, 4]$, the estimate of Proposition 3.1 follows using (3.1) and Lemma 3.2. If not, the phase of (3.1) is not stationary w.r.t. τ and we proceed by integrations by parts which give at most $O(h^\infty/t)$. \square

Proof. (Proof of Lemma 3.2) We apply the stationary phase with respect to z in the integral (3.2): let $r = 1 + x_Q$ and set $z = z_Q + \tilde{z} \sqrt{1 + r^2 - 2r \cos(y_Q - y)}$. As $r \geq \sqrt{2}$, this is well defined and $dz/d\tilde{z} = \phi(x_Q, y - y_Q, 0, 1)$. As $\phi(x_Q, y - y_Q, z - z_Q, 1) = \phi(x_Q, y - y_Q, 0, 1) \sqrt{1 + \tilde{z}^2}$ the phase of $I_{0,gl}$ becomes $\tau(z_Q \gamma - \sqrt{1 - \gamma^2}(-y\tilde{\alpha} + \Gamma_0(\tilde{\alpha}, s)) + \phi(x_Q, y - y_Q, 0, 1)(\tilde{z}\gamma - \sqrt{1 + \tilde{z}^2}))$ and its critical point with respect to \tilde{z} satisfies $\tilde{z} = \frac{\gamma}{\sqrt{1 - \gamma^2}}$. As, in case j large, this value is large, we renormalize \tilde{z} by taking $\tilde{z} = \sqrt{\frac{w^2}{1 - \gamma^2} - 1}$; as such, the critical point is $w = 1$ and the second order derivative of the phase equals $\tau \phi(x_Q, y - y_Q, 0, 1) \sqrt{1 - \gamma^2}$. The stationary phase in w yields a factor $\tau^{-1/2} \times (1 - \gamma^2)^{-\frac{1}{2} - \frac{1}{4}}$ and the symbol $\tau^{1+5/6} \frac{b_\partial(1-\gamma^2)^{5/12}}{\phi(x_Q, y - y_Q, z - z_Q, 1)}$ becomes $\tau^{1+1/3} (1 - \gamma^2)^{\frac{5}{12} - \frac{3}{4}} \frac{\tilde{b}_\partial(y, z_Q, \alpha, \gamma, \tau)}{\phi^{1/2}(x_Q, y - y_Q, 0, 1)}$, where \tilde{b}_∂ has main contribution b_∂ . We obtain

$$\begin{aligned} I_{0,gl}(Q, Q_0, \tau) &= \tau^{\frac{4}{3}} \int e^{i\tau(z_Q \gamma - \sqrt{1-\gamma^2}(-y\tilde{\alpha} + \Gamma_0(\tilde{\alpha}, s) + \phi(x_Q, y - y_Q, 0, 1)))} \\ &(1 - \gamma^2)^{-1/3} \frac{\tilde{b}_\partial(y, Q, \tilde{\alpha}, \gamma, \tau)}{\phi^{1/2}(x_Q, y - y_Q, 0, 1)} \frac{f(\alpha, \gamma, \tau) \psi_0(1 - \gamma^2) \chi_\varepsilon(\tilde{\alpha})}{(s^2 - 1)^{1/4} A_+(\tau^{\frac{2}{3}} \zeta_0(\alpha, \gamma))} d\tilde{\alpha} d\gamma dy. \end{aligned} \quad (3.3)$$

The phase $\phi(x_Q, y - y_Q, 0, 1)$ has two degenerate critical points of order exactly two at $y = y_Q \pm \arccos(1/r)$, where $r = 1 + x_Q$. Near $y_Q - \arccos(1/r)$, its first order derivative equals -1 , hence for y near this point the phase of $I_{0,gl}$ is non-stationary w.r.t. y and repeated integrations by parts yield $O(\frac{\tau^{-\infty}}{\sqrt{s^2 - 1}})$. Let $y_c := y_Q + \arccos(1/r)$. Notice that, if $y \in [0, 2\pi)$ is sufficiently close to y_* on the support of $I_{0,gl}$ (say $|y - y_*| \leq \frac{\pi}{16}$) and is such that $|y - y_c| \geq \frac{\pi}{8}$, then $1 - \tilde{\alpha}$ has to be bounded from below by a fixed constant there where the phase of $I_{0,gl}$ is stationary w.r.t. y . Taking ε smaller if necessary, it follows that for such value of y outside a small, fixed neighborhood of y_c , $\tilde{\alpha}$ cannot belong to the support of $\chi_\varepsilon(\tilde{\alpha})$. We are reduced to studying the integral (3.3) for $|y - y_c| \leq \frac{\pi}{8} < 1$. Let $\varepsilon_1 > 0$ be small enough. We study separately the cases $|y - y_c| \leq \tau^{-1/3 + \varepsilon_1}$ and $\tau^{-1/3 + \varepsilon_1} \lesssim |y - y_c| \leq \frac{\pi}{8}$; to do that, we introduce a smooth cut-off χ_0 supported in $[-2, 2]$ and equal to 1 on $[-3/2, 3/2]$ and split $I_{0,gl} = I_{0,gl}^{\chi_0} + I_{0,gl}^{1-\chi_0}$, where $I_{0,gl}^{\chi_0}$ has the form (3.3) with additional cut-off $\chi((y - y_c)\tau^{1/3 - \varepsilon_1})$.

3.1.1. *Case* $\tau^{-1/3+\epsilon_1} \leq |y-y_c| \leq \frac{\pi}{8}$: *study of* $I_{1,gl}^{1-\chi_0}$. We set $\tilde{\alpha} = \tilde{\alpha}(\beta, \tau) := 1 - \tau^{-2/3}\beta$: as on the support of $\chi_{\epsilon_1}(\tilde{\alpha})$ we have $1 - \tilde{\alpha} \lesssim \epsilon$, it follows that $\tau^{-2/3}\beta \lesssim \epsilon$. This choice of coordinates is motivated by the behavior of the Airy factor $A_+(\tau^{2/3}\zeta_0(\alpha, \gamma))$: as $\tau^{2/3}\zeta_0(\alpha, \gamma) = \tau^{2/3}\alpha^{2/3}\tilde{\zeta}(\frac{\sqrt{1-\gamma^2}}{\alpha}) = \tau^{2/3}\sqrt{1-\gamma^2}^{2/3}\tilde{\alpha}^{2/3}\tilde{\zeta}(\frac{1}{\tilde{\alpha}})$, then

$$\tau\alpha(-\tilde{\zeta})^{3/2}(\frac{1}{\tilde{\alpha}}) = \sqrt{2}\sqrt{1-\gamma^2}\beta^{3/2}(1 + O(\tau^{-2/3}\beta)), \quad (3.4)$$

where we used Lemma 2.5. As such, for $(\sqrt{2}\sqrt{1-\gamma^2}\beta^{3/2})^{2/3}$ large enough, $A_+(\tau^{2/3}\zeta_0(\alpha, \gamma))$ does oscillate, while for $(\sqrt{2}\sqrt{1-\gamma^2}\beta^{3/2})^{2/3}$ bounded it may be brought into the symbol. Write $1 = \chi_0(\beta) + (1-\chi_0)(\beta)$. On the support of $1-\chi_0(\beta)$ the Airy factor may oscillate and the phase function of $I_{0,gl}^{1-\chi_0}$ equals $z_Q\gamma - \sqrt{1-\gamma^2}\varphi$, where we have set

$$\varphi(y, \tilde{\alpha}, r) := -y\tilde{\alpha} + \Gamma_0(\tilde{\alpha}, s) + \phi(x_Q, y - y_Q, 0, 1) - \frac{2}{3}(-\tilde{\zeta})^{3/2}(\frac{1}{\tilde{\alpha}}). \quad (3.5)$$

With φ defined in (3.5) we have

$$I_{0,gl}^{1-\chi_0}(Q, Q_0, \tau) = \tau^{\frac{4}{3}-\frac{2}{3}} \int e^{-i\tau(z_Q\gamma - \sqrt{1-\gamma^2}\varphi)} \tilde{\chi}_\epsilon(1 - \tau^{-2/3}\tilde{\beta})(1 - \chi_0)((y - y_c)\tau^{1/3-\epsilon_1}) \\ \beta^{1/4}(1 - \gamma^2)^{\frac{1}{12}-\frac{1}{3}}(\tilde{b}f)(y, \beta, \gamma, \tau) \times \frac{\psi_0(1 - \gamma^2)}{(s^2 - 1)^{1/4}\phi^{1/2}(x_Q, y - y_Q, 0, 1)} dy d\gamma d\beta, \quad (3.6)$$

where the factor $\beta^{1/4}(1 - \gamma^2)^{1/12}$ comes from the Airy term A_+^{-1} (using (3.4)).

Lemma 3.3. *Let* $y = y_c + Y$, *where* $y_c = y_Q + \arccos(1/r)$. *There exists a unique change of variables* $Y \mapsto \sigma$ *which is smooth and satisfying* $\frac{dY}{d\sigma} \notin \{0, \infty\}$ *such that, for* $\tilde{\zeta}$ *given by Lemma 2.5, we have*

$$-(y_c + Y)\tilde{\alpha} + \phi(x_Q, y - y_Q + Y, 0, 1) = \frac{\sigma^3}{3} + \sigma\tilde{\alpha}^{2/3}\tilde{\zeta}(\frac{1}{\tilde{\alpha}}) + \tilde{\Gamma}(\tilde{\alpha}, r), \quad (3.7)$$

and where $\tilde{\Gamma}(\tilde{\alpha}, r) := \sqrt{r^2 - 1} - y_c\tilde{\alpha} + \frac{(1-\tilde{\alpha})^2}{2\sqrt{r^2-1}}(1 + O(1 - \tilde{\alpha}))$.

Proof. We proceed exactly as in the proof of Lemma 2.10 (where now $x = 0$ and s is replaced by r). As y_c is the degenerate critical point of order 2 of ϕ , there exist a smooth change of variable $Y \rightarrow \sigma$ and smooth phase functions $\zeta^\#$ and $\tilde{\Gamma}$ such that the LHS term in (3.7) reads as $\frac{\sigma^3}{3} + \sigma\zeta^\#(\tilde{\alpha}, r) + \tilde{\Gamma}(\tilde{\alpha}, r)$. Exactly as in Lemma 2.10 we obtain that $\zeta^\# = \tilde{\alpha}^{2/3}\tilde{\zeta}(\frac{1}{\tilde{\alpha}})$. It remains to determine $\tilde{\Gamma}$. The two critical points satisfy

$$r \cos(\arccos(1/r) + Y_\pm) = \tilde{\alpha}^2 \mp \sqrt{r^2 - \tilde{\alpha}^2}\sqrt{1 - \tilde{\alpha}^2}.$$

We have as before $\phi(x_Q, y_c + Y_\pm, 0, 1) = \sqrt{r^2 - \tilde{\alpha}^2} \pm \sqrt{1 - \tilde{\alpha}^2}$. As $\cos(\arccos(1/r) + Y) = \sin(\arcsin(1/r) - Y)$ we use the computations from Lemma 2.10 to determine $\arcsin(1/r) - \frac{1}{2}(Y_+ + Y_-)$. As $-y_c = -y_Q - \arccos(1/r) = -y_Q - \frac{\pi}{2} + \arcsin(1/r)$ we obtain $\tilde{\Gamma}(\tilde{\alpha}, r) = -(y_Q + \frac{\pi}{2})\tilde{\alpha} + \Gamma_0(\tilde{\alpha}, r)$ where $\Gamma_0(\tilde{\alpha}, r)$ is the same as in Lemma 2.10 and compute the derivatives of this new function at $\tilde{\alpha} = 1$ using those of Γ_0 as follows

$$\tilde{\Gamma}(1, r) = \sqrt{r^2 - 1} - y_c, \tilde{\Gamma}'(1, r) = -y_c, \tilde{\Gamma}''(1, r) = \frac{1}{\sqrt{r^2 - 1}}, \tilde{\Gamma}^{(k)} = \frac{c_k}{\sqrt{r^2 - 1}}(1 + O(\frac{1}{\sqrt{r^2 - 1}})).$$

□

Using the changes of variable $y \rightarrow y_c + Y$, $Y \rightarrow \sigma$ from Lemma 2.10 yields

$$\varphi(y, \tilde{\alpha}, r) = \frac{\sigma^3}{3} + \sigma\tilde{\alpha}^{2/3}\tilde{\zeta}(\frac{1}{\tilde{\alpha}}) + \tilde{\Gamma}(\tilde{\alpha}, r) + \Gamma_0(\tilde{\alpha}, s) - \frac{2}{3}(-\tilde{\zeta})^{3/2}(\frac{1}{\tilde{\alpha}}).$$

Let $y = y_c + Y$, $Y \rightarrow \sigma$ as in the Lemma 3.3 and set moreover $\sigma = \tau^{-1/3}w$: then $\tau^{\epsilon_1} \lesssim |w|$ on the support of the symbol (and if $|\tau^{-1/3}w| \geq \frac{\pi}{4}$ the integral defining $I_{0,gl}^{1-\chi_0}$ is $O(\tau^{-\infty})$). We apply the stationary phase in w near the critical points : let χ be a smooth cut-off supported in a fixed neighborhood of 1 and equal to 1 near 1 and set $\chi_\pm := \chi(\pm \frac{\sqrt{2\beta}}{w})$, $\tilde{\alpha} = 1 - \tau^{-2/3}\beta$; let also $\bar{\chi} := 1 - \chi_+ - \chi_-$. Write

$$I_{0,gl}^{1-\chi_0}(Q, Q_0, \tau) = \sum_{\chi \in \{\chi_\pm, \bar{\chi}\}} I_{0,gl}^{1-\chi_0, \chi},$$

where $I_{0,gl}^{1-\chi_0, \chi}$ are given by (3.6) with additional cutoffs $\chi(\frac{\sqrt{2\beta}}{w})$.

Lemma 3.4. *For w in a small, fixed neighborhood of $\pm\sqrt{2\beta}$, we have*

$$I_{0,gl}^{1-\chi_0,\chi_\pm}(Q, Q_0, \tau) = \tau^{4/3-2/3-1/3} \int e^{-i\tau(z_Q\gamma - \sqrt{1-\gamma^2}\varphi_\pm)} \chi_\varepsilon(1 - \tau^{-2/3}\beta) \Sigma_\pm(\beta, \gamma, \tau) (1 - \gamma^2)^{\frac{1}{12} - \frac{1}{3} - \frac{1}{4}} \\ \times \frac{(1 - \chi_0)(Y(\tau^{-1/3}w)\tau^{1/3-\varepsilon_1})\psi_0(1 - \gamma^2)}{(s^2 - 1)^{1/4}\phi^{1/2}(x_Q, \arccos(1/r) + Y(\tau^{-1/3}w_\pm), 0, 1)} d\gamma d\beta, \quad (3.8)$$

where $\varphi_\pm := \mp\frac{2}{3}(-\tilde{\zeta})^{3/2}(\frac{1}{\alpha}) + \tilde{\Gamma}(\tilde{\alpha}, r) + \Gamma_0(\tilde{\alpha}, s) - \frac{2}{3}(-\tilde{\zeta})^{3/2}(\frac{1}{\alpha})|_{\tilde{\alpha}=1-\tau^{-2/3}\beta}$. Here Σ_\pm are asymptotic expansions with parameter $\tau^{-\varepsilon_1}$ and main contribution $\frac{dY}{d\sigma} \frac{\beta^{1/4}}{\sqrt{|w_\pm|}} \chi_\pm(\frac{\sqrt{2\beta}}{w_\pm}) \Sigma(y_c + \tau^{-1/3}w_\pm, \beta, \tau)$.

Proof. The critical points are $w_\pm := \pm\tau^{1/3}\sqrt{-\tilde{\zeta}(\frac{1}{1-\tau^{-2/3}\beta})}$ which gives $w_\pm = \pm\sqrt{2\beta}(1 + O(\sqrt{2\tau^{-2/3}\beta}))$. As $|w| \geq \tau^{\varepsilon_1}$ on the support of $(1 - \chi_0)(Y(\tau^{-1/3}w)\tau^{1/3-\varepsilon_1})$ and $\frac{w}{\sqrt{2\beta}} \sim \pm 1$ on the support of the symbol, we also have $\sqrt{2\beta} \gtrsim \tau^{\varepsilon_1}$. At w_\pm , the second order derivative of the phase equals $\partial_w^2(\tau\sqrt{1-\gamma^2}\varphi)|_{w_\pm} = \sqrt{1-\gamma^2}w_\pm(1 + O(\tau^{-1/3}w_\pm))$. As $\sqrt{1-\gamma^2} \geq 1/8$ on the support of ψ_0 and $|w| \geq \tau^{\varepsilon_1}$ on the support of $(1 - \chi_0)(w\tau^{-\varepsilon_1})$, it follows that $\sqrt{1-\gamma^2} \times w_\pm \gtrsim \tau^{\varepsilon_1}$ and the stationary phase applies at w_\pm with a parameter larger than τ^{ε_1} . The exponent of the factor $(1 - \gamma^2)$ is $1/12 - 1/3 - 1/4 = -1/2$. The factor $\tau^{-1/3}$ before the integral (3.8) comes from the change of variables $\sigma \rightarrow w$. (For $1 \leq j \leq j(s, h)$, replace τ by τ^{-j}). \square

We now consider the integral $I_{0,gl}^{1-\chi_0,\bar{\chi}}(Q, Q_0, \tau)$ whose symbol is supported for $\frac{w^2}{2\beta} \notin [1/2, 3/2]$.

Lemma 3.5. *The stationary phase applies in γ with large parameter τ and yields*

$$I_{0,gl}^{1-\chi_0,\bar{\chi}}(Q, Q_0, \tau) = \tau^{\frac{4}{3}-1-\frac{1}{2}} \int e^{-i\tau\sqrt{\varphi^2+z_Q^2}} \chi_\varepsilon(1 - \tau^{-2/3}\beta) (1 - \chi_0)(|w|\tau^{-\varepsilon_1}) \\ \beta^{1/4} \bar{\chi}\left(\frac{\sqrt{2\beta}}{w}\right) \Sigma(y, \beta, \tau) \left(\frac{\varphi^2}{\varphi^2+z_Q^2}\right)^{\frac{1}{2}} \frac{1}{\varphi^{1/2}} \times \frac{\psi_0\left(\frac{\varphi^2}{\varphi^2+z_Q^2}\right)}{(s^2 - 1)^{1/4}\phi^{1/2}(x_Q, y - y_Q, 0, 1)} d\beta dy, \quad (3.9)$$

where Σ is an asymptotic expansion with small parameter τ^{-1} and main contribution $b_{\partial f}$.

Proof. The critical point satisfies : $\gamma_c = -z_Q/\sqrt{\varphi^2+z_Q^2}$ and at γ_c , the second order derivative of the phase equals $\frac{\varphi}{\sqrt{1-\gamma_c^2}} = \varphi \times \frac{\sqrt{\varphi^2+z_Q^2}}{\varphi^3} \geq \varphi$ and its critical value equals $-\sqrt{z_Q^2+\varphi^2}$. In order to show that the stationary phase applies we will show that $\varphi \geq \sqrt{s^2-1}$. From Lemmas 3.3 and 2.10 we have $\tilde{\Gamma}(\tilde{\alpha}, r) + \Gamma_0(\tilde{\alpha}, s) = \sqrt{r^2-1} + \sqrt{s^2-1} + (y_* - y_c)\tilde{\alpha} + O(1 - \tilde{\alpha})$. As y is close to y_* on the support of the symbol $I_{1,gl}$ ($|y - y_*| \leq \frac{\pi}{16}$) and $|y - y_c| \leq \frac{\pi}{8}$ (these constants may be shrunk if necessary), then $|y_c - y_*| \leq \frac{3\pi}{16} < \frac{5}{8}$ while $\sqrt{r^2-1} \geq \sqrt{2}\sqrt{\sqrt{2}-1} \geq \frac{4}{5}$. Moreover, on the support of $\chi_\varepsilon(\tilde{\alpha})$ we have $|1 - \tilde{\alpha}| \lesssim \varepsilon$ so we conclude taking ε_1 small enough compared to ε_0 . The stationary phase yields (3.9). The exponent of $\left(\frac{\varphi^2}{\varphi^2+z_Q^2}\right)$ equals $\frac{1}{12} - \frac{1}{3} + \frac{3}{4}$, where the last term comes from the second order derivative. \square

Corollary 3.6. *We have $I_{0,gl}^{1-\chi_0,\bar{\chi}}(Q, Q_0, \tau) = O(\tau^{-\infty}/t)$. Moreover, modulo $O(\tau^{-\infty}/t)$,*

$$I_{0,gl}^{1-\chi_0,\chi_\pm}(Q, Q_0, \tau) = \tau^{4/3-1-1/2} \int e^{-i\tau\sqrt{\varphi_\pm^2+z_Q^2}} \chi_{\varepsilon_1}(1 - \tau^{-2/3}\beta) \\ \frac{\tilde{\Sigma}_\pm(\beta, \tau)}{\varphi_\pm^{1/2}} \times \frac{\check{\psi}_0\left(\frac{\varphi_\pm}{\sqrt{\varphi_\pm^2+z_Q^2}}\right)}{(s^2 - 1)^{1/4}\phi^{1/2}(x_Q, y_c - y_Q + Y(\tau^{-1/3}w_\pm), 0, 1)} d\gamma d\beta, \quad (3.10)$$

where $\check{\psi}_0(\cdot) = (\cdot)^{1/2}\tilde{\chi}_1$ and $\tilde{\Sigma}_-$ is a classical symbol with main contribution $\Sigma(\beta, \gamma_c, \tau)$.

Proof. Using Lemma 3.5, $I_{0,gl}^{1-\chi_0,\bar{\chi}}(Q, Q_0, \tau)$ of the form (3.9) : with symbol supported for w far from w_\pm : repeated integrations by parts yield $O(\tau^{-\infty}/t)$ where the factor $1/t$ is obtained from the symbol $(\varphi\sqrt{s^2-1})^{-1/2} \lesssim 1/\sqrt{s^2-1} \lesssim 1/t$ as $2\sqrt{s^2-1} \geq \varphi \geq \sqrt{s^2-1}$, as $t \sim \sqrt{\varphi^2+z_Q^2} \leq 8\varphi$ on the support of ψ_0 and $r \geq \sqrt{2}$. To obtain (3.10) we use the proof of Lemma 3.5 to (3.8) for the \pm signs. \square

Using the Corollary, we obtain $I_{0,gl}^{1-\chi_0}(Q, Q_0, \tau) = \sum_{\pm} I_{0,gl}^{1-\chi_0, \chi_{\pm}} + O(\tau^{-\infty}/t)$, where $I_{0,gl}^{1-\chi_0, \chi_{\pm}}$ are given in (3.10). We are left with the integration with respect to β in the integrals (3.10) whose symbols $(1 - \chi_0)(\sqrt{\beta}\tau^{-\epsilon_1})\chi_{\epsilon_1}(1 - \tau^{-2/3}\beta)$ are supported for $\beta \gtrsim \tau^{2\epsilon_1}$ and $\tau^{-2/3}\beta \lesssim \epsilon_1$. As β takes values in a large interval, we consider separately dyadic intervals where $\beta \sim 2^{2k}$ and then sum all the contributions. Let $\tilde{\chi}$ supported near 1 and equal to 1 on $[\frac{3}{4}, \frac{5}{4}]$ such that

$$(1 - \chi_0)(\sqrt{\beta}\tau^{-\epsilon_1})\chi_{\epsilon_1}(1 - \tau^{-2/3}\beta) \sum_k \tilde{\chi}(\beta 2^{-2k}) = (1 - \chi_0)(\sqrt{\beta}\tau^{-\epsilon_1})\chi_{\epsilon_1}(1 - \tau^{-2/3}\beta). \quad (3.11)$$

On the support of $(1 - \chi_0)(\sqrt{\beta}\tau^{-\epsilon_1})\chi_{\epsilon_1}(1 - \tau^{-2/3}\beta)$ we have $\tau^{2\epsilon_1} < \beta \lesssim \epsilon_1\tau^{2/3}$ and for each k in the previous sum, $\tilde{\chi}(\beta 2^{-2k})$ localize at $\beta \sim 2^{2k}$. The sum is thus taken for $\epsilon_1 \log_2(\tau) \leq k < \frac{1}{3} \log_2(\tau)$. Recall that $\varphi|_{\pm} = \varphi|_{w_{\pm}}$ where $\varphi_- = \tilde{\Gamma}(\tilde{\alpha}, r) + \Gamma_0(\tilde{\alpha}, s)$ and $\varphi_+ = \varphi_- - \frac{4}{3}(-\tilde{\zeta})^{3/2}(\frac{1}{\tilde{\alpha}})$. We deal separately with the \pm signs. Let $I_{0,gl}^{1-\chi_0, \chi_{\pm}, k}$ denote the integrals in (3.10) with additional cutoff $\tilde{\chi}(\beta 2^{-2k})$. Using (3.11) we have

$$I_{0,gl}^{1-\chi_0, \chi_{\pm}} = \sum_{k=\epsilon_1 \log_2 \tau}^{(\log_2 \tau)/3} I_{0,gl}^{1-\chi_0, \chi_{\pm}, k}.$$

Lemma 3.7. *There exists a constant $C = C_+(\varepsilon)$ such that $|I_{0,gl}^{1-\chi_0, \chi_+}| \leq \sum_{k=\epsilon_1 \log_2 \tau}^{(\log_2 \tau)/3} |I_{0,gl}^{1-\chi_0, \chi_+, k}| \leq C_+(\varepsilon)/t$.*

Proof. At $w_+ = \sqrt{2\beta}(1 + O(\sqrt{\tau^{-2/3}\beta}))$, the phase φ_+ is stationary when $\tau^{1/3}(y_c - y_*) = 2\sqrt{2\beta}(1 + O(\sqrt{\tau^{-2/3}\beta}))$. Let $\beta = 2^{2k}\Xi$ on the support of $\tilde{\chi}(\beta 2^{-2k})$, with $\Xi \in [1/2, 3/2]$. As

$$\begin{aligned} \partial_{\Xi}(\tau\sqrt{\varphi_+^2 + z_Q^2}) &= \frac{\varphi_+}{\sqrt{\varphi_+^2 + z_Q^2}} \frac{\partial\beta}{\partial\Xi} (\tau\partial_{\beta}\varphi_+) \\ &= \frac{\varphi_+}{\sqrt{\varphi_+^2 + z_Q^2}} 2^{2k+k} \left(\frac{\tau^{1/3}(y_c - y_*)}{2^k} - 2\sqrt{2\Xi}(1 + O(\sqrt{\tau^{-2/3}\beta})) \right), \end{aligned} \quad (3.12)$$

the phase is stationary for $\Xi \sim 1$ only when $\frac{\tau^{1/3}(y_c - y_*)}{2^k} \sim 2\sqrt{2}$; as $\frac{\varphi_+}{\sqrt{\varphi_+^2 + z_Q^2}} \geq 1/8$ on the support of ψ_0 and as $2^{3k} \geq \tau^{3\epsilon_1/2}$, it follows that, for $|\frac{\tau^{1/3}(y_c - y_*)}{2^k} - 2\sqrt{2}| \geq 4$ and $\Xi \in [1/2, 3/2]$, repeated integrations by parts yield a contribution $O(\tau^{-\infty}/t)$. We deduce that there are at most a finite number of values of k for which the phase may be stationary; for such k the stationary phase applies at the critical point $2\sqrt{2\Xi} \sim \frac{\tau^{1/3}(y_c - y_*)}{2^k}$ as, there, the second order derivative equals $-\frac{\varphi_+}{\sqrt{\varphi_+^2 + z_Q^2}} 2^{3k} \times \frac{\sqrt{2}}{\sqrt{\Xi}}$ and $\Xi \sim 1$. For all such k , the stationary phase yields a factor $2^{2k} \times 2^{-3k/2} \times \sqrt{\varphi_+^2 + z_Q^2}^{1/2} / \varphi_+^{1/2}$, where the exponent $2k$ comes from the change of variables and the exponent $2^{-3k/2}$ from the second order derivative at $\Xi \sim 1$. As $2^{2k} \leq \tau^{2/3}$, the sum over all such k yields at most $2^{k/2} \leq \tau^{1/6}$ and the exponent $1/6$ is canceled by the exponent of $\tau^{4/3-2/3-1/2-1/3}$ from $I_{0,gl}^{1-\chi_0, \chi_+}$. We conclude using that $(\varphi_+ \sqrt{s^2 - 1})^{-1/2} \leq C_+(\varepsilon)/t$, where $C_+(\varepsilon)$ depends only on ε . \square

Lemma 3.8. *There exists a constant $C = C_-(\varepsilon)$ such that $\sum_{k=\epsilon_1 \log_2 \tau}^{(\log_2 \tau)/3} |I_{0,gl}^{1-\chi_0, \chi_-, k}(Q, Q_0, t)| \lesssim C_-(\varepsilon)/t$.*

Proof. We have $\varphi_- = \tilde{\Gamma}(\tilde{\alpha}, r) + \Gamma_0(\tilde{\alpha}, s)$ hence, for $\tilde{\alpha} = 1 - \tau^{-2/3}\beta$, we have

$$\begin{aligned} \tau\varphi_- &= \tau \left(\sqrt{r^2 - 1} + \sqrt{s^2 - 1} - (y_c - y_*)\tilde{\alpha} \right. \\ &\quad \left. + \frac{(1 - \tilde{\alpha})^2}{2} \left(\frac{1}{\sqrt{r^2 - 1}}(1 + O(1 - \tilde{\alpha})) + \frac{1}{\sqrt{s^2 - 1}}(1 + O(1 - \tilde{\alpha})) \right) \right) \\ \tau\partial_{\beta}\varphi_- &= \tau^{1/3}(y_c - y_*) + \tau^{-1/3}\beta \left(\frac{1}{\sqrt{r^2 - 1}}(1 + O(\tau^{-2/3}\beta)) + \frac{1}{\sqrt{s^2 - 1}}(1 + O(\tau^{-2/3}\beta)) \right). \end{aligned} \quad (3.13)$$

At $\beta = 2^{2k}\Xi$ we have $\partial_{\Xi}(\tau\sqrt{\varphi_-^2 + z_Q^2}) = \frac{\varphi_-}{\sqrt{\varphi_-^2 + z_Q^2}} 2^{2k} \partial_{\beta}(\tau\varphi_-)|_{\beta=2^{2k}\Xi}$ hence

$$\begin{aligned} \partial_{\Xi}(\tau\sqrt{\varphi_-^2 + z_Q^2}) &= \frac{2^{4k}\tau^{-1/3}\varphi_-}{\sqrt{\varphi_-^2 + z_Q^2}} \left(\frac{\tau^{2/3}}{2^{2k}}(y_c - y_*) + \Xi \left(\frac{1}{\sqrt{r^2 - 1}}(1 + O(\tau^{-2/3}2^{2k})) + \frac{1}{\sqrt{s^2 - 1}}(1 + O(\tau^{-2/3}2^{2k})) \right) \right), \\ \partial_{\Xi}^2(\tau\sqrt{\varphi_-^2 + z_Q^2})|_{\partial_{\beta}\varphi_- = 0} &= \frac{2^{4k}\tau^{-1/3}\varphi_-}{\sqrt{\varphi_-^2 + z_Q^2}} \left(\frac{1}{\sqrt{r^2 - 1}}(1 + O(\tau^{-2/3}2^{2k})) + \frac{1}{\sqrt{s^2 - 1}}(1 + O(\tau^{-2/3}2^{2k})) \right). \end{aligned}$$

As $s \geq r$ and for $\frac{\varphi_-}{\sqrt{\varphi_-^2 + z_Q^2}}$ on the support of ψ_0 we obtain a lower bound for the second order derivative of $\frac{2^{4k}\tau^{-1/3}}{\sqrt{r^2 - 1}}$. From now on we can proceed as in the case of φ_+ : if $\frac{2^{4k}\tau^{-1/3}}{\sqrt{r^2 - 1}} \geq \tau^\epsilon$ for some $\epsilon > 0$, we apply the stationary phase if moreover $\frac{\tau^{2/3}\sqrt{r^2 - 1}}{2^{2k}}(y_* - y_c) \sim 1$: the last condition reduces the number of such k to at most three values for which we find

$$|I_{0,gl}^{1-\chi_0, \chi-, k}(Q, Q_0, t)| \lesssim \frac{\tau^{-1/6}}{t} \times \frac{2^{2k}}{\phi^{1/2}} \left(\frac{\varphi_-}{\sqrt{\varphi_-^2 + z_Q^2}} \right)^{3/4 - 1/2} \times \left(\frac{2^{4k}\tau^{-1/3}}{\sqrt{r^2 - 1}} \right)^{-1/2} \sim 1/t, \quad (3.14)$$

where we used that $(\varphi\sqrt{s^2 - 1})^{-1/2} \lesssim 1/t$ and $\phi \geq r - 1$ to obtain $\frac{(r^2 - 1)^{1/4}}{\phi^{1/2}} \lesssim 1$. If $\frac{\tau^{2/3}\sqrt{r^2 - 1}}{2^{2k}}(y_* - y_c) \notin [1/4, 4]$, repeated integrations by parts yield a $O(\tau^{-\infty}/t)$ contribution (and we conclude using that $2^k \lesssim \tau^{1/3}$).

Fix $M > 4$ large enough and consider $2^{4k}\tau^{-1/3}\frac{1}{\sqrt{r^2 - 1}} \in [M^2, \tau^{\epsilon'}]$ for some $\epsilon' > 0$. As this parameter is large, we still may apply the stationary phase but we need to verify that the remainders are sufficiently small and that we can bound their sum. There is still a finite number of k for which the phases may be stationary. At the critical points Ξ_c , the stationary phase applies and we obtain, for all $N \geq 1$,

$$\begin{aligned} I_{0,gl}^{1-\chi_0, \chi-, k}(Q, Q_0, t) &= \tau^{-1/6} e^{-i\tau\sqrt{\varphi_-^2 + z_Q^2}} \frac{\tilde{\Sigma}_-(2^{2k}\Xi_c, \tau)}{\varphi_-^{1/2}(s^2 - 1)^{1/4} \phi^{1/2}(x_Q, \arccos(1/r) + \tau^{-1/3}w_-, 0, 1)} \frac{2^{2k} \times |\partial_{\Xi}^2(\tau\sqrt{\varphi_-^2 + z_Q^2})|^{-1/2}}{\phi^{1/2}(x_Q, \arccos(1/r) + \tau^{-1/3}w_-, 0, 1)} \\ &\quad + O\left(\left(\frac{2^{4k}\tau^{-1/3}}{\sqrt{r^2 - 1}} \right)^{-N} \tau^{-1/6} \times \frac{2^{2k}}{\sqrt{s^2 - 1}(r - 1)^{1/4}} \right), \quad (3.15) \end{aligned}$$

where the main contribution of $I_{0,gl}^{1-\chi_0, \chi-, k}(Q, Q_0, t)$ in the first line still satisfies (3.14) and where the remainder in the second line is $O\left(\left(\frac{2^{4k}\tau^{-1/3}}{4\sqrt{r^2 - 1}} \right)^{-N}/t \right)$. In the second line we used $\phi \geq (r - 1)^{1/2}$. The bounds for the remainders follow using $\sup |\partial_{\Xi}^2(\tau\sqrt{\varphi_-^2 + z_Q^2})| \geq \frac{2^{4k}\tau^{-1/3}}{\sqrt{r^2 - 1}}$. Notice that, taking one derivative of the cutoff $\chi_{\varepsilon}(\tau^{-2/3}2^{2k}\Xi) = \chi(\tau^{-2/3}2^{2k}\Xi/\varepsilon)$ yields a factor $\tau^{-2/3}2^{2k}/\varepsilon$ but, as $\Xi \sim 1$ on the support of $\chi(\beta 2^{-2k}) = \chi(\Xi)$, on the support of $\chi(\tau^{-2/3}2^{2k}\Xi/\varepsilon)\chi(\Xi)$ we have $\tau^{-2/3}2^{2k}/\varepsilon \lesssim 1$ hence for $M > 4$ sufficiently large this factor doesn't change the contribution of the remainder. For all k s.t. the phase is not stationary, integration by parts yields a contribution of at most

$$\tau^{-1/6} \frac{2^{2k}}{t(r - 1)^{1/2}} \times (2^{-4k}\tau^{1/3}\sqrt{r^2 - 1})^{N+1} = \frac{1}{t} \times (2^{-2k}\tau^{1/6}\sqrt{r^2 - 1}^{1/2}) \times (2^{-4k}\tau^{1/3}\sqrt{r^2 - 1})^N$$

for all $N \geq 0$. Let $N = 0$ and sum over k with $2^{4k}\tau^{-1/3}\frac{1}{\sqrt{r^2 - 1}} \in [M^2, \tau^{\epsilon'}]$, then

$$\frac{1}{t} \left(\sum_{M^2 \leq 2^{4k}\tau^{-1/3}\frac{1}{\sqrt{r^2 - 1}} \leq \tau^{\epsilon'}} 2^{-2k} \right) \times \tau^{1/6}\sqrt{r^2 - 1}^{1/2} \lesssim 1/(Mt).$$

Let now k such that $\tau^{\epsilon_1} \lesssim 2^k$ and $2^{4k}\tau^{-1/3}\frac{1}{\sqrt{r^2 - 1}} \leq M^2$ for some large, fixed $M > 1$. We bound each $I_{0,gl}^{1-\chi_0, \chi-, k}$ by $\tau^{-1/6}\frac{2^{2k}}{t(r - 1)^{1/4}} \lesssim M/t$ using $2^{2k} \leq M\tau^{1/6}\sqrt{r^2 - 1}^{1/2}$ and conclude. \square

Remark 3.9. In the two previous Lemmas, the bounds for $I_{0,gl}^{1-\chi_0, \chi_{\pm}}$ come with additional factors $(\frac{\varphi_{\pm}}{\sqrt{\varphi_{\pm}^2 + z_Q^2}})^{1/4}$. This is useful to keep in mind for the case when $1 - \gamma^2$ behaves like 2^{-2j} .

For β on the support of $\chi_0(\beta)$, using (3.4), the Airy factor can be brought in the symbol. The phase of $I_{0,gl}^{1-\chi_0}$ equals $\tau(z_Q\gamma - \sqrt{1-\gamma^2}\varphi_0)$, where $\varphi_0 := -(y_c - y_* + \tau^{-1/3}w)(1 - \tau^{-2/3}\beta) + \sqrt{s^2 - 1} + \phi(x_Q, \arccos(1/r) + \tau^{-1/3}w, 0, 1) \geq \sqrt{s^2 - 1}$. As $\sqrt{1-\gamma^2} \times w \geq \tau^{\epsilon_1}$ on the support of $\psi_0(1-\gamma^2)(1-\chi_0)(w\tau^{-\epsilon_1})$, it follows that the phase is non-stationary in w as $\beta \leq 2 \ll \tau^{2\epsilon_1} \lesssim w^2/2$ and we integrate by parts to obtain $O(\tau^{-\infty}/t)$.

3.1.2. *Case $|y-y_c| \leq 2\tau^{-1/3+\epsilon_1}$: study of $I_{0,gl}^{\chi_0}$.* Let $y = y_c + \tau^{-1/3}w$, with $|w| \leq \tau^{\epsilon_1}$. As $\partial_w \phi(x_Q, \arccos(1/r) + \tau^{-1/3}w, 0, 1) = \tau^{-1/3} \left(1 - \tau^{-2/3}w^2/2(1 + O(\tau^{-1/3}w))\right)$, the derivative w.r.t. w of phase of $I_{0,gl}^{\chi_0}$ equals

$$\tau^{-1/3} \sqrt{1-\gamma^2} \left(1 - \tau^{-2/3}\beta - 1 + \tau^{-2/3}w^2/2(1 + O(\tau^{-1/3}w))\right) = \sqrt{1-\gamma^2}(-\beta + w^2/2(1 + O(\tau^{-1/3}w))),$$

hence, for $\beta \geq \tau^{2\epsilon_1}$ we perform repeated integrations by parts to obtain a $O(\tau^{-\infty}/t)$ contribution (using that the support in w, β is bounded). We introduce $\chi_0(\beta\tau^{-2\epsilon_1})$ into the symbol of $I_{0,gl}^{\chi_0}$ without changing its contribution modulo $O(\tau^{-\infty}/t)$ terms. If we introduce moreover a cutoff $\chi_0(\beta)$ supported for $\beta \leq 2$, the Airy factor doesn't oscillate and may be brought into the symbol : in this case the phase of $I_{0,gl}^{\chi_0}$ is given by

$$\tau(z_Q\gamma - \sqrt{1-\gamma^2}(-(y_c - y_* + \tau^{-1/3}w)(1 - \tau^{-2/3}\beta) + \sqrt{s^2 - 1} + \phi(x_Q, \arccos(1/r) + \tau^{-1/3}w, 0, 1))).$$

Let $\varphi_0 := -(y_c - y_* + \tau^{-1/3}w)(1 - \tau^{-2/3}\beta) + \sqrt{s^2 - 1} + \phi(x_Q, \arccos(1/r) + \tau^{-1/3}w, 0, 1)$, then $\varphi_0 \geq \sqrt{s^2 - 1}$ and the stationary phase w.r.t. γ applies exactly as before. The critical point γ satisfies $z_Q = \frac{\gamma}{\sqrt{1-\gamma^2}}\varphi_0$.

The contribution of $I_{0,gl}^{\chi_0}(Q, Q_0, \tau)$ is of the form (3.9) where moreover $\beta \leq 2$ and $|w| \leq \tau^{\epsilon_1}$. We then obtain

$$|I_{0,gl}^{\chi_0}(Q, Q_0, \tau)| \lesssim \frac{\tau^{1/6-1/3}}{\varphi_0^{1/2}(s^2-1)^{1/4}} \times \tau^{\epsilon_1}, \quad (3.16)$$

where the exponents $1/6 - 1/3$ come from (3.9) and the change of variable $y = y_c + \tau^{-1/3}w$, and the factor τ^{ϵ_1} from the size of the support in w . Let now $\beta \in [3/2, \tau^{2\epsilon_1}]$ on the support of $(1 - \chi_0(\beta))\chi_0(\beta\tau^{2\epsilon_1})$, when the Airy factor does oscillate. We also have $\varphi \geq \sqrt{s^2 - 1}$ and the stationary phase w.r.t γ applies. The corresponding contribution of $I_{0,gl}^{\chi_0}(Q, Q_0, \tau)$ may be bounded as in (3.16) but with an additional factor $\tau^{2\epsilon_1}$ arising from the support wr.t. $\beta \leq \tau^{2\epsilon_1}$. Taking $\epsilon_1 < 1/18$ allows to conclude. \square

3.2. Dispersive bounds when $d(Q, \partial\Omega) \leq \sqrt{2} - 1 \leq d(Q_0, \partial\Omega)$. Let $1 \leq r \leq \sqrt{2} \leq s$ and let $0 \leq j \leq j(s, h)$. We proceed as in [7, Section 3.3] to obtain directly the form of the reflected wave, which may be done using the Melrose and Taylor parametrix as the observation point Q is close to the boundary; formula (2.4) becomes useless since $d(Q, \partial\Omega)$ may be arbitrarily small. By Proposition 2.8 we are reduced to prove $|\sum_{j=0}^{j(s,h)} \int \chi(h\tau)\tau e^{it\tau} I_j(Q_0, Q, \tau) d\tau| \leq \frac{C}{h^{2t}}$ for a constant independent of Q_0 and Q , where we set

$$I_j(\tau, Q_0, Q) := \tau \int e^{i\tau(y\alpha + z\gamma)} \left(a_j A_+(\tau^{2/3}\zeta) + b_j \tau^{-1/3} A'_+(\tau^{2/3}\zeta) \right) \frac{A(\tau^{2/3}\zeta_0)}{A_+(\tau^{2/3}\zeta_0)} \psi_j(1-\gamma^2) \widehat{F}_{j,\tau}(\alpha, \gamma) d\alpha d\gamma$$

obtained as done in the last part of Section 2.1.

Lemma 3.10. *There exists a constant $C > 0$, such that, for all Q in a small neighborhood of \mathcal{C}_{Q_0} , $|y - y_*| \leq \frac{\pi}{16}$ and $t \sim \text{dist}(Q_0, \partial\Omega) + \text{dist}(Q, \partial\Omega)$ the following holds $|\sum_{j=0}^{j(s,h)} I_j(Q_0, Q, \tau)| \leq \frac{C}{t}$.*

The Lemma follows exactly as in [7, Lemma 3.25] for all $j \leq j(s, h)$ as the observation point Q is located near a glancing point of the boundary (notice that, in the case Q far from $\partial\Omega$, the geometry of the obstacle was important and the approach to obtain dispersive bounds in the case of the exterior of the cylinder was different from the one in the exterior of a ball; when Q is near $\partial\Omega$ the same arguments hold in both cases so we do not reproduce the proof here. Moreover, all stationary arguments hold for $j \leq j(s, h)$).

3.3. Dispersive bounds for the "non-glancing" part, $d(Q_0, \partial\Omega) \geq \sqrt{2} - 1$. Let $s \geq \sqrt{2} - 1$ as before.

We let $u_{j,he,h}^\#(Q, Q_0, t) := \int_{\partial\Omega} \frac{\partial_x u_{j,he,h}^+(P, Q_0, t - |Q-P|)}{4\pi|Q-P|} d\sigma(P)$, where $\partial_x u_{j,he}^+|_{\partial\Omega}$ has been defined in (2.23).

Proposition 3.11. *There exists $C = C(\varepsilon) > 0$ such that for all $t > h$, the following holds uniformly with respect to Q, Q_0 such that $s \geq r \geq \sqrt{2}$ (where $s = 1 + x_{Q_0}$, $r = 1 + x_Q$):*

$$\sum_{j=0}^{j(s,h)} |u_{j,he,h}^\#(Q, Q_0, t)| \leq \frac{C}{h^2 t}.$$

Proof. Using (2.23), it follows that the phase function of $u_{j,he,h}^\#(Q, Q_0, t)$ is $\tau(t - \Phi)$ where $\Phi := |Q - P| + |P - Q_0|$ and the symbol is $\frac{\tau^2}{|P-Q||P-Q_0|} \sigma_{j,he}(y, z, s, \tau)$ with $\sigma_{j,he}$ a classical symbol of order 0 with respect to τ supported for P with coordinates $(x, y, z)_P = (0, y, z)$ such that $s(\sin y_* - \sin y) \geq c(\varepsilon)|P - Q_0|$ and $2^{-j}(-z) \sim \phi(0, y, 0, s)$. In the following it will be convenient to work with the coordinates (r, θ, z) (instead of (x, y, z)). Recall that we set $r = 1 + x$, $\theta = \frac{\pi}{2} - y$. In these coordinates, the support conditions for $\sigma_{j,he}$ become $s(\cos \theta_* - \cos \theta) \geq c(\varepsilon)|P - Q_0|$, $\theta_* = \arccos(1/s)$. We compute the derivative of the phase Φ , where

$$\Phi := |Q - P| + |Q_0 - P| = \tilde{\phi}(1, \theta - \theta_Q, r_Q, z - z_Q) + \tilde{\phi}(1, \theta, s, z),$$

where now $P = (\cos \theta, \sin \theta, z) \in \mathbb{R}^3$, $Q_0 = (s, 0, 0)$ and $Q = (r_Q \cos \theta_Q, r_Q \sin \theta_Q, z_Q)$ and where $\tilde{\phi}$ is defined in (1.9). Let $r = r_Q$. The critical points satisfy $\partial_\theta \Phi = \partial_z \Phi = 0$, which is equivalent to

$$\begin{cases} \frac{z}{\tilde{\phi}(1, \theta, s, z)} + \frac{z - z_Q}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} = 0, \\ \frac{s \sin \theta}{\tilde{\phi}(1, \theta, s, z)} + \frac{r \sin(\theta - \theta_Q)}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} = 0. \end{cases} \quad (3.17)$$

We aim at applying the stationary phase with respect to both θ and z . We evaluate the second order derivatives of Φ at $\nabla_{\theta,z} \Phi = 0$. The second order derivative of Φ satisfies

$$\partial_{z,z}^2 \Phi|_{\partial_z \Phi=0} = \left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right) \left(1 - \frac{z^2}{\tilde{\phi}^2(1, \theta, s, z)} \right). \quad (3.18)$$

Next, as $\partial_{\theta,z}^2 \Phi = -\left(\frac{zs \sin \theta}{\tilde{\phi}^3(1, \theta, s, z)} + \frac{(z - z_Q)r \sin(\theta - \theta_Q)}{\tilde{\phi}^3(1, \theta - \theta_Q, r, z - z_Q)} \right)$, we obtain, using the system (3.17),

$$\partial_{\theta,z}^2 \Phi|_{\nabla_{\theta,z} \Phi=0} = -\left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right) \frac{zs \sin \theta}{\tilde{\phi}^2(1, \theta, s, z)}. \quad (3.19)$$

Finally, we compute

$$\partial_{\theta,\theta}^2 \Phi = \frac{s \cos \theta}{\tilde{\phi}(1, \theta, s, z)} - \frac{s^2 \sin^2 \theta}{\tilde{\phi}^3(1, \theta, s, z)} + \frac{r \cos(\theta - \theta_Q)}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} - \frac{r^2 \sin^2(\theta - \theta_Q)}{\tilde{\phi}^3(1, \theta - \theta_Q, r, z - z_Q)}. \quad (3.20)$$

To evaluate $\partial_{\theta,\theta}^2 \Phi|_{\nabla_{\theta,z} \Phi=0}$ we need a refined analysis of the critical points. Using both equations in (3.17) gives $\frac{s \sin \theta}{\psi(s, \theta)} = -\frac{r \sin(\theta - \theta_Q)}{\psi(r, \theta - \theta_Q)}$, where $\psi(s, \theta) = \sqrt{1 - 2s \cos \theta + s^2}$, and hence $\theta \in [\theta_Q - \pi, \theta_Q]$. Taking the square in the last equality in (3.17), subtracting 1 and then using the first in (3.17) yields $\frac{s \cos \theta - 1}{\tilde{\phi}(1, \theta, s, z)} = \pm \frac{(r \cos(\theta - \theta_Q) - 1)}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)}$. Depending on the sign, we separate three situations :

Different signs. Consider first the case $\frac{s \cos \theta - 1}{\tilde{\phi}(1, \theta, s, z)} = -\frac{(r \cos(\theta - \theta_Q) - 1)}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)}$ when

$$\frac{s \cos \theta}{\tilde{\phi}(1, \theta, s, z)} + \frac{r \cos(\theta - \theta_Q)}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} = \frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)}.$$

We find

$$\partial_{\theta,\theta}^2 \Phi|_{\nabla_{\theta,z} \Phi=0} = \left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right) \left(1 - \frac{s^2 \sin^2 \theta}{\tilde{\phi}^2(1, \theta, s, z)} \right). \quad (3.21)$$

Using (3.18), (3.21), (3.19), the determinant of the Hessian matrix equals

$$\left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right)^2 \times \frac{(s \cos \theta - 1)^2}{\tilde{\phi}^2(1, \theta, s, z)} \Big|_{\nabla_{\theta,z} \Phi=0} \quad (3.22)$$

and on the support of the symbol $\sigma_{1,he}$ the second factor in (3.22) takes values in $[c^2(\varepsilon_1), 1]$. The unique critical point w.r.t. z reads as $z_c = z_Q \times \frac{\psi(s,\theta)}{\psi(s,\theta) + \psi(r,\theta - \theta_Q)}$.

Lemma 3.12. *When $\tau \times \left(\frac{1}{\tilde{\phi}(1,\theta,s,z)} + \frac{1}{\tilde{\phi}(1,\theta - \theta_Q,r,z - z_Q)} \right) \geq M$ for some $M > 1$ large enough, the usual stationary phase applies for θ, z near the critical points and yields $|\mathcal{F}(u_{j,he,h}^\#)(Q, Q_0, \tau)| \lesssim \frac{\tau}{t}$ when $t \sim \tilde{\phi}(1, \theta_c - \theta_Q, r, z_c - z_Q) + \tilde{\phi}(1, \theta_c, s, z_c)$. For z, θ outside a fixed neighborhood of the critical points the previous estimate still holds.*

Proof. We let $j = 0$ for simplicity. When $\tau \times \left(\frac{1}{\tilde{\phi}(1,\theta,s,z)} + \frac{1}{\tilde{\phi}(1,\theta - \theta_Q,r,z - z_Q)} \right) \geq \tau^\epsilon$ for some $\epsilon > 0$, the stationary phase obviously applies with large parameter $\gtrsim \tau^\epsilon$: then $\mathcal{F}(u_{0,he,h}^\#)(Q, Q_0, \tau)$ takes the form

$$\begin{aligned} \mathcal{F}(u_{0,he,h}^\#)(Q, Q_0, \tau) &= \frac{\tau^{2-1} e^{i\tau(t-\Phi)_{z_c, \theta_c}} \tilde{\sigma}_{0,he}(\theta, z, s, \tau)}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q) \tilde{\phi}(1, \theta, s, z)} \left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right)^{-1} \Big|_{z_c, \theta_c} \\ &\quad + O\left(\frac{\tau^{-\infty}}{\tilde{\phi}(1, \theta_c - \theta_Q, r, z_c - z_Q) + \tilde{\phi}(1, \theta_c, s, z_c)} \right) \end{aligned} \quad (3.23)$$

for some new symbol $\tilde{\sigma}_{0,he}$ which reads as an asymptotic expansion with main contribution $\sigma_{j,he}$ and small parameter $\lesssim \tau^{-\epsilon'}$. As the main contribution of $\mathcal{F}(u_{0,he,h}^\#)(Q, Q_0, \tau)$ can be bounded by $\frac{\tau \tilde{\sigma}_{0,he}(\theta, z, s, \tau)}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q) + \tilde{\phi}(1, \theta, s, z)}$, for $t \sim \tilde{\phi}(1, \theta_c - \theta_Q, r, z_c - z_Q) + \tilde{\phi}(1, \theta_c, s, z_c)$ this allows to conclude using the integration w.r.t. τ . For t that doesn't satisfy this condition we conclude by integrations by parts, finite speed of propagation and support properties of the symbol. If we replace $\tau^{\epsilon'}$ by some large constant M , the main contribution of $\mathcal{F}(u_{0,he,h}^\#)(Q, Q_0, \tau)$ can be bounded in the same way, but we need to bound the remainder terms as follows

$$\frac{\tau^2}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q) \tilde{\phi}(1, \theta, s, z)} \times \tau^{-1} \left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right) M^{-N}$$

for all $N \geq 1$, which is enough to conclude. For $j \leq j(s, h)$ we conclude in the same way.

Let now z, θ outside a fixed neighborhood of the critical points. If moreover $|z| \geq 2t$, the phase $\tau(t - \Phi)$ is not stationary w.r.t. τ ; let $|z| \leq 2t$ such that $|\frac{z}{z_c} - 1| \geq c$ for some fixed constant $c > 0$. If $\tau \frac{|z_Q|}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \geq M_1$ for some large $M_1 > 1$, then we make repeated integrations by parts as $\tau \partial_z \Phi = \frac{z_Q}{\tilde{\phi}_2(1, \theta - \theta_Q, r, z - z_Q)} (\frac{z}{z_c} - 1)$. Let $\tau \frac{|z_Q|}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} < M_1$. As $\tau \partial_z \Phi = \tau \left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right) z - \tau \frac{z_Q}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)}$, then if $\tau \partial_z \Phi \geq M_2$ for some large constant $M_2 > 1$, repeated integrations by parts allow to conclude; if, instead, $\tau \partial_z \Phi \leq M_2$ then

$$|z| \leq \left(\frac{M_2}{\tau} + \frac{|z_Q|}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right) \frac{1}{\left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right)}$$

and we directly obtain, using the size of the support of the integrand (z, θ) (with θ bounded)

$$|\mathcal{F}(u_{0,he,h}^\#)(Q, Q_0, \tau)| \lesssim \frac{\tau^2}{\tilde{\phi}_1 \tilde{\phi}_2} \frac{M_1 + M_2}{\tau} \times \frac{\tilde{\phi}_1 \tilde{\phi}_2}{\tilde{\phi}_1 + \tilde{\phi}_2} \lesssim \frac{1}{t},$$

where $\tilde{\phi}_1 = \tilde{\phi}(1, \theta, s, z)$ and $\tilde{\phi}_2 = \tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)$. Similar arguments hold for all $j \leq j(s, h)$. \square

Lemma 3.13. *When $\tau \times \left(\frac{1}{\tilde{\phi}(1,\theta,s,z)} + \frac{1}{\tilde{\phi}(1,\theta - \theta_Q,r,z - z_Q)} \right) \leq M$ estimate (3.23) still holds for $t \sim \tilde{\phi}(1, \theta_c - \theta_Q, r, z_c - z_Q) + \tilde{\phi}(1, \theta_c, s, z_c)$.*

Proof. For $|z/z_c - 1| \geq c$ we may proceed as in the second part of the proof of the previous lemma. Let therefore $z/z_c \in [1/4, 4]$ and make the change of variables $z = z_c \Xi$. Then

$$\tau \partial_\Xi \Phi = \tau z_c \partial_z \Phi|_{z=z_c \Xi} = \tau z_c \times \frac{z_Q}{\tilde{\phi}_2} (\Xi - 1) = \tau z_c^2 \frac{\tilde{\phi}_1}{\tilde{\phi}_2} \frac{1}{\tilde{\phi}_1 + \tilde{\phi}_2} (\Xi - 1).$$

Using (3.18), we obtain $\tau \partial_{\Xi}^2 \Phi|_{\Xi=1} = \tau z_c^2 \partial_z^2 \Phi|_{z=z_c \Xi} = \tau \left(\frac{1}{\phi_1} + \frac{1}{\phi_2} \right) z_c^2 \frac{\psi_1^2}{\phi_1^2}$, where, from the support properties of the symbol, the last factor is bounded from below by a fixed constant. If $\tau \left(\frac{1}{\phi_1} + \frac{1}{\phi_2} \right) z_c^2 \geq M$, we apply the stationary phase near $\Xi = 1$ only with respect to Ξ (and not with θ) as in the previous lemma and, using that θ belongs to a compact set, we find the following uniform bound

$$|\mathcal{F}(u_{0,he,h}^\#)(Q, Q_0, \tau)| \lesssim \frac{\tau^2}{\tilde{\phi}_1 \tilde{\phi}_2} \times z_c \times \tau^{-1/2} z_c^{-1} \left(\frac{1}{\tilde{\phi}_1} + \frac{1}{\tilde{\phi}_2} \right)^{-1/2} \lesssim \frac{\tau}{\tilde{\phi}_1 + \tilde{\phi}_1} \times \left(\tau \left(\frac{1}{\tilde{\phi}_1} + \frac{1}{\tilde{\phi}_2} \right) \right)^{1/2} \quad (3.24)$$

and we conclude using the hypothesis $\tau \times \left(\frac{1}{\tilde{\phi}(1,\theta,s,z)} + \frac{1}{\tilde{\phi}(1,\theta-\theta_Q,r,z-z_Q)} \right) \leq M$. Same for all $j \leq j(s, h)$. \square

Same sign, P in the illuminated regime of Q_0, Q . Consider now the situation $\frac{s \cos \theta - 1}{\psi(s, \theta)} = \frac{r \cos(\theta - \theta_Q) - 1}{\psi(r, \theta - \theta_Q)}$. The formula (3.18) remains unchanged and $\partial_{z,z}^2 \Phi$ is strictly positive. Moreover, from the support condition of $\sigma_{1,he}$ we have $s \cos \theta > 1$ then $r \cos(\theta - \theta_Q) > 1$ and in (3.20) we obtain a lower bound for the sum of the first and third terms at the critical points as follows :

$$\frac{s \cos \theta}{\tilde{\phi}(1, \theta, s, z)} + \frac{r \cos(\theta - \theta_Q)}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \geq \frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)}$$

and we can proceed exactly as in the previous case.

Remark 3.14. Notice that the positivity condition $s \cos \theta > 1$ is equivalent to $\cos \theta > \cos \theta_*$, where $\theta_* = \arccos(1/s)$, which in turn assures that the point P belongs to the illuminated region of Q (as $\theta < \theta_*$). When both conditions hold ($\cos \theta > 1/s$ and $\cos(\theta - \theta_Q) > 1/r$), the point P belongs to the illuminated regions from Q_0 and Q . In fact, the line Q_0Q is tangent to the boundary when $\arccos(1/s) + \arccos(1/r) = \theta_Q$: if $P \in \partial\Omega$ is such that the cosine of the angle between QO and OP is larger than $1/r$, then the point Q belongs to the illuminated regime of Q_0 . As such, the previous case when $\pm(s \cos \theta - 1) > 0$ and $\pm(1 - r \cos(\theta - \theta_Q)) > 0$ corresponds to points P which belong to the illuminated regime of only one of the two points Q_0 and Q . In the last case $s \cos \theta - 1 < 0$ and $1 - r \cos(\theta - \theta_Q) < 0$ that will be dealt with in the remaining of this section, P does not belong to the illuminated regions of Q_0, Q .

Same sign, P in the shadow regime of Q_0, Q . In this case we do not have a lower bound for the determinant of the Hessian matrix as before. Replacing $\frac{r \cos(\theta - \theta_Q)}{\phi_2} = -\frac{1}{\phi_1} + \frac{1}{\phi_2} + \frac{s \sin \theta}{\phi_1}$ in the expression (3.20) yields the following form for the determinant of the Hessian matrix at this critical point :

$$\left(\frac{1}{\tilde{\phi}_1} + \frac{1}{\tilde{\phi}_2} \right) \times \frac{(1 - s \cos \theta)}{\tilde{\phi}_1} \times \left| \left(\frac{1}{\tilde{\phi}_1} + \frac{1}{\tilde{\phi}_2} \right) \frac{(1 - s \cos \theta)}{\tilde{\phi}_1} - 2 \frac{\psi_1^2}{\tilde{\phi}_1^2} \right|_{\nabla_{\theta,z} \Phi = 0}. \quad (3.25)$$

Lemma 3.15. At $\nabla_{\theta,z} \Phi = 0$ the following holds

$$\begin{aligned} \frac{\psi_2^2}{\tilde{\phi}_2^2} - \frac{1}{\tilde{\phi}_2} \frac{(1 - r \cos(\theta - \theta_Q))}{\tilde{\phi}_2} &= \frac{r(r - \cos(\theta - \theta_Q))}{\tilde{\phi}_2^2} \geq \frac{r(r - 1)}{\tilde{\phi}_2^2}. \\ \frac{\psi_1^2}{\tilde{\phi}_1^2} - \frac{1}{\tilde{\phi}_1} \frac{(1 - s \cos \theta)}{\tilde{\phi}_1} &= \frac{s(s - \cos \theta)}{\tilde{\phi}_2^2} \geq \frac{s(s - 1)}{\tilde{\phi}_1^2}. \end{aligned}$$

The lemma is a direct computation. Taking the sum of the terms in the left hand side and using that $\tilde{\phi}_j = \psi_j \sqrt{1 + \frac{z_Q^2}{(\psi_1 + \psi_2)^2}}$ for $j \in \{1, 2\}$ and $\frac{(1 - s \cos \theta)}{\tilde{\phi}_1} = \frac{(1 - r \cos(\theta - \theta_Q))}{\tilde{\phi}_2}$ yields, at $\nabla_{\theta,z} \Phi = 0$,

$$\left(2 \frac{\psi_1^2}{\tilde{\phi}_1^2} - \left(\frac{1}{\tilde{\phi}_1} + \frac{1}{\tilde{\phi}_2} \right) \frac{(1 - s \cos \theta)}{\tilde{\phi}_1} \right) \geq \frac{s(s - 1)}{\tilde{\phi}_1^2} + \frac{r(r - 1)}{\tilde{\phi}_2^2},$$

which further induces the following lower bound for the determinant of the Hessian matrix

$$\left(\frac{1}{\tilde{\phi}_1} + \frac{1}{\tilde{\phi}_2} \right) \times \frac{(1 - s \cos \theta)}{\tilde{\phi}_1} \times \left(\frac{s(s - 1)}{\tilde{\phi}_1^2} + \frac{r(r - 1)}{\tilde{\phi}_2^2} \right).$$

From now on we may proceed as in the proof of the first case, applying the stationary phase when this determinant is sufficiently large and obtaining bounds using the size of the intervals of integration when the stationary phase fails to apply. Thus, we obtain the equivalent of Lemma 3.12

Lemma 3.16. *When $\tau \times \left(\frac{1}{\tilde{\phi}(1, \theta, s, z)} + \frac{1}{\tilde{\phi}(1, \theta - \theta_Q, r, z - z_Q)} \right)^{1/2} \left(\frac{s(s-1)}{\tilde{\phi}_1^2} + \frac{r(r-1)}{\tilde{\phi}_2^2} \right)^{1/2} \geq M$ for some large $M > 1$, the usual stationary phase applies for θ, z near the critical points and yields $|\mathcal{F}(u_{0,he,h}^\#)(Q, Q_0, \tau)| \lesssim \frac{\tau}{t}$ for $t \sim \tilde{\phi}(1, \theta_c - \theta_Q, r, z_c - z_Q) + \tilde{\phi}(1, \theta_c, s, z_c)$. For z, θ outside a fixed neighborhood of the critical points the previous estimate still holds.*

Proof. The main contribution of $\mathcal{F}(u_{0,he,h}^\#)(Q, Q_0, \tau)$ after applying the stationary phase is bounded by

$$\frac{\tau^2}{\tilde{\phi}_1 \tilde{\phi}_2} \times \tau^{-1} \left(\frac{1}{\tilde{\phi}_1} + \frac{1}{\tilde{\phi}_2} \right)^{-1/2} \left(\frac{s(s-1)}{\tilde{\phi}_1^2} + \frac{r(r-1)}{\tilde{\phi}_2^2} \right)^{-1/2} \lesssim \frac{\tau}{\tilde{\phi}_1 + \tilde{\phi}_2} \times \sqrt{\frac{1}{\tilde{\phi}_1} + \frac{1}{\tilde{\phi}_2}} \left(\frac{\tilde{\phi}_1}{4s} + \frac{\tilde{\phi}_2}{4r} \right).$$

On the support of $\sigma_{1,he}$ we obtain the desired estimates. We let the other situations to the reader. \square

When $1 \leq j \leq j(s, h)$ the phase is the same, only the symbol comes with non positive powers of 2^j : to sum them up, notice that the phase is stationary in τ only when $t \sim |z| \sim 2^j s$, hence for a finite number of j . \square

4. HIGH-FREQUENCY CASE. PARAMETRIX AND DISPERSIVE ESTIMATES FOR $d(Q, \partial\Omega) < \sqrt{2} - 1$ AND $d(Q_0, \partial\Omega) < \sqrt{2} - 1$, OR FOR $d(Q, \partial\Omega) \geq \sqrt{2} - 1$ AND $\sqrt{1 - \gamma^2} \sim 2^{-j}$ WITH $\tau 2^{-3j} d(Q, \partial\Omega) \lesssim 1$

In this section both Q and Q_0 are close to the boundary and $t > 0$. For convenience, we will assume this time that $s \leq r \leq \sqrt{2}$. Denote $\mathcal{R}(Q, Q_0, \tau)$ the outgoing solutions of the Helmholtz equation $(\tau^2 + \Delta)w = \delta_{Q_0}$, $w|_{\partial\Omega} = 0$ with $Q_0 = (s, 0, 0)$ where we recall that, in cylindrical coordinates, $\Delta = \partial_r^2 + \frac{\partial_r}{r} + \frac{\partial_\vartheta^2}{r^2} + \partial_z^2$. Then the solution of the wave equation with initial condition $(u_0, u_1) = (\delta_{Q_0}, 0)$ is given by

$$u(Q, Q_0, t) = \int_0^\infty e^{it\tau} \mathcal{R}(Q, Q_0, \tau) \frac{d\tau}{\pi}. \quad (4.1)$$

For a given $w_0(r, \theta, z)$, the solution to the inhomogeneous equation $(\tau^2 + \Delta)w = w_0$ reads as

$$w(\tau, r, \theta, z) = \int_{\mathbb{R}} e^{iz\vartheta} \sum_{n \in \mathbb{Z}} e^{in\theta} \int_1^\infty G_n(r, \tilde{r}, \kappa(\vartheta, \tau)) \tilde{r}^2 \hat{w}_0(\tilde{r}, n, \vartheta) d\tilde{r} d\vartheta,$$

where the kernel G_n is symmetric w.r.t. r, \tilde{r} and, for $r \geq \tilde{r}$, it is given by

$$\begin{aligned} G_n(r, \tilde{r}, \kappa) &= \frac{\pi}{2i} (r\tilde{r})^{-\frac{1}{2}} \left(J_n(\tilde{r}\kappa) - \frac{J_n(\kappa)}{H_n(\kappa)} H_n(\tilde{r}\kappa) \right) H_n(r\kappa), \\ &= \frac{\pi}{4i} (r\tilde{r})^{-\frac{1}{2}} \left(\overline{H}_n(\tilde{r}\kappa) - \frac{\overline{H}_n(k)}{H_n(k)} H_n(\tilde{r}\kappa) \right) H_n(r\kappa). \end{aligned} \quad (4.2)$$

Here $J_n(z) = \frac{1}{2}(H_n(z) + \overline{H}_n(z))$ denotes the Bessel function and $\kappa(\vartheta, \tau) := \sqrt{\tau^2 - \vartheta^2}$. As n is an integer, $H_{-n}(z) = (-1)^n H_n(z)$, therefore $G_n = G_{-n}$. Taking $w_0 = \delta_{Q_0}$, $Q_0 = (s, 0, 0)$ and $s \leq r$ yields $\tilde{r} = s$ and

$$\mathcal{R}(Q, Q_0, \tau) = s^2 \int_{\mathbb{R}} e^{iz\vartheta} \sum_{n \in \mathbb{Z}} e^{in\theta} G_{|n|}(r, s, \kappa(\vartheta, \tau)) d\vartheta.$$

Let $\psi_0, \psi \in C_0^\infty$ as in Section 1.0.1 such that ψ_0 is equal to 1 on $[1/81, 1]$, and to 0 on $[0, 1/100]$, $\psi \in C_0^\infty(1/4, 4)$ is equal to 1 near 1 is such that $1 - \psi_0(\beta) = \sum_{j \geq 1} \psi(2^{2j}\beta)$ and $0 \leq \psi_0, \psi \leq 1$, and set

$$\mathcal{R}_j(Q, Q_0, \tau) = s^2 \int_{\mathbb{R}} e^{iz\vartheta} \sum_{n \in \mathbb{Z}} e^{in\theta} \psi(2^{2j}(1 - (\vartheta/\tau)^2)) G_{|n|}(r, s, \kappa(\vartheta, \tau)) d\vartheta,$$

for $j \geq 1$; for $j = 0$, replace ψ by $\psi_0(1 - \gamma^2)$.

Lemma 4.1. *Fix $0 < h_0 < 1$ small enough and let $h \leq h_0$. Let $\chi \in C_0^\infty(1/2, 2)$ valued in $[0, 1]$ and equal to 1 on $[\frac{3}{4}, \frac{3}{2}]$. There exist a constant $C > 0$ such that for all $1 \leq s \leq r \leq \sqrt{2}$ and all $t > 0$, we have*

$$I(Q, Q_0, h) := \int_0^\infty e^{it\tau} \chi(h\tau) \mathcal{R}(Q, Q_0, \tau) d\tau \leq \frac{C}{h^2 t}. \quad (4.3)$$

Moreover, for $j \geq j(r, h)$ with $j(r, h)$ defined in Definition 2.3, we have $\sum_{j \geq j(r, h)} I^j(Q, Q_0, \tau) \leq \frac{C}{h^2 t}$, where

$$I^j(Q, Q_0, h) := \int_0^\infty e^{it\tau} \chi(h\tau) \mathcal{R}_j(Q, Q_0, \tau) d\tau. \quad (4.4)$$

In the remaining of this section we prove Lemma 4.1. Let $\kappa = \kappa(\vartheta, \tau) = \sqrt{\tau^2 - \vartheta^2}$ and set

$$G_n^+(r, s, \kappa) = \frac{\pi}{4i\sqrt{rs}} \overline{H}_n(s\kappa) H_n(r\kappa), G_n^-(r, s, \kappa) := \frac{\pi}{4i\sqrt{rs}} \frac{\overline{H}_n(\kappa)}{H_n(\kappa)} H_n(s\kappa) H_n(r\kappa). \quad (4.5)$$

Substitute (4.5) in (4.2) and denote \mathcal{R}^\pm and $I^\pm(Q, Q_0, \tau)$ the corresponding contributions, respectively, so that $I = I^+ + I^-$. Let $\chi_0 \in C_0^\infty(-2, 2)$ valued in $[0, 1]$ and equal to 1 on $[-1, 1]$ and $\chi_\pm(\ell) := (1 - \chi_0(\ell)) \mathbf{1}_{\pm\ell > 0}$. Consider $n \neq 0$ and write $I^\pm = \sum_{* \in \{0, \pm\}} I_{\chi_*}^\pm$, where, for $\chi_* \in \{\chi_0, \chi_\pm\}$, $n \geq 1$ and $j \geq 0$ we define

$$I_{\chi_*}^{\pm, n, j} := \int_0^\infty e^{it\tau} \chi(h\tau) \int e^{iz\vartheta} \chi_* \left(\left(\frac{\sqrt{\tau^2 - \vartheta^2}}{n} - 1 \right) / \varepsilon \right) s^2 \psi(2^{2j}(1 - (\vartheta/\tau)^2)) G_n^\pm(r, s, \kappa(\vartheta, \tau)) d\vartheta d\tau \quad (4.6)$$

and set $I_{\chi_*}^\pm = \sum_{j \geq 0} \sum_{n \in \mathbb{N} \setminus \{0\}} (e^{in\theta} + e^{-in\theta}) I_{\chi_*}^{\pm, n, j}$ for some small $\varepsilon > 0$. Then $I^j = 2 \sum_{* \in \{0, \pm\}} \sum_n \cos(n\theta) I_{\chi_*}^{\pm, n, j}$. Then $\sqrt{\tau^2 - \vartheta^2}/n < 1 - \varepsilon$, $\sqrt{\tau^2 - \vartheta^2}/n \in [1 - 2\varepsilon, 1 + 2\varepsilon]$ and $\sqrt{\tau^2 - \vartheta^2}/n > 1 + \varepsilon$ on the support of χ_-, χ_0, χ_+ .

In the following, we look for upper bounds for $|I_{\chi_*}^{\pm, n, j}|$ first when $s \leq r \leq \sqrt{2}$, then for $r \geq \sqrt{2}$ and $j \geq j(r, h)$ and check that the sums over n, j remain bounded by $C/(h^2 t)$ for some uniform constant $C > 0$ independent of the parameters. We may assume that $n \geq n_0$ for some large n_0 , as, for bounded values, the result is trivial. We start with the main part $I_{\chi_+}^\pm$ which corresponds to values $\rho := (\sqrt{\tau^2 - \vartheta^2})/n \geq 1 + \varepsilon$. Let $\vartheta = \tau\gamma$ then $\rho = \tau\sqrt{1 - \gamma^2}/n \geq 1 + \varepsilon$. Let $\tilde{\rho} \in \{\rho, r\rho, s\rho\}$. With Φ_+ given in Lemma 6.1 we get from (6.3)

$$H_n(n\tilde{\rho}) \sim_{1/n} 2e^{-\frac{i\pi}{3}} \left(\frac{4\tilde{\zeta}(\tilde{\rho})}{1 - \rho^2} \right)^{\frac{1}{4}} n^{-\frac{1}{3}} A_+(n^{\frac{2}{3}} \tilde{\zeta}(\tilde{\rho})) \left[\sum_{j \geq 0} \left(a_j + n^{-\frac{4}{3}} \Phi_+(n^{\frac{2}{3}} \tilde{\zeta}(\tilde{\rho})) b_j \right) (-n^{\frac{2}{3}} \tilde{\zeta}(\tilde{\rho}))^{-3j/2} \right], \quad (4.7)$$

$$A_+(n^{\frac{2}{3}} \tilde{\zeta}(\tilde{\rho})) \sim_{1/n} n^{-\frac{1}{6}} (-\tilde{\zeta}(\tilde{\rho}))^{-\frac{1}{4}} e^{-i\frac{2}{3}n(-\tilde{\zeta}(\tilde{\rho}))^{\frac{3}{2}}} \left(1 + O((-n^{\frac{2}{3}} \tilde{\zeta}(\tilde{\rho}))^{-1}) \right), \text{ if } n^{\frac{2}{3}} \tilde{\zeta}(\tilde{\rho}) > 2.$$

On the support of the cut-off functions in (4.6) for $* = +$, the symbol of $I_{\chi_+}^{\pm, n, j}$ becomes

$$J_{\chi_+}^{\pm, n, j}(r, s, \rho) := n^{-2 \times \frac{1}{3} - 2 \times \frac{1}{6}} \frac{s^2 \chi_+((\tau\sqrt{1 - \gamma^2}/n - 1)/\varepsilon)}{(rs)^{\frac{1}{2}} ((r\rho)^2 - 1)^{\frac{1}{4}} ((s\rho)^2 - 1)^{\frac{1}{4}}} \Sigma_\pm(r, s, \rho, n) \tau \chi(h\tau) \psi(2^{2j}(1 - \gamma^2)), \quad (4.8)$$

where Σ_\pm are asymptotic expansions with small parameter n^{-1} and with main contribution obtained as a product of a_0 in (6.3) and σ_0 in (6.1) hence elliptic. The phase functions of $I_{\chi_+}^{\pm, n, j}$, denoted ϕ_n^\pm , read as

$$\tau\phi_n^\pm := t\tau + z\gamma\tau - n \left(f_0(r, \rho) \mp f_0(s, \rho) \right), \quad f_0(r, \rho) := \frac{2}{3} (-\tilde{\zeta}(r\rho))^{\frac{3}{2}} - \frac{2}{3} (-\tilde{\zeta}(\rho))^{\frac{3}{2}}, \quad (4.9)$$

where we recall $\rho = \frac{\tau\sqrt{1 - \gamma^2}}{n}$. The phases ϕ_n^\pm of $I_{\chi_+}^{\pm, n, j}$ are stationary when $\nabla_{\tau, \gamma}(\tau\phi_n^\pm) = 0$, that is

$$\partial_\tau(\tau\phi_n^\pm) = t + z\gamma - \frac{n}{\tau} (f_1(r, \rho) \mp f_1(s, \rho)), \quad \tau\partial_\gamma\phi_n^\pm = \tau \left(z + \frac{\gamma}{\sqrt{1 - \gamma^2}} \frac{(f_1(r, \rho) \mp f_1(s, \rho))}{\rho} \right), \quad (4.10)$$

where $f_1(r, \rho) := \sqrt{(r\rho)^2 - 1} - \sqrt{\rho^2 - 1}$ and where the derivative of f_0 is obtained from Lemma 2.5.

Lemma 4.2. *There exists $C > 0$ so that for all $\sqrt{2} \geq r \geq s \geq 1$ the following holds $\sum_{n \geq n_0, j \geq 0} |I_{\chi_+}^{\pm, n, j}| \leq \frac{C}{h^2 t}$. For $r \geq s$ with $r \geq \sqrt{2}$ and for $j(r, h)$ given in Definition 2.3 we also have $\sum_{n \geq n_0, j \geq j(r, h)} |I_{\chi_+}^{\pm, n, j}| \leq \frac{C}{h^2 t}$.*

Proof. We focus on $I_{\chi_+}^-, n, j$. Let $\varphi := 2^j \sqrt{1 - \gamma^2}$, then $\varphi \in (1/2, 2)$ on the support of $\psi(2^{2j}(1 - \gamma^2)) = \psi(\varphi^2)$ and $|\gamma| \geq 1/4$ (when $j = 0$ there is no need to change variables). Let $\phi_{n, j}^- := \phi_n^-|_{\gamma = \sqrt{1 - 2^{-2j}\varphi^2}}$ for $\varphi \sim 1$. As $r \geq s$ and $\rho \geq 1 + \varepsilon$, the factor depending on r, s, ρ in (4.11) is uniformly bounded by $1/\rho$.

Let first $1 < s \leq r \leq \sqrt{2}$ and $t \sim |z|$. If $2^{-2j}|z| \gtrsim 1$ then $\tau|\partial_\varphi\phi_{n,j}^-| = \tau|\frac{\partial\gamma}{\partial\varphi}\partial_\gamma\phi_{n,j}^-| \sim \tau 2^{-2j}|z| \gtrsim 1/h$: repeated integrations by parts yield $O(h^N(2^{-2j}|z|)^{-N})$ for all $N \geq 1$, hence for small r we find

$$|I_{\chi_+}^{-,n,j}(Q, Q_0, h)| \lesssim \frac{1}{h^2} \times \frac{2^{-2j}}{n} \times \frac{s^2}{(rs)^{1/2}} \frac{h^N(2^{-2j}|z|)^{-N}}{((r\rho)^2 - 1)^{1/4}((s\rho)^2 - 1)^{1/4}}. \quad (4.11)$$

Take $N = 1$, then $\sum_{n < 2^{-j}/h} \sum_{2^{2j} \lesssim |z|} 2^{-2j}/(n\rho) \times h(2^{2j}/|z|) \leq \frac{1}{|z|} \sum_{2^{2j} \lesssim |z|} 2^j \times 2^{-j}/h \leq \frac{h \log(1/h)}{|z|}$ where we used $(n\rho)^{-1} = 2^j h$, $n < 2^{-j}/h$ and $j \leq \log_2(1/h)$. Same computation with $N \geq 1$ yields $\sum_{n, 2^{2j} \lesssim t} |I_{\chi_+}^{-,n,j}| \lesssim \frac{O(h^N) \log(1/h)}{h^{2t}}$. For $2^{-2j}|z| \leq 1$ we have again, $|I_{\chi_+}^{-,n,j}| \lesssim \frac{1}{h^2} \frac{2^{-2j}}{n\rho}$ and $\sum_{n < 2^{-j}/h} \sum_{2^{2j} \geq |z|} 2^{-2j}/(n\rho) \lesssim \sum_{2^{2j} \geq |z|} 2^{-2j+j} h \times (2^{-j}/h) \leq \sum_{2^{2j} \geq |z|} 2^{-2j} \leq 1/|z| \sim 1/t$. If $t/|z| \notin [1/2, 2]$, repeated integrations by parts in τ yield the same kind of bounds with additional factors h^N for all $N \geq 1$.

Let now $r \geq \sqrt{2}$ and $s \leq r$ such that $\tau 2^{-3j}r \leq 1$; since the phase is stationary w.r.t. γ when $|z| \sim 2^j r$, it follows that, if $\tau 2^{-4j}|z| \geq 4$, we may integrate by parts in φ (in which case the remainders may be dealt with as before) to conclude. Let therefore $\tau 2^{-4j}|z| \leq 4$. We notice that when $t \geq 4(|z| + 2^j r)$ the phase $\tau\phi_{n,j}^-$ is not stationary in τ : in this case we integrate by parts in τ and obtain an upper bound for $|I_{\chi_+}^{-,n,j}|$ of the form (4.11) but with $h^N(2^{-2j}|z|)^{-N}$ replaced by $(h/t)^N$. For $N \geq 1$ gives $\sum_{n < 2^{-j}/h, j \geq j(r,h)} |I_{\chi_+}^{-,n,j}| \lesssim \frac{h^N}{h^{2t}} \sum_{n < 2^{-j}/h, j \leq \log(1/h)} \frac{2^{-2j}}{n\rho} \leq \frac{h^N \log(1/h)}{h^{2t}}$ (where we didn't use that $j \geq j(r,h)$).

Let $|z| \sim 2^j r$ and $t \leq 4(|z| + 2^j r) \sim 4|z|$, then we have again $|I_{\chi_0, \chi_0, n}^{-, \chi_+, j}| \lesssim \frac{1}{h^2} \frac{2^{-2j}}{n\rho}$ and we are left to estimate the sum over $j \geq 1$ satisfying $\tau 2^{-4j}|z|, \tau 2^{-3j}r \lesssim 1$. If moreover $2^{-2j}|z| \leq 1$, we find

$$\sum_{1 < 2^{-j}/(nh), j \geq j(r,h)} |I_{\chi_+}^{-,n,j}| \lesssim \frac{1}{h^2} \sum_{n \leq 2^{-j}/h, 2^{-2j}|z| \leq 1} \frac{2^{-2j}}{n} \times nh 2^j = \frac{1}{h^2} \sum_{2^{-2j} \leq 1/|z|} h 2^{-2j+j} \times \frac{2^{-j}}{h} \lesssim \frac{1}{h^2 t}. \quad (4.12)$$

When $2^{-2j}|z| \gtrsim 1$ we bound from below $\frac{\tilde{\tau}}{h} \partial_\varphi^2 \phi_{n,j}^-|_{\partial_\varphi \phi_{n,j}^- = 0} \gtrsim \frac{2^{-2j}|z|}{h}$. The stationary phase yields

$$I_{\chi_+}^{-,n,j} = \frac{1}{h^2} \int e^{i\tilde{\tau}\phi_{n,j}^-} \chi(\tilde{\tau}) \frac{(\tilde{\tau}/h)^{-1/2}}{\sqrt{\partial_\varphi^2 \phi_{n,j}^-|_{\partial_\varphi \phi_{n,j}^- = 0}}} \left(\tilde{J}_{\chi_+}^{-,n,j}(r, s, \frac{\tilde{\tau}}{2^j nh}) + h 2^{-j} O((2^{-2j}|z|/h)^{-\infty}) \right) d\tilde{\tau}, \quad (4.13)$$

where $\tilde{J}_{\chi_+}^{-,n,j}(r, s, \frac{\tilde{\tau}}{2^j nh})$ is the symbol with main contribution $J_{\chi_+}^{-,n,j}$ introduced in (4.8) and where $h 2^{-j}$ comes from the factors $2^{-2j} \times \frac{1}{n} \times \frac{nh}{2^{-j}}$ of the symbol. In order to uniformly bound the sum of (4.13), notice that the phase is stationary when $t \sim |z| \sim 2^j r$. As $2^{-4j}|z| \lesssim h$, then $|z|^{1/2} \leq h^{1/2} 2^{2j}$ and

$$|I_{\chi_+}^{-,n,j}| \lesssim \frac{1}{h^2 t} \times \frac{h^{1/2}|z|^{1/2} 2^{-2j+j}}{n} \times nh 2^j \leq \frac{1}{h^2 t} \times h^2 2^{2j}, \quad h^2 \sum_{n_0 \leq n, 2^j < 1/(hn)} 2^{2j} \leq h \sum 2^j \lesssim, \quad (4.14)$$

where we used that $n \leq 2^{-j}/h$ on the support of χ_+ . Notice that the condition $j \geq j(r,h)$ was particularly useful here in order to obtain the sharp bounds in (4.14). In the same way one may deal with $I_{\chi_+}^{+,n,j}$ and obtain similar bounds. The proof of the Lemma is achieved. \square

Next, we turn to $I_{\chi_0}^{\pm, n, j}$ whose symbols are supported for $\frac{\tau\sqrt{1-\gamma^2}}{n} \in [1 - 2\varepsilon, 1 + 2\varepsilon]$. For each $j \geq 1$, it will be convenient to take $\tau 2^{-j}\varphi = n + n^{1/3}w$: on the support of the symbol of $I_{\chi_0}^{\pm, n, j}$ we now have $wn^{-2/3} \in [-2\varepsilon, +2\varepsilon]$ and $2^{-j}/h \sim n \geq 1$ as $\tau \sim 1/h$. Write again $1 = \sum_{* \in \{0, \pm\}} \chi_*(w)$ where $\chi_\pm(\ell) = (1 - \chi_0)(\ell) 1_{\pm\ell > 0}$ and denote $I_{\chi_0, \chi_*}^{\pm, n, j}$ the corresponding integrals (defined as in (4.6) but with additional cutoffs $\chi_*(n^{2/3}(\frac{\sqrt{\tau^2 - \vartheta^2}}{n} - 1))$). We deal separately with the cases $w > 1$, $|w| \leq 2$ and $w < -1$.

Lemma 4.3. *For $1 < s \leq r \leq \sqrt{2}$ we have $\sum_{n \geq n_0, j \geq 1} |I_{\chi_0, \chi_*}^{\pm, n, j}| \lesssim \frac{1}{h^2 t}$, $* \in \{0, +\}$. For $r \geq s$ with $r \geq \sqrt{2}$ and $j(r,h)$ as in Definition 2.3 we have $\sum_{n \geq n_0, j \geq j(r,h)} |I_{\chi_0, \chi_*}^{\pm, n, j}| \lesssim \frac{1}{h^2 t}$, $* \in \{0, +\}$.*

Proof. On the support of $\chi_+(w)$ we may proceed in a similar way as in Lemma 4.2 as the same asymptotic expansions hold for the Hankel factors; as the computations are similar (modulo the change of variable w.r.t.

τ) we focus on $I_{\chi_0, \chi_0}^{-, n, j}$ with symbol $\chi_0(w)$. The expansion (4.7) still holds (with the simpler form (6.5)): when $n^{2/3}(-\tilde{\zeta}(\tilde{\rho})) < 2$ (with $\tilde{\rho} \in \{\rho, r\rho, s\rho\}$), the Airy factors don't oscillate and may be brought into the symbol. Let first $1 < s \leq r \leq \sqrt{2}$ when the last inequality holds. The phase of $I_{\chi_0, \chi_0}^{-, n, j}$ equals $\tau(t + z\sqrt{1 - 2^{-2j}\varphi^2})$, and taking $\tau = \frac{2^j}{\varphi}(n + n^{1/3}w)$ we are reduced to obtaining uniform bounds for

$$\frac{s^2}{(rs)^{1/2}} \int e^{i\frac{2^j}{\varphi}(n+n^{1/3}w)(t+z\sqrt{1-2^{-2j}\varphi^2})} \chi(h\frac{2^j}{\varphi}(n+n^{1/3}w)) \frac{2^j}{\varphi}(n+n^{1/3}w) \psi(\varphi) \chi_0(w) n^{-2/3+1/3} \frac{2^{-2j+j}}{\varphi} d\varphi dw, \quad (4.15)$$

where the factors $2^{-2j+j} \frac{n^{1/3}}{\varphi}$ come from $\gamma \rightarrow \varphi$, $\tau \rightarrow w$ and where $n \sim \frac{2^{-j}}{h}$. For $t \lesssim h^{-1/3}$, the sum of all contributions of the form (4.15) may be bounded as follows

$$\sum_{j, n} |I_{\chi_0, \chi_0}^{-, n, j}| \leq \sum_{j, n \sim 2^{-j}/h} n^{2/3} \sim \sum_{j, h^{1/3}2^{j/3} \leq 1} (2^{-j}/h)^{5/3} \leq \frac{h^{1/3}}{h^2} \lesssim \frac{1}{h^{2t}}. \quad (4.16)$$

For $t \gtrsim h^{-1/3}$ satisfying $t \geq 2|z|$, the phase is non-stationary w.r.t. w ; integrations by parts with the large parameter $2^j n^{1/3} \sim 2^{2j/3}/h^{1/3}$ yield a contribution $O((2^j n^{1/3}/|t|)^{-N})$ for all $N \geq 1$ and we conclude. For $h^{-1/3} \lesssim t \leq 4|z|$ we have $\frac{1}{|z|} \leq \frac{4}{t}$ and we apply the stationary phase in both w, φ : let $\phi_{0, n, j}^- := \frac{2^j}{\varphi}(n + n^{1/3}w)(t + z\sqrt{1 - 2^{-2j}\varphi^2})$ then $\partial_w^2 \phi_{0, n, j}^- = 0$ and the determinant of the Hessian matrix equals $(\partial_{w, \varphi}^2 \phi_{0, n, j}^-)^2 \sim (2^{-j} n^{1/3} |z|)^2$ for $\varphi \sim 1$. If $2^{-j} n^{1/3} |z| \geq h^{-\epsilon}$ for some small $\epsilon > 0$, we find, for small r, s ,

$$\sum_{j, n \geq n_0} |I_{\chi_0, \chi_0}^{-, n, j}| \leq \sum_{j, n \sim 2^{-j}/h} \frac{n^{2/3}}{2^{-j} n^{1/3} |z|} \lesssim \sum_{j, 2^j < 1/h} \frac{2^j}{|z|} \times (2^{-j}/h)^{4/3} (1 + O(h^\infty)) \lesssim \frac{h^{2/3}}{h^{2t}}.$$

If $2^{-j} n^{1/3} |z| \leq 2h^{-\epsilon}$ then we bound the sum of $|I_{\chi_0, \chi_0}^{-, n, j}|$ as in (4.16) by $\sum_{j, n \sim 2^{-j}/h} n^{2/3}$ and use that $2^{-j/3}/h^{1/3} \sim n^{1/3} \leq 2^{j+1} h^{-\epsilon}/|z|$ which gives $\sum_{j, n \sim 2^{-j}/h} n^{2/3} \leq \frac{1}{h^{2|z|}} h^{2/3-\epsilon} \sum_j 2^{j+1-4j/3} \lesssim \frac{h^{2/3-\epsilon}}{h^{2t}}$.

Let now $r \geq \sqrt{2}$ such that $n^{2/3}(-\tilde{\zeta}(r\rho)) > 1$. If moreover $n^{2/3}(-\tilde{\zeta}(s\rho)) > 1$, then both Airy factors $A(n^{2/3}\tilde{\zeta}(r\rho))$, $A(n^{2/3}\tilde{\zeta}(s\rho))$ do oscillate and we may proceed as with $\chi_+(w)$ (the only differences with the case χ_+ are the absence of the phase functions of $\overline{H}_n(n\rho)/H_n(n\rho)$, which means replacing $f_1(r, \rho)$ by $\sqrt{(r\rho)^2 - 1}$, and also the fact that the factor depending on r, s, ρ in (4.14) may not be bounded but at most $n^{1/3}$). Consider $n^{2/3}(-\tilde{\zeta}(s\rho)) \leq 2$, then $|H_n(n\rho)| \sim n^{-1/3}$ and the symbol of $I_{\chi_0, \chi_0}^{-, n, j}$ becomes

$$J_{\chi_0, \chi_0}^{-, n, j}(r, s, \rho) := n^{-2 \times \frac{1}{3} - \frac{1}{6}} \frac{s^2 \Sigma_0(r, s, \rho, n)}{(rs)^{1/2} ((r\rho)^2 - 1)^{\frac{1}{4}}} \psi(\varphi) \chi_0(w) \chi(h\frac{2^j}{\varphi}(n + n^{1/3}w)) \frac{2^j}{\varphi}(n + n^{1/3}w) 2^{-2j} \frac{2^j n^{1/3}}{\varphi}$$

where the elliptic symbol Σ_0 is an asymptotic expansion with small parameter n^{-1} and with main contribution obtained as a product of a_0 in (6.3), σ_0 in (6.1) and $H_n(n\rho) n^{-1/3} \times \frac{\overline{H}_n(n\rho)}{H_n(n\rho)}$. The factor $((r\rho)^2 - 1)^{-1/4}$ is always bounded by $n^{1/6}$. The factors $2^j(n + n^{1/3}w) \times 2^{-2j} \times 2^j n^{1/3}/\varphi$ occur from the changes of variables $\vartheta \rightarrow \tau\gamma$, $\gamma \rightarrow \varphi$, $\tau \rightarrow w$. If $t \lesssim h^{-1/3}$ we conclude as in (4.16). Let $t \geq h^{-1/3}$. The phase $\phi_{0, n, j}^- := \tau(t + z\sqrt{1 - 2^{-2j}\varphi^2}) - \frac{2}{3}n(-\tilde{\zeta}(r\rho))^{3/2}$ is not stationary for z such that $\tau 2^{-j}|z| \sim 2^j n \times 2^{-2j}|z| \sim 2^{-j} n |z| \geq h^{-\epsilon}$ for some small $\epsilon > 0$ and we perform repeated integrations by parts to conclude. If $2^{-j} n |z| \leq 2h^{-\epsilon}$, then for $|t| \geq 4|z|$ we integrate by parts, while for $|t| \leq 4|z|$ we use $2^{-j} n |z| \sim 2^{-2j}|z|/h \leq 2h^{-\epsilon}$, $2^{-j}/h \geq 1$ to obtain

$$\sum_{j \geq 1, n \sim 2^{-j}/h} |I_{\chi_0, \chi_0}^{-, n, j}| \leq \sum_{n \sim 2^{-j}/h, 2^{-2j} \leq h^{1-\epsilon}/|z|} n^{2/3} \leq \frac{h^{1/3}}{h^2} \sum_{h^2 \leq 2^{-2j} \leq 2h^{1-\epsilon}/|z|} 2^{-2j+j/3} < \frac{h^{1-\epsilon}}{h^{2t}}.$$

Let $r \geq \sqrt{2}$ and $j \geq j(r, h)$: for s such that $n^{2/3}(-\tilde{\zeta}(s\rho)) \leq 2$ we conclude as before (with an additional factor $1/r$ in the symbol). For $n^{2/3}(-\tilde{\zeta}(s\rho)) \geq 1$, the situation is similar to the one of χ_+ dealt with before. \square

Lemma 4.4. *For $1 < s \leq r \leq \sqrt{2}$ we have $\sum_{n \geq n_0, j \geq 1} |\sum_{\pm} I_{\chi_0, \chi_{\pm}}^{\pm, n, j}| \lesssim \frac{1}{h^{2t}}$. For $r \geq s$ with $r \geq \sqrt{2}$ and $j(r, h)$ given in Definition 2.3, we also have $\sum_{n \geq n_0, j \geq j(r, h)} |\sum_{\pm} I_{\chi_0, \chi_{\pm}}^{\pm, n, j}| \lesssim \frac{1}{h^{2t}}$.*

Proof. Recall that $1 - \gamma^2 = 2^{-2j}\varphi^2$, with $\varphi \sim 1$ on the support of ψ , and $\rho = \tau\sqrt{1 - \gamma^2}/n = 1 + n^{-2/3}w$: as $w < -1$ on the support the symbol of $I_{\chi_0, \chi_-}^{\pm, n, j}$ then $\rho \in [1 - \varepsilon, 1 - n^{-2/3}]$. It will be convenient to use the representation of G_n in terms of Bessel functions J_n instead of H_n , hence the first line in (4.2). We estimate

$$\begin{aligned} & \frac{s^2}{(rs)^{1/2}} \sum_{n \geq 1} e^{in\theta} \sum_{j \geq 1} \int e^{i\frac{2j(n+n^{1/3}w)}{\varphi}(t-z\sqrt{1-2^{-2j}\varphi^2})} \chi\left(\frac{2^j h(n+n^{1/3}w)}{\varphi}\right) \frac{J_n(n(1+n^{-\frac{2}{3}}w))}{H_n(n(1+n^{-\frac{2}{3}}w))} \\ & \times \psi(\varphi)\chi_-(w)H_n(nr(1+n^{-\frac{2}{3}}w))H_n(ns(1+n^{-\frac{2}{3}}w)) \frac{2^j}{\varphi} (n+n^{1/3}w)2^{-2j+j}n^{\frac{1}{3}}dwd\varphi. \end{aligned} \quad (4.17)$$

The Bessel function $J_n(n\rho)$ is given by (6.4). The factor J_n/H_n corresponds to the quotient $\frac{A}{A_+}(n^{\frac{2}{3}}\tilde{\zeta}(\rho)) = e^{-2i\pi/3} + e^{-\frac{4}{3}|n|\tilde{\zeta}(\rho)^{\frac{3}{2}}}$ (see Lemma 6.1). On the support of the cut-offs of $I_{\chi_0, \chi_-}^{-, n, j}$, its symbol has the form

$$J_{\chi_0, \chi_-}^{-, n, j} := \frac{s^2}{(rs)^{1/2}} n^{-2/3} \left(\frac{4\tilde{\zeta}(r\rho)}{1-(r\rho)^2}\right)^{1/4} \left(\frac{4\tilde{\zeta}(s\rho)}{1-(s\rho)^2}\right)^{1/4} A_+(n^{2/3}\tilde{\zeta}(r\rho))A_+(n^{2/3}\tilde{\zeta}(s\rho))e^{-\frac{4}{3}n\tilde{\zeta}(\rho)^{\frac{3}{2}}}\Sigma_-, \quad (4.18)$$

for some symbol Σ_- of order 0. In the case of $I_{\chi_0, \chi_-}^{+, n, j}$ one should replace $A_+(n^{2/3}\tilde{\zeta}(s\rho))$ by $A(n^{2/3}\tilde{\zeta}(s\rho))$ and remove the exponential decreasing factor. When $n^{2/3}|\tilde{\zeta}(r\rho)|, n^{2/3}|\tilde{\zeta}(s\rho)| < 2$ we can proceed exactly as for $I_{\chi_0, \chi_0}^{-, n, j}$ with $r-1, s-1 \lesssim n^{-2/3}$ small. Assume $n^{2/3}\tilde{\zeta}(s\rho) \geq n^{2/3}\tilde{\zeta}(r\rho) \geq 1$ with $\rho \leq 1 - \frac{1}{n^{2/3}} < 1$, then

$$J_{\chi_0, \chi_-}^{-, n, j} := \frac{s^2}{(rs)^{1/2}} \frac{n^{-\frac{1}{3}-\frac{1}{6}}}{(1-(r\rho)^2)^{\frac{1}{4}}} \frac{n^{-\frac{1}{3}-\frac{1}{6}}}{(1-(s\rho)^2)^{\frac{1}{4}}} e^{-\frac{4}{3}n\tilde{\zeta}(\rho)^{\frac{3}{2}} + \frac{2}{3}n\tilde{\zeta}(s\rho)^{\frac{3}{2}} + \frac{2}{3}n\tilde{\zeta}(r\rho)^{\frac{3}{2}}}\Sigma_-. \quad (4.19)$$

As we are assuming $\tilde{\zeta}(r\rho), \tilde{\zeta}(s\rho) > 0$, we have, using Lemma 2.5, $\frac{2}{3}\tilde{\zeta}(s\rho)^{3/2} - \frac{2}{3}\tilde{\zeta}(r\rho)^{3/2} = -\int_{\rho}^{s\rho} \frac{\sqrt{1-w^2}}{w}dw \leq 0$.

The phase function of $I_{\chi_0, \chi_-}^{\pm, n, j}$ is $\tau(t+z\gamma)$ and the factor $\frac{s^2}{(rs)^{1/2}(1-(s\rho)^2)^{\frac{1}{4}}(1-(r\rho)^2)^{\frac{1}{4}}}$ is at most $n^{1/3}$ when $1-s\rho \sim 1-r\rho \sim n^{-2/3}$, while for $r, s \geq 2$ this term is uniformly bounded by 1. From now one can proceed as in the case of $I_{\chi_0, \chi_0}^{-, n, j}$ as on the support of $\chi(h\tau)$ we still have $n \sim 2^{-j}/h$, the phase is stationary for $t \sim |z|$ and for $2^{-j}n|z| \geq h^{-\varepsilon}$ we integrate by parts, while for $2^{-j}n|z| \leq 2h^{-\varepsilon}$ we conclude as done previously. \square

Lemma 4.5. *For $1 < s \leq r \leq \sqrt{2}$ we have $\sum_{n \geq n_0, j \geq 1} |\sum_{\pm} I_{\chi_{\pm}}^{\pm, n, j}| \lesssim \frac{1}{h^2 t}$. For $r \geq s$ with $r \geq \sqrt{2}$ and $j(r, h)$ as in Definition 2.3, we also have $\sum_{n \geq n_0, j \geq j(r, h)} |\sum_{\pm} I_{\chi_{\pm}}^{\pm, n, j}| \lesssim \frac{1}{h^2 t}$.*

Proof. On the support of $I_{\chi_{\pm}}^{-, n, j}$ we have $\rho = \frac{\tau\sqrt{1-\gamma^2}}{n} \leq 1 - \varepsilon$. The symbol of $I_{\chi_{\pm}}^{-, n, j}$ has also the form (4.18). For small $r, s \leq \sqrt{2}$ and $\varepsilon > 2(\sqrt{2}-1)$, we write $1-r\rho = 1-r+r(1-\rho)$ to deduce that, if $\rho \leq 1 - \varepsilon$, then the symbol (4.18) takes the form (4.19) where the factor $\frac{s^2}{(rs)^{1/2}(1-(s\rho)^2)^{\frac{1}{4}}(1-(r\rho)^2)^{\frac{1}{4}}}$ is uniformly bounded by a constant depending only on ε and we conclude as before. When r, s are large (and $\tau 2^{-2j}|z| \leq M, \tau 2^{-j}r \leq M$ for large $M > 1$), we separate the possible situations: the only new one is the case $n^{2/3}(1-r\rho), n^{2/3}(1-s\rho) \geq 1$ and r such that $r < 1/\rho \leq 1/(1-\varepsilon)$, in which case $\frac{s^2}{(rs)^{1/2}(1-(s\rho)^2)^{\frac{1}{4}}(1-(r\rho)^2)^{\frac{1}{4}}} \leq rn^{1/3} \leq \frac{n^{1/3}}{(1-\varepsilon)}$. In this case we have additional decay from the exponential factors and conclude as before. \square

5. SMALL FREQUENCY CASE

Let $\tau \leq 1/h_0$ for some fixed $h_0 > 0$, small enough. We use again the parametrix in terms of Bessel functions introduced in Section 4 and keep the same notations. We split $I = I^+ + I^-$, and for $n \geq 1$ large enough, $I^{\pm} = \sum_{* \in \{0, \pm\}} I_{\chi_*}^{\pm}$, with $I_{\chi_*}^{\pm}$ introduced as a sum of $I_{\chi_*}^{\pm, n, j}$ given in (4.6) where $\chi(h\tau)$ is replaced by $\tilde{\chi}(\tau)$ supported for $\tau \leq 2/h_0$. Take $n_0 = 4/h_0$. We aim at proving that $|\sum_{\pm} I_{\chi_*}^{\pm}| \lesssim C(h_0)/t$.

- On the support of $I_{\chi_*}^{\pm}$, $* \in \{+, 0\}$, and for $n \geq n_0$ we have $n_0 \leq n \leq \frac{\tau\sqrt{1-\gamma^2}}{1-\varepsilon} < \frac{4}{h_0} = n_0$.
- On the support of $I_{\chi_*}^{\pm}$, $* \in \{+, 0\}$, and for $1 \leq n \leq n_0$ as $\sqrt{1-\gamma^2} \sim 2^{-j}$ and $\tau \leq 2/h_0$, only a finite number of j such that $2^j \leq 1/(h_0(1-\varepsilon))$ may contribute. For each j, n on this finite set, the symbols of $I_{\chi_*}^{\pm}$ are bounded and their phase may oscillate only for large t or large $|z|$. If t is bounded then if r or $|z|$ are larger than $\max\{4t, M\}$ for some $M > 1$ large enough, integrations by parts allow to conclude (using that the sum is finite); if $|z|, r \leq 4t$ each integral is bounded and we obtain $|I_{\chi_*}^{\pm}| \lesssim C(h_0)$.

If t be sufficiently large, then if $t/(|z| + 2^j r) \notin [1/8, 8]$, integrations by parts yields a contribution $O(1/t^N)$ for each pair (j, n) on the support of $I_{\chi_*^\pm}^{n,j}$. If $t/(|z| + 2^j r) \in [1/8, 8]$, we separate the cases $2^{-2j}|z| \geq M$ for some large M , when we apply the stationary phase in $\varphi = 2^j \sqrt{1 - \gamma^2}$ and we conclude as in (4.14) or $2^{-2j}|z| \leq M$, when we bound directly as in (4.12).

- On the support of $I_{\chi_-}^\pm$ we have $n \geq \tau \sqrt{1 - \gamma^2}/(1 - \varepsilon)$, hence the sum over n is unbounded but as $n \gg \tau \sqrt{1 - \gamma^2}$ we may use (6.7) and conclude.

6. APPENDIX

6.1. Airy functions. For $w \in \mathbb{C}$, the Airy function is defined as follows : $A(w) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{i(s^3/3 + sw)} ds$. Let $A_\pm(w) := A(e^{\mp 2i\pi/3} w)$, then $A_-(w) = \overline{A_+(\overline{w})}$ and $A(w) = e^{i\pi/3} A_+(w) + e^{-i\pi/3} A_-(w)$. Moreover, $A_\pm(w), A'_\pm(w)$ are not zero for any $w \in \mathbb{R}$, while all the zeros of $A(w)$ and $A'(w)$ are real and non positive. We say that $f(w)$ admits an asymptotic expansion for $w \rightarrow 0$ if there exists $(c_j)_{j \in \mathbb{N}}$ such that for any $j \geq 0$ we have $\lim_{w \rightarrow 0} w^{-j-1}(f(w) - \sum_0^j c_i w^i) = c_{j+1}$. We write $f(w) \sim_w \sum_j c_j w^j$.

Lemma 6.1. *Let $\Sigma(w) := (A_+(w)A_-(w))^{1/2}$, then $\Sigma(z) = |A_+(w)| = |A_-(w)|$ is real, monotonic increasing in w and nowhere vanishing. We let $\mu(w) := \frac{1}{2i} \log(\frac{A_-(w)}{A_+(w)})$ for $w < -1/4$. Then $A_\pm(w) = \Sigma(w)e^{\mp i\mu(w)}$. For $w < -1$, the following asymptotic expansions hold*

$$\Sigma(w) \sim_{\frac{1}{w}} (-w)^{-\frac{1}{4}} \sum_{j \geq 0} \sigma_j (-w)^{-\frac{3j}{2}}, \quad \mu(w) \sim_{\frac{1}{w}} \frac{2}{3} (-w)^{\frac{3}{2}} \sum_{j \geq 0} e_j (-w)^{-\frac{3j}{2}}, \quad \sigma_0 = \frac{1}{2\sqrt{\pi}}, \quad e_0 = 1. \quad (6.1)$$

The Airy quotient $\Phi_+(w) = \frac{A'_+(w)}{A_+(w)} = \frac{\Sigma'(w)}{\Sigma(w)} - i\mu'(w)$ satisfies everywhere $\Phi'_+(w) = w - \Phi_+^2(w)$. In particular $\Phi'_+(w)$ is bounded on $(-\infty, -1)$ and $\Phi_+(w) \sim_{\frac{1}{w}} (-w)^{\frac{1}{2}} \sum_{j \geq 0} d_j (-w)^{-\frac{3j}{2}}$, $d_0 = 1$, for $(-w) > 1$ large.

For $w > 1$, the functions $A_\pm(w)$ grow exponentially $A_\pm(w) = \Sigma_\pm(w)e^{\frac{2}{3}w^{3/2}}$, where Σ_\pm are classical symbols of order $-1/4$ and we have $\frac{A_-(w)}{A_+(w)} + e^{2i\pi/3} = O(w^{-\infty})$ when $w \rightarrow \infty$ and $\frac{A_-(w)}{A_+(w)} \sim_{\frac{1}{w}} e^{2i\mu(w)}$ when $w \rightarrow -\infty$. Moreover the Airy function $A(w)$ decays exponentially for $w > 1$, $A(w) \sim_{\frac{1}{w}} |w|^{-\frac{1}{4}} e^{-\frac{2}{3}w^{3/2}}$.

6.2. Bessel and Hankel functions. The Hankel function $H_\nu(z)$ is a solution to the Bessel's equation $w^2 H''_\nu(w) + w H'_\nu(w) + (w^2 - \nu^2) = 0$. The couple $\{H_\nu(w), \overline{H}_\nu(w)\}$ is a fundamental system of solutions for the Bessel equation. The real and imaginary part of $H_\nu(w)$, denoted $J_\nu(w)$ and $Y_\nu(w)$ respectively, are the usual Bessel function of the first and second type. The Hankel function of order ν is defined by ([1, (9.1.25)])

$$H_\nu(w) = \int_{-\infty}^{+\infty - i\pi} e^{w \sinh t - \nu t} dt. \quad (6.2)$$

For large positive order ν and $w = \nu\rho$, the Hankel functions have the following expansions that hold *uniformly* with respect to ρ in the sector $|\arg(\rho)| < \pi - \varepsilon$, where $\varepsilon > 0$ is an arbitrary number [1, (9.3.37)]:

$$H_\nu(\nu\rho) = 2e^{-\frac{i\pi}{3}} \left(\frac{-4\tilde{\zeta}(\rho)}{\rho^2 - 1} \right)^{\frac{1}{4}} \left(\nu^{-\frac{1}{3}} A_+(\nu^{\frac{2}{3}} \tilde{\zeta}(\rho)) \left(\sum_{j \geq 0} a_j(\tilde{\zeta}) \nu^{-2j} \right) + \nu^{-\frac{5}{3}} A'_+(\nu^{\frac{2}{3}} \tilde{\zeta}(\rho)) \left(\sum_{j \geq 0} b_j(\tilde{\zeta}) \nu^{-2j} \right) \right), \quad (6.3)$$

$$J_\nu(\nu\rho) = 2e^{-\frac{i\pi}{3}} \left(\frac{-4\tilde{\zeta}(\rho)}{\rho^2 - 1} \right)^{\frac{1}{4}} \left(\nu^{-\frac{1}{3}} A(\nu^{\frac{2}{3}} \tilde{\zeta}(\rho)) \left(\sum_{j \geq 0} a_j(\tilde{\zeta}) \nu^{-2j} \right) + \nu^{-\frac{5}{3}} A'(\nu^{\frac{2}{3}} \tilde{\zeta}(\rho)) \left(\sum_{j \geq 0} b_j(\tilde{\zeta}) \nu^{-2j} \right) \right). \quad (6.4)$$

Here $a_j(\tilde{\zeta}), b_j(\tilde{\zeta})$ are given in [1, (9.3.40)] and $\tilde{\zeta}(\rho)$ is provided in Lemma 2.5 (see [1, (9.3.38),(9.3.39)]). When $\rho = 1 + \nu^{-2/3}v$, $v = O(1)$, $w = \nu\rho = \nu + \nu^{1/3}v$, these formulas reduce to (see [1, (9.3.23),(9.3.24)])

$$H_\nu(\nu + \nu^{1/3}v) = \frac{2^{1/3}}{\nu^{1/3}} A_+(-2^{1/3}v) \left(1 + \sum_{j \geq 1} \tilde{a}_j(v) \nu^{-2j/3} \right) + \frac{2^{2/3}}{\nu} A'_+(-2^{1/3}v) \left(\sum_{j \geq 0} \tilde{b}_j(v) \nu^{-2j/3} \right), \quad (6.5)$$

$$J_\nu(\nu + \nu^{1/3}v) = \frac{2^{1/3}}{\nu^{1/3}} A(-2^{1/3}v) \left(1 + \sum_{j \geq 1} \tilde{a}_j(v) \nu^{-2j/3} \right) + \frac{2^{2/3}}{\nu} A'(-2^{1/3}v) \left(\sum_{j \geq 0} \tilde{b}_j(v) \nu^{-2j/3} \right). \quad (6.6)$$

where \tilde{a}_j, \tilde{b}_j are polynomials in v given in [1, (9.3.25),(9.3.26)].

Remark 6.2. *The formulas (6.3), (6.4) are among the deepest and most important results in the theory of Bessel functions. In order to prove (6.4) starting from (6.2) one may chose a suitable contour that yields $J_\nu(\nu\rho) = (2\pi)^{-1} \int e^{i\nu\phi(\rho,t)} dt$ with $\phi(\rho,t) = \rho \sin t - t$; for ν large enough and for $\rho > 1$, the critical point $t(\rho) := \arccos(1/\rho)$ is real and the critical value equals $\phi(\rho, t(\rho)) = \sqrt{\rho^2 - 1} - \arccos(1/\rho) = \frac{2}{3}(-\tilde{\zeta})^{3/2}$, where $\tilde{\zeta}(\rho)$ is defined as in (2.7). As the phase function of $A(\nu^{2/3}\tilde{\zeta})$ equals $\nu(s^3 + s\tilde{\zeta})$ and has critical points $s^2 = -\tilde{\zeta}$ and critical values $\pm\frac{2}{3}(-\tilde{\zeta})^{3/2}$, one obtains (6.4) by stationary phase (see [13] for details).*

When the order is much larger than the argument $n \gg w$, (6.3), (6.4) reduce to (see [1, (9.3.1)])

$$J_n(w) = \sqrt{\frac{1}{2\pi n}} \left(\frac{ew}{2n}\right)^n \left(1 + O\left(\frac{|w|}{n}\right)\right), \quad Y_n(w) = -\sqrt{\frac{1}{2\pi n}} \left(\frac{ew}{2n}\right)^{-n} \left(1 + O\left(\frac{|w|}{n}\right)\right), \quad n \gg 1. \quad (6.7)$$

As we consider cylindrical coordinates we deal only with $\nu = n \in \mathbb{Z}$: in view of the well-known relations $H_{-n}(w) = (-1)^n H_n(w)$ (see [1, (9.1.6)]), we may consider only non negative values of n in our discussion.

REFERENCES

- [1] M.Abramowitz and I.A.Stegun. *Handbook of mathematical functions, with formulas, graphs, and mathematical tables..* Edited by Milton Abramowitz and Irene A. Stegun. Dover publications Inc., New York, 1966.
- [2] C.Chester, B.Friedman, F.Ursell. *An extension of the method of steepest descents.* Proc. Cambridge Philos. Soc., 53:599-611,1957
- [3] J.Ginibre and G.Velo The global Cauchy problem for the nonlinear Schrödinger equation revisited Ann. Inst. H. Poincaré Anal. Non Linéaire, 2(4):309-327, 1985
- [4] J. Ginibre and G. Velo. The global Cauchy problem for the nonlinear Klein-Gordon equation. *Math. Z.*, 189(4):487–505, 1985.
- [5] J.Ginibre and G.Velo. Generalized Strichartz inequalities for the wave equation Partial differential operators and mathematical physics (Holzhau, 1994), vol. 78 of Open Theory Adv. Appl. p 153-160, Birkhäuser, Basel, 1995
- [6] L. Hörmander. *The Analysis of Linear Partial Differential Operators III.* Springer, 1994
- [7] O.Ivanovici and G.Lebeau. Dispersion for the wave and Schrödinger equations outside strictly convex obstacles and counterexamples. preprint <https://arxiv.org/pdf/2012.08366.pdf>
- [8] H.Lindblad and Ch. Sogge. On existence and scattering with minimal regularity for semi-linear wave equation *J.Funct. Anal.*, 130(2):357-426, 1995
- [9] L.Kapitanski. Some generalizations of the Strichartz-Brenner inequality *Algebra i Analiz*, 1(3):127-159, 1989
- [10] M.Keel and T.Tao. Endpoint Strichartz estimates *Amer. J. Math.*, 120(5):955-980, 1998
- [11] L. Meas. *Dispersive estimates for the wave equation inside cylindrical convex domains* *Comptes Rendus Mathématique*, vol.355, issue 2, 161–165 (2017)
- [12] R. Melrose, M.Taylor. *Boundary problems for the wave equations with grazing and gliding rays*, 1987.
- [13] F.Olver. *Asymptotics and Special functions.* Academic Press, New York, 1974
- [14] H. F. Smith. A parametrix construction for wave equations with $C^{1,1}$ coefficients. *Ann. Inst. Fourier (Grenoble)*, 48(3):797–835, 1998.
- [15] H. F. Smith and Ch. D. Sogge. L^p regularity for the wave equation with strictly convex obstacles. *Duke Math. J.*, 73(1):97–153, 1994.
- [16] H. F. Smith and Ch. D. Sogge. On the critical semilinear wave equation outside strictly convex obstacles. *J. Amer. Math. Soc.*, 8(4):879–916, 1995.
- [17] R.Strichartz. Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equation *Duke Math. J.* 44(3):705-714, 1977
- [18] D.Tataru. Strichartz estimates for second order hyperbolic operators with non-smooth coefficients III *J.Amer.Math.Soc.*, 15(2):419-442 (electronic), 2002
- [19] M. Taylor. *Partial Differential Equations II.* Springer, 1996.
- [20] M. Zworski. High frequency scattering by a convex obstacle. *Duke Math. J.*, 61(2):545–634, 1990.

*SORBONNE UNIVERSITÉ, CNRS, LJLL, F-75005 PARIS, FRANCE

Email address: felice.iandoli@sorbonne-universite.fr, oana.ivanovici@sorbonne-universite.fr