# Homomorphisms of (n, m)-graphs with respect to generalized switch

SAGNIK SEN<sup>a</sup>, ÉRIC SOPENA<sup>b</sup>, S TARUNI<sup>a</sup>

- (a) Indian Institute of Technology Dharwad, India.
  - (b) Université de Bordeaux, LaBRI, France.

#### Abstract

The study of homomorphisms of (n, m)-graphs, that is, adjacency preserving vertex mappings of graphs with n types of arcs and m types of edges was initiated by Nešetřil and Raspaud [Journal of Combinatorial Theory, Series B 2000]. Later, some attempts were made to generalize the switch operation that is popularly used in the study of signed graphs, and study its effect on the above mentioned homomorphism.

In this article, we too provide a generalization of the switch operation on (n, m)graphs, which to the best of our knowledge, encapsulates all the previously known generalizations as special cases. We approach to study the homomorphism with respect to the switch operation axiomatically. We prove some fundamental results that are essential tools in the further study of this topic. In the process of proving the fundamental results, we have provided yet another solution to an open problem posed by Klostermeyer and MacGillivray [Discrete Mathematics 2004]. We also prove the existence of a categorical product for (n, m)-graphs on with respect to a particular class of generalized switch which implicitly uses category theory. This is a counter intuitive solution as the number of vertices in the Categorical product of two (n, m)-graphs on p and q vertices has a multiple of pq many vertices, where the multiple depends on the switch. This solves an open question asked by Brewster in the PEPS 2012 workshop as a corollary. We also provide a way to calculate the product explicitly, and prove general properties of the product. We define the analog of chromatic number for (n, m)-graphs with respect to generalized switch and explore the interrelations between chromatic numbers with respect to different switch operations. We find the value of this chromatic number for the family of forests using group theoretic notions.

**Keywords:** colored mixed graphs, switching, homomorphisms, categorical product, chromatic number.

#### 1 Introduction

A graph homomorphism, that is, an edge-preserving vertex mapping of a graph G to a graph H, also known as an H-coloring of G, was introduced as a generalization of

coloring [15]. It allows us to unify certain important constraint satisfaction problems, especially related to scheduling and frequency assignments, which are otherwise expressed as various coloring and labeling problems on graphs [15]. Thus the notion of graph homomorphism manages to capture a wide range of important applications in an uniform setup. When viewed as an operation on the set of all graphs, it induces rich algebraic structures: a quasi order (and a partial order), a lattice, and a category [15].

The study of graph homomorphism can be characterized into three major branches:

- (i) The study of various application motivated optimization problems are modeled using graph homomorphisms. These usually involve finding an H having certain prescribed properties such that every member of a graph family  $\mathcal{F}$  is H-colorable [14, 27, 32].
- (ii) The study of the algorithmic aspects of the *H*-coloring problem, including characterizing its dichotomy, and finding exact (polynomial), approximation or parameterized algorithms for the hard problems [6, 11, 12].
- (iii) The study of the algebraic structures that gets induced by the notion of graph homomorphisms [15].

Unsurprisingly, these three areas of research have interdependencies and connections. The notion of graph homomorphisms, initially introduced for undirected and directed graphs, later got extended to 2-edge-colored graphs [1], k-edge-colored graphs [13] and (n,m)-graphs [24]. These graphs, due to their various types of adjacencies, manages to capture complex relational structures and are useful for mathematical modeling. For instance, the Query Evaluation Problem (QEP) in graph database (the immensely popular databases that are now used to handle highly interrelated data in social networks like Facebook, Twitter, etc.), is modeled on homomorphisms of (n,m)-graphs [2, 4].

From a theoretical point of view, apart from being the generalization of the well-studied graph homomorphisms of oriented, signed, 2-edge-colored, and k-edge-colored graphs, homomorphisms of (n, m)-graphs is known to have connections with some topics in graph theory as well as other mathematical disciplines. For instance, connections with acyclic coloring, harmonious coloring, and flows are pointed out in [24], [1], and [5], respectively. On the other hand, connections with the study of binary predicate logic, and Coxeter groups are established in [24] and [1]. It is worth mentioning that in a recent work by Borodin, Kim, Kostochka, West [5], the then best approximation of Jagear's conjecture for planar graphs was established by showing that every planar graph with girth at least  $\frac{20t-2}{3}$  has circular chromatic number at most  $2+\frac{1}{t}$ . However, this follows as a corollary of the main result of the paper [5] which is a theorem on homomorphisms of (n, m)- graphs.

Thus, indeed the study of homomorphisms of (n, m)-graphs is a significant area of research. However, as there are not many known well-structured (n, m)-graphs, the study of its homomorphisms becomes extremely difficult. In contrast, there are known well-structured oriented, and signed graphs such as the Paley tournaments, signed Paley graphs, and Tromp constructions [20, 25]. In an other work [9], we have implemented the theory from this work (related to the switch operation) to construct some well-structured (0,3) and (1,1)-graphs and used them to prove upper bounds for (n,m)-chromatic number of partial 2-trees.

In recent times, researchers have started further extending the graph homomorphism studies by exploring the effect of switch operation on homomorphisms. Notably, homomorphisms of signed graphs, which are essentially obtained by observing the effect of the switch operation on 2-edge-colored graphs, has gained immense popularity [21, 23, 25, 29] in recent times due to its strong connection with the graph minor theory. Also, graph homomorphism with respect to some other switch-like operations, such as, push operation on oriented graphs [18], cyclic switch on k-edge-colored graphs [7], and switching (n,m)-graphs with respect to Abelian groups of special type (which does not allow switching an edge to an arc or vice versa) [8, 19] and switching (n,m)-graphs with respect to non-Abelian group [10] to list a few, has been recently studied. Naturally, all three main branches of research listed above in the context of graph homomorphism are also explored for the above-mentioned extensions and variants. However, in comparison, the global algebraic structure is a less explored branch for the extensions.

**Organization:** In this article, we are going to introduce a homomorphism with respect to a generalized switch operation (using a group  $\Gamma$ ) on (n, m)-graphs which, in particular, allows arcs to become edges and vice-versa. We call it a  $\Gamma$ -homomorphism whose detailed definition is deferred to the next section. In particular, it is possible to view the set of all (n, m)-graphs as a category with  $\Gamma$ -homomorphism playing the role of morphism. However, in this article, we have refrained from using the language of category theory as much as possible, and have used the language of graph theory instead.

- In Section 2 we introduce the notion of  $\Gamma$ -homomorphisms of (n, m)-graphs and prove some basic results. In particular, we show that the switch operation defined by us is (strictly) more general than the switch operation defined by Leclerc, MacGillivray, and Warren [19].
- In Section 3 we study algebraic properties of  $\Gamma$ -homomorphisms and explore their relation with (n, m)-homomorphisms. In particular, we prove a generalized version of an open problem due to Klostermeyer and MacGillivray [18]. Also, we show that the important notion of "core" is well-defined in this setup (category).
- In Section 4 we establish the existence of categorical product and co-product of (n, m)-graphs under  $\Gamma$ -homomorphism. Furthermore, we prove some fundamental properties of these categorical products and co-products. It is worth mentioning that proving the existence of categorical product is non-trivial. Interestingly, given two (n, m)-graphs G and H of order m and n, respectively, their categorical product is a graph on  $|\Gamma|mn$  vertices.
- In Section 5 we define and study the Γ-chromatic number for the family of forests, where the Γ-chromatic number is defined using Γ-homomorphism and the detailed definition is deferred to Section 5. This result generalizes Theorem 1.1 of [24].
- In Section 6 we conclude our work and propose possible future directions of works on this topic.

## 2 Homomorphisms of (n, m)-graphs and generalized switch

Throughout this article, we will follow the standard graph theoretic, algebraic and category theory notions from West [30], Artin [3], and Jacobson [16], respectively.

An (n, m)-graph G is a graph with vertex set V(G), arc set A(G) and edge set E(G), where each arc is colored with one of the n colors from  $\{2, 4, \dots, 2n\}$  and each edge is colored with one of the m colors from  $\{2n+1, 2n+2, \dots, 2n+m\}$ . In particular, if there is an arc of color i from u to v, we say that v is an i-neighbor of u, or equivalently, u is an i-neighbor of v. Furthermore, if there is an edge of color j between u and v, then we say that u (resp., v) is a j-neighbor of v (resp., u). For convenience, we use the following convention throughout this article: if u is an i-neighbor of v, then we say v is an  $\bar{i}$ -neighbor of v. In particular, it is worth noting that,  $\bar{i} = i$ .

Let  $\Gamma \subseteq S_{2n+m}$ , where  $S_{2n+m}$  is the permutation group on  $A_{n,m} = \{1, 2, \dots, 2n, 2n + 1, \dots, 2n + m\}$ . A  $\sigma$ -switch at a vertex v of an (n, m)-graph is to change its incident arcs and edges in such a way that its t-neighbors become  $\sigma(t)$ -neighbors for all  $t \in A_{n,m}$  where  $\sigma \in \Gamma$ . To  $\Gamma$ -switch a vertex v of an (n, m)-graph is to apply a  $\sigma$ -switch on v for some  $\sigma \in \Gamma$ . An (n, m)-graph G' obtained by a sequence of  $\Gamma$ -switches performed on the vertices of G is a  $\Gamma$ -equivalent graph of G.

In the very first work on (n, m)-graphs, Nešetřil and Raspaud [24] in 2000, extended the notion of graph homomorphisms to homomorphisms of (n, m)-graphs<sup>1</sup>, this generalization for particular cases implies the study related to homomorphisms of oriented, signed and k-edge colored graphs [1, 7, 21, 23, 27, 28].

**Definition 2.1.** Let G and H be two (n,m)-graphs. An (n,m)-homomorphism of G to H is a vertex mapping  $\phi: V(G) \to V(H)$  satisfying the following: for any  $u, v \in V(G)$ , if u is a t-neighbor of v, then f(u) is a t-neighbor of f(v) in H, for some  $v \in A_{n,m}$ .

We now extend the Definition 2.1 to homomorphisms of (n, m)-graphs with  $\Gamma$ -switch.

**Definition 2.2.** A  $\Gamma$ -homomorphism of G to H is a function  $f:V(G) \to V(H)$  such that there exists a  $\Gamma$ -equivalent graph G' of G satisfying the following: for any  $u, v \in V(G) = V(G')$ , if u is a t-neighbor of v in G' then f(u) is a t-neighbor of f(v) in G' then this by  $G \xrightarrow{\Gamma} H$ .

A Γ-isomorphism of G to H is a bijective Γ-homomorphism whose inverse is also a Γ-homomorphism. We denote this by  $G \equiv_{\Gamma} H$ . Observe that if  $\Gamma = \langle e \rangle$  is the singleton group with the identity element e, then our Γ-homomorphism definition becomes the same as homomorphism of (n, m)-graphs.

A related work on homomorphism with respect to a switch operation on (n, m)-graphs has been studied [19] in which, an Abelian subgroup of  $S_m \otimes (S_2 \wr S_n)$  acts on the vertices of (n, m)-graph such that the edges switch color with edges and the arcs switch color with arcs. Formally, Let  $\phi \in S_m$ ,  $\psi \in S_n$ , and  $\pi = (p_1, p_2, \dots, p_n) \in (\mathbb{Z}_2)^n$ , for an ordered triple

<sup>&</sup>lt;sup>1</sup>In their work, (n, m)-graphs were termed as colored mixed graphs and (n, m)-homomorphisms of (n, m)-graphs as colored homomorphism of colored mixed graphs. Also, in [5], (n, m)-graphs are referred as s-graphs

 $\gamma = (\phi, \psi, \pi)$ , an LMW-switch<sup>2</sup> [19] at a vertex v is said to be the process of transforming G into an (n, m)-graph  $G^{v,\gamma}$  where edges with color i incident to v changes to edges of color  $\phi(i)$ , arcs of color j incident to v, changes to arcs with color  $\psi(j)$  with the orientation reversed if and only if  $p_j = 1$  in  $\pi$ . This definition is a natural extension to the definitions given in [7, 18] for switching or pushing in the case of (n, m) = (1, 0), (0, m) graphs. In this paper, we study a more generalized switch operation on (n, m) graphs which also captures LMW-switch in particular. There also has been studies of  $\Gamma$  switch-homomorphisms of (n, m)-graphs when  $\Gamma$  is non-Abelian [17]. Here, we will restrict ourselves to Abelian subgroups  $\Gamma$  of  $S_{2n+m}$  unless otherwise stated.

Let u, v be any two adjacent vertices of an (n, m)-graph G. A switch-commutative group  $\Gamma \subseteq S_{2n+m}$  is such that for any  $\sigma, \sigma' \in \Gamma$ , by performing  $\sigma$ -switch on u and  $\sigma'$ -switch on v, the adjacency between u and v changes in the same way irrespective of the order of the switches. Observe that, this property not only depends on the group but also depends on the values of n and m. A  $\Gamma$ -switch, where  $\Gamma$  is a switch-commutative group, in general, is called a commutative switch.

**Theorem 2.3.** Every LMW-switch operation is a commutative switch operation. Moreover, there exists infinitely many commutative switches which are not LMW-switch.

Proof. Let G be an (n, m)-graph. Consider an Abelian subgroup  $\Gamma \subseteq S_m \otimes (S_2 \wr S_n)$ . For two vertices  $u, v \in V(G)$ , and for  $\sigma_u(t) = (\phi_u(t), \psi_u(t), \pi_u(t))$  applied on u and  $\sigma_v(t) = (\phi_v(t), \psi_v(t), \pi_v(t))$  applied on v, where  $\sigma_u, \sigma_v \in \Gamma$ . It is enough to prove that this action is a commutative switch operation. That is, to show that, the adjacency between u and v changes in the same way irrespective of the order in which the switch is applied.

Suppose v is a t-neighbor of u. If t is the color of an edge, and as  $\Gamma$  is Abelian,  $\sigma_u(\sigma_v(t)) = \sigma_u(\sigma_v(t))$  as  $\phi_u$  and  $\phi_v$  are responsible to change the color of the edge, which is a symmetric relation between u and v, and they commute.

If t is a color of an arc, then after applying the switches, the color of the arc changes as per the functions  $\psi_u$  and  $\psi_v$  and it does not matter in which order the switches are applied as the color of the arc (not the direction) is a symmetric relation between u and v, and as  $\Gamma$  is commutative. On the other hand, the change in the direction of the arc is determined by  $\pi_u(t).\pi_v(t)$ . To be precise, if  $\pi_u(t).\pi_v(t) = 0$  then the direction does not change, and if  $\pi_u(t).\pi_v(t) = 1$  then the direction changes. This completes the proof.

We give an example of a commutative switch which is not an LMW-switch. Consider (1,1)-graph G. Let  $C_3 = \{e, \sigma, \sigma^2\} \subset S_3$ , where,

$$\sigma(t) = \begin{cases} 2, & \text{if } t = 1\\ 3, & \text{if } t = 2\\ 1, & \text{if } t = 3 \end{cases}$$

For two adjacent vertices say  $u, v \in V(G)$ ,  $\sigma$  applied on u and then  $\sigma$  applied on v yields the same result as that of  $\sigma$  applied on v first and then on u. Thus,  $\Gamma$  is a commutative switch whereas it is clearly not a LMW-switch as an arc (color 2) is switched to edge

<sup>&</sup>lt;sup>2</sup>No particular name was given for this switch in [19]. We use the initials of the author names for convenience.

(color 3) under this operation. Further, we can extend this example to (n, n)-graph, for any  $n \in \mathbb{N}$  and  $\Gamma \subsetneq S_{3n}$ , where  $\Gamma = C_3 \oplus C_3 \oplus C_3 \oplus C_3 \oplus C_3$ , we have n-types of arc and n-types of edges, where i-th  $C_3$  permuting ith type of arc or edge respectively.

#### 3 Basic algebraic properties

Let  $\Gamma \subseteq S_{2n+m}$  be a group and let G be an (n, m)-graph with set of vertices  $\{v_1, v_2, \cdots, v_k\}$ . Let  $G^*$  be the (n, m)-graph having vertices of the type  $v_i^{\sigma}$  where  $i \in \{1, 2, \cdots, k\}$  and  $\sigma \in \Gamma$ . Also a vertex  $v_i^{\sigma}$  is a t-neighbor of  $v_j^{\sigma'}$  in  $G^*$  if and only if  $v_i$  is a t-neighbor of  $v_j$  in G where  $i, j \in \{1, 2, \cdots, k\}$  and  $\sigma, \sigma' \in \Gamma$ . The  $\Gamma$ -switched graph denoted by  $\rho_{\Gamma}(G)$  of G is the (n, m)-graph obtained from  $G^*$  by performing a  $\sigma$ -switch on  $v_i^{\sigma}$  for all  $i \in \{1, 2, \cdots, k\}$  and  $\sigma \in \Gamma$ .

**Lemma 3.1.** If  $\Gamma$  is switch-commutative group, then we have  $\overline{\sigma_v(\overline{\sigma_u(t)})} = \sigma_u(\overline{\sigma_v(\overline{t})})$ , for  $\sigma_u, \sigma_v \in \Gamma$  and  $t \in A_{n,m}$ .

Proof. Let  $u, v \in V(G)$ , and let v be a t-neighbor of u. Suppose  $\sigma_u$  is applied on u. Then we have, v is a  $\sigma_u(t)$ -neighbor of u, then by our convention, we have u is a  $\overline{\sigma_u(t)}$ -neighbor of v. Now we apply  $\sigma_v$  on v, we get, u is a  $\sigma_v(\overline{\sigma_u(t)})$ -neighbor of v, this implies, v is a  $\overline{\sigma_v(\overline{\sigma_u(t)})}$  neighbor of u.

On the other hand, as, v is a t-neighbor of u, we have, u is a  $\overline{t}$ -neighbor of v. Now we apply,  $\sigma_v$  first on v, therefore, u is a  $\sigma_v(\overline{t})$ -neighbor of v, and this is same as v is a  $\overline{\sigma_v(\overline{t})}$ -neighbor of u, we apply  $\sigma_u$  on u, we have, v is a  $\sigma_u(\overline{\sigma_v(\overline{t})})$ -neighbor of u.

Since  $\Gamma$  is switch-commutative, we get,

$$\overline{\sigma_v(\overline{\sigma_u(t)})} = \sigma_u(\overline{\sigma_v(\overline{t})})$$

Hence, the proof.

Corollary 3.2. Let G be an (n, m)-graph and let  $\Gamma$  be a switch-commutative group. Suppose, v is a t-neighbor of  $\underline{u}$  in G, and  $\sigma_u$ ,  $\sigma_v \in \Gamma$  is applied on u, v respectively. Then after the switches, v becomes a  $\sigma_v(\overline{\sigma_u(t)})$ -neighbor of u or equivalently, v is a  $\sigma_u(\overline{\sigma_v(\overline{t})})$ -neighbor of u.

**Theorem 3.3.** The  $\Gamma$ -switched graph  $\rho_{\Gamma}(G)$  is well defined for all (n, m)-graph G if and only if  $\Gamma$  is a switch-commutative group.

Proof. As,  $\Gamma$  is a switch-commutative group, it is clear that  $\rho_{\Gamma}(G)$  is well defined for all (n,m)-graphs G. Suppose,  $\Gamma$ -switched graph  $\rho_{\Gamma}(G)$  is well defined for all (n,m)-graphs G, Let v be a t-neighbor of u for some  $u,v\in V(G)$ . Then by definition,  $v^{\sigma_j}$  is a  $\sigma_j(\overline{\sigma_i(t)})$ -neighbor of  $u^{\sigma_i}$  in  $\rho_{\Gamma}(G)$ . As,  $\rho_{\Gamma}(G)$  is well defined, the order in which we switch the vertices should not matter, which forces  $\Gamma$  to be a switch-commutative group.  $\square$ 

This  $\Gamma$ -switch graph helps build a bridge between  $\langle e \rangle$ -homomorphism and  $\Gamma$ -homomorphism of two (n, m)-graphs. We prove a useful property of a switch commutative group.

**Proposition 3.4.** Let G and H be two (n,m)-graphs. We have  $G \xrightarrow{\Gamma} H$  if and only if  $G \xrightarrow{\langle e \rangle} \rho_{\Gamma}(H)$ , where  $\Gamma$  is a switch-commutative group.

*Proof.* For the "only if" part of the proof, suppose  $f: G \xrightarrow{\Gamma} H$ . Thus,  $f: G' \xrightarrow{\langle e \rangle} H$  for some  $G' \equiv_{\Gamma} G$ . Let  $S \subset V(G)$  be such that the vertices  $g_i$  of G in S are switched with  $\sigma_i$  to get G'.

Consider,  $\phi: G \to \rho_{\Gamma(H)}$  such that,

$$\phi(g_i) = \begin{cases} f(g_i) & \text{if } g_i \notin V(G) \setminus S \\ f(g_i)^{\sigma_i^{-1}} & \text{if } g_i \in S. \end{cases}$$

We prove that  $\phi$  is an  $\langle e \rangle$ -homomorphism.

- (i) If  $g_i \in S$  and  $g_j \in V(G) \setminus S$  and let  $g_j$  be a t-neighbor of  $g_i$  in G, As,  $\sigma_i$  is applied on  $g_i$ , we have,  $g_j$  to be a  $\sigma_i(t)$ -neighbor of  $g_i$  in G'. As f is a  $\langle e \rangle$ -homomorphism,  $f(g_j)$  is  $\sigma_i(t)$ -neighbor of  $f(g_i)$  in H. From the definition of  $\Gamma$ -switched graph,  $(f(g_j))$  is  $\sigma_i^{-1}(\sigma_i(t))$ -neighbor of  $f(g_i)^{\sigma_i^{-1}}$  in  $\rho_{\Gamma}(H)$ . As,  $\Gamma$  is Abelian,  $\sigma_i^{-1}(\sigma_i(t)) = t$ , Hence,  $\phi(g_j)$  is a t-neighbor of  $\phi(g_i)$  in  $\rho_{\Gamma}(H)$ .
- (ii) If  $g_i, g_j \in S$ , and let  $g_j$  be a t-neighbor of  $g_i$  in G. As  $g_i, g_j \in S$ , let  $\sigma_i, \sigma_j$  applied on  $g_i, g_j$  respectively. Thus by Corollary 3.2,

$$g_j$$
 is a  $\overline{\sigma_i(\overline{\sigma_j(t)})}$ -neighbor of  $g_i$  in  $G'$ .

As f is an  $\langle e \rangle$ -homomorphism, we have,

$$f(g_j)$$
 is a  $\sigma_i(\overline{\sigma_j(t)})$ -neighbor of  $f(g_i)$  in  $H$ .

By the definition of  $\rho_{\Gamma}(G)$ ,

$$f(g_j)^{\sigma_j^{-1}}$$
 is  $\overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(\overline{\sigma_i(\overline{\sigma_j(t)})})})}$ -neighbor of  $f(g_i)^{\sigma_i^{-1}}$  in  $\rho_{\Gamma}(H)$ .

Thus, it is enough to prove,

$$f(g_j)^{\sigma_j^{-1}}$$
 is  $\overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(\overline{\sigma_i(\overline{\sigma_j(t)})})})} = t$ -neighbor of  $f(g_i)^{\sigma_i^{-1}}$  in  $\rho_{\Gamma(H)}$ .

Due to Lemma 3.1, we have,

$$\overline{\sigma_i(\overline{\sigma_j(t)})} = \sigma_j(\overline{\sigma_i(\overline{t})})$$

which implies

$$\overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(\overline{\sigma_i(\overline{\sigma_j(t)})})})} = \overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(\sigma_j(\overline{\sigma_i(\overline{t})}))})} = \overline{\sigma_i^{-1}(\sigma_i(\overline{t}))} = \overline{\overline{t}} = t.$$

(iii) If  $g_i, g_j \in V(G) \setminus S$ , then  $\phi(g_i) = f(g_i)$  is a t-neighbor of  $\phi(g_j) = f(g_j)$ , as f is a homomorphism.

For the "if" part of the proof, suppose  $f: G \xrightarrow{\langle e \rangle} \rho_{\Gamma}(H)$  be a  $\langle e \rangle$ -homomorphism. Let  $g_1, g_2, \dots, g_p$  be the vertices of G, and let  $h_1, h_2, \dots, h_q$  be the vertices of H. Also, let  $f(g_i) = h_j^{\sigma_i}$  for some  $\sigma_i \in \Gamma$ . In that case, we define the function  $\varphi: V(G') \to V(H)$  as follows:

$$\varphi(g_i) = h_j$$
.

where we obtain G' by performing a  $\sigma_i^{-1}$ -switch on the vertex  $g_i$  of G. Note that it is enough to show  $\varphi$  is a  $\langle e \rangle$ -homomorphism of G' to H.

Let  $g_j^{\sigma_i^{-1}}$  be a t-neighbor of  $g_i^{\sigma_j^{-1}}$  in G'. We have to show that  $\varphi(g_j^{\sigma_i^{-1}}) = h_i$  (say) is a t-neighbor of  $\varphi(g_i^{\sigma_j^{-1}}) = h_j$  (say) in H.

$$g_i^{\sigma_i^{-1}}$$
 be a t-neighbor of  $g_i^{\sigma_j^{-1}}$  in  $G'$ .

By corollary 3.2, we have,

$$g_j$$
 is a  $\overline{\sigma_i^{-1}(\overline{\sigma_i^{-1}}(t))}$ -neighbor of  $g_i$  in  $G$ .

As f is a  $\langle e \rangle$ -homomorphism, we have,

$$f(g_j)$$
 is a  $\overline{\sigma_j^{-1}(\overline{\sigma_i^{-1}}(t))}$ -neighbor of  $f(g_i)$  in  $\rho_{\Gamma(H)}$ .

That is,

$$h_i^{\sigma_j}$$
 is a  $\overline{\sigma_j^{-1}(\overline{\sigma_i^{-1}}(t))}$ -neighbor of  $h_j^{\sigma_i}$  in  $\rho_{\Gamma(H)}$ .

By corollary 3.2, we have,

$$h_i$$
 is a  $\sigma_j(\overline{\sigma_i(\overline{\sigma_j^{-1}(\overline{\sigma_i^{-1}}(t))})})$ -neighbor of  $h_j$  in  $H$ .

Thus, it is enough to prove,

$$\sigma_j(\overline{\sigma_i(\overline{\sigma_j^{-1}(\overline{\sigma_i^{-1}}(t))})}) = t.$$

By Lemma 3.1, we have,

$$\overline{\sigma_i(\overline{\sigma_j(t)})} = \sigma_j(\overline{\sigma_i(\overline{t})})$$

which implies

$$\overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(\overline{\sigma_i(\overline{\sigma_j(t)})})})} = \overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(\sigma_j(\overline{\sigma_i(\overline{t})}))})} = \overline{\sigma_i^{-1}(\sigma_i(\overline{t}))} = \overline{\overline{t}} = t.$$

Thus,  $h_i = \varphi(g_j^{\sigma_i^{-1}})$  is a t-neighbor of  $h_j = \varphi(g_i^{\sigma_j^{-1}})$  in H.

**Theorem 3.5.** Let G and H be (n,m)-graphs. Then,  $G \equiv_{\Gamma} H$  if and only if  $\rho_{\Gamma}(G) \equiv_{\langle e \rangle} \rho_{\Gamma}(H)$ , where  $\Gamma$  is a switch-commutative group.

*Proof.* For the "only if" part of the proof, suppose :  $G \equiv_{\Gamma} H$ . Let G' be a  $\Gamma$ -equivalent graph of G such that  $f: G' \xrightarrow{\langle e \rangle} H$  is an  $\langle e \rangle$ -isomorphism. Suppose that G has vertices  $g_1, g_2, \dots, g_p$ , and G' is obtained by performing  $\tau_i$ -switch on  $g_i$ , for some  $\tau_i \in \Gamma$ . For any  $g_i^{\sigma_j} \in V(\rho_{\Gamma}(G))$  we define the function  $\phi: V(\rho_{\Gamma}(G)) \to V(\rho_{\Gamma}(H))$  as follows:

$$\phi(g_i^{\sigma_j}) = \left( f(g_i^{\tau_i^{-1}}) \right)^{\sigma_j}.$$

Next we prove that  $\phi$  is a  $\langle e \rangle$ -isomorphism of  $\rho_{\Gamma}(G)$  to  $\rho_{\Gamma}(H)$ . Let  $g_j^{\sigma_i}$  be a t-neighbor of  $g_i^{\sigma'_j}$  in  $\rho_{\Gamma(G)}$ , then by Corollary 3.2,

$$g_j$$
 is a  $\overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(t)})}$  – neighbor of  $g_i$  in  $G$ .

$$g_i^{\tau_j}$$
 is a  $\overline{\tau_j(\overline{\tau_i}(\overline{\sigma_i^{-1}}(\overline{\sigma_j^{-1}}(t))))}$  neighbor of  $g_i^{\tau_i}$  in  $G'$ .

As f is a  $\langle e \rangle$ -isomorphism,

$$f(g_j^{\tau_j})$$
 is a  $\overline{\tau_j(\overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(t)})})}$  neighbor of  $f(g_i^{\tau_i})$  in  $H$ .

Now,

$$(f(g_j^{\tau_j}))^{\tau_j^{-1}} \text{ is a } \overline{\tau_j^{-1}(\tau_i^{-1}(\overline{\tau_i(\overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(t)})}))))} \text{ neighbor of } f(g_i^{\tau_i})^{\tau_i^{-1}} \text{ in } \rho_{\Gamma}(H).$$

Now,

$$\phi(g_j^{\sigma_i}) = \left(f(g_j^{\tau_j^{-1}})\right)^{\sigma_i} \text{ is a } \overline{\sigma_i(\overline{\sigma_j(\tau_j^{-1}(\overline{\tau_i(\overline{\sigma_i^{-1}(\overline{\sigma_j^{-1}(t)})}))}))) \text{ neighbor of } } \phi(g_i^{\sigma_j}) = \left(f(g_i^{\tau_i^{-1}})\right)^{\sigma_j}.$$

By repeated application of Lemma 3.1 on (1), (2) and (3), and as  $\Gamma$  is Abelian, we get,

$$\overline{\sigma_{i}(\sigma_{j}(\tau_{j}^{-1}(\tau_{i}^{-1}(\overline{\tau_{j}(\overline{\sigma_{i}^{-1}(\overline{\sigma_{j}^{-1}(\overline{\sigma_{j}^{-1}(t)})})))))))))}))))} = t$$
(1)
(2)
(3)

Therefore,  $\phi(g_i) = (f(g_i)^{\tau_i^{-1}})^{\sigma_j}$  is t-neighbor of  $(f(g_k)^{\tau_k^{-1}})^{\sigma_l}$  in  $\rho_{\Gamma(H)}$ . Thus,  $\phi$  a  $\langle e \rangle$ isomorphism of  $\rho_{\Gamma}(G)$  to  $\rho_{\Gamma}(H)$ .

For the "if" part of the proof, suppose  $\rho_{\Gamma}(G) \equiv_{\langle e \rangle} \rho_{\Gamma}(H)$  and we have to show  $G \equiv_{\Gamma} H$ . Assume  $g_1, g_2, \dots, g_p$  be the vertices of G. A sequence of vertices in  $\rho_{\Gamma}(G)$  of the form  $(g_1^{\sigma_1}, g_2^{\sigma^2}, \cdots, g_p^{\sigma_p})$  is a representative sequence of G in  $\rho_{\Gamma}(G)$ , where  $\sigma_i \in \Gamma$  is any element for  $i \in \{1, 2, \dots, p\}$  (repetition of elements among  $\sigma_i$  is allowed here).

Given an  $\langle e \rangle$ -isomorphism  $\psi : \rho_{\Gamma}(G) \xrightarrow{\langle e \rangle} \rho_{\Gamma}(H)$  and a representative sequence S of G in  $\rho_{\Gamma}(G)$ , define the set

$$Y_{S,\psi} = \{v^{\sigma} \mid \psi(v^{\sigma}) = (\psi(v))^{\sigma} \text{ where } v \in S \text{ and } \sigma \in \Gamma\}.$$

Let  $Y_{S^*,\varphi}$  be the set satisfying the property  $|Y_{S^*,\varphi}| \ge |Y_{S,\psi}|$  where S varies over all representative sequences and  $\psi$  varies over all  $\langle e \rangle$ -isomorphisms.

We will show that  $Y_{S^*,\varphi} = V(\rho_{\Gamma}(G))$ . We will prove by the method of contradiction. Thus, let us assume the contrary, that is, let  $Y_{S^*,\varphi} \neq V(\rho_{\Gamma}(G))$ . This implies that there exists a  $v^{\sigma}$ , for some  $v \in S^*$  and some  $\sigma \in \Gamma$  such that  $\varphi(v^{\sigma}) \neq (\varphi(v))^{\sigma}$ . Next let us define the function

$$\hat{\varphi}(x) = \begin{cases} \varphi(x) & \text{if } x \neq g, v^{\sigma}, \\ \varphi(g) & \text{if } x = v^{\sigma}, \\ \varphi(v^{\sigma}) & \text{if } x = g, \end{cases}$$

where  $g = \varphi^{-1}(\varphi(v)^{\sigma}) \in \rho_{\Gamma}(G)$ .

Next we are going to show that  $\hat{\varphi}$  is an  $\langle e \rangle$ -isomorphism of  $\rho_{\Gamma}(G)$  and  $\rho_{\Gamma}(H)$ . So, we need to show that x is a t-neighbor of y in  $\rho_{\Gamma}(G)$  if and only if  $\hat{\varphi}(x)$  is a t-neighbor of  $\hat{\varphi}(y)$  in  $\rho_{\Gamma}(H)$ . Notice that, it is enough to check this for x = g and  $x = v^{\sigma}$  while y varies over all vertices of  $\rho_{\Gamma}(G)$ . We will separately handle the exceptional case when  $x = v^{\sigma}$  and y = g first.

- (i) If  $x = v^{\sigma}$  and y = g, then  $\varphi(v)$  and  $\varphi(v)^{\sigma}$  are non-adjacent by the definition of  $\rho_{\Gamma}(G)$ . Thus,  $v = \varphi^{-1}(\varphi(v))$  and  $g = \varphi^{-1}(\varphi(v)^{\sigma})$  are non-adjacent. Hence  $x = v^{\sigma}$  and y = g are also non-adjacent. On the other hand, this implies that  $\hat{\varphi}(x) = \varphi(g)$  and  $\hat{\varphi}(y) = \varphi(v^{\sigma})$  are non-adjacent.
- (ii) If  $x = v^{\sigma}$  and  $y \neq g$ , then y is a t-neighbour of x in  $\rho_{\Gamma}(G)$  if and only if  $\varphi(y)$  is a t-neighbor of  $\varphi(x)$  in  $\rho_{\Gamma}(H)$ , as  $\varphi$  is an  $\langle e \rangle$ -isomorphism. Observe that  $\hat{\varphi}(x) = \varphi(g) = \varphi(v)^{\sigma}$  as  $g = \varphi^{-1}(\varphi(v)^{\sigma})$ , and  $\hat{\varphi}(y) = \varphi(y)$ . Since  $x = v^{\sigma}$ , y is a  $\sigma^{-1}(t)$ -neighbor of v in  $\rho_{\Gamma}(G)$ , if and only if  $\varphi(y)$  is a  $\sigma^{-1}(t)$ -neighbor of  $\varphi(v)$  in  $\rho_{\Gamma}(H)$ , if and only if,  $\varphi(y) = \hat{\varphi}(y)$  is a t-neighbour of  $\varphi(v)^{\sigma} = \hat{\varphi}(x)$  in  $\rho_{\Gamma}(H)$ .
- (iii) If x=g and  $y\neq v^{\sigma}$ , then y is a t-neighbor of x in  $\rho_{\Gamma}(G)$  if and only if,  $\varphi(y)$  is a t-neighbor of  $\varphi(x)=\varphi(g)=\varphi(v)^{\sigma}$  in  $\rho_{\Gamma}(H)$ , as  $g=\varphi^{-1}(\varphi(v)^{\sigma})$ . The previous statement holds if and only if  $\varphi(y)$  is a  $\sigma^{-1}(t)$ -neighbor of  $\varphi(v)$  in  $\rho_{\Gamma}(H)$  if and only if y is a  $\sigma^{-1}(t)$ -neighbor of v in  $\rho_{\Gamma}(G)$  if and only if y is a t-neighbor of v in  $\rho_{\Gamma}(G)$  if and only if  $\varphi(y)=\hat{\varphi}(y)$  is a t-neighbor of  $\varphi(v)=\hat{\varphi}(x)$  in  $\varphi_{\Gamma}(H)$ .

However, now we have  $|Y_{S^*,\varphi}| < |Y_{S^*,\hat{\varphi}}|$ . This is a contradiction to the definition of  $Y_{S^*,\varphi}$ , and hence  $Y_{S^*,\varphi} = V(\rho_{\Gamma}(G))$ .

Let  $v_1, v_2$  be two distinct vertices in  $S^*$ . If  $\varphi(v_1)^{\sigma} = \varphi(v_2)$  for any  $\sigma \in \Gamma$ , then  $\varphi(v_1^{\sigma}) = \varphi(v_2)$ . This implies  $v_1^{\sigma} = v_2$  because  $\varphi$  is a bijection. However, this is not possible as  $v_1, v_2$  are distinct vertices from a representative sequence  $S^*$  of G in  $\rho_{\Gamma}(G)$ . Hence,  $\varphi(v_1)^{\sigma} \neq \varphi(v_2)$  for any  $v_1, v_2 \in S^*$ . That means,  $\varphi(S^*) = R$  is a representative

sequence of H in  $\rho_{\Gamma}(H)$ . Thus, note that  $\langle e \rangle$ -isomorphism restricted to the induced subgraph  $\rho_{\Gamma}(G)[S^*]$  is also an  $\langle e \rangle$ -isomorphism to the induced subgraph  $\rho_{\Gamma}(H)[R]$ . That is,  $\rho_{\Gamma}(G)[S^*] \equiv_{\langle e \rangle} \rho_{\Gamma}(H)[R]$ . As  $\langle e \rangle \subseteq \Gamma$ , this also means  $\rho_{\Gamma}(G)[S^*] \equiv_{\Gamma} \rho_{\Gamma}(H)[R]$ .

On the other hand, as  $S^*$  and R are representative sequences of G and H, respectively, we have  $\rho_{\Gamma}(G)[S^*] \equiv_{\Gamma} G$  and  $\rho_{\Gamma}(H)[R] \equiv_{\Gamma} H$ . Thus we are done by composing the  $\Gamma$ -isomorphisms.

The above result generalizes results by Brewster and Graves [7] (see Theorem 12) and Sen [26] (see Theorem 3.4). Additionally, it (re)proves an open problem given by Klostermeyer and MacGillivray [18] (see Open Problem 2 in the conclusion) by restricting the result to (n, m) = (1, 0) where  $\Gamma$  is the group in which the only non-identity element simply reverses the direction of the arcs.

The next result follows from the fundamental theorem of finite abelian groups.

**Theorem 3.6.** Let  $\Gamma_1$  be a switch-commutative subgroup of  $S_{2n+m}$ . Let  $\Gamma_2 \subseteq \Gamma_1$ . If  $p^2 \nmid |\Gamma_1|$  for any prime p, then  $\rho_{\Gamma_1}(G) \equiv_{\langle e \rangle} \rho_{\Gamma_1/\Gamma_2}(\rho_{\Gamma_2}(G))$ .

Proof. Since  $\Gamma_1$  is a finite Abelian group,  $\Gamma_1/\Gamma_2$  and  $\Gamma_2$  both are normal subgroups of  $\Gamma_1$ , As,  $p^2 \nmid |\Gamma_1|$ , we have  $\Gamma_1/\Gamma_2 \times \Gamma_2 \equiv \Gamma_1$ . Thus along with the fact that  $\Gamma_2$  and  $\Gamma_1/\Gamma_2$  are normal subgroups, we have  $\Gamma_2 \cdot \Gamma_1/\Gamma_2 = \Gamma_1$ . Observe that, every element  $\sigma \in \Gamma_1$ , can be uniquely written as  $\alpha.\beta$ , where  $\alpha \in \Gamma_1/\Gamma_2$ ,  $\beta \in \Gamma_2$ . Now let G be an (n, m)-graph. We prove,  $f: \rho_{\Gamma_1}(G) \to \rho_{\Gamma_1/\Gamma_2}(\rho_{\Gamma_2}(G))$  is an isomorphism. Consider,

$$f: V(\rho_{\Gamma_1}(G)) \to V(\rho_{\Gamma_1/\Gamma_2}(\rho_{\Gamma_2}(G)),$$
  
$$f(u^{\sigma}) = (u^{\alpha})^{\beta}.$$

where  $\alpha . \beta = \sigma$ .

Suppose,  $v^{\sigma_j}$  be a t-neighbor  $u^{\sigma_i}$  in  $\rho_{\Gamma_1}(G)$  for some i, j if and only if  $v^{\alpha_j,\beta_j}$  is a t-neighbor of  $u^{\alpha_i,\beta_i}$ , where  $\sigma_i = \alpha_i.\beta_i$  and  $\sigma_j = \alpha_j.\beta_j$ . As  $\Gamma_2$  and  $\Gamma_1/\Gamma_2$  are Abelian, every  $\sigma \in \Gamma_1$ , can be uniquely represented as  $\alpha.\beta$ , where  $\alpha \in \Gamma_1/\Gamma_2$  and  $\beta \in \Gamma_2$ .

A Γ-core of an (n, m)-graph G is a subgraph H of G such that  $G \xrightarrow{\Gamma} H$ , whereas H does not admit a Γ-homomorphism to any of its proper subgraphs.

**Theorem 3.7.** The core of an (n, m)-graph G is unique up to  $\Gamma$ -isomorphism.

*Proof.* Let  $H_1$  and  $H_2$  be two  $\Gamma$ -cores of G. We have to show that  $H_1$  and  $H_2$  are  $\Gamma$ -isomorphic.

Note that, there exist  $\Gamma$ -homomorphisms  $f_1: G \xrightarrow{\Gamma} H_1$  and  $f_2: G \xrightarrow{\Gamma} H_2$  as  $H_1, H_2$  are  $\Gamma$ -cores. Moreover, there exists the inclusion  $\Gamma$ -homomorphisms  $i_1: H_1 \xrightarrow{\Gamma} G$  and  $i_2: H_2 \xrightarrow{\Gamma} G$ .

Now consider the composition  $\Gamma$ -homomorphism  $f_2 \circ i_1 : H_1 \xrightarrow{\Gamma} H_2$ . Note that it must be a surjective vertex mapping. Not only that, for any non-adjacent pair u, v of vertices in  $H_1$ , the vertices  $(f_2 \circ i_1)(u)$  and  $(f_2 \circ i_1)(v)$  are non-adjacent in  $H_2$ . The reason is that, if the above two conditions are not satisfied, then the composition  $\Gamma$ -homomorphism  $f_2 \circ i_1 \circ f_1 : G \xrightarrow{\Gamma} H_2$  can be considered as a  $\Gamma$ -homomorphism to a proper subgraph of

 $H_2$ . This will contradict the fact that  $H_2$  is a  $\Gamma$ -core. Therefore,  $f_2 \circ i_1$  is a bijective  $\Gamma$ -homomorphism whose inverse is also a  $\Gamma$ -homomorphism. In other words,  $f_2 \circ i_1$  is a  $\Gamma$ -isomorphism.

Due to the above theorem, it is possible to define the  $\Gamma$ -core of G and let us denote it by  $core_{\Gamma}(G)$ . Notice that, this is the analogue of the fundamental algebraic concept of core in the study of graph homomorphism.

### 4 Categorical products

Taking the set of (n, m)-graphs as objects and their  $\Gamma$ -homomorphisms as morphisms, one can consider the category of (n, m)-graphs with respect to  $\Gamma$ -homomorphism. In this section, we study whether products and co-products exist in this category or not. We would also like to remark that the existence of categorical product and co-product will not only contribute in establishing the category of (n, m)-graphs with respect to  $\Gamma$ -homomorphism as a richly structured category, but it will also show that the lattice of (n, m)-graphs induced by  $\Gamma$ -homomorphisms is a distributive lattice with the categorical products and co-products playing the roles of join and meet, respectively. Moreover, categorical product was useful in proving the density theorem [15] for undirected and directed graphs. Thus, it is not wrong to hope that it may become useful to prove the analogue of the density theorem in our context. It is worth commenting that the the idea to prove the existence of categorical products in this context generalizes the idea of the same in the context of signed graphs from [22].

Before proceeding further with the results, let us recall what categorical product and co-product mean in our context. Let G, H be two (n, m)-graphs and let  $\Gamma \subseteq S_{2n+m}$  be an Abelian group.

The categorical product of G and H with respect to  $\Gamma$ -homomorphism is an (n,m)-graph P having two projection mappings of the form  $f_g: P \xrightarrow{\Gamma} G$  and  $f_h: P \xrightarrow{\Gamma} H$  satisfying the following universal property: if any (n,m)-graph P' admit  $\Gamma$ -homomorphisms  $\phi_g: P' \xrightarrow{\Gamma} G$  and  $\phi_h: P' \xrightarrow{\Gamma} H$ , then there exists a unique  $\Gamma$ -homomorphism  $\varphi: P' \xrightarrow{\Gamma} P$  such that  $\phi_g = f_g \circ \varphi$  and  $\phi_h = f_h \circ \varphi$ .

The categorical co-product of G and H with respect to  $\Gamma$ -homomorphism is an (n,m)-graph C along with the two inclusion mappings of the form  $i_g: G \xrightarrow{\Gamma} C$  and  $i_h: H \xrightarrow{\Gamma} C$  satisfying the following universal property: if for any (n,m)-graph C' there are  $\Gamma$ -homomorphisms  $\phi_g: G \xrightarrow{\Gamma} C'$  and  $\phi_h: H \xrightarrow{\Gamma} C'$ , then there exists a unique  $\Gamma$ -homomorphism  $\varphi: C \xrightarrow{\Gamma} C'$  such that  $\phi_g = \varphi \circ i_g$  and  $\phi_h = \varphi \circ i_h$ .

Let G, H be two (n, m)-graphs and let  $\Gamma \subseteq S_{2n+m}$  be a switch-commutative group. Then  $G \times_{\langle e \rangle} H$  denotes the (n, m)-graph on set of vertices  $V(G) \times V(H)$  where (u, v) is a t-neighbor of (u', v') in  $G \times_{\langle e \rangle} H$  if and only if u is a t-neighbor of u' in G and v is a t-neighbor of v' in H. Moreover, the (n, m)-graph  $G \times_{\Gamma} H$  is the subgraph of  $\rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(H)$  induced by the set of vertices

$$X = \{(u^{\sigma}, v^{\sigma}) : (u, v) \in V(G) \times V(H) \text{ and } \sigma \in \Gamma\}.$$

**Theorem 4.1.** The categorical product of (n,m)-graphs G and H with respect to  $\Gamma$ -homomorphism exists and is  $\Gamma$ -isomorphic to  $G \times_{\Gamma} H$ , where  $\Gamma$  is a switch-commutative group.

Proof. Let  $(G \times_{\Gamma} H)'$  be Γ-switched graph of  $G \times_{\Gamma} H$ , where we apply  $\sigma^{-1}$  on  $(u^{\sigma}, v^{\sigma}) \in V(G \times_{\Gamma} H)$ . Thus, we define  $f_g(u^{\sigma}, v^{\sigma}) = u$  and  $f_h(u^{\sigma}, v^{\sigma}) = v$  as the two projections. Observe that  $f_g$  and  $f_h$  are  $\langle e \rangle$ -homomorphisms of  $(G \times_{\Gamma} H)'$  to G and H, respectively. If there exists an (n, m)-graph P' such that,  $\phi_g : P' \xrightarrow{\Gamma} G$  and  $\phi_h : P' \xrightarrow{\Gamma} H$ , then define  $\phi : P' \xrightarrow{\Gamma} G \times_{\Gamma} H$  such that  $\phi(p) = (\phi_g(p), \phi_h(p))$ . From the definition of  $\phi$ , we have  $\phi_g = f_g \circ \phi$  and  $\phi_h = f_h \circ \phi$ . Note that this is the unique way we can define  $\phi$  which satisfies the universal property from the definition of products. Thus,  $G \times_{\Gamma} H$  is indeed the categorical product of G and H with respect to  $\Gamma$ -homomorphism once we prove its uniqueness up to  $\Gamma$ -isomorphism.

Suppose  $P_1$  with projection mappings  $f_g$ ,  $f_h$  and  $P_2$  with projection mappings  $\phi_g$ ,  $\phi_h$  be two (n,m)-graphs that satisfy the universal properties of categorical product of G and H, then there exists  $\varphi: P_1 \xrightarrow{\Gamma} P_2$  and  $\varphi': P_2 \xrightarrow{\Gamma} P_1$  with  $\phi_g \circ \varphi = f_g$ ,  $\phi_h \circ \varphi = f_h$  and  $f_g \circ \varphi' = \phi_g$ ,  $f_h \circ \varphi' = \phi_h$ . Now consider the composition,  $\varphi' \circ \varphi: P_1 \xrightarrow{\Gamma} P_1$ . As,

$$f_a \circ (\varphi' \circ \varphi) = (f_a \circ \varphi') \circ \varphi = \phi_a \circ \varphi = f_a.$$

Similarly,

$$f_h \circ (\varphi' \circ \varphi) = (f_h \circ \varphi') \circ \varphi = \phi_h \circ \varphi' = f_h.$$

From this we can conclude that  $\varphi' \circ \varphi$  is a identity homomorphism on  $P_1$ . Similarly  $\varphi \circ \varphi'$  must be the identity homomorphism on  $P_2$ . Thus implying,  $\varphi' = \varphi^{-1}$  is an  $\Gamma$ -isomorphism of  $P_2$  and  $P_1$ .

In PEPS 2012 workshop, Brewster had asked whether Categorical product exists for signed graphs or not. The above theorem answers this question in affirmative as a special case (with respect to  $\Gamma$ -homomorphisms of (0,2)-graphs, where  $\Gamma$  is non-trivial).

Corollary 4.2. For any (n,m)-graphs G and H, and a switch-commutative group  $\Gamma \subseteq S_{2n+m}$ , we have  $\rho_{\Gamma}(G \times_{\Gamma} H) \equiv_{\langle e \rangle} \rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(H)$ .

*Proof.* The proof follows from the Theorems 3.5 and Theorem 4.1.  $\Box$ 

Let G + H denotes the disjoint union of the (n, m)-graphs G and H.

**Theorem 4.3.** The categorical co-product of (n,m)-graphs G and H with respect to  $\Gamma$ -homomorphism exists and is  $\Gamma$ -isomorphic to G+H, where  $\Gamma$  is any subgroup of  $S_{2n+m}$ .

Proof. Consider the inclusion mapping  $i_g: G \xrightarrow{\Gamma} G + H$ , and  $i_h: H \xrightarrow{\Gamma} G + H$ . Suppose there exists an (n,m)-graph C and there are Γ-homomorphisms  $\phi_g: G \xrightarrow{\Gamma} C$  and  $\phi_h: H \xrightarrow{\Gamma} C$ , then there exists a Γ-homomorphism  $\varphi: G + H \xrightarrow{\Gamma} C$  such that

$$\varphi(x) = \begin{cases} \phi_g(x) & \text{if } x \in V(G), \\ \phi_h(x) & \text{if } x \in V(H). \end{cases}$$

Observe that,  $\phi_g = \varphi \circ i_g$  and  $\phi_h = \varphi \circ i_h$ . Note that such a  $\varphi$  is unique.

Suppose we have P with  $\Gamma$ -homomorphisms  $f_g, f_h$  and P' with  $\Gamma$ -homomorphisms  $\phi_g, \phi_h$  satisfying the universal property of categorical co-product of G and H, then there exists  $\Gamma$ -homomorphisms,  $\varphi: P \xrightarrow{\Gamma} P'$  and  $\varphi': P' \xrightarrow{\Gamma} P$  with  $\phi_g = \varphi \circ f_g, \phi_h = \varphi \circ f_h$  and  $f_g = \varphi' \circ \phi_g, f_h = \varphi' \circ \phi_h$ . Now consider the composition,  $\varphi' \circ \varphi: P \xrightarrow{\Gamma} P$ . As,

$$(\varphi' \circ \varphi) \circ f_g = \varphi' \circ (\varphi \circ f_g) = \varphi' \circ \phi_g = f_g$$

Similarly,

$$(\varphi' \circ \varphi) \circ f_h = \varphi' \circ (\varphi \circ f_h) = \varphi' \circ \phi_h = f_h$$

Therefore, we should have  $\varphi' \circ \varphi$  to be the identity mapping on P. Similarly  $\varphi \circ \varphi'$  must be the identity mapping on P. Thus implying,  $\varphi' = \varphi^{-1}$  is an  $\Gamma$ -isomorphism of P' and P. Therefore, we have, the categorical co-product of (n, m)-graphs G and H with respect to  $\Gamma$ -homomorphism is G + H.

Thus both categorical product and co-product exists with respect to  $\Gamma$ -homomorphism when  $\Gamma$  is a switch commutative group. Furthermore, the usual algebraic identities hold with respect to these operations too.

**Corollary 4.4.** For any (n,m)-graphs G and H, and a switch-commutative group  $\Gamma \subseteq S_{2n+m}$ , we have  $\rho_{\Gamma}(G+H) \equiv_{\langle e \rangle} \rho_{\Gamma}(G) + \rho_{\Gamma}(H)$ .

*Proof.* The proof follows from the Theorems 3.5 and Theorem 4.3.

**Theorem 4.5.** For any (n, m)-graphs G, H, K and  $\Gamma$ , a switch-commutative group, We have the following.

- (i)  $G \times_{\Gamma} H \equiv_{\Gamma} H \times_{\Gamma} G$ ,
- (ii)  $G \times_{\Gamma} (H \times_{\Gamma} K) \equiv_{\Gamma} (G \times_{\Gamma} H) \times_{\Gamma} K$ ,
- (iii)  $G \times_{\Gamma} (H + K) \equiv_{\Gamma} (G \times_{\Gamma} H) + (G \times_{\Gamma} K).$

Proof. (i) Consider the mapping  $\phi: G \times_{\Gamma} H \to H \times_{\Gamma} G$ , such that,  $\phi(g^{\sigma}, h^{\sigma'}) = (h^{\sigma'}, g^{\sigma})$  We show that  $\phi$  is a Γ-isomorphism. Let  $(g_1^{\sigma_1}, h_1^{\sigma_1}), (g_2^{\sigma_2}, h_2^{\sigma_2}) \in V(G \times_{\Gamma} H)$ , suppose,  $(g_2^{\sigma_2}, h_2^{\sigma_2})$  be a t-neighbor of  $(g_1^{\sigma_1}, h_1^{\sigma_1})$  in  $G \times_{\Gamma} H$ , which implies,  $g_2^{\sigma_2}$  is a t-neighbor of  $g_1^{\sigma_1}$  in G and  $h_2^{\sigma_2}$  is a t-neighbor of  $h_1^{\sigma_1}$  in H, and hence,  $(h_2^{\sigma_2}, g_2^{\sigma_2})$  be a t-neighbor of  $(h_1^{\sigma_1}, g_1^{\sigma_1})$  in  $H \times_{\Gamma} G$ . Thus  $\phi$  is a Γ-isomorphism.

(ii) Observe that when  $\Gamma = \langle e \rangle$ . the function

$$\phi(g,(h,k)) = ((g,h),k)$$

is a  $\langle e \rangle$ -isomorphism of  $G \times_{\Gamma} (H \times_{\Gamma} K)$  to  $(G \times_{\Gamma} H) \times_{\Gamma} K$  where  $g \in G$ ,  $h \in H$ , and  $k \in K$ .

Next we will prove it for general  $\Gamma$ . Notice that by Theorem 4.1, we have

$$\rho_{\Gamma}(G \times_{\Gamma} (H \times_{\Gamma} K)) = \rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(H \times_{\Gamma} K) = \rho_{\Gamma}(G) \times_{\langle e \rangle} (\rho_{\Gamma}(H) \times_{\langle e \rangle} \rho_{\Gamma}(K)) \qquad (1)$$

and

$$\rho_{\Gamma}((G \times_{\Gamma} H) \times_{\Gamma} K) = \rho_{\Gamma}(G \times_{\Gamma} H) \times_{\langle e \rangle} \rho_{\Gamma}(K) = (\rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(H)) \times_{\langle e \rangle} \rho_{\Gamma}(K). \tag{2}$$

Since we have already proved that our equality holds for  $\Gamma = \langle e \rangle$ , we obtain,

$$\rho_{\Gamma}(G) \times_{\langle e \rangle} (\rho_{\Gamma}(H) \times_{\langle e \rangle} \rho_{\Gamma}(K)) \equiv_{\langle e \rangle} (\rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(H)) \times_{\langle e \rangle} \rho_{\Gamma}(K).$$

Therefore by equations (1) and (2) we have

$$\rho_{\Gamma}(G \times_{\Gamma} (H \times_{\Gamma} K)) \equiv_{\langle e \rangle} \rho_{\Gamma}((G \times_{\Gamma} H) \times_{\Gamma} K).$$

By Theorem 3.5 we have

$$G \times_{\Gamma} (H \times_{\Gamma} K) \equiv_{\Gamma} (G \times_{\Gamma} H) \times_{\Gamma} K.$$

This concludes the proof.

(iii) When  $\Gamma = \langle e \rangle$  consider the function

$$\phi(g, x) = (g, x)$$

where  $g \in G$  and if  $x \in (H+K)$ . However, here if  $x \in H$ , then the image  $(g,x) \in G \times_{\Gamma} H$  and if  $x \in K$ , then the image  $(g,x) \in G \times_{\Gamma} K$ . Observe that  $\phi$  is a  $\langle e \rangle$ -isomorphism of  $G \times_{\Gamma} (H+K)$  to  $(G \times_{\Gamma} H) + (G \times_{\Gamma} K)$ .

Next we will prove it for general  $\Gamma$ . Notice that by Theorem 4.1, we have

$$\rho_{\Gamma}(G \times_{\Gamma} (H + K)) = \rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(H + K) = \rho_{\Gamma}(G) \times_{\langle e \rangle} (\rho_{\Gamma}(H) + \rho_{\Gamma}(K)) \tag{3}$$

and

$$\rho_{\Gamma}((G \times_{\Gamma} H) + (G \times_{\Gamma} K)) = \rho_{\Gamma}(G \times_{\Gamma} H) + \rho_{\Gamma}(G \times_{\Gamma} K)$$

$$= (\rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(H)) + (\rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(K)). \tag{4}$$

Since we have already proved that our equality holds for  $\Gamma = \langle e \rangle$ , we obtain,

$$\rho_{\Gamma}(G) \times_{\langle e \rangle} (\rho_{\Gamma}(H) + \rho_{\Gamma}(K)) \equiv_{\langle e \rangle} (\rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(H)) + (\rho_{\Gamma}(G) \times_{\langle e \rangle} \rho_{\Gamma}(K)).$$

Therefore by equations (3) and (4) we have

$$\rho_{\Gamma}(G \times_{\Gamma} (H + K)) \equiv_{\langle e \rangle} \rho_{\Gamma}((G \times_{\Gamma} H) + (G \times_{\Gamma} K)).$$

By Theorem 3.5 we have

$$G \times_{\Gamma} (H + K) \equiv_{\Gamma} (G \times_{\Gamma} H) + (G \times_{\Gamma} K).$$

This concludes the proof.

Remark 4.6. The existence of product and co-product in the category of (n,m)-graphs with  $\Gamma$ -homomorphism playing the role of morphism shows the richness of the category. Moreover, it also shows that the  $\Gamma$ -homomorphism order, that is an order defined by  $G \leq H$  when  $G \xrightarrow{\Gamma} H$  defines a lattice on the set of all  $\Gamma$ -cores. Here the (cores of the) categorical products and coproducts play the roles of meet and join, respectively.

#### 5 Chromatic number

We know that the ordinary chromatic number of a simple graph G can be expressed as the minimum |V(H)| such that G admits a homomorphism to H. The analogue of this definition is a popular way for defining chromatic number of other types of graphs, namely, oriented graphs, k-edge-colored graphs, (n, m)-graphs, signed graphs, push graphs, etc. Here also, we can follow the same. The  $\Gamma$ -chromatic number of an (n, m)-graph is given by,

$$\chi_{\Gamma:n,m}(G) = \min\{|V(H)| : G \xrightarrow{\Gamma} H\}.$$

Moreover, for a family  $\mathcal{F}$  of (n, m)-graphs, the  $\Gamma$ -chromatic number is given by,

$$\chi_{\Gamma:n,m}(\mathcal{F}) = \max\{\chi_{\Gamma:n,m}(G) : G \in \mathcal{F}\}.$$

Let  $\Gamma \subseteq S_{2n+m}$  be an Abelian group acting on the set  $A_{n,m}$ . For  $x \in A_{n,m}$ , we call the set,  $\operatorname{Orb}_x = \{\sigma(x) : \sigma \in \Gamma\}$  as orbit of x. A consistent group  $\Gamma \subset S_{2n+m}$  is such that each orbit induced by  $\Gamma$  acting on the set  $A_{n,m}$  contains  $\overline{i}$  if and only if it contains i for  $i \in \{1, 2, \dots, 2n\}$ . Notice that, these orbits form a partition on the set  $A_{n,m}$  as the relation,  $x \sim y$  whenever  $x = \sigma(y)$  for some  $\sigma \in \Gamma$ , is an equivalence relation.

Next we establish a useful observation.

**Theorem 5.1.** Let G be any (n,m)-graph,  $\Gamma$  be a switch commutative group, then we have,

$$\chi_{\Gamma:n,m}(G) \le \chi_{n,m}(G) \le |\Gamma| \cdot \chi_{\Gamma:n,m}(G).$$

*Proof.* As  $\Gamma$ -homomorphism is, in particular, an  $\langle e \rangle$ -homomorphism, the first inequality holds. The second inequality follows from Proposition 3.4.

An immediate corollary follows.

Corollary 5.2. Let G be an (n, m)-graph and let  $\Gamma_1, \Gamma_2 \subseteq S_{2n+m}$  be two switch-commutative groups. Then we have

$$\frac{\chi_{\Gamma_1:n,m}(G)}{|\Gamma_2|} \le \chi_{\Gamma_2:n,m}(G) \le |\Gamma_1| \cdot \chi_{\Gamma_1:n,m}(G).$$

**Proposition 5.3.** Let  $\Gamma \subseteq S_{2n+m}$  be a consistent group, G be an (n, m)-graph, and G' be a  $\Gamma$ -equivalent graph of G. If a vertex v is a t-neighbor of u in G, then v must be a  $\sigma(t)$  neighbor of u in G' for some  $\sigma \in \Gamma$ .

*Proof.* Observe that, it is enough to prove the statement assuming G' is obtained from G by performing a  $\sigma$ -switch on v.

By Lemma 3.1, in G', v is a  $\overline{\sigma(\overline{t})}$ -neighbor of u. Note that,  $\overline{t} \in Orb_t$  as  $\Gamma$  is a consistent group. Since  $\overline{t} \in Orb_t$ , we have  $\sigma(\overline{t}) \in Orb_{\overline{t}} = Orb_t$ . Therefore,  $\overline{\sigma(\overline{t})} \in Orb_t$  as  $\Gamma$  is consistent.

Next we focus on studying the  $\Gamma$ -chromatic number of (n, m)-forests.

**Theorem 5.4.** Let  $\mathcal{F}$  be the family of (n,m)-forests and let k be the number of orbits of  $A_{n,m}$  with respect to the action of  $\Gamma$ . Then,

$$\chi_{\Gamma:n,m}(\mathcal{F}) \leq \begin{cases} k+2 & \text{if } k \text{ is even,} \\ k+1 & \text{if } k \text{ is odd.} \end{cases}$$

Moreover, equality holds if  $\Gamma$  is consistent.

*Proof.* We will start by proving the upper bound. First assume that k is odd. In this case consider the complete graph  $K_{k+1}$ . We will construct a complete (n, m)-graph having  $K_{k+1}$  as its underlying graph. As (k+1) is even, we know that  $K_{k+1}$  can be decomposed into  $\frac{k-1}{2}$  edge-disjoint Hamiltonian cycles and a perfect matching.

Let  $\{\alpha_1, \alpha_2, \dots \alpha_k\}$  be the representatives of the k orbits. If  $v_0v_1 \dots v_tv_0$  is the  $i^{th}$  Hamiltonian cycle from the decomposition, then assign adjacencies to the edges of it in such a way that  $v_j$  is a  $\alpha_{2j-1}$ -neighbor of  $v_{j-1}$  and  $v_{j+1}$  is a  $\alpha_{2j}$ -neighbor of  $v_j$ , for  $j \in \{0, 1, \dots, \frac{k-1}{2}\}$  where the + operation of the indices is considered modulo (k+1). Moreover, let  $u_0w_0, u_1w_1, \dots, u_{\frac{k-1}{2}}w_{\frac{k}{2}}$  be the edges of the perfect matching from the decomposition mentioned above. Assign adjacencies to these edges in such a way that  $u_j$  is a  $\alpha_k$ -neighbor of  $w_j$  for all  $j \in \{0, 1, \dots, \frac{k-1}{2}\}$ . With a little abuse of notation, we denote the so obtained (n, m)-graph by  $K_{k+1}$  itself.

Notice that, every vertex of  $K_{k+1}$  has an  $\alpha_i$ -neighbor for every  $i \in \{1, 2, \dots, k\}$ . We claim that every (n, m)-forest admit a  $\Gamma$ -homomorphism to  $K_{k+1}$ . If not, then there exists a minimal (with respect to number of vertices) counter-example F that does not admit  $\Gamma$ -homomorphism to  $K_{k+1}$ . Let u be a leaf, having v as its neighbor, of F, then  $F \setminus \{u\}$  is no longer a minimal counter-example, thus it admits a  $\Gamma$ -homomorphism f to  $K_{k+1}$ . That means, there exists a  $\Gamma$ -equivalence (n, m)-graph  $F' \setminus \{u\}$  such that f is a  $\langle e \rangle$ -homomorphism of it to  $K_{k+1}$ . Also assume that, F' is such a  $\Gamma$ -equivalent (n, m)-graph of F, that v is a  $\alpha_i$ -neighbor of u for some  $i \in \{1, 2, \dots, k\}$ . This is possible as one can switch the vertex v to make the adjacency of u with v match the corresponding orbit's representative. Now, we extend f to an  $\langle e \rangle$ -homomorphism of F' to  $K_{k+1}$  by mapping v to the  $\alpha_i$ -neighbor f(u) in  $K_{k+1}$ . That means, there exists a  $\Gamma$ -homomorphism of F to  $K_{k+1}$ . This contradicts the minimality of F. Hence every (n, m)-forest admits a  $\Gamma$ -homomorphism to  $K_{n,m}$ .

Secondly, assume that k is even. Note that, if there were (k+1) orbits instead, then by what we have proved above, it would be possible to show that every (n, m)-forest admits a  $\Gamma$ -homomorphism to an (n, m)-graph having  $K_{k+2}$  as underlying graph. Therefore, assuming a dummy orbit we are done with this case too.

Next we will prove the tightness of the upper bound when  $\Gamma$  is consistent. Let  $\{\alpha_1, \alpha_2, \cdots \alpha_k\}$  be the representatives of the k orbits. For odd values of k, consider the star (n, m)-graph S on (k + 1) vertices: the central vertex v having k neighbors  $v_1, v_2, \cdots, v_k$ . Let  $v_i$  be a  $\alpha_i$ -neighbor of v for all  $i \in \{1, 2, \cdots, k\}$ . As, no matter how we switch the vertices of S, the vertex v will have k distinctly adjacent neighbors. Therefore we have  $\chi_{\Gamma:n,m}(S) \geq k+1$ , and thus  $\chi_{\Gamma:n,m}(\mathcal{F}) = k+1$  when k is odd and  $\Gamma$  is consistent.

For even values of k, consider a rooted tree T of height two in which every vertex, other than the leaves, has exactly one  $\alpha_i$ -neighbor for  $i \in \{1, 2, \dots, k\}$ . Suppose T admits

a  $\Gamma$ -homomorphism f to an (n, m)-graph H. Let r be the root of T. If H has (k+1) vertices, then the images of the vertices from N[r] under f will be a spanning subgraph in H. Furthermore, notice that each vertex of N[r] has at least one  $\beta_i$ -neighbor, where  $\beta_i$  belongs to the  $i^{th}$  orbit. Thus their images should also have the same property, that is each of them must have at least one  $\beta_i$ -neighbor, where  $\beta_i$  belongs to the  $i^{th}$  orbit. However, as H has only (k+1) vertices, each of its vertices are forced to have exactly one  $\beta_i$ -neighbor, where  $\beta_i$  belongs to the  $i^{th}$  orbit. Now if we restrict ourselves to only the neighbors whose type is from a particular orbit, that must give us a perfect matching, which is impossible as (k+1) is odd. Therefore, H must have at least (k+2) vertices which implies the lower bound.

The above result implies the upper bound of Theorem 1.1 of [24].

### 6 Concluding remarks

In this article, we introduced a generalized switch operation on (n, m)-graphs and studied their basic algebraic properties. Naturally, this topic generates a lot of interesting open questions, especially, in an effort of extending the known results in the domain of graph homomorphisms. We list a few of them here.

- (i) Is it possible to generalize the notion of exponential graphs (see section 2.4 in [15]) in the category of (n, m)-graphs with respect to  $\Gamma$ -homomorphism where  $\Gamma$  is switch-commutative?
- (ii) Is it possible to obtain the analogue of the Density Theorem (see Theorem 3.30 in [15]) in this context? At present, such an analogue is unknown even for (0, 2)-graphs.
- (iii) Finding the  $\Gamma$ -chromatic number for other graph families like planar graphs, partial 2-trees, outerplanar graphs, cycles, grids, graphs having bounded maximum degree, etc. can be other directions of study in this set up.
- (iv) Studying the (n, m)-cycles may further lead us to finding a characterization of  $\Gamma$ -equivalent graphs, similar to what Zaslavsky [31] did for signed graphs.
- (v) If  $\Gamma$  is not switch-commutative, then does the categorical product exist?
- (vi) Given a pre-decided (n, m)-graph H, a switch-commutative group  $\Gamma$  and an input (n, m)-graph G, the decision problem "does G admit a  $\Gamma$ -homomorphism to H?" is in NP due to Proposition 3.4. Can we characterize the full dichotomy of this problem?

As a remark, it is worth mentioning that using the notion of generalized switch (implicitly), it was possible to improve the existing upper bounds of the  $\langle e \rangle$ -chromatic number of (n,m)-partial 2-trees where 2n+m=3 [9]. Therefore, it will not be surprising if  $\Gamma$ -homomorphism becomes useful as a technique to establish bounds for (n,m)-chromatic number of graphs.

**Acknowledgement:** This work is partially supported by French ANR project "HOSI-GRA" (ANR-17-CE40-0022), IFCAM project "Applications of graph homomorphisms" (MA/IFCAM/18/39), SERB-MATRICS "Oriented chromatic and clique number of planar graphs" (MTR/2021/000858).

#### References

- [1] N. Alon and T. H. Marshall. Homomorphisms of edge-colored graphs and coxeter groups. *Journal of Algebraic Combinatorics*, 8(1):5–13, 1998.
- [2] R. Angles, M. Arenas, P. Barceló, A. Hogan, J. Reutter, and D. Vrgoč. Foundations of modern query languages for graph databases. *ACM Computing Surveys (CSUR)*, 50(5):1–40, 2017.
- [3] M. Artin. Algebra. Birkhäuser, 1998.
- [4] L. Beaudou, F. Foucaud, F. Madelaine, L. Nourine, and G. Richard. Complexity of conjunctive regular path query homomorphisms. In *Conference on Computability in Europe*, pages 108–119. Springer, 2019.
- [5] O. Borodin, S.-J. Kim, A. Kostochka, and D. West. Homomorphisms from sparse graphs with large girth. *Journal of Combinatorial Theory, Series B*, 90(1):147–159, 2004.
- [6] R. C. Brewster, F. Foucaud, P. Hell, and R. Naserasr. The complexity of signed graph and edge-coloured graph homomorphisms. *Discrete Mathematics*, 340(2):223– 235, 2017.
- [7] R. C. Brewster and T. Graves. Edge-switching homomorphisms of edge-coloured graphs. *Discrete mathematics*, 309(18):5540–5546, 2009.
- [8] R. C. Brewster, A. Kidner, and G. MacGillivray. k-colouring (m, n)-mixed graphs with switching.  $arXiv\ preprint\ arXiv:2203.08070,\ 2022.$
- [9] S. Das, A. Lahiri, S. Nandi, S. Sen, and S. Taruni. An update on (n, m)-chromatic numbers. CoRR, abs/2306.08069, 2023.
- [10] C. Duffy, G. MacGillivray, and B. Tremblay. Switching *m*-edge-coloured graphs using non-abelian groups. *arXiv preprint arXiv:2207.12528*, 2022.
- [11] T. Feder, P. Hell, and B. Mohar. Acyclic homomorphisms and circular colorings of digraphs. SIAM Journal on Discrete Mathematics, 17(1):161–169, 2003.
- [12] F. Foucaud, H. Hocquard, D. Lajou, V. Mitsou, and T. Pierron. Graph modification for edge-coloured and signed graph homomorphism problems: Parameterized and classical complexity. *Algorithmica*, 84(5):1183–1212, 2022.
- [13] G. Guśpiel and G. Gutowski. Universal targets for homomorphisms of edge-colored graphs. *Journal of Combinatorial Theory, Series B*, 127:53–64, 2017.

- [14] P. Hell and J. Nešetřil. On the complexity of H-coloring. *Journal of Combinatorial Theory*, Series B, 48(1):92–110, 1990.
- [15] P. Hell and J. Nesetril. *Graphs and homomorphisms*, volume 28. OUP Oxford, 2004.
- [16] N. Jacobson. Basic algebra I. Courier Corporation, 2012.
- [17] A. Kidner. Gamma-Switchable 2-Colourings of (m, n)-Mixed Graphs. PhD thesis, 2021.
- [18] W. F. Klostermeyer and G. MacGillivray. Homomorphisms and oriented colorings of equivalence classes of oriented graphs. *Discrete Mathematics*, 274(1-3):161–172, 2004.
- [19] E. Leclerc, G. MacGillivray, and J. M. Warren. Switching (m, n)-mixed graphs with respect to abelian groups.  $arXiv\ preprint\ arXiv:2110.01576$ , 2021.
- [20] T. H. Marshall. Homomorphism bounds for oriented planar graphs. *J. Graph Theory*, 55(3):175–190, 2007.
- [21] R. Naserasr, E. Rollová, and É. Sopena. Homomorphisms of signed graphs. *Journal of Graph Theory*, 79(3):178–212, 2015.
- [22] R. Naserasr, S. Sen, and E. Sopena. The homomorphism order of signed graphs. Journal of Combinatorial Mathematics and Combinatorial Computing, 116(4):169–182, 2020.
- [23] R. Naserasr, E. Sopena, and T. Zaslavsky. Homomorphisms of signed graphs: An update. *European Journal of Combinatorics*, 91:103222, 2021.
- [24] J. Nešetřil and A. Raspaud. Colored homomorphisms of colored mixed graphs. *Journal of Combinatorial Theory, Series B*, 80(1):147–155, 2000.
- [25] P. Ochem, A. Pinlou, and S. Sen. Homomorphisms of 2-edge-colored triangle-free planar graphs. *J. Graph Theory*, 85(1):258–277, 2017.
- [26] S. Sen. On homomorphisms of oriented graphs with respect to the push operation. *Discrete Mathematics*, 340(8):1986–1995, 2017.
- [27] E. Sopena. The chromatic number of oriented graphs. *Journal of Graph Theory*, 25(3):191–205, 1997.
- [28] É. Sopena. Homomorphisms and colourings of oriented graphs: An updated survey. *Discrete Mathematics*, 339(7):1993–2005, 2016.
- [29] E. Sopena, E. Rollová, and R. Naserasr. Homomorphisms of planar signed graphs to signed projective cubes. *Discrete Mathematics & Theoretical Computer Science*, 15, 2013.
- [30] D. B. West. *Introduction to graph theory*, volume 2. Prentice hall Upper Saddle River, 2001.

- [31] T. Zaslavsky. Signed graphs. Discrete Applied Mathematics, 4(1):47–74, 1982.
- [32] X. Zhu. Uniquely H-colorable graphs with large girth. *Journal of Graph Theory*, 23(1):33–41, 1996.