

THE REVERSE MATHEMATICS OF CAC FOR TREES

JULIEN CERVELLE, WILLIAM GAUDELIER, AND LUDOVIC PATEY

Abstract. *CAC for trees* is the statement asserting that any infinite subtree of $\mathbb{N}^{<\mathbb{N}}$ has an infinite path or an infinite antichain. In this paper, we study the computational strength of this theorem from a reverse mathematical viewpoint. We prove that *CAC for trees* is robust, that is, there exist several characterizations, some of which already appear in the literature, namely, the statement *SHER* introduced by Dorais et al. [8], and the statement $\text{TAC} + \text{B}\Sigma_2^0$ where *TAC* is the tree antichain theorem introduced by Conidis [6]. We show that *CAC for trees* is computationally very weak, in that it admits probabilistic solutions.

§1. Introduction. In this paper we study the computability-theoretic strength of the statement *CAC for trees*, which is a variation on the well-studied chain-antichain theorem (*CAC*). It turns out *CAC for trees* has different characterizations, making it a robust notion, suitable for future studies in reverse mathematics.

We are going to use two frameworks: reverse mathematics and computable reduction. For a good and more complete introduction to reverse mathematics, see Simpson [18], or Hirschfeldt [12] which also covers the computable reduction and classical results on Ramsey’s theorem.

Reverse mathematics is a foundational program which seeks to determine the optimal axioms to prove “ordinary” theorems. It itself uses the

This project started as the study of Ramsey-like theorems for 3-variable forbidden patterns. The attempt to prove Corollary 6.13 naturally led to the study of the *SHER* principle, already defined by Dorais and al. [8]. Thanks to multiple personal communications with François Dorais, we realized that the *SHER* principle is closely related to trees, and more precisely, equivalent to the Chain-Antichain principle for trees, a principle studied by Binns et al. in [3]. We later realized that *SHER* is also equivalent to $\text{TAC} + \text{B}\Sigma_2^0$, where *TAC* is an antichain principle for completely branching c.e. trees, defined by Conidis [6]. Some of the results are therefore independent rediscoveries of some theorems from [3, 6], but in a more unified setting. The authors are thankful to Chris Conidis, François Dorais and Alberto Marcone for interesting comments and discussions. The authors are also thankful to the anonymous referee for his careful reading and his numerous improvement suggestions. The authors were partially supported by grant ANR “ACTC” #ANR-19-CE48-0012-01.

framework of subsystems of second-order arithmetic, with a base theory called RCA_0 , which informally captures “computable mathematics”.

Computable reduction makes precise the idea of being able to computably solve a problem P using another problem Q . A problem is defined as a Π_2^1 formula in the language of second-order arithmetic, thought to be of the form $\forall X(\psi(X) \implies \exists Y\varphi(X, Y))$, where ψ and φ are arithmetical formulas. An instance of a problem is a set X verifying $\psi(X)$, and a solution of an instance X is a set Y such that $\varphi(X, Y)$. With this formalism, we say that “ P is computably reducible to Q ”, and we write $P \leq_c Q$ if for any instance I of P , there is an I -computable instance \widehat{I} of Q , such that, for any solution \widehat{S} of Q , there is an $\widehat{S} \oplus I$ -computable solution S of P .

The early study of reverse mathematics has seen the emergence of four subsystems of second-order arithmetic, linearly ordered by the provability relation, such that most of the ordinary theorems are either provable in RCA_0 , or equivalent in RCA_0 to one of them. These subsystems, together with RCA_0 , form the “Big Five”. Among the theorems studied in reverse mathematics, Ramsey’s theorem for pairs RT_2^2 plays an important role, since it is historically the first example of a natural statement which does not satisfy this empirical observation. The theorems we study in this paper are all consequences of RT_2^2 .

1.1. A chain-antichain theorem for trees. Among the consequences of Ramsey’s theorem for pairs, the chain-antichain theorem received a particular focus in reverse mathematics.

DEFINITION 1.1 (CAC, chain-antichain theorem). Every infinite partial order has either an infinite chain or an infinite antichain.

CAC was first studied in [13] by Hirschfeldt and Shore, following a question raised by Cholak, Jockusch and Slaman in [4, Question 13.8] asking whether or not $\text{CAC} \implies \text{RT}_2^2$ over RCA_0 , for which they proved the answer is negative (Corollary 3.12). The reciprocal $\text{RCA}_0 \vdash \text{RT}_2^2 \implies \text{CAC}$ is easier to obtain, by defining a coloring such that $\{x, y\}$ has color 1 if its elements are comparable, and 0 otherwise. Any homogeneous set for this coloring is either a chain or an antichain, depending on its color.

In this article, we focus on the special case where the order is the predecessor relation $<$ on a tree.

DEFINITION 1.2 (CAC for trees). Every infinite subtree of $\mathbb{N}^{<\mathbb{N}}$ has an infinite path or an infinite antichain.

This statement was first introduced by Binns et al. in [3], where the authors showed that every infinite computable tree must have either an infinite computable chain or an infinite Π_1^0 antichain. Furthermore, they

showed that these bounds are optimal, by constructing an infinite computable tree which has no infinite Σ_1^0 chain or antichain. They also showed that $\text{RCA}_0 + \text{WKL} \not\vdash \text{CAC}$ for binary trees (see Definition 2.1).

1.2. Ramsey-like theorems. In [17], Patey identified a formal class of theorems, encompassing several statements surrounding Ramsey's theorem. Indeed, many of them are of the form “for every coloring $f : [\mathbb{N}]^n \rightarrow k$ avoiding some set of forbidden patterns, there exists an infinite set $H \subseteq \mathbb{N}$ avoiding some other set of forbidden patterns (relative to f)”

For example the Erdős-Moser theorem (EM) asserts that, “for any coloring $f : [\mathbb{N}]^2 \rightarrow 2$, there exists an infinite set $H \subseteq \mathbb{N}$ which is transitive for f ”, i.e. $\forall i < 2, \forall x < y < z \in H, f(x, y) = i \wedge f(y, z) = i \implies f(x, z) = i$. In other terms, we want H to avoid the patterns that would make it not transitive for f , i.e. $f(x, y) = i \wedge f(y, z) = i \wedge f(x, z) = 1 - i$ for any $i < 2$. Another example comes from ADS which is equivalent over RCA_0 (see [13, Theorem 5.3]) to the statement “for any transitive coloring $f : [\mathbb{N}]^2 \rightarrow 2$ (i.e. avoiding certain patterns), there exists an infinite set $H \subseteq \mathbb{N}$ which is f -homogeneous”. With these definitions, one sees that RT_2^2 is equivalent to $\text{EM} + \text{ADS}$ over RCA_0 , since EM takes any coloring and “turns it into” a transitive one, and ADS takes any transitive coloring and finds an infinite set which is homogeneous for it.

Forbidden patterns on 3 variables and 2 colors are generated by the following three basic patterns:

- (1) $f(x, y) = i \wedge f(y, z) = i \wedge f(x, z) = 1 - i$
- (2) $f(x, y) = i \wedge f(y, z) = 1 - i \wedge f(x, z) = i$
- (3) $f(x, y) = 1 - i \wedge f(y, z) = i \wedge f(x, z) = i$

Avoiding them respectively leads to **transitivity**, **semi-ancestry**, and **semi-heredity** (for the color i). Each of them generates two ramsey-like statements, one restricting the input coloring, and one restricting the output infinite set, namely “for any 2-coloring of pairs avoiding the forbidden pattern, there exists an infinite homogeneous set” and “for any 2-coloring of pairs, there exists an infinite set $H \subseteq \mathbb{N}$ which avoids the forbidden pattern”. We now survey the known results about these three patterns.

Transitivity. The statement “for any 2-coloring of pairs, there exists an infinite set which is transitive for some color” is a weaker version of EM. The Erdős-Moser theorem was proven to be strictly weaker than Ramsey's theorem for pairs over RCA_0 by Lerman, Solomon and Towsner [16, Corollary 1.16]. On the other hand, the statement “for any 2-coloring of pairs which is transitive for some color, there exists an infinite homogeneous set” is equivalent to CAC (see [13, Theorem 5.2]), which is also known to be strictly weaker than RT_2^2 over RCA_0 (see Hirschfeldt and Shore [13, Corollary 3.12]).

Semi-ancestry. The statement “for any 2-coloring of pairs which has semi-ancestry for some color, there exists an infinite homogeneous set” is a consequence of the statement STRIV, defined by Dorais et al. [8, Statement 5.12]), because a 2-coloring is semi-trivial if and only if it has semi-ancestry. And STRIV itself is equivalent to $\forall k, \text{RT}_k^1$ (see the remark below its definition). The statement “for any 2-coloring of pairs, there exists an infinite set which has semi-ancestry for some color” is equivalent to RT_2^2 (see Proposition 6.11).

Semi-heredity. The statement “for any 2-coloring of pairs which is semi-hereditary for some color, there exists an infinite homogeneous set” is the statement SHER, which was first introduced by Dorais et al. [8, Statement 5.11]. In Section 6, we will show that it is equivalent to CAC for trees. Finally, the statement “for any 2-coloring of pairs, there exists an infinite set which is semi-hereditary for some color” is equivalent to RT_2^2 (see Corollary 6.13).

1.3. Notation. Let the symbols $\exists^\infty x$ and $\forall^\infty x$ be abbreviations for $\forall y, \exists x > y$ and $\exists y, \forall x > y$ respectively. In particular they are the dual of each other.

Given $x, y \in \mathbb{N} \cup \{\pm\infty\}$, we define $\llbracket x, y \rrbracket := \{z \in \mathbb{N} \mid z \geq x \wedge z \leq y\}$, an inequality being strict when its respective bracket is flipped, e.g. $\llbracket x, y \rrbracket := \{z \in \mathbb{N} \mid z \geq x \wedge z < y\}$. As in set theory, an integer n can also represent the set of all integers strictly smaller to it, i.e. $\llbracket 0, n \rrbracket$. Moreover $\langle -, \dots, - \rangle$ represents a bijection from \mathbb{N}^n to \mathbb{N} (for some n), which verifies $\forall x, y \in \mathbb{N}, \langle x_1, \dots, x_k \rangle \geq \max\{x_1, \dots, x_k\}$ and which is increasing on each variable. Given a set $A \subseteq \mathbb{N}$, we denote by χ_A its **indicator function**.

A **string** is a finite sequence of integers, and the set of all strings is denoted $\mathbb{N}^{<\mathbb{N}}$. In particular, a **binary string** is a finite sequence of 0 and 1, and the set of all binary strings is denoted $2^{<\mathbb{N}}$. The **length** of a given string $\sigma : n \rightarrow \mathbb{N}$ is the integer $|\sigma| := n$. The **empty string** $\emptyset \rightarrow \mathbb{N}$ is denoted ε . Given two strings σ and τ of respective length ℓ and m , we define their **concatenation** as the finite sequence $\sigma \cdot \tau : \ell + m \rightarrow \mathbb{N}$ which, given $j < \ell + m$, associates $\sigma(j)$ if $j < \ell$, and $\tau(j - \ell)$ otherwise. So we usually write a string $\sigma : n \rightarrow \mathbb{N}$ as $\sigma_0 \cdot \dots \cdot \sigma_{n-1}$, where $\forall j < n, \sigma_j := \sigma(j)$. We define the partial order on strings “is **prefix of**”, denoted \prec , by $\sigma \prec \tau \iff |\sigma| < |\tau| \wedge \forall j < |\sigma|, \sigma(j) = \tau(j)$, and denote by \preceq its reflexive closure. When two strings σ and τ are **incomparable** for \prec , we write $\sigma \perp \tau$.

A **tree** T is a subset of $\mathbb{N}^{<\mathbb{N}}$ which is downward closed for \prec , i.e. $\forall \sigma \in T, \tau \prec \sigma \implies \tau \in T$. A **binary tree** is a subset of $2^{<\mathbb{N}}$ which is downward closed for \prec . A subset $S \subseteq T$ of a tree is a **chain** when it is linearly ordered for \prec , and it is an **antichain** when its elements are pairwise incomparable for \prec .

1.4. Organization of the paper. In Section 2, we prove the robustness of CAC for trees in reverse mathematics and over the computable reduction, by proving its equivalence with several variants of the statement. In Section 3, we provide a probabilistic proof of CAC for trees, and give a precise analysis of this proof in terms of DNC functions. In Section 4 we show that both ADS and EM imply CAC for trees. In Section 5 we prove there is a computable instance of TAC whose solutions are all of hyperimmune degree, with almost explicit witness. In particular we show a computable instance of TAC can avoid a uniform sequence of Δ_2^0 sets. In Section 6, we show the equivalence between CAC for trees and SHER. In Section 7, we explore the relations between stable versions of previously mentioned statements, namely CAC for trees, ADS and SHER, leading to the result that CAC for stable c.e. trees admits low solutions. Finally, in Section 8, we conclude the paper with remaining open questions and summary diagrams.

§2. Equivalent definitions. In this section, we study some variations of CAC for trees, and prove they are all equivalent. We also study TAC and show that $\text{TAC} + \text{B}\Sigma_2^0$ is equivalent to CAC for trees. We start by defining these statements.

DEFINITION 2.1 (CAC for c.e. (binary) trees). Every infinite c.e. (binary) subtree of $\mathbb{N}^{<\mathbb{N}}$ has an infinite path or an infinite antichain.

Remark 2.2. In the context of reverse mathematics “being c.e.” is a notion that is relative to the model considered, i.e. an object is c.e. when it can be approximated in a c.e. manner by objects from the model.

DEFINITION 2.3 (Completely branching tree). A node σ of a tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$ is a **split node** when there is $n_0, n_1 \in \mathbb{N}$ such that $\forall i < 2, \sigma \cdot n_i \in T$. In particular, if T is a binary tree, then σ is a split node when both $\sigma \cdot 0 \in T$ and $\sigma \cdot 1 \in T$. A tree $T \subseteq 2^{<\mathbb{N}}$ is **completely branching** when, for any of its node σ , if σ is not a leaf then it is a split node.

The following statement was introduced by Conidis [6] (personal communication), motivated by the reverse mathematics of commutative noetherian rings.

DEFINITION 2.4 (TAC [6], tree antichain theorem). Any infinite c.e. subtree of $2^{<\mathbb{N}}$ which is completely branching, contains an infinite antichain.

Conidis proved that TAC follows from CAC over RCA_0 , and constructed an instance of TAC, all of whose solutions are of hyperimmune degree. In particular, $\text{RCA}_0 + \text{WKL} \not\equiv \text{TAC}$. Now we can proceed with the proof of the equivalence.

THEOREM 2.5. *The following statements are equivalent over RCA_0 and computable reduction:*

- (1) CAC for trees
- (2) CAC for c.e. trees
- (3) CAC for c.e. binary trees
- (4) TAC + $\text{B}\Sigma_2^0$

PROOF. (2) \implies (1) and (2) \implies (3) are immediate. (3) \implies (4) is Proposition 2.6 and Proposition 2.7. (4) \implies (2) is Proposition 2.8. (1) \implies (2) is Proposition 2.9. \dashv

We shall see in Section 3 that the use of $\text{B}\Sigma_2^0$ is necessary in the equivalence above, as TAC does not imply $\text{B}\Sigma_2^0$ over RCA_0 .

PROPOSITION 2.6. $\text{RCA}_0 \vdash \text{CAC for c.e. binary trees} \implies \text{TAC and TAC} \leq_c \text{CAC for c.e. binary trees}.$

PROOF. Let $T \subseteq 2^{<\mathbb{N}}$ be an infinite completely branching c.e. tree. By the statement CAC for c.e. binary trees, either there is an infinite antichain, or an infinite path P . In the former case, we are done. In the latter case, using the fact that T is completely branching, the set $\{\sigma \cdot (1-i) \mid \sigma \cdot i \prec P\}$ is an infinite antichain of T . \dashv

PROPOSITION 2.7. $\text{RCA}_0 \vdash \text{CAC for c.e. binary trees} \implies \forall k, \text{RT}_k^1$
and $\forall k, \text{RT}_k^1 \leq_c \text{CAC for c.e. binary trees}.$

PROOF. Let $f : \mathbb{N} \rightarrow k$ be a coloring, there are two possibilities. Either $\exists i < k, \exists^\infty x, f(x) = i$, in which case there is an infinite computable f -homogeneous set. Otherwise $\forall i < k, \forall^\infty x, f(x) \neq i$, in which case we define an infinite binary c.e. tree T , via a strictly increasing sequence of computable trees $(T_j)_{j \in \mathbb{N}}$ defined by $T_0 := \{0^i \mid i < k\}$ and $T_{s+1} := T_s \cup \{0^{f(s)} \cdot 1^{m+1}\}$, where m is the number of $x < s$ such that $f(x) = f(s)$.

Every antichain in T is of size at most k , thus, by CAC for c.e. binary trees, T must contain an infinite path, and so $\exists i < k, \exists^\infty x, f(x) = i$, which is a contradiction. \dashv

PROPOSITION 2.8. $\text{RCA}_0 \vdash \text{TAC} + \text{B}\Sigma_2^0 \implies \text{CAC for c.e. trees and CAC for c.e. trees} \leq_c \text{TAC}$

PROOF. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be an infinite c.e. tree. We can deal with two cases directly: when T has a node with infinitely many immediate children, as they contain a computable infinite antichain; and when T has finitely many split nodes, in which case it has finitely many paths P_0, \dots, P_{k-1} , which are all computable. Moreover one of them is infinite, as otherwise they would all be finite, i.e. $\forall i < k, \exists s, \forall n > s, n \notin P_i$, and thus their union would be finite since $\text{B}\Pi_1^0$ (which is equivalent to $\text{B}\Sigma_2^0$) yields $\exists b, \forall i < k, \exists s < b, \forall n > s, n \notin P_i$. But since $T = \bigcup_{i < k} P_i$, this would lead to a contradiction.

A **split triple** of T is a triple $(\mu, n_0, n_1) \in T \times \mathbb{N} \times \mathbb{N}$ such that $\mu, \mu \cdot n_0, \mu \cdot n_1 \in T$. In particular, μ is a split node in T .

Idea. The general idea is to build greedily a completely branching c.e. tree S by looking for split triples in T , and mapping them to split nodes in S . This correspondence is witnessed by an injective function $f : S \rightarrow T$ that will be constructed alongside S . The main difficulty is that, since T is c.e., a split node ρ can be discovered after μ even though $\rho \prec \mu$, which means that we will not be able to ensure that S can be embedded in T . In particular, f will not be a tree morphism. However, the only property that needs to be ensured is that for every infinite antichain A of S , the set $f(A)$ will be an infinite antichain of T . To guarantee this, the function f needs to verify

$$(*) \quad \forall \sigma, \nu \in S, \sigma \perp \nu \implies f(\sigma) \perp f(\nu)$$

During the construction, we are going to associate to each node $\sigma \in S$ a set $N_\sigma \subseteq T$, which might decrease in size over time ($N_\sigma^0 \supseteq N_\sigma^1 \supseteq \dots$), with the property that at every step s , the elements of $\{N_\sigma^s : \sigma \in S\}$ are pairwise disjoint, and their union contains cofinitely many elements of T . The role of N_σ is to indicate that “if a split triple is found in N_σ , then the nodes in S , associated via f , must be above σ ”.

Construction. Initially, $N_\varepsilon^0 := T$, $S := \{\varepsilon\}$ and $f(\varepsilon) := \varepsilon$.

At step s , suppose we have defined a finite, completely branching binary tree $S \subseteq 2^{<\mathbb{N}}$, and for every $\sigma \in S$, a set $N_\sigma^s \subseteq T$ such that $\{N_\sigma^s : \sigma \in S\}$ forms a partition of T minus finitely many elements. Moreover, assume we have defined a mapping $f : S \rightarrow T$.

Search for a split triple (μ, n_0, n_1) in $\bigcup_{\sigma \in S} N_\sigma^s$. Let $\sigma \in S$ be such that $\mu \in N_\sigma^s$. Let τ be any leaf of S such that $\tau \succeq \sigma$ (for example pick the left-most successor of σ). Add $\tau \cdot 0$ and $\tau \cdot 1$ to S , and set $f(\tau \cdot i) = \mu \cdot n_i$ for each $i < 2$. Note that S is still completely branching.

Then, split N_σ^s into three disjoint subsets N_σ^{s+1} , $N_{\tau \cdot 0}^{s+1}$, $N_{\tau \cdot 1}^{s+1}$ as follows: (for $i < 2$) $N_{\tau \cdot i}^{s+1} := \{\rho \in N_\sigma^s \mid \rho \succ \mu \cdot n_i\}$ and $N_\sigma^{s+1} := \{\rho \in N_\sigma^s \mid \forall i < 2, \rho \perp \mu \cdot n_i\}$. Note that these sets do not form a partition of N_σ^s as we missed the nodes in $\{\rho \in N_\sigma^s \mid \rho \preceq \mu\}$, fortunately there are only finitely many of them. Lastly, set $N_\nu^{s+1} := N_\nu^s$ for every $\nu \in S - \{\sigma, \tau \cdot 0, \tau \cdot 1\}$.

Verification. First, let us prove that at any step s , $\bigcup_{\sigma \in S} N_\sigma^s$ contains infinitely many split triples, which ensures that the search always terminates. Note that $T - \bigcup_{\sigma \in S} N_\sigma^s$ is a c.e. subset of $\bigcup_{\sigma \in S} \{\rho \mid \rho \preceq \sigma\}$, hence exists by bounded Σ_1 comprehension, which follows from RCA_0 . Moreover, by assumption, T is a finitely branching c.e. tree, which means that every $\rho \in T - \bigcup_{\sigma \in S} N_\sigma^s$ belongs to a bounded number of split triples of T . By $\text{B}\Sigma_1^0$ (which follows from RCA_0), the number of split triples in T which involve a node from $T - \bigcup_{\sigma \in S} N_\sigma^s$ is bounded. Since by assumption, T

contains infinitely many split triples, $T - \bigcup_{\sigma \in S} N_\sigma^s$ must contain infinitely many of them.

Second, we prove by induction on s that $\forall s \in \mathbb{N}, \forall \sigma \neq \nu \in S, N_\sigma^s \perp N_\nu^s$, i.e. $\forall \mu \in N_\sigma^s, \forall \rho \in N_\nu^s, \mu \perp \rho$. At step 0, the assertion is trivially verified. At step s , suppose we found the split triple (μ, n_0, n_1) in the set N_σ^s , and that $\forall i < 2, f(\tau \cdot i) = \mu \cdot n_i$ where $\tau \succ \sigma$. Since μ was found in N_σ^s , the latter is split into $N_{\tau \cdot 0}^{s+1}, N_{\tau \cdot 1}^{s+1}$ and N_σ^{s+1} , the other sets remain identical. By construction, and because they are all subsets of N_σ^s , the assertion holds.

We now prove (*), consider $\sigma, \nu \in S$ such that $\sigma \perp \nu$. WLOG suppose ν was added to S sooner than σ , more precisely $f(\nu)$ appeared (as child in a split triple) at step s in some set N_ν^s , so N_ν^{s+1} contains $f(\nu)$ by construction. Since σ was added to S after ν , there exists $\rho \in S$ such that $f(\sigma) \in N_\rho^{s+1}$. By contradiction $\rho \neq \sigma$ holds, as otherwise $f(\sigma) \in N_\tau^{s+1}$, and so σ would extend τ by construction of S . Thus by using the previous assertion, we deduce $f(\sigma) \perp f(\nu)$. \dashv

PROPOSITION 2.9. $\text{RCA}_0 \vdash \text{CAC for trees} \implies \text{CAC for c.e. trees and CAC for c.e. trees} \leq_c \text{CAC for trees}$.

PROOF. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be a c.e. tree. We define the computable tree $S \subseteq \mathbb{N}^{<\mathbb{N}}$ by $\langle n_0, s_0 \rangle \cdot \dots \cdot \langle n_{k-1}, s_{k-1} \rangle \in S$ if and only if for all $j < k$, s_j is the smallest integer such that $n_0 \cdot \dots \cdot n_j \in T[s_j]$, where $T[s_j]$ is the approximation of T at stage s_j .

By CAC for trees, there is an infinite chain (resp. antichain) in S , and by forgetting the second component of each string, we obtain an infinite chain (resp. antichain) in T . \dashv

Before finishing this section, we introduce a set version of the principle TAC, which is more convenient to manipulate than TAC. Indeed, when working with TAC, the downward-closure of the tree is not relevant, and we naturally end up taking an infinite computable subset of the tree rather than working with the c.e. tree. This motivates the following definitions.

DEFINITION 2.10. A set $X \subseteq 2^{<\mathbb{N}}$ is **completely branching** if for every $\sigma \in 2^{<\mathbb{N}}$, $\sigma \cdot 0 \in X$ iff $\sigma \cdot 1 \in X$.

Note that the above definition is compatible with the notion of completely branching tree.

DEFINITION 2.11 (SAC, set antichain theorem). Every infinite completely branching set $X \subseteq 2^{<\mathbb{N}}$ has an infinite antichain.

The set antichain theorem is equivalent to the tree antichain theorem, as shows the following lemma.

LEMMA 2.12. $\text{RCA}_0 \vdash \text{SAC} \iff \text{TAC}$.

PROOF. SAC \implies TAC. Let $T \subseteq 2^{<\mathbb{N}}$ be an infinite, completely branching c.e. tree. Let $S \subseteq T$ be an infinite computable, completely branching set. By SAC, there is an infinite antichain $A \subseteq X$. In particular, A is an antichain for T .

SAC \iff TAC. Let $S \subseteq 2^{<\mathbb{N}}$ be an infinite computable, completely branching set. One can define an infinite computable tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$ by letting $\sigma \in T$ iff for every $n < |\sigma|$, $\sigma(n)$ codes for a binary string $\tau_n \in S$, such that for every $n < |\sigma| - 1$, $\tau_n \prec \tau_{n+1}$, and there is no string in S strictly between τ_n and τ_{n+1} . The tree T is such that every chain in T codes for a chain in S , and every antichain in T codes for an antichain in S . We can see T as an instance of CAC for trees. Moreover, since S is completely branching, then T has infinitely many split triples, so the proof of Proposition 2.8 applied to this instance T of CAC for trees does not use BS_2^0 . Thus there is either an infinite chain for T , or an infinite antichain for S . With the appropriate decoding, we obtain an infinite antichain for S . \dashv

§3. Probabilistic proofs of SAC. The restriction of CAC to trees yields a strictly weaker statement from the viewpoint the arithmetical bounds in the arithmetic hierarchy. Indeed, by Herrmann [11, Theorem 3.1], there is a computable partial order with no Δ_2^0 infinite chain or antichain, while by Binns et al. in [3, Theorem 6.2], every infinite computable tree must have either an infinite computable chain or an infinite Π_1^0 antichain. In this section, we go one step further in the study of the weakness of CAC for trees by proving that SAC admits probabilistic solutions. Before this, we prove two technical lemmas.

LEMMA 3.1 (RCA_0). *Let $S \subseteq 2^{<\mathbb{N}}$ be an infinite completely branching set. Then for every n , there exists an antichain of size n .*

PROOF. By finite Ramsey's theorem for pairs and 2 colors (which holds in RCA_0), there exists some $p \in \mathbb{N}$ such that for every 2-coloring of $[p]^2$, there exists a homogeneous set of size n . Since S is infinite, there exists a subset $P \subseteq S$ of size p . By choice of p , there exists a subset $Q \subseteq P$ of size n such that either Q is a chain, or an antichain. In the latter case, we are done. In the former case, since S is completely branching, the set $\{\hat{\sigma} \mid \sigma \in Q\} \subseteq S$ is an antichain, where $\hat{\sigma}$ is the string obtained from σ by flipping its last bit. \dashv

LEMMA 3.2 (RCA_0). *Let $S \subseteq 2^{<\mathbb{N}}$ be an infinite completely branching set. Then for every antichain $A \subseteq S$, for all but at most one $\sigma \in A$, the set $S_\sigma := \{\tau \in S \mid \sigma \perp \tau \text{ and } |\tau| > |\sigma|\}$ is infinite and completely branching.*

PROOF. First, since $S \subseteq 2^{<\mathbb{N}}$ completely branching, then for every $\sigma \in 2^{<\mathbb{N}}$, the set S_σ is completely branching. Suppose for the sake of contradiction that there exists two strings $\sigma, \rho \in A$ such that S_σ and S_ρ are both finite. Then pick any $\tau \in S - (S_\sigma \cup S_\rho)$ with $|\tau| > \max(|\sigma|, |\rho|)$. It follows that $\sigma \prec \tau$ and $\rho \prec \tau$, and thus that σ and ρ are comparable, contradicting the fact that A is an antichain. \dashv

PROPOSITION 3.3. *The measure of the oracles computing a solution for a computable instance of SAC is 1.*

Remark 3.4. The proof of the above proposition is carried out purely as a computability statement, hence we have access to as much induction as needed.

PROOF. Let $S \subseteq 2^{<\mathbb{N}}$ be an infinite computably branching set. We are going to build a decreasing sequence of infinite completely branching sets of strings $S_0 \supseteq S_1 \supseteq \dots$, with $S_0 := S$, together with finite antichains $A_i \subseteq S_i$ (for $i \in \mathbb{N}$), in order to have an infinite antichain $A := \{\sigma_i \mid i \in \mathbb{N}\}$ where $\sigma_i \in A_i$.

This construction will work with positive probability, and since the class of oracles computing a solution to the instance S is invariant under Turing equivalence, this implies that this class has measure 1. Indeed, by Kolmogorov's 0-1 law, every measurable Turing-invariant class has either measure 0 or 1.

First, let $S_0 := S$. At step k , assume the sets $S_0 \supseteq S_1 \supseteq \dots \supseteq S_k$ and A_0, \dots, A_{k-1} have been defined, as well as the finite antichain $\{\sigma_0, \dots, \sigma_{k-1}\}$, such that $\forall \tau \in S_k, \forall i < k, \sigma_i \perp \tau$.

Search computably for a finite antichain $A_k \subseteq S_k$ of size 2^{k+2} . If found, pick an element $\sigma_k \in A_k$ at random. Then define $S_{k+1} := \{\tau \in S_k \mid \sigma_k \perp \tau \text{ and } |\tau| > |\sigma_k|\}$ for the next step.

If the procedure never stops, it yields an infinite antichain $A := \{\sigma_i \mid i \in \mathbb{N}\}$ thanks to the definition of the sets $(S_i)_{i < k}$. Assuming that S_k is an infinite completely branching set, Lemma 3.1 ensures that A_k will be found.

However, if at any point, S_k is not an infinite completely branching set, then at some point t we will not be able to find a large enough A_t in it. If this happens, since S_{k+1} is completely determined by S_k and σ_k , it means that we have chosen some "bad" $\sigma_k \in A_k$. Luckily, by Lemma 3.2, there is at most one element of this kind in A_k . Thus, if we pick σ_k at random in A_k , we have at most $\frac{1}{|A_k|} = \frac{1}{2^{k+2}}$ chances for this case to happen. So the overall probability that this procedure fails is less than $\sum_{k \geq 0} \frac{1}{2^{k+2}} = \frac{1}{2}$. Hence we found an antichain with positive probability. \dashv

Very few theorems studied in reverse mathematics admit a probabilistic proof. Proposition 3.3 provides a powerful method for separating the

statement **CAC for trees** from many theorems in reverse mathematics. In what follows, **AMT** stands for the Atomic Model Theorem, studied by Hirschfeldt, Shore and Slaman [14], **COH** is the cohesiveness principle, defined by Cholak, Jockusch and Slaman [4, Statement 7.7], and **RWKL** is the Ramsey-type Weak König’s lemma, defined by Flood [10, Statement 2] under the name **RKL**.

COROLLARY 3.5. *Over RCA_0 , **CAC for trees** implies none of **AMT**, **COH** and **RWKL**.*

PROOF. These three statements have a computable instance such that the measure of the oracles computing a solution is 0, see Astor et al. [1]. \dashv

The argument of Proposition 3.3 can be formalized over RCA_0 to yield the following result.

DEFINITION 3.6 (2-RAN). For every sequence of uniformly Π_2^0 binary trees T_0, T_1, \dots such that, for every n , $\mu([T_n]) > 1 - 2^{-n}$, there is some n and some set X such that $X \in [T_n]$.

PROPOSITION 3.7. $\text{RCA}_0 \vdash \text{2-RAN} \implies \text{SAC}$.

PROOF. For every n , consider the construction of Proposition 3.3, where the antichain A_k is of size 2^{n+k+1} instead of 2^{k+2} . For each k , let $\sigma_k \in A_k$ be the unique “bad” choice (if it exists), that is, which makes the set S_{k+1} finite, and let τ_k be the string of length $n+k+1$ corresponding to the binary representation of the rank of σ_k in A_k for some fixed order on binary strings. Then one can compute σ_k from τ_k and the finite set A_k . Note that τ_k is undefined when σ_k does not exist.

Consider the Σ_2^0 class $\mathcal{U}_n := \{X \in 2^{\mathbb{N}} \mid \exists k, \tau_k \prec X\} = \bigcup_k [\tau_k]$. It verifies

$$\mu(\mathcal{U}_n) \leq \sum_{\substack{k \geq 0 \\ \sigma_k \text{ exists}}} \mu([\tau_k]) \leq \sum_{k \geq 0} \frac{1}{2^{n+k+1}} = 2^{-n}$$

Let T_n be a Π_2^0 tree such that $[T_n] = 2^{\mathbb{N}} - \mathcal{U}_n$. We can now consider the sequence of trees $(T_n)_{n \in \mathbb{N}}$. By 2-RAN, there is some n and some $X \in [T_n]$. For any instance of **SAC**, find a solution by running the construction given in Proposition 3.3 with the help of X to avoid the potential “bad” choice in each A_k . \dashv

COROLLARY 3.8. *Over RCA_0 , **SAC** (and therefore **TAC**) implies none of $\text{B}\Sigma_2^0$ and **CAC for trees**.*

PROOF. Slaman [unpublished] proved that 2-RAN does not imply $\text{B}\Sigma_2^0$ over RCA_0 ; The result can also be found in [2]. The corollary follows from $\text{RCA}_0 \vdash \text{SAC} \implies \text{TAC}$ (Lemma 2.12) and $\text{RCA}_0 \vdash \text{TAC} + \text{B}\Sigma_2^0 \iff \text{CAC for trees}$ (Theorem 2.5). \dashv

We are now going to refine Proposition 3.3 by proving that some variant of DNC is sufficient to compute a solution of SAC.

DEFINITION 3.9 (Diagonally non-computable function). A function $f : \mathbb{N} \rightarrow \mathbb{N}$ is **diagonally non-computable relative to X** (or $\text{DNC}(X)$) if for every e , $f(e) \neq \Phi_e^X(e)$. Whenever f is dominated by a function $h : \mathbb{N} \rightarrow \mathbb{N}$, then we say that f is $\text{DNC}_h(X)$. A Turing degree is $\text{DNC}_h(X)$ if it contains a $\text{DNC}_h(X)$ function.

The following lemma gives a much more convenient way to work with $\text{DNC}_h(X)$ functions.

LEMMA 3.10 (Folklore). *Let A, X be subsets of \mathbb{N} . The following are equivalent:*

- (1) *A is of degree $\text{DNC}_h(X)$ for some computable (primitive recursive) function $h : \mathbb{N} \rightarrow \mathbb{N}$.*
- (2) *A computes a function $g : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that*

$$\forall e, n, |W_e^X| \leq n \implies g(e, n) \notin W_e^X$$

and which is dominated by a computable function $b : \mathbb{N}^2 \rightarrow \mathbb{N}$, i.e.

$$\forall e, n, g(e, n) < b(e, n)$$

PROOF. (2) \implies (1). Let $i : \mathbb{N} \rightarrow \mathbb{N}$ be a computable (primitive recursive) function such that for any $e \in \mathbb{N}$ and $B \subseteq \mathbb{N}$ we have $\Phi_{i(e)}^B(x) \downarrow \iff x = \Phi_e^B(e)$. Thus

$$W_{i(e)}^B = \begin{cases} \{\Phi_e^B(e)\} & \text{if } e \in B' \\ \emptyset & \text{otherwise} \end{cases}$$

From there, define the A -computable function $f : \mathbb{N} \rightarrow \mathbb{N}$ by $f : e \mapsto g(i(e), 1)$. It is $\text{DNC}(X)$ because $g(i(e), 1) \notin W_{i(e)}^X$ since $|W_{i(e)}^X| \leq 1$. Moreover, f is dominated by the computable function $e \mapsto b(i(e), 1)$, because b computably dominates g .

(1) \implies (2). Let f be a $\text{DNC}_h(X)$ function computed by A . Given the pair e, n , we describe the process that defines $g(e, n)$.

Construction. For each $i < n$, we compute the code $u(e, i)$ of the X -computable function which, on any input, looks for the i^{th} element of W_e^X . If it finds such an element, then it interprets it as an n -tuple $\langle k_0, \dots, k_{n-1} \rangle$ and returns the value k_i . If it never finds such an element, then the function diverges. Finally we define $g : e, n \mapsto \langle f(u(e, 0)), \dots, f(u(e, n-1)) \rangle$

Verification. First, since f is dominated by h , and since the function $\langle -, \dots, - \rangle$ computing an n -tuple is increasing on each variable, we can dominate g with the computable function

$$b : e, n \mapsto \langle h(u(e, 0)), \dots, h(u(e, n-1)) \rangle$$

Now, by contradiction, suppose g does not satisfy (2), i.e. suppose there exists e, n such that $|W_e^X| \leq n$ but $g(e, n) \in W_e^X$. Because W_e^X has fewer than n elements, we can suppose $g(e, n)$ is the i^{th} one for a some $i < n$. Thus the function $\Phi_{u(e,i)}^X$ is constantly equal to k_i where $g(e, n) = \langle k_0, \dots, k_{n-1} \rangle$, in particular $\Phi_{u(e,i)}^X(u(e, i)) = k_i$. But we also have

$$g(e, n) = \langle f(u(e, 0)), \dots, f(u(e, n-1)) \rangle$$

implying $f(u(e, i)) = k_i = \Phi_{u(e,i)}^X(u(e, i))$, which is impossible as f is supposed to be $\text{DNC}_h(X)$. \dashv

We are now ready to prove the following proposition. Conidis [6] independently proved the same statement for TAC with a similar construction. Note that by the equivalence of TAC + $\text{B}\Sigma_2^0$ with CAC for trees, Conidis result implies Proposition 3.11.

PROPOSITION 3.11. *Let $S \subseteq \mathbb{N}^{<\mathbb{N}}$ be an instance of SAC. Every set X of degree $\text{DNC}_h(\emptyset')$, with h a computable function, computes a solution of S .*

Remark 3.12. Once again, as in the case of Proposition 3.3, the proof here is carried purely as a computability statement, we have access to as much induction as we need.

PROOF. First, since X is of degree DNC_h for a computable function h , by Lemma 3.10, it computes a function $g : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that $\forall e, n, |W_e^{\emptyset'}| \leq n \implies g(e, n) \notin W_e^{\emptyset'}$ and which is dominated by a computable function $b : \mathbb{N}^2 \rightarrow \mathbb{N}$.

The idea of this proof is the same as in Proposition 3.3, but this time we are going to use g to avoid selecting the potential “bad” element in each finite antichain, i.e. the element which is incompatible with only finitely many strings. For any finite set $A \subseteq S$, let $\psi_A : \mathbb{N} \rightarrow S$ be a bijection such that $\psi_A(\llbracket 0, |A| \rrbracket) = A$.

The procedure is the following. Initially, $S_0 := S$. At step k , assume $S_k \subseteq S$ has been defined. To find the desired antichain A_k we use the fixed point theorem to find an index e_k such that $\Phi_{e_k}^{\emptyset'}(n)$ is the procedure that halts if it finds an antichain $A \subseteq S_k$ whose size is greater than $b(e_k, 1)$ and $\psi_A(n) \in A$, and finds (using \emptyset') an integer m such that $\forall \ell > m, \psi_A(\ell) \succ \psi_A(n)$.

Define $A_k := A$. By choice of A and e_k ,

$$W_{e_k}^{\emptyset'} = \begin{cases} \{\psi_A^{-1}(\rho)\} & \text{if } A_k \text{ has a bad element } \rho \\ \emptyset & \text{otherwise} \end{cases}$$

Finally we can define $\sigma_k := \psi_A(g(e_k, 1))$. Indeed since $|W_{e_k}^{\emptyset'}| \leq 1$ by construction, $g(e_k, 1) \notin W_{e_k}^{\emptyset'}$. Moreover $\sigma_k \in A_k$, because $g(e_k, 1) < b(e_k, 1) < |A_k|$. This implies that σ_k is not a bad element of A_k , in other words the set $S_{k+1} := \{\tau \in S_k \mid \tau \perp \sigma_k \text{ and } |\tau| > |\sigma_k|\}$ is infinite. \dashv

§4. ADS and EM. Ramsey's theorem for pairs admits a famous decomposition into the Ascending Descending Sequence theorem (ADS) and the Erdős-Moser theorem (EM) over RCA_0 . As mentioned in the introduction, both statements are strictly weaker than RT_2^2 . Actually, these statements are generally thought of as decomposing Ramsey's theorem for pairs into its disjunctiveness part with ADS, and its compactness part with EM. Indeed, the standard proof of ADS is disjunctive, and does not involve any notion of compactness, while the proof of EM is non-disjunctive and implies RWKL, which is the compactness part of RT_2^2 .

ADS and EM are relatively disjoint, in that they are only known to have the hyperimmunity principle as common consequence, which is a particularly weak principle. In this section however, we show that CAC for trees follows from both ADS and EM over RCA_0 . We shall see in Section 5 that CAC for trees implies the hyperimmunity principle.

The following proposition was proved by Dorais (personal communication) for SHER and independently by Conidis [6] for TAC. We adapted the proof of Dorais to obtain the following proposition.

PROPOSITION 4.1. $\text{RCA}_0 \vdash \text{ADS} \implies \text{CAC for trees and CAC for trees} \leq_c \text{ADS}$

PROOF. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be an infinite tree. Define the total order $<_0$ over T by $\sigma <_0 \tau \iff \sigma \prec \tau \vee (\sigma \perp \tau \wedge \sigma(d) <_{\mathbb{N}} \tau(d))$ where $d := \min\{k \in \mathbb{N} \mid \sigma(k) \neq \tau(k)\}$. By ADS, there is an infinite ascending or descending sequence (σ_i) for $(T, <_0)$.

If it is descending, there are two possibilities. Either $\forall^\infty i, \sigma_i \not\prec \sigma_{i+1}$, which means we eventually have an infinite \prec -decreasing sequence of strings, which is impossible. Or $\exists^\infty i, \sigma_i \perp \sigma_{i+1}$, in which case we designate by (h_k) the sequence $(\sigma_{\ell_k+1})_{k \in \mathbb{N}}$ of all such σ_{i+1} , and we show that it is an antichain of T .

To do so, it suffices to prove by induction on m that $\forall m > 0, \forall k, h_k \not\prec h_{k+m}$. When $m = 1$, due to how (σ_i) is structured, we have $\forall k, h_k \not\prec \sigma_{\ell_k} \perp h_{k+1}$. We now consider h_k and $h_{k+(m+1)}$. By induction hypothesis, $h_k \not\prec h_{k+m}$ and $h_{k+m} \not\prec h_{k+(m+1)}$. Moreover since $h_k >_0 h_{k+m} >_0 h_{k+(m+1)}$, we know there are minima d and e such that $h_k(d) > h_{k+m}(d)$ and $h_{k+m}(e) > h_{k+(m+1)}(e)$. Now $e \leq d$ implies $h_{k+(m+1)}(e) < h_{k+m}(e) = h_k(e)$, and $e > d$ implies $h_{k+(m+1)}(d) = h_{k+m}(d) < h_k(d)$; in any case $h_k \not\prec h_{k+(m+1)}$.

Now if the sequence (σ_i) is ascending, we again distinguish two possibilities. Either $\forall^\infty i, \sigma_i \not\prec \sigma_{i+1}$, which means we eventually obtain an infinite path of the tree. Or $\exists^\infty i, \sigma_i \perp \sigma_{i+1}$, in which case we work in the same fashion as in the descending case (designate by (h_k) the sequence $(\sigma_{\ell_k})_{k \in \mathbb{N}}$ of all such σ_i , and show by induction on m that $\forall m > 0, \forall k, h_k \not\prec h_{k+m}$). \dashv

PROPOSITION 4.2. $\text{RCA}_0 \vdash \text{EM} \implies \text{CAC for trees and CAC for trees} \leq_c \text{EM}$

PROOF. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be an infinite tree. We first define a computable bijection $\psi : \mathbb{N} \rightarrow T$. To do so, let $\varphi : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N}$ be the bijection $x_0 \cdot \dots \cdot x_{n-1} \mapsto p_0^{x_0} \times \dots \times p_{n-1}^{x_{n-1}} - 1$ where p_k is the k^{th} prime number. The elements of the sequence $(\varphi^{-1}(n))_{n \in \mathbb{N}}$ that are in T form a subsequence denoted $(s_n)_{n \in \mathbb{N}}$, and the function $\psi : \mathbb{N} \rightarrow T$ is defined by $n \mapsto s_n$.

Note that, by construction, the range of ψ is infinite and computable. Moreover, if $\sigma \prec \tau \in T$, then $\varphi(\sigma) < \varphi(\tau)$, hence $\psi^{-1}(\sigma) < \psi^{-1}(\tau)$. Also note that the range of ψ is not necessarily a tree.

Let $f : [\mathbb{N}]^2 \rightarrow 2$ be the coloring defined by $f(\{x, y\}) = 1$ iff $x <_{\mathbb{N}} y$ and $\psi(x) \prec \psi(y)$ coincide. By EM, there is an infinite transitive set $S \subseteq \mathbb{N}$, i.e. $\forall i < 2, \forall x < y < z \in S, f(x, y) = f(y, z) = i \implies f(x, z) = i$

Note that if there are $x < y \in S$ such that $f(x, y) = 0$, then $\forall z > y \in S, f(x, z) = 0$. Indeed given $x < y < z \in S$ such that $f(x, y) = 0$, either $f(y, z) = 0$, and so by transitivity we have $f(x, z) = 0$; or $f(y, z) = 1$, but in that case $f(x, z) \neq 1$ because it is impossible to simultaneously have $\psi(y) \prec \psi(z)$, $\psi(x) \prec \psi(z)$ and $\psi(x) \perp \psi(y)$.

Now two cases are possible. Either $\exists^\infty j \in \mathbb{N}, f(s_j, s_{j+1}) = 0$, so consider the infinite set A made of all such s_j . Thanks to the previous property, A is f -homogeneous for the color 0, and so $\psi(A)$ is an infinite antichain. Or $\forall^\infty j \in \mathbb{N}, f(s_j, s_{j+1}) = 1$, so there is a large enough $k \in \mathbb{N}$ such that $\psi(s_k) \prec \psi(s_{k+1}) \prec \dots$, i.e. we found an infinite path. \dashv

§5. TAC, lowness and hyperimmunity. Binns et al. in [3] and Conidis [6] respectively studied the reverse mathematics of CAC for trees and TAC. Since CAC for trees is computably equivalent to $\text{TAC} + \text{B}\Sigma_2^0$ and this equivalence also holds in reverse mathematics, the analysis of CAC for trees and TAC is very similar. For example, Binns et al. [3, Theorem 6.4] proved that for any fixed low set L , there is a computable instance of CAC for trees with no L -computable solution, while Conidis [6] proved the existence of a computable instance of TAC whose solutions are all of hyperimmune degree. In this section, we prove a general statement regarding TAC (Theorem 5.1) and show that it encompasses both results.

THEOREM 5.1. *Let (A_n) be an uniform sequence of infinite Δ_2^0 sets. There is a computable instance of TAC such that no A_n is a solution.*

PROOF. First, for any n , let e_n be the index of A_n , i.e. $\Phi_{e_n}^{\emptyset'} = A_n$. We also write $A_n[s] := \Phi_{e_n}^{\emptyset'[s]}[s]$.

Idea. We are going to construct a tree $T \subseteq 2^{<\mathbb{N}}$, such that for each $n \in \mathbb{N}$, there is $\sigma_n \in A_n$ verifying $\sigma_n \notin T$ or $\sigma_n \in T \wedge \forall^\infty \tau \in T, \sigma_n \prec \tau$. These

requirements are respectively denoted \mathcal{R}_n and \mathcal{S}_n , and A_n cannot be an infinite antichain of T if one of them is met.

The sequence (σ_n) is constructed via a movable marker procedure, with steps s and sub-steps $e < s$. At each step s we are going to manipulate an approximation σ_n^s of σ_n , and variables $\widehat{\sigma}_n^s$ that will help us keep track of which requirement is satisfied by σ_n^s .

Construction. At the beginning of each step s , let T_s be the approximation of the tree T defined by $T_s := T_{s-1} \cup \{\tau_s \cdot 0, \tau_s \cdot 1\}$ where τ_s is the leftmost (for example) leaf of T_{s-1} such that $\tau_s \succ \widehat{\sigma}_{s-1}^s$. For $s = 0$, we let $T_0 := \{\varepsilon\}$.

At step s , sub-step e , let σ_e^s be the string whose code is the smallest in the uniformly computable set $\{\tau \in A_e[s] \upharpoonright_s \mid (\tau \in T_s \wedge \tau \succ \widehat{\sigma}_{e-1}^s) \vee \tau \notin T_s\}$ with $\widehat{\sigma}_{-1}^s := \varepsilon$ and σ_e^s is undefined when the set is empty.

Besides, define $\widehat{\sigma}_e^s := \begin{cases} \sigma_e^s & \text{if } \sigma_e^s \in T_s \text{ (and therefore } \sigma_e^s \succ \widehat{\sigma}_{e-1}^s) \\ \widehat{\sigma}_{e-1}^s & \text{otherwise} \end{cases}$

Verification. By induction on e , we prove that $\sigma_e := \lim_s \sigma_e^s$ exists and is an element of A_e , also we prove $\widehat{\sigma}_e := \lim_s \widehat{\sigma}_e^s$ exists, and σ_e satisfies \mathcal{R}_e or \mathcal{S}_e .

Suppose we reached a step r such that for all $e' < e$ the values of $\sigma_{e'}^r$ and $\widehat{\sigma}_{e'}^r$ have stabilized. And thus, for any step $s > r$, as $\tau_s \succ \widehat{\sigma}_{s-1}^s \succ \widehat{\sigma}_{e-1}^s = \widehat{\sigma}_{e-1}$, the tree will always be extended with nodes above $\widehat{\sigma}_{e-1}$, implying only a finite part of the tree is not above $\widehat{\sigma}_{e-1}$.

Now suppose k is the smallest code of a string τ such that $(\tau \in T \wedge \tau \succ \widehat{\sigma}_{e-1}) \vee \tau \notin T$. Such a string exists because A_e is infinite, whereas the set of strings in T that are below $\widehat{\sigma}_{e-1}$ is not. If $\tau \in T$, then $\exists x, \forall y \succ x, \tau \in T_y$, otherwise define $x := 0$. Since A_e is Δ_2^0 , there exists $s \geq \max\{k+1, r, x\}$ such that $A_e[s] \upharpoonright_{k+1}$ has stabilized i.e. $\forall t > s, A_e[t] \upharpoonright_{k+1} = A_e[s] \upharpoonright_{k+1}$. Thus $\sigma_e^s = \tau$ because $\tau \in A_e \upharpoonright_{k+1} = A_e[s] \upharpoonright_{k+1} \subseteq A_e[s] \upharpoonright_s$. This ensures that for any $t > s$, $\sigma_e^t = \tau$, i.e. $\sigma_e = \tau$.

Finally, we distinguish two cases. Either $\sigma_e \in T$ and so $\exists t, \sigma_e^t \in T_t$, thus $\forall u > t, \widehat{\sigma}_e^u = \sigma_e^u$. So \mathcal{S}_e is satisfied, as cofinitely many nodes of T will be above $\widehat{\sigma}_e = \sigma_e$. Or $\sigma_e \notin T$, in which case, either $\forall t, \sigma_e^t \notin T_t$, implying $\forall t, \widehat{\sigma}_e^t := \widehat{\sigma}_{e-1}^t$ and thus \mathcal{R}_e is satisfied. \dashv

We now show how Theorem 5.1 relates to the result of Binns et al. in [3, Theorem 6.4], that is, the existence, for any fixed low set L , of a computable instance of CAC for trees with no L -computable solution.

LEMMA 5.2. *For any low set P , the sequence of infinite P -computable sets is uniformly Δ_2^0 .*

PROOF. Since $P' \leq_T \emptyset'$ we can \emptyset' -compute the function

$$f(e, x) = \begin{cases} \Phi_e^P(x) & \text{when } \forall y \leq x, \Phi_e^P(y) \downarrow \text{ and } \exists y > x, \Phi_e^P(y) \downarrow = 1 \\ 1 & \text{otherwise} \end{cases}$$

Now let $A_e := \{f(e, x) \mid x \in \mathbb{N}\}$. If Φ_e^P is total and infinite then A_e is equal to it, so it is P -computable. Otherwise A_e is cofinite, and in particular it is infinite and P -computable. \dashv

We are now ready to state the result of Binns et al. in [3, Theorem 6.4], but for TAC.

COROLLARY 5.3. *For any low set P , there exists a computable instance of TAC with no P -computable solution.*

PROOF. Given P , we can use Lemma 5.2 to obtain a uniform sequence, on which we apply Theorem 5.1. \dashv

The previous corollary is very useful to show that $\text{RCA}_0 + \text{WKL} \not\leq \text{TAC}$, since there exists a model of $\text{RCA}_0 + \text{WKL}$ below a low set. The following corollary will be useful to prove that the result of Binns et al. in [3, Theorem 6.4] implies the result of Conidis.

COROLLARY 5.4. *There exist a PA degree P and an instance of TAC with no P -computable solution.*

PROOF. It follows from the existence of a low PA degree by the low basis theorem, see [15, Corollary 2.2]. \dashv

The next proposition has two purposes. First, it will be used to show the existence of a computable instance of TAC whose solutions are all of hyperimmune degree (see Theorem 5.6). Second, it shows that, for any such instance, one can choose two specific functionals to witness this hyperimmunity, without loss of generality (see Corollary 5.8).

PROPOSITION 5.5. *Let T be an instance of TAC. For any set P of PA degree, if T has no P -computable solution, then for any solution $(\sigma_n)_{n \in \mathbb{N}}$, the function $t_{T, (\sigma_n)_{n \in \mathbb{N}}} : n \mapsto \min\{t \mid \sigma_n \in T[t]\}$ or $\ell_{(\sigma_n)_{n \in \mathbb{N}}} : n \mapsto |\sigma_n|$ is hyperimmune.*

PROOF. By contraposition, suppose there exists a solution $(\sigma_n)_{n \in \mathbb{N}}$ such that $t_{T, (\sigma_n)_{n \in \mathbb{N}}}$ and $\ell_{(\sigma_n)_{n \in \mathbb{N}}}$ are computably dominated by t and ℓ respectively. Then the set

$$\left\{ (\tau_n)_{n \in \mathbb{N}} \mid \begin{array}{l} (\tau_n)_{n \in \mathbb{N}} \text{ is an infinite antichain of } T \\ t_{T, (\tau_n)_{n \in \mathbb{N}}} \leq t \text{ and } \ell_{(\tau_n)_{n \in \mathbb{N}}} \leq \ell \end{array} \right\}$$

is a non-empty Π_1^0 class. It is non-empty because $(\sigma_n)_{n \in \mathbb{N}}$ belongs to it, and to show it is a Π_1^0 class, it can be written as

$$\left\{ (\tau_n)_{n \in \mathbb{N}} \left| \begin{array}{l} \forall m < n, \tau_n \perp \tau_m \\ \forall n, \tau_n \in T[t(n)] \\ |\tau_n| \leq \ell(n) \end{array} \right. \right\}$$

Thanks to ℓ , the number of elements at each level n of the tree associated to this class is computably bounded by $2^{\ell(n)}$, thus it can be coded by a Π_1^0 class of $2^{\mathbb{N}}$. Finally since P is of PA degree, it computes an element of any Π_1^0 class of the Cantor space, hence the result. \dashv

Combining Corollary 5.4 and Proposition 5.5, we obtain the following theorem from Conidis [6].

THEOREM 5.6 (Conidis [6]). *There is a computable instance of TAC such that each solution is of hyperimmune degree.*

PROOF. Let P be of low PA degree. By using Corollary 5.3 we get a computable instance T of TAC with no P -computable solution. Thus, by using Proposition 5.5 we deduce that, for any solution (σ_n) , its function $t_{T,(\sigma_n)_{n \in \mathbb{N}}}$ or $\ell_{(\sigma_n)_{n \in \mathbb{N}}}$ is hyperimmune. And $(\sigma_n)_{n \in \mathbb{N}}$ computes both, since T is computable ; meaning it is of hyperimmune degree. \dashv

COROLLARY 5.7. $\text{RCA}_0 \vdash \text{TAC} \implies \text{HYP}$

In his direct proof of Theorem 5.6, Conidis [6] constructed computable instance of TAC and two functionals Φ, Ψ such that for every solution H , either Φ^H or Ψ^H is hyperimmune. Interestingly, Proposition 5.5 can be used to show that Φ and Ψ can be chosen to be $t_{T,-}$ and ℓ_- , without loss of generality.

COROLLARY 5.8. *For any instance T of TAC whose solutions are all of hyperimmune degree, at least one of the function $t_{T,-}$ or ℓ_- is a witness.*

PROOF. Let T be an instance of TAC whose solutions are all of hyperimmune degree, and let $(\sigma_n)_{n \in \mathbb{N}}$ be such a solution. By contradiction, if we suppose $t_{T,(\sigma_n)_{n \in \mathbb{N}}}$ and $\ell_{(\sigma_n)_{n \in \mathbb{N}}}$ are both computably dominated, then by Proposition 5.5, T has a P -computable solution. If we choose P to be computably dominated, then it cannot compute a solution of hyperimmune degree, hence a contradiction. \dashv

Note that for every (computable or not) instance of TAC, there is a solution $(\sigma_n)_{n \in \mathbb{N}}$ such that $\ell_{(\sigma_n)_{n \in \mathbb{N}}}$ is dominated by the identity function, by picking any path, and building an antichain along it.

§6. SHER. We have seen in Section 4 that CAC for trees follows from both ADS and EM over RCA_0 . The proof of CAC for trees from ADS used only one specific property of the partial order $(T, <)$, that we shall refer to

as *semi-heredity*. Dorais and al. [8] introduced the principle SHER, which is the restriction of Ramsey's theorem for pairs to semi-hereditary colorings. In this section, we show that the seemingly artificial principle SHER turns out to be equivalent to the rather natural principle CAC for trees. This equivalence can be seen as more step towards the robustness of CAC for trees.

DEFINITION 6.1 (Semi-heredity). A coloring $f : [\mathbb{N}]^2 \rightarrow 2$ is **semi-hereditary** for the color $i < 2$ if

$$\forall x < y < z, f(x, z) = f(y, z) = i \implies f(x, y) = i$$

The name "semi-heredity" comes from the contraposition of the previous definition $\forall x < y < z, f(x, y) = 1 - i \implies f(x, z) = 1 - i \vee f(y, z) = 1 - i$

DEFINITION 6.2 (SHER, [8]). For any semi-hereditary coloring f , there exists an infinite f -homogeneous set.

The first proposition consists essentially of noticing that, given a set of strings $T \subseteq \mathbb{N}^{<\mathbb{N}}$, the partial order (T, \prec) behaves like a semi-hereditary coloring. The whole technicality of the proposition comes from the definition of an injection $\psi : \mathbb{N} \rightarrow T$ with some desired properties.

PROPOSITION 6.3. $\text{RCA}_0 \vdash \text{SHER} \implies \text{CAC for c.e. trees and CAC for c.e. trees} \leq_c \text{SHER}$

PROOF. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be an infinite c.e. tree. First, let $\varphi : \mathbb{N}^{<\mathbb{N}} \rightarrow \mathbb{N}$ the bijection $x_0 \cdot \dots \cdot x_{n-1} \mapsto p_0^{x_0} \times \dots \times p_{n-1}^{x_{n-1}} - 1$ where p_k is the k^{th} prime number. Define $\psi : \mathbb{N} \rightarrow T$ by letting $\psi(n)$ be the least $\sigma \in T$ (in order of apparition) such that $\phi(\sigma)$ is bigger than $\phi(\psi(0)), \phi(\psi(1)), \dots, \phi(\psi(n-1))$. Note that, by construction, the range of ψ is infinite and computable. Moreover, if $\sigma \prec \tau$, then $\varphi(\sigma) < \varphi(\tau)$, hence $\psi^{-1}(\sigma) < \psi^{-1}(\tau)$. Also note that the range of ψ is not necessarily a tree.

Let $f : [\mathbb{N}]^2 \rightarrow 2$ be the coloring defined by $f(\{x, y\}) = 1$ iff $x \prec_{\mathbb{N}} y$ and $\psi(x) \prec \psi(y)$ coincide. Let us show that f is semi-hereditary for the color 1. Suppose we have $x < y < z$ and that $f(x, z) = f(y, z) = 1$, i.e. letting $\sigma := \psi(x), \tau := \psi(y), \rho := \psi(z)$ then we have $\sigma \prec \rho$ and $\tau \prec \rho$, thus either $\sigma \prec \tau$ or $\tau \prec \sigma$. But since $x < y$, i.e. $\psi^{-1}(\sigma) < \psi^{-1}(\tau)$, only $\sigma \prec \tau$ can hold due to the above note, meaning $f(x, y) = 1$.

By SHER applied to f , there is an infinite f -homogeneous set H . If it is homogeneous for the color 0, then the set $\psi(H)$ corresponds to an infinite antichain of T . Likewise, if it is homogeneous for the color 1, then the set $\psi(H)$ is an infinite path of T . \dashv

We now prove the converse of the previous proposition.

DEFINITION 6.4 (Weak homogeneity). Given a coloring $f : [\mathbb{N}]^2 \rightarrow k$, a set $A := \{a_0 < a_1 < \dots\} \subseteq \mathbb{N}$ is **weakly-homogeneous** for the color $i < k$ if $\forall j, f(a_j, a_{j+1}) = i$

Before proving Proposition 6.6, we need a technical lemma.

LEMMA 6.5 (RCA₀). *Let $f : [\mathbb{N}]^2 \rightarrow 2$ be a semi-hereditary coloring for the color $i < 2$. For every infinite set $A := \{a_0 < a_1 < \dots\}$ which is weakly-homogeneous for the color i , there is an infinite f -homogeneous subset $B \subseteq A$.*

PROOF (DORAIS). We first show that any a_j falls in one of these two categories:

1. $\forall k > j, f(a_j, a_k) = i$
2. $\exists \ell > j, (\forall k \in \llbracket j, \ell \rrbracket, f(a_j, a_k) = i \wedge \forall k \geq \ell, f(a_j, a_k) = 1 - i)$

Indeed, for any $\ell > j$ such that $f(a_j, a_\ell) = i$, by semi-heredity, $f(a_j, a_{\ell-1}) = i$. So with a finite induction we get $\forall k \in \llbracket j, \ell \rrbracket, f(a_j, a_k) = i$.

There are now two possibilities. Either there are finitely many a_j of type 2, and so by removing these elements, the resulting set is f -homogeneous for the color i . Otherwise there are infinitely many a_j of type 2, in which case one can define an infinite f -homogeneous subset for color $1 - i$ using $\mathbf{B}\Sigma_1^0(A)$ as follows: due to the observation above, “ a_j is of type 2” is equivalent to the $\Sigma_1^0(A)$ formula $\exists \ell > j, f(a_j, a_\ell) = 1 - i$. Thus, given a finite set of type 2 elements $\{a_{j_0}, \dots, a_{j_{k-1}}\}$, by $\mathbf{B}\Sigma_1^0(A)$ there is $b > \max\{j_0, \dots, j_{k-1}\}$, and so j_k is defined as the smallest integer j_k such that $j_k \geq b$ and $f(a_{j_{k-1}}, a_{j_k}) = 1 - i$.

⊣

PROPOSITION 6.6. $\text{RCA}_0 \vdash \text{CAC for trees} \implies \text{SHER and}$
 $\text{SHER} \leq_c \text{CAC for trees}$

PROOF. Let $f : [\mathbb{N}]^2 \rightarrow 2$ be a semi-hereditary coloring for the color $i < 2$. We begin by constructing a tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$ defined as $T := \{\sigma_n \mid n \in \mathbb{N}\}$, where σ_n is the unique string which is:

1. strictly increasing (as a function), with last element n
2. weak-homogeneous for the color i , i.e. $\forall k < |\sigma_n| - 1, f(\sigma_n(k), \sigma_n(k + 1)) = i$
3. maximal as a weak-homogeneous set, i.e. $\forall y < \sigma_n(0), f(y, \sigma_n(0)) = 1 - i$ and $\forall j < |\sigma_n| - 1, \forall y \in \llbracket \sigma_n(j), \sigma_n(j + 1) \rrbracket, f(\sigma_n(j), y) = 1 - i \vee f(y, \sigma_n(j + 1)) = 1 - i$

To ensure existence, unicity and that T is a tree, we prove σ_n is obtained via the following effective procedure. Start with the string n . If the string $s_0 \cdot \dots \cdot s_m$ has been constructed, then look for the biggest integer $j < s_0$ such that $f(j, s_0) = i$. If there is none, the process ends. Else, the process is repeated with the string $j \cdot s_0 \cdot \dots \cdot s_m$.

The string obtained is maximal by construction. It is unique, because at each step, if there are two (or more) integers $j_0 < j_1$ smaller than s_0 and such that $f(j_0, s_0) = f(j_1, s_0) = i$, then by semi-heredity we have $f(j_0, j_1) =$

1. This means we will eventually add j_0 after j_1 . In particular, the string contains all the $j < n$ such that $f(j, n) = i$. Moreover this shows that T is a tree, since the procedure is the same at any point during construction.

Now we can apply CAC for trees to T , leading to two possibilities. Either there is an infinite path, which is a weakly-homogeneous set for the color i thanks to condition 2. And so apply Lemma 6.5 to obtain a f -homogeneous set for the color i .

Or there is an infinite antichain, which is of the form $(\sigma_{n_j})_{j \in \mathbb{N}}$. Let us show the set $H := \{n_j \mid j \in \mathbb{N}\}$ is f -homogeneous for the color $1-i$. Indeed, if $f(n_s, n_t) = i$ for some $s < t$, then $n_s \in \sigma_{n_t}$, since σ_{n_t} contains all the elements $y < n_t$ such that $f(y, n_t) = i$. But then $\sigma_{n_s} \prec \sigma_{n_t}$, contradicting the fact that $(\sigma_{n_j})_{j \in \mathbb{N}}$ is an antichain. \dashv

We end this section by studying RT_2^2 with respect to 3-variables forbidden patterns. As explained in the introduction, there are three basic 3-variables forbidden patterns, yielding the notions of semi-heredity, semi-ancestry and semi-transitivity, respectively. These forbidden patterns induce Ramsey-like statements of the form “for any 2-coloring of pairs, there exists an infinite set which avoids some kind of forbidden patterns”. This statement applied to semi-transitivity yields a consequence of the Erdős-Moser theorem, known to be strictly weaker than Ramsey’s theorem for pairs over RCA_0 . We now show that the two remaining forbidden patterns yield statements equivalent to RT_2^2 . This completes the picture of the reverse mathematics of Ramsey-like theorems for 3-variable forbidden patterns.

DEFINITION 6.7 (Semi-ancestry). A coloring $f : [\mathbb{N}]^2 \rightarrow 2$ has **semi-ancestry** for the color $i < 2$ if

$$\forall x < y < z, f(x, y) = f(x, z) = i \implies f(y, z) = i$$

Before proving RT_2^2 from the Ramsey-like statement about semi-ancestry over RCA_0 , we need to prove that this statement implies $\text{BS}\Sigma_2^0$. This is done by proving the following principle.

DEFINITION 6.8 (D_2^2). Every Δ_2^0 set admits an infinite subset in it or its complement.

PROPOSITION 6.9. *The statement “for any 2-coloring of pairs, there exists an infinite set which has semi-ancestry for some color” implies D_2^2 over RCA_0 .*

PROOF. Let A be a Δ_2^0 set whose approximations are $(A_t)_{t \in \mathbb{N}}$. We define the coloring $f(x, y) := \chi_{A_y}(x)$, and use the statement of the proposition to obtain an infinite set B that has semi-ancestry for some color.

If B has semi-ancestry for the color 1, then $\forall x < y < z \in B, x \in A_y \wedge x \in A_z \implies y \in A_z$. Now either $B \subseteq \overline{A}$ and we are done. Or $\exists x \in A \cap B$, which

means $\forall^\infty y \in B, x \in A_y$, implying that $\forall^\infty y > x \in B, \forall z > y \in B, y \in A_z$ by semi-ancestry, i.e. $\forall^\infty y > x \in B, y \in A$. So we can compute a subset H of B which is infinite and such that $H \subseteq A$.

This argument also works when B has semi-ancestry for the color 0, we just need to switch A and \overline{A} , as well as \in and \notin , when needed. \dashv

COROLLARY 6.10. *The statement “for any 2-coloring of pairs, there exists an infinite set which has semi-ancestry for some color” implies $\mathbf{B}\Sigma_2^0$ over \mathbf{RCA}_0 .*

PROOF. Immediate, since $\mathbf{RCA}_0 \vdash \mathbf{D}_2^2 \implies \mathbf{B}\Sigma_2^0$, see [5, Theorem 1.4]. \dashv

PROPOSITION 6.11. *The statement “for any 2-coloring of pairs, there exists an infinite set which has semi-ancestry for some color” implies \mathbf{RT}_2^2 over \mathbf{RCA}_0 and over the computable reduction.*

PROOF. Let $f : [\mathbb{N}]^2 \rightarrow 2$ be a coloring. We can apply the statement to obtain an infinite set A which has semi-ancestry for the color i , i.e. $\forall x < y < z \in A, f(x, y) = i \wedge f(x, z) = i \implies f(y, z) = i$. There are two possibilities. Either there exists $a \in A$ such that $\exists^\infty b > a \in A, f(a, b) = i$, in which case all such elements b form an infinite f -homogeneous set due to the property of A . Otherwise any $a \in A$ verifies $\forall^\infty b > a, f(a, b) = 1 - i$, i.e. all the elements of A have a limit color equal to $1 - i$ for the coloring $f \upharpoonright_{[A]^2}$. Thus we can use $\mathbf{B}\Sigma_2^0$ (Corollary 6.10) to compute an infinite homogeneous set (see [9, Proposition 6.2]). \dashv

The proof that the Ramsey-like statement about semi-heredity implies Ramsey’s theorem for pairs is indirect, and uses the Ascending Descending Sequence principle.

PROPOSITION 6.12. *The statement “for any 2-coloring of pairs, there exists an infinite set which is semi-hereditary for some color” implies \mathbf{ADS} over \mathbf{RCA}_0 and over the computable reduction.*

PROOF. Let $\mathcal{L} = (\mathbb{N}, <_{\mathcal{L}})$ be a linear order. Let $f : [\mathbb{N}]^2 \rightarrow 2$ be the coloring defined by $f(\{x, y\}) = 1$ iff $<_{\mathcal{L}}$ and $<_{\mathbb{N}}$ coincide on $\{x, y\}$. By the statement of the proposition, there is an infinite set H on which the coloring is semi-hereditary for some color i .

Before continuing, note that if there is a pair $x < y \in H$ such that $f(x, y) = 1 - i$ then $\forall z > y \in H, f(x, z) = 1 - i$. Indeed, either $f(y, z) = 1 - i$, in which case by transitivity of $<_{\mathcal{L}}$ we have $f(x, z) = 1 - i$. Or $f(y, z) = i$, in which case by semi-heredity $f(x, z) = 1 - i$, because otherwise it would imply that $f(x, y) = i$.

Now if $\exists^\infty x \in H, \exists y > x \in H, f(x, y) = 1 - i$, then we construct an infinite decreasing sequence by induction. Suppose the sequence $x_0 >_{\mathcal{L}} \dots >_{\mathcal{L}} x_{n-1}$ has already be constructed, and we have $m \in H$ such that $f(x_{n-1}, m) = 1 - i$, then we look for a $\{x < y\}$ such that $x > m$ and $f(x, y) = 1 - i$. By

the previous remark we have that $f(x_{n-1}, x) = 1 - i$, i.e. $x_{n+1} >_{\mathcal{L}} x$, so we extend the sequence with x and redefine $m := y$.

Else $\forall^\infty x \in H, \forall y > x \in H, f(x, y) = i$, and so getting rid of finitely many elements yields an infinite increasing sequence. \dashv

COROLLARY 6.13. *The statement “for any 2-coloring of pairs, there exists an infinite set which is semi-hereditary for some color” implies RT_2^2 over RCA_0 .*

PROOF. This comes from the fact that $\text{RT}_2^2 \iff S + \text{SHER}$ by definition (with S denoting the statement in question). And we have $\text{RCA}_0 \vdash S \implies \text{SHER}$ by using Proposition 6.12, Proposition 4.1 and Proposition 6.6. \dashv

Remark 6.14. Let S denote the statement “for any 2-coloring of pairs, there exists an infinite set which is semi-hereditary for some color”. The proof that S implies RT_2^2 over RCA_0 involves two applications of S . The first one to obtain an infinite set over which the coloring is semi-hereditary, and a second one to solve **SHER** using the fact that S implies **ADS**, which itself implies **SHER**. It is unknown whether RT_2^2 is computably reducible to S .

§7. Stable counterparts: SADS and CAC for stable c.e. trees. Cholak, Jockusch and Slaman [4] made significant progress in the understanding of Ramsey’s theorem for pairs by dividing the statement into a stable and a cohesive part.

DEFINITION 7.1. A coloring $f : [\mathbb{N}]^2 \rightarrow k$ is **stable** if for every $x \in \mathbb{N}$, $\lim_y f(x, y)$ exists. A linear order $\mathcal{L} = (\mathbb{N}, <_{\mathcal{L}})$ is **stable** if it is of order type $\omega + \omega^*$.

We call SRT_k^2 and **SADS** the restriction of RT_k^2 and **ADS** to stable colorings and stable linear orders, respectively. Given a linear order $\mathcal{L} = (\mathbb{N}, <_{\mathcal{L}})$, the coloring corresponding to the order is stable iff the linear order is of order type $\omega + \omega^*$, or $\omega + k$ or $k + \omega^*$. In particular, SRT_2^2 implies **SADS** over RCA_0 .

In this section, we study the stable counterparts of **CAC for trees** and **SHER**, and prove that they are equivalent over RCA_0 . We show **SADS** implies **CAC for stable c.e. trees** over RCA_0 . It follows in particular that every computable instance of **CAC for stable c.e. trees** admits a low solution.

DEFINITION 7.2 (Stable tree, Dorais [7]). A tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$ is **stable** when for every $\sigma \in T$ either $\forall^\infty \tau \in T, \sigma \perp \tau$ or $\forall^\infty \tau \in T, \sigma \not\perp \tau$

Note that any stable finitely branching tree admits a unique path.

PROPOSITION 7.3. $\text{RCA}_0 \vdash \text{SADS} \implies \text{CAC for stable c.e. trees}$

Remark 7.4. In the proof below, we use the statement that $\text{RCA}_0 \vdash \text{SADS} \implies \text{B}\Sigma_2^0$. This is because $\text{RCA}_0 \vdash \text{SADS} \implies \text{PART}$ ([13, Proposition 4.6]) and $\text{RCA}_0 \vdash \text{PART} \iff \text{B}\Sigma_2^0$ ([5, Theorem 1.2]).

PROOF. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be an infinite c.e. tree which is stable.

Consider the total order $<_0$, defined by $\sigma <_0 \tau \iff \sigma \prec \tau \vee (\sigma \perp \tau \wedge \sigma(d) < \tau(d))$ where $d := \min\{y \mid \sigma(y) \neq \tau(y)\}$. We show that it is of type $\omega + \omega^*$, i.e.

$$\forall \sigma \in T, (\forall^\infty \tau \in T \sigma <_0 \tau) \vee (\forall^\infty \tau \in T, \tau <_0 \sigma)$$

So let $\sigma \in T$, there are two possibilities. Either $\forall^\infty \tau \in T, \sigma \not\prec \tau$, meaning we even have $\forall^\infty \tau \in T, \sigma \prec \tau$, which directly implies $\forall^\infty \tau \in T, \sigma <_0 \tau$.

Or $\forall^\infty \tau \in T, \sigma \perp \tau$. In this case, we consider all the nodes τ successors of a prefix of σ but not prefix of σ , there are finitely many of them, because there are finitely many prefixes of σ and no infinitely-branching node (WLOG, as otherwise there would be a computable infinite antichain). So we can apply the pigeon-hole principle, by using $\text{B}\Sigma_2^0$, to deduce that there is a certain τ which has infinitely many successors.

Moreover, by stability of T there cannot be another such node. Indeed, by contradiction, if there were two such nodes τ and τ' , then we would have that $\exists^\infty \eta \in T, \eta \perp \tau$, because the successors of τ' are incomparable to τ . And since τ already verifies $\exists^\infty \eta \in T, \eta \succ \tau$, contradicting the stability of T .

Therefore we have that $\forall^\infty \eta \in T, \eta \succ \tau$, and so depending on whether $\tau <_0 \sigma$ or $\sigma <_0 \tau$, we obtain that $\forall^\infty \eta \in T, \eta <_0 \sigma$ or $\forall \eta \in T, \sigma <_0 \eta$ respectively. From there we can apply SADS and the proof is exactly like in Proposition 4.1. \dashv

COROLLARY 7.5. CAC for stable c.e. trees admits low solutions.

PROOF. This comes from the fact that any instance of SADS has a low solution, as proven in [13, Theorem 2.11]. \dashv

The proof that SHER follows from CAC for stable trees will use $\text{B}\Sigma_2^0$. We therefore first prove that CAC for stable trees implies $\text{B}\Sigma_2^0$ over RCA_0 .

LEMMA 7.6. $\text{RCA}_0 \vdash \text{CAC for stable trees} \implies \forall k, \text{RT}_k^1$

PROOF. Let $f : \mathbb{N} \rightarrow k$ be a coloring. There are two possibilities: Either $\exists i < k, \exists^\infty x, f(x) = i$, in which case there is an infinite computable f -homogeneous set. Otherwise $\forall i < k, \forall^\infty x, f(x) \neq i$, in which case we consider the infinite tree

$$T := \{\sigma \in \text{Inc} \mid \forall y \leq \max \sigma, f(y) = f(\max \sigma) \implies y \in \text{ran} \sigma\}$$

where Inc is the set of strictly increasing strings of $\mathbb{N}^{<\mathbb{N}}$.

By hypothesis, for every node $\sigma \in T, \forall^\infty \tau \in T, \sigma \perp \tau$, otherwise T would have an infinite T -computable path. Thus, T is stable. Moreover, every

antichain is of size at most k , thus T is a stable tree with no infinite path and no infinite antichain, contradicting CAC for stable trees. \dashv

PROPOSITION 7.7. *Under RCA_0 the statement CAC for stable trees implies SHER for stable colorings*

PROOF. Let $f : [\mathbb{N}]^2 \rightarrow 2$ be a stable coloring, semi-hereditary for the color i . We distinguish two cases. Either there are finitely many integers with limit color i , meaning we can ignore them and use $\text{B}\Sigma_2^0$ (Lemma 7.6) to compute an infinite homogeneous set (see [9, Proposition 6.2]). Otherwise there are infinitely many integers whose limit color is i , in which case we use the same proof as in Proposition 6.6, but we must prove the tree T we construct is stable. So let σ_n be a node of this tree.

Suppose first n has limit color i . Let p be sufficiently large so that $f(n, p) = i$. As explained in Proposition 6.6, σ_p contains all the integers $m < p$ such that $f(m, p) = i$. It follows that $n \in \sigma_p$. Moreover, if $n \in \sigma_p$, then $\sigma_n \preceq \sigma_p$. Thus, $\forall^\infty p, \sigma_n \prec \sigma_p$.

Suppose now n has limit color $1 - i$, then since there are infinitely many integers with limit color i , there is one such that $p > n$. In particular, σ_p verifies $\forall^\infty \tau \in T, \sigma_p \prec \tau$. Thus, if $\sigma_n \prec \sigma_p$ then $\forall^\infty \tau \in T, \sigma_n \prec \tau$, and if $\sigma_n \perp \sigma_p$ then $\forall^\infty \tau \in T, \sigma_n \perp \tau$. \dashv


PROPOSITION 7.8. *Under RCA_0 the statement SHER for stable colorings implies CAC for stable c.e. trees*

PROOF. Let $T \subseteq \mathbb{N}^{<\mathbb{N}}$ be an infinite stable c.e. tree. The proof is the same as in Proposition 6.3, but we must verify that the coloring $f : [\mathbb{N}]^2 \rightarrow 2$ defined is stable. Given $x \in \mathbb{N}$, we claim that $\exists i < 2, \forall^\infty y f(x, y) = i$. Since T is stable, either $\forall^\infty y, \psi(x) \not\prec \psi(y)$ or $\forall^\infty y, \psi(x) \perp \psi(y)$ holds. In the first case $\forall^\infty y, f(x, y) = 1$, and in the second one $\forall^\infty y, f(x, y) = 0$. Thus the coloring is stable, and the proof can be carried on. \dashv

COROLLARY 7.9. *The following are equivalent over RCA_0 :*

- (1) CAC for stable trees
- (2) CAC for stable c.e. trees
- (3) SHER for stable colorings

§8. Conclusion. The following figure sums up the implications proved in this paper. All implications hold both in RCA_0 and over the computable reduction.



We have established in Theorem 2.5 the equivalence between $TAC + B\Sigma_2^0$ and other statements.

Question 8.1. What is the first-order part of TAC ?

Recall that, by Corollary 5.3, for every fixed low set X , there is a computable instance of TAC with no X -computable solution. By computable equivalence, this property also holds for CAC for trees. It is however unknown whether Corollary 5.3 can be improved to defeat all low sets simultaneously.

Question 8.2. Does every computable instance of CAC for trees admit a low solution?

Note that by Corollary 7.5, any negative witness to the previous question would yield a non-stable tree.

We have also seen by Proposition 3.11 that for every computable instance T of CAC for trees, every computably bounded DNC function relative to \emptyset' computes a solution to T . The natural question would be whether we can improve this result to any DNC function relative to \emptyset' .

Question 8.3. Is there some X such that for every computable instance T of CAC for trees, every DNC function relative to X computes a solution to T ?

Note that in the case of a computable set X , the answer is negative, as there exist DNC functions of low degree.

Finally, recall from Remark 6.14 the following question:

Question 8.4. Is RT_2^2 computably reducible to the statement “for any 2-coloring of pairs, there exists an infinite set which is semi-hereditary for some color”?

REFERENCES

- [1] ERIC P. ASTOR and LAURENT BIENVENU and DAMIR DZHAFAROV and LUDOVIC PATEY and PAUL SHAFER and REED SOLOMON and LINDA BROWN WESTRICK, *The weakness of typicality*, In preparation.
- [2] D BELANGER, C CHONG, W LI, W WANG, and Y YANG, *Where pigeonhole principles meet König lemmas*, *Transactions of the AMS*, vol. 374 (2021), pp. 8275–8303.
- [3] STEPHEN BINNS, BJØRN KJOS-HANSEN, MANUEL LERMAN, JAMES H. SCHMERL, and REED SOLOMON, *Self-embeddings of computable trees*, (2014).
- [4] PETER A. CHOLAK, CARL G. JOCKUSCH, and THEODORE A. SLAMAN, *On the strength of Ramsey’s theorem for pairs*, *The Journal of Symbolic Logic*, vol. 66 (2001), no. 01, pp. 1–55.
- [5] C. CHONG, STEFFEN LEMPP, and YUE YANG, *On the role of the collection principle for Σ_2^0 -formulas in second-order reverse mathematics*, *Proceedings of the American Mathematical Society*, vol. 138 (2010), no. 3, pp. 1093–1100.
- [6] CHRIS J. CONIDIS, *Computability and combinatorial aspects of minimal prime ideals in noetherian rings*, Submitted.
- [7] FRANÇOIS G. DORAIS, *On a theorem of hajnal and surányi*, 2012.
- [8] FRANÇOIS G. DORAIS, DAMIR D. DZHAFAROV, JEFFRY L. HIRST, JOSEPH R. MILETI, and PAUL SHAFER, *On uniform relationships between combinatorial problems*, *Trans. Amer. Math. Soc.*, vol. 368 (2016), no. 2, pp. 1321–1359.
- [9] DAMIR D. DZHAFAROV, DENIS R. HIRSCHFELDT, and SARAH C. REITZES, *Reduction games, provability, and compactness*, 2020.
- [10] STEPHEN FLOOD, *Reverse mathematics and a Ramsey-type König’s lemma*, *The Journal of Symbolic Logic*, vol. 77 (2012), no. 4, pp. 1272–1280.
- [11] E. HERRMANN, *Infinite chains and antichains in computable partial orderings*, *J. Symbolic Logic*, vol. 66 (2001), no. 2, pp. 923–934.
- [12] DENIS R. HIRSCHFELDT, *Slicing the truth*, Lecture Notes Series. Institute for Mathematical Sciences. National University of Singapore, vol. 28, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2015, On the computable and reverse mathematics of combinatorial principles, Edited and with a foreword by Chitat Chong, Qi Feng, Theodore A. Slaman, W. Hugh Woodin and Yue Yang.
- [13] DENIS R. HIRSCHFELDT and RICHARD A. SHORE, *Combinatorial principles weaker than Ramsey’s theorem for pairs*, *The Journal of Symbolic Logic*, vol. 72 (2007), no. 1, pp. 171–206.
- [14] DENIS R. HIRSCHFELDT, RICHARD A. SHORE, and THEODORE A. SLAMAN, *The atomic model theorem and type omitting*, *Transactions of the American Mathematical Society*, vol. 361 (2009), no. 11, pp. 5805–5837.
- [15] CARL G. JOCKUSCH and ROBERT IRVING SOARE, π_1^0 classes and degrees of theories, *Transactions of the American Mathematical Society*, vol. 173 (1972), pp. 33–56.
- [16] MANUEL LERMAN, REED SOLOMON, and HENRY TOWNSNER, *Separating principles below Ramsey’s theorem for pairs*, *Journal of Mathematical Logic*, vol. 13 (2013), no. 02, p. 1350007.
- [17] LUDOVIC PATEY, *Ramsey-like theorems and moduli of computation*, 2019.
- [18] STEPHEN G. SIMPSON, *Subsystems of Second Order Arithmetic*, Cambridge University Press, 2009.

UNIV PARIS EST CRETEIL, LACL, F-94010 CRETEIL, FRANCE

E-mail: julien.cervelle@u-pec.fr

URL: <https://jc.lacl.fr>

UNIV PARIS EST CRETEIL, LACL, F-94010 CRETEIL, FRANCE

E-mail: william.gaudelier@gmail.com

CNRS, ÉQUIPE DE LOGIQUE

UNIVERSITÉ DE PARIS

PARIS, FRANCE

E-mail: ludovic.patey@computability.fr

URL: <http://ludovicpatey.com>