Spin alignment of vector mesons from quark dynamics in a rotating medium

Minghua Wei*

Institute of Modern Physics, Fudan University, Shanghai 200433, China

Mei Huang[†]

School of Nuclear Science and Technology, University of Chinese Academy of Sciences, Beijing 100049, China

(Dated: March 6, 2023)

Vorticities in heavy-ion collisions (HICs) are supposed to induce spin alignment and polarization phenomena of quarks and mesons. In this work, we analyze the spin alignment of vector mesons ϕ and ρ induced by rotation from quark dynamics in the framework of the Nambu-Jona-Lasinio (NJL) model. The rotating angular velocity induces mass splitting of spin components for vector ϕ, ρ mesons $M_{\phi,\rho}(\Omega) \simeq M_{\phi,\rho}(\Omega = 0) - s_z \Omega$. This behavior contributes to the spin alignment of vector mesons ϕ, ρ in an equilibrium medium and naturally explains the negative deviation of $\rho_{00} - 1/3$ for vector mesons. Incidentally, the positive deviation of $\rho_{00} - 1/3$ under the magnetic field can also be easily understood from quark dynamics.

I. INTRODUCTION

Relativistic heavy-ion collision experiments provide a platform for studying quantum chromodynamics (QCD) matter under extreme conditions. It has been expected that guark gluon plasma (QGP) can be created through heavy ion collisions (HICs) [1]. In a specific case, a noncentral collision, QGP is created with a large orbital angular momentum (OAM) at a range of $10^4 - 10^5\hbar$ [2–4]. Meanwhile, a strong magnetic field can reach the magnitude of $10m_{\pi}^2$ at the initial time of the HICs and evolves with time [5-7]. The magnitude of the magnetic field decays severely while the averaged angular velocity still maintains its magnitude along the axis which is perpendicular to the reaction plane [3]. Therefore, the effects of a rotating medium will play significant roles on QCD phase diagram, dilepton production rate and spin polarization.

Spin alignment of vector meson ϕ and K^{*0} has been one of the intriguing topics in HICs. Experimental evidence suggests that spin density matrix element ρ_{00} has a remarkable deviation from 1/3 [8, 9]. ALICE collaboration has measured ρ_{00} for K^{*0} and ϕ meson at $\sqrt{s_{NN}} = 2.76$ TeV in Pb-Pb collisions [8], and a negative deviation from 1/3 at lower transverse momentum has been reported. STAR collaboration had measured ρ_{00} between $\sqrt{s_{NN}} = 11.5$ and 200 GeV in Au-Au collisions [9]. In this collision energy range, a positive deviation form 1/3 is reported for ρ_{00} of vector meson ϕ , which can be explained by a ϕ -meson field [10].

Both the magnetic field and vorticities are expected to contribute to the spin alignment. In Ref.[11], the effect of magnetic field on spin alignment has been studied. In this work, we introduce the effects of global rotation on quark matter. Generally, the rotation has an inhibition effect on quark and anti-quark pairing [12], which is different from Magnetic Catalysis (MC) [13–16]. As a consequence, chiral condensate will disappear as the angular velocity grows. Since the chiral condensate is an order parameter for chrial phase transition, its behaviors demonstrate that first order phase transition occurs at low temperature regions and crossover occurs in high temperature regions. The characteristics imply an analogy between the rotation and chemical potential [17, 18]. However, it is found that the rotation behaves beyond an effective chemical potential in dilepton production [19] and the rotation enhances the dilepton production rate and induces the ellipticity of lepton pairs.

It is also worth emphasizing that previous studies have assumed an infinite size without boundary conditions [3, 17, 20]. The boundary conditions on a finite size system should be taken into account for a rotating system[21–24], and inhomogeneous chiral condensation will be developed [25, 26]. For example, no-flux or MIT bag boundary conditions are applied in a spherical or cylindrical system [27–29]. The choice of boundary conditions will influence the behavior of chiral condensate near the surface. Particularly, when no-flux boundary condition is applied in a cylindrical system, chiral condensate will almost keep a constant in the inner part of the cylinder but get enhanced like a Gibbs phenomenon near the surface [21].

In QCD phase diagram studies, it is also possible to consider inhomogeneous rotation with boundary conditions. Chiral condensate is enhanced near the area where the angular velocity changes severely, and this phenomenon is called centrifugal effect [25]. Furthermore, chiral condensate reveals the dynamical mass of quarks. Besides the light quarks u and d, heavy flavor quarks, like s,c, are studied in electromagnetic and rotational fields [30]. Corresponding ϕ , D mesons are expected to

^{*} weiminghua@fudan.edu.cn

[†] huangmei@ucas.ac.cn

be influenced by rotation as well. However, it is complicated to describe the freeze-out of heavy particles from the QGP [31]. Therefore, we only consider light-flavor vector meson modes excited and annihilated in a rotating medium.

External fields, such as magnetic field and vortical field, will induce the mass splitting of spin-1 vector mesons. In recent years, much attention has been paid to the magnetic field effect of vector mesons, such as ρ , ϕ . For example, in case of charged ρ mesons, mass splitting behaves like:

$$M_{\rho^{\pm}}(eB) = M_{\rho^{\pm}}(0) \mp \kappa eB \tag{1}$$

where κ is a coefficient. For point-like particle, $\kappa = 1$; when quark-antiquark loop effect is considered, $\kappa \neq 1$. Similarly, ρ meson under rotation also exhibits a mass splitting phenomenon [32]:

$$M_{\rho}(\Omega) = M_{\rho}(0) - s_z \Omega \tag{2}$$

This relation indicates that vector mesons tend to occupy the $s_z = +1$ state. In this paper, a similar result can be obtained for vector meson ϕ .

Recently, the mass splitting and spin alignment of vector meson ϕ has been investigated under the magnetic field [11]. Another important contribution from the rotation on the spin alignment should also be taken into account. Therefore, in this work, we calculate the rotational contribution on matrix element ρ_{00} .

This paper is organized as follows. In section II, quantum field theory in a rotating frame is introduced. Based on it, the quark propagator and the self-energy are modified by uniform rotation. Consequently, masses of vector mesons are obtained by random phase approximation. In section III, the numerical result of quark mass, meson mass and spin alignment will be presented. Finally, the conclusion and summary will be given in section IV.

II. FORMALISM

A. Quantum field theory in a uniformly rotating frame

In quantum field theory, a Lorentz vector field $V^a(x)$ can be described by tetard $e_a(x) = e_a^{\mu}(x) \frac{\partial}{\partial x^{\mu}}$ in curved space-time. In this section, Latin letters, a, b = 0, 1, 2, 3, stand for Lorentz indices. And Greek letters, $\mu, \nu = 0, 1, 2, 3$, stand for coordinate indices. Parallel transport of $V^a(x)$ acts as follows:

$$V^{a}(x+dx) = V^{a}(x) + \Gamma_{ab\mu}V^{b}dx^{\mu}, \qquad (3)$$

where $\Gamma_{ab\mu}$ is called spin connection. Metric compatibility ensures $\Gamma_{ab\mu} = -\Gamma_{ba\mu}$, and non-torsion condition provides a relation between the tetard and the spin connection:

$$\Gamma^{ab}_{\mu} = \frac{1}{2} \begin{bmatrix} e^{a\nu} \left(\partial_{\mu} e^{b}_{\nu} - \partial_{\nu} e^{b}_{\mu} \right) - e^{b\mu} \left(\partial_{\mu} e^{a}_{\nu} - \partial_{\nu} e^{a}_{\mu} \right) \\ -e^{a\rho} e^{bl} \left(\partial_{\rho} e_{c\sigma} - \partial_{\sigma} e_{c\rho} \right) e^{c}_{\mu} \end{bmatrix}.$$

$$\tag{4}$$

For spinor field, corresponding spinor connection is given by $\Gamma_{\mu} = \sigma^{ab}\Gamma_{ab\mu}$, where σ^{ab} is spinor representation of Lorentz group. In the co-moving frame of the QGP, free fermions are described by modified Dirac equation:

$$[i\bar{\gamma}^{\mu}(\partial_{\mu}+\Gamma_{\mu})-M_{f}]\psi=0, \qquad (5)$$

where M_f is the mass of a fermion and $\bar{\gamma}^{\mu} = e^{\mu}_a \gamma^a$ satisfies $\{\bar{\gamma}_{\mu}, \bar{\gamma}_{\nu}\} = g^{\mu\nu}$.

Particularly, in a uniformly rotating frame, the tetard can be described by

$$e^a_\mu = \delta^a_\mu + \delta^a_i \delta^0_\mu v_i, \quad e^\mu_a = \delta^\mu_a - \delta^0_a \delta^\mu_i v_i, \tag{6}$$

where $\vec{v} = \vec{\Omega} \times \vec{x}$ gives the velocity in the lab frame. We use the capital Greek letter Ω to represent the magnitude of angular velocity. If we substitute Eq.(6) into Eq.(4), non-zero terms of spin connection will be expressed as following:

$$\Gamma_{ij0} = \frac{1}{2} \left(\partial_i v_j - \partial_j v_i \right), \\ \Gamma_{0ij} = -\frac{1}{2} \left(\partial_i v_j + \partial_j v_i \right), \\ \Gamma_{0ij} = -\frac{1}{2} \left(\partial_i v_j + \partial_j v_i \right), \\ \Gamma_{0i0} = -\frac{1}{2} \left(v_j \partial_i v_j + v_j \partial_j v_i \right).$$

$$(7)$$

Finally, in a uniformly rotating frame, Dirac equation can be rewritten by [33]

$$[i\gamma^a\partial_a + \gamma^0\Omega\hat{J}_z - M_f]\psi = 0.$$
(8)

Here, z-axis is chosen as the direction of the rotation and it is perpendicular to the reaction plane. The total angular momentum \hat{J}_z can be expressed as $\hat{J}_z = \hat{L}_z + \hat{S}_z$ where \hat{L}_z is the orbital angular momentum and $\hat{S}_z = \frac{1}{2} \begin{pmatrix} \hat{\sigma}_z & 0\\ 0 & \hat{\sigma}_z \end{pmatrix}$ contributes to the spin part.

For solving modified Dirac equation Eq.(8), Ref.[12] has given the solutions in the cylindrical coordinates where a position in space-time is labeled by $\tilde{r} = (t, r, \theta, z)$. We can write down a complete set of commuting operators: the Hamiltonian \hat{H} , the longitudinal momentum \hat{k}_z , the square of transverse momentum \hat{k}_t^2 , the total angular momentum \hat{J}_z , and the helicity operator on transverse plan \hat{h}_t . The eigenstates for fermion and anti-fermion [12] are given by:

$$U = \sqrt{\frac{E_k + M_f}{4E_k}} e^{ik_z z} e^{in\theta} \begin{pmatrix} J_n(k_t r) \\ s e^{i\theta} J_{n+1}(k_t r) \\ \frac{k_z - isk_t}{E_k + M_f} J_n(k_t r) \\ \frac{-sk_z + ik_t}{E_k + M_f} e^{i\theta} J_{n+1}(k_t r) \end{pmatrix},$$
(9)

$$V = \sqrt{\frac{E_k + M_f}{4E_k}} e^{-ik_z z} e^{in\theta} \begin{pmatrix} \frac{k_z - isk_t}{E_k + M_f} J_n\left(k_t r\right) \\ \frac{sk_z - ik_t}{E_k + M_f} e^{i\theta} J_{n+1}\left(k_t r\right) \\ J_n\left(k_t r\right) \\ -s e^{i\theta} J_{n+1}\left(k_t r\right) \end{pmatrix},$$
(10)

where $J_n(k_t r)$ is the *n*-th order Bessel function and $n \in \mathbb{Z}$ stands for the quantum number of angular momentum.

In Eq.(9) and Eq.(10), k_t and k_z are the eigenvalues of transverse and longitudinal momentum, respectively, and $s = \pm 1$ is the eigenvalue of the helicity operator \hat{h}_t . Besides, E_k is defined by $E_k \equiv \sqrt{k_z^2 + k_t^2 + M_f^2}$ and the energy is given by $E = \pm E_k - (n + 1/2)\Omega$.

Based on the solutions of Dirac equation, one can write down the quark propagator by definition: $S(\tilde{r};\tilde{r}') = \langle 0 | T\psi(\tilde{r})\bar{\psi}(\tilde{r}') | 0 \rangle$. As a standard procedure, $\psi(\tilde{r})$ and $\bar{\psi}(\tilde{r}')$ are expanded by Eq.(9) and Eq.(10), and a explicit form can be obtained as following:

$$S\left(\tilde{r};\tilde{r}'\right) = \frac{1}{(2\pi)^2} \sum_{n} \int \frac{dk_0}{2\pi} \int k_t dk_t \int dk_z \frac{e^{in(\theta-\theta')} e^{-ik_0(t-t')+ik_z(z-z')}}{\left[k_0 + \left(n + \frac{1}{2}\right)\Omega\right]^2 - k_t^2 - k_z^2 - M_f^2 + i\epsilon} \\ \times \left\{ \left[\left[k_0 + \left(n + \frac{1}{2}\right)\Omega\right]\gamma^0 - k_z\gamma^3 + M_f \right] \left[J_n\left(k_tr\right)J_n\left(k_tr'\right)\mathcal{P}_+ + e^{i\left(\theta-\theta'\right)}J_{n+1}\left(k_tr\right)J_{n+1}\left(k_tr'\right)\mathcal{P}_- \right] \right]^{(11)} - i\gamma^1 k_t e^{i\theta} J_{n+1}\left(k_tr\right)J_n\left(k_tr'\right)\mathcal{P}_+ - \gamma^2 k_t e^{-i\theta'}J_n\left(k_tr\right)J_{n+1}\left(k_tr'\right)\mathcal{P}_- \right\}.$$

Here, we have simplified the expression by projection operators $\mathcal{P}_{\pm} = \frac{1}{2} (1 \pm i\gamma^1 \gamma^2)$. In Appendix A, we will present another procedure for obtaining the quark propagator in a rotating medium.

Nevertheless, Eq.(11) is the result under the infinite size approximation. One can find a finite size version with a boundary condition (see Ref.[21]). The transverse momentum k_t is discrete and its integral is replaced by the summation of series. In this manuscript, firstly, we have to calculate the spectral functions of vector mesons in a rotating medium. So, we apply the infinite size approximation to avoid the tedious summation of the series. This compromise will cause the violation of causality in the case of large angular velocity. In this manuscript, we study the phenomena and quantities at a fixed radius $r = 0.1 \text{GeV}^{-1}$ and in a range of angular velocity from $\Omega = 0 \text{ GeV}$ to $\Omega = 0.4 \text{ GeV}$ so that the velocity of a fixed point is smaller than the speed of light, i.e. $\Omega r < 1$.

B. The 3-flavor NJL model

In order to study the microscopic properties of vector mesons, we use the Nambu-Jona-Lasinio(NJL) model to estimate the strong interaction[34]. The 3-flavor NJL model is required to investigate ϕ meson, which contains s quark [35]. The Lagrangian of the 3-flavor NJL model is given as following:

$$\mathcal{L}_{3\mathrm{NJL}} = \psi [i\bar{\gamma}^{\mu} (\partial_{\mu} + \Gamma_{\mu}) - m_{f}]\psi + G \sum_{a=0}^{8} \left[\left(\bar{\psi} \lambda^{a} \psi \right)^{2} + \left(\bar{\psi} i \gamma_{5} \lambda^{a} \psi \right)^{2} \right] - K \left[\det \bar{\psi} \left(1 + \gamma_{5} \right) \psi + \det \bar{\psi} \left(1 - \gamma_{5} \right) \psi \right],$$
(12)

where K is the coupling constant of six-fermion interaction, G_s and G_V are coupling constants of four-fermion interaction for scalar and vector channels respectively. In the 3-flavor NJL model, $\psi = (\psi_u, \psi_d, \psi_s)$ is a Dirac spinor which contains u, d and s quarks. Correspondingly, λ_a are the Gell-Man matrices and m_f is the current quark mass with different flavors.

The NJL model is an effective model which only contains quarks. The rotation affects the quark dynamics through the non-zero term of spinor connection Γ^{μ} . Furthermore, by applying the mean field approximation, the Lagrangian will be rewritten as:

$$\mathcal{L}_{\rm MF} = \sum_{f=u,d,s} \bar{\psi}_f \left[i \bar{\gamma}_\mu (\partial_\mu + \Gamma_\mu) - M_f \right] \psi_f - 2G_S \sum_{f=u,d,s} \sigma_f^2 + 4K \sigma_u \sigma_d \sigma_s$$
(13)

where σ_f is the chiral condensate $\sigma_f \equiv \langle \bar{\psi}_f \psi_f \rangle$ for a specific flavor. And M_f is dynamical quark mass which is given by

$$M_f \equiv m_f - 4G_S \sigma_f + 2K \prod_{f' \neq f} \sigma_{f'}.$$
 (14)

In a uniformly rotating medium, the spinor connection Γ^{μ} has been estimated in section II A. We assume the direction of rotation is parallel to the z-axis. By applying the standard procedure in finite-temperature field theory [36], the grand potential for quarks with specific flavor is shown as follows:

$$\Omega_{f}(r) = \frac{N_{c}}{8\pi^{2}}T\sum_{n}\int dk_{t}^{2}\int dk_{z} \left[J_{n}\left(k_{t}r\right)^{2} + J_{n+1}\left(k_{t}r\right)^{2}\right] \\ \times \left[E_{k}/T + \ln\left(1 + e^{-\left(E_{k} - \left(n + \frac{1}{2}\right)\Omega\right)/T}\right) + \ln\left(1 + e^{-\left(E_{k} + \left(n + \frac{1}{2}\right)\Omega\right)/T}\right)\right].$$
(15)

Consequently, the total grand potential is

$$\Omega_{\rm tot}(r) = \sum_{f=u,d,s} \left(2G_S \sigma_f^2 - \Omega_f \right) + 4K \sigma_u \sigma_d \sigma_s.$$
(16)

Here, $N_c = 3$ is the degeneracy of color and T is the temperature of the medium. We can obtain the dynamical

quark mass M_f and chiral condensate σ_f by solving the gap equations:

$$\frac{\partial \Omega_{\rm tot}}{\partial \sigma_f} = 0, \quad \frac{\partial^2 \Omega_{\rm tot}}{\partial \sigma_f^2} > 0. \tag{17}$$

In principle, if other parameters are fixed, the dynamical quark mass $M_f(r)$ will be a function of radius r. Currently, most of studies have assumed that $M_f(r)$ is changed smoothly, i.e. $\partial M_f(r)/\partial r \simeq 0$. Such assumption is called local density approximation(LDA)[12, 17, 25]. In this manuscript, we use the LDA and choose a fixed radius $r = 0.1 \text{GeV}^{-1}$. Then, $M_f(\Omega)$ will be evaluated numerically in section III A.

C. Vector meson mass splitting under the rotation

In the NJL model, the vector meson ϕ is treated as a $s\bar{s}$ bound state or a resonance state, and it can be constructed by quark-antiquark scattering[34]. In the random phase approximation (RPA), vector meson propagator can be obtained by summation of quark loops and one loop polarization function is given by

$$\Pi^{\mu\nu}(q) = -i \int d^4 \tilde{r} \operatorname{Tr}_{sfc}[i\gamma^{\mu}S(0;\tilde{r})i\gamma^{\nu}S(\tilde{r};0)]e^{iq\cdot\tilde{r}}.$$
 (18)

Here, Tr_{sfc} stands for the trace in the spinor, flavor and color spaces. Since ϕ meson is purely constituted by s quark, $S(0; \tilde{r})$ stands for the s quark propagator with the dynamical mass M_s given by mean field approximation. The polarization function of ϕ meson is supposed to be modified in a rotating medium.

Currently, we focus on the mesons which remain at rest in the rotating frame, i.e. $\vec{q} = 0$. In this case, polarization vectors are given by

$$\epsilon_1^{\mu} = \frac{1}{\sqrt{2}}(0, 1, i, 0),$$

$$\epsilon_2^{\mu} = \frac{1}{\sqrt{2}}(0, 1, -i, 0),$$

$$b^{\mu} = (0, 0, 0, 1)$$
(19)

where ϵ_1^{μ} and ϵ_2^{μ} are the right and left-hand polarization vectors respectively. Longitudinal polarization vector is parallel to the direction of rotation. Correspondingly, the projection operators are given by

$$P_1^{\mu\nu} = -\epsilon_1^{\mu} \epsilon_1^{\nu}, (s_z = -1 \text{ for } \phi \text{ meson }),$$

$$P_2^{\mu\nu} = -\epsilon_2^{\mu} \epsilon_2^{\nu}, (s_z = +1 \text{ for } \phi \text{ meson }),$$

$$L^{\mu\nu} = -b^{\mu} b^{\nu}, (s_z = 0 \text{ for } \phi \text{ meson }).$$
(20)

As we have assumed $\vec{q} = 0$, nonzero elements of the polarization functions can be read in the following matrix:

$$\Pi_{\phi}^{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \Pi^{11} & \Pi^{12} & 0 \\ 0 & \Pi^{21} & \Pi^{22} & 0 \\ 0 & 0 & 0 & \Pi^{33} \end{pmatrix}.$$
 (21)

The explicit expressions of matrix elements are shown in Ref.[19]. This tensor can be decomposed by projection operators in Eq.(20) as following:

$$\Pi_{\phi}^{\mu\nu} = A_{-1,-1}P_1^{\mu\nu} + A_{11}P_2^{\mu\nu} + A_{00}L^{\mu\nu}, \qquad (22)$$

Similarly, the vector meson propagator can be decomposed as following:

$$D^{\mu\nu}_{\phi}(q) = D_{-1,-1}(q)P^{\mu\nu}_1 + D_{11}(q)P^{\mu\nu}_2 + D_{00}(q)L^{\mu\nu}.$$
 (23)

Here, coefficients D_{λ} are obtained by RPA summation and expressed by:

$$D_{\lambda\lambda}(q) = \frac{4G_V}{1 + 4G_V A_{\lambda\lambda}},\tag{24}$$

where

$$A_{-1,-1} = -(\Pi_{11} - i\Pi_{12}), \quad (s_z = -1 \text{ for } \phi \text{ meson }),$$

$$A_{11} = -\Pi_{11} - i\Pi_{12}, \quad (s_z = +1 \text{ for } \phi \text{ meson }),$$

$$A_{00} = -\Pi_{33}, \quad (s_z = 0 \text{ for } \phi \text{ meson }).$$

(25)

Then we can obtain the corresponding spectrum functions for different spin components, which takes the following form:

$$\xi_{\lambda\lambda}(\omega) \equiv \frac{1}{\pi} \operatorname{Im} D_{\lambda\lambda}(\omega) = \frac{(4G_V)^2 \operatorname{Im} A_{\lambda\lambda}(\omega)}{\pi \left\{ \left[1 + 4G_V \operatorname{Re} A_{\lambda\lambda}(\omega) \right]^2 + \left[4G_V \operatorname{Im} A_{\lambda\lambda}(\omega) \right]^2 \right\}}$$
(26)

where ω is the energy of meson and we set the momentum to $\vec{q} = 0$.

D. spin alignment of vector meson ϕ

Heavy-ion collisions will create an ensemble of particles under extreme conditions. Particularly, we consider vector meson ϕ with spin-1. For a fixed direction, a normalized spin state of a ϕ meson is labeled by $|\lambda\rangle$ with $\lambda = 1, 0, -1$. Spin density operator ρ is defined by

$$\rho = \sum_{\lambda\lambda'} \rho_{\lambda\lambda'} \left| \lambda \right\rangle \left\langle \lambda' \right|, \qquad (27)$$

Here, $\rho_{\lambda\lambda'}$ comprises a 3 × 3 spin density matrix as following:

$$\rho_{\lambda\lambda'} = \begin{pmatrix} \rho_{11} & \rho_{10} & \rho_{1,-1} \\ \rho_{01} & \rho_{00} & \rho_{1,-1} \\ \rho_{-1,1} & \rho_{-1,0} & \rho_{-1,-1} \end{pmatrix}, \qquad (28)$$

An ensemble of ϕ mesons, for which spin information is described by $\rho_{\lambda\lambda'}$, will decay to

$$\phi \to K^+ + K^-. \tag{29}$$

In this process, daughter particles will have an angular distribution[37]:

$$\frac{dN}{d\Omega^*} = \frac{3}{4\pi} \left\{ \cos^2 \theta \rho_{00} + \sin^2 \theta \left(\rho_{11} + \rho_{-1-1} \right) / 2 - \sin 2\theta \left(\cos \phi \operatorname{Re} \rho_{10} - \sin \phi \operatorname{Im} \rho_{10} \right) / \sqrt{2} + \sin 2\theta \left(\cos \phi \operatorname{Re} \rho_{-10} + \sin \phi \operatorname{Im} \rho_{-10} \right) / \sqrt{2} - \sin^2 \theta \left[\cos(2\phi) \operatorname{Re} \rho_{1-1} - \sin(2\phi) \operatorname{Im} \rho_{1-1} \right] \right\}.$$
(30)

From experimental data of θ -distribution, ρ_{00} can be obtained as a coefficient of angular distribution. Since experiments can measure the value of $\rho_{\lambda\lambda'}$, theoretical studies should explain the results of spin alignment and evaluate $\rho_{\lambda\lambda'}$ qualitatively. Magnetic field and vorticity are two factors which are taken into account in many models. The approaches are generalized as follows:

B,
$$\Omega$$
 (quarks) $\xrightarrow{\text{influence}} \rho_{\lambda\lambda'}(\phi) \xrightarrow{\text{determine}} \frac{dN}{d\Omega^*} (K^+, K^-)$
(31)

Theoretical studies are interested in the first step of Eq.(31). In a real heavy-ion collision process, the evolution is complicated, and the history will influence $\rho_{\lambda\lambda'}$ after the freeze out. One possible method is the quark coalescence model. However, it is unable to take the medium effects into account in the quark coalescence model. As we mentioned in previous sections, quark mass and meson mass will be modified in a rotating medium.

In this investigation, we aim at a uniformly rotating medium and the created ϕ mesons are in global equilibrium. In this case, the particle number density $\overline{\rho}_{\lambda\lambda'}(\mathbf{q})$ can be expressed by

$$\overline{\rho}_{\lambda\lambda'}(\mathbf{q}) = \int d\omega \, \frac{2\omega}{e^{\omega/T} - 1} \xi_{\lambda\lambda'}(\omega, \mathbf{q}) \,, \qquad (32)$$

where $\xi_{\lambda\lambda'}(\omega, \mathbf{k})$ is the spectral function given in Eq.(26). At the one loop level, the spectral function in a rotating medium is calculated in Ref.[19], and we present it in Appendix B for convenience. Generally, the spectral function is the imaginary part of the full propagator:

$$\xi_{\lambda\lambda'}(q) \equiv \frac{1}{\pi} \text{Im} D_{\lambda\lambda'}(q) \,. \tag{33}$$

Particularly, if the rotation axis is chosen as the quantized direction, $\xi_{\lambda\lambda'}(\omega, \mathbf{q})$ and $\overline{\rho}_{\lambda\lambda'}(\mathbf{q})$ will become diagonal:

$$\overline{\rho}_{\lambda\lambda'}(\mathbf{q}) = \begin{pmatrix} \overline{\rho}_{11} & 0 & 0\\ 0 & \overline{\rho}_{00} & 0\\ 0 & 0 & \overline{\rho}_{-1,-1} \end{pmatrix} .$$
(34)

Therefore we divide the spectral function into two part: a delta function part and a continuum part, i.e.

$$\xi_{\lambda\lambda}(\omega,\mathbf{q}) = \delta\left(\omega^2 - M_{\phi,\lambda}^2\right) + \xi_{\lambda\lambda}^*(\omega,\mathbf{q}) \qquad (35)$$

where $\xi_{\lambda\lambda}^{*}(\omega, \mathbf{q})$ is the continuum part of spectral function and $M_{\phi,\lambda}$ is the vector meson mass for different spin components. Correspondingly the particle number density $\bar{\rho}_{\lambda\lambda}(\mathbf{q})$ takes the form of:

$$\bar{\rho}_{\lambda\lambda}(\mathbf{q}) = \frac{1}{\exp\left(M_{\phi,\lambda}/T\right) - 1} + \int d\omega \frac{2\omega\xi_{\lambda\lambda}^*(\omega, \mathbf{q})}{\exp(\omega/T) - 1} \quad (36)$$

In experiment measurement, spin alignment is evaluated by matrix element ρ_{00} which can be expressed by:

$$\rho_{00}(\mathbf{q}) \equiv \frac{\overline{\rho}_{00}(\mathbf{q})}{\sum_{\lambda=0,\pm 1} \overline{\rho}_{\lambda\lambda}(\mathbf{q})} \,. \tag{37}$$

In the non-rotating case, ϕ mesons with different spin components have the same mass $M_{\phi}(s_z = 0, \pm 1)$, which leads to $\rho_{00} = 1/3$. In a finite angular velocity, the mass of ϕ meson with $s_z = +1$ component decreases linearly thus ϕ mesons tend to occupy the $s_z = +1$ state. As a consequence, spin polarization and spin alignment can be obtained in our formalism.

In this section, Eqs.(32-37) are momentum-dependent. However, in this investigation, we evaluate the spectral functions and spin alignment with $\mathbf{q} = \mathbf{0}$, which means vector mesons are staying at rest in the rotating frame. In general, Ref.[11] has revealed the spin density matrix for arbitrary measuring direction which is characterized by Euler angles (α , β , γ). The explicit form is:

$$\overline{\rho}_{\lambda\lambda'}(\mathbf{0};\alpha,\beta,\gamma) = \sum_{\lambda_1,\lambda_2} R_{\lambda\lambda_1}(\alpha,\beta,\gamma) \overline{\rho}_{\lambda_1\lambda_2}(\mathbf{0}) R_{\lambda_2\lambda'}^{-1}(\alpha,\beta,\gamma) . \quad (38)$$

Here, $R_{\lambda\lambda'}(\alpha, \beta, \gamma)$ is the spin-1 representation of the rotation. As a result, the spin alignment only depends on Euler angles β , and the explicit form is:

$$\rho_{00}(\mathbf{0};\alpha,\beta,\gamma) = \frac{\overline{\rho}_{00}(\mathbf{0})\cos^2\beta + \overline{\rho}_{11}(\mathbf{0})\sin^2\beta}{\overline{\rho}_{00}(\mathbf{0}) + \overline{\rho}_{11}(\mathbf{0}) + \overline{\rho}_{-1,-1}(\mathbf{0})}.$$
 (39)

In Section III C, numerical results will be presented for $\beta = 0$, and the following formula is used:

$$\rho_{00}^{\Omega}(\mathbf{0}) \equiv \frac{\bar{\rho}_{00}(\mathbf{0})}{\bar{\rho}_{00}(\mathbf{0}) + \bar{\rho}_{11}(\mathbf{0}) + \bar{\rho}_{-1,-1}(\mathbf{0})} \,. \tag{40}$$

III. NUMERICAL RESULT

Since a rotating system has broken the Lorentz symmetry, it is not necessary to use Pauli-Villas regularization. In fact, the Bessel function will not reflect the oscillation behavior in the cut-off energy scale Λ . Therefore we choose a hard cut-off function:

$$f_{\Lambda} = \frac{\Lambda^{2*10}}{\Lambda^{2*10} + p^{2*10}} \tag{41}$$

where $\Lambda = 620.411 \text{MeV}$ is the energy scale for cut-off. And the corresponding coupling constants are: $G_S = \frac{1.710}{\Lambda^2}$, $G_V = 0.67 * \frac{1.710}{\Lambda^2}$ and $K = \frac{12035}{1000\Lambda^5}$. Current quark masses are: $m_u = m_d = 5.5 \text{ MeV}$, $m_s = 135.433 \text{ MeV}$.




FIG. 1. Dynamical quark masses as functions of angular velocity at temperature T = 10 MeV. Red lines stand for mass of s quark and the blue lines stand for light quark u and d.




FIG. 2. Dynamical quark masses as functions of angular velocity at temperature T = 150 MeV. Red lines stand for mass of s quark and the blue lines stand for light quark u and d.

A. Dynamical quark mass

As mentioned in previous sections, chiral condensate will be suppressed under rotation. In FIG. 1, at an almost zero temperature, T = 10MeV, rotation induces a first order transition at angular velocity $\Omega_c = 0.713$ GeV. Both M_u and M_d stay at a constant mass when the angular velocity is below Ω_c . From Eq.(14), we know that dynamical quark masses are determined by the chiral condensates σ_u , σ_d and σ_s . In fact, $\sigma_u = \sigma_d$ and drops at Ω_c while σ_s still varies smoothly. Consequently, M_s has a small jump at Ω_c and then decrease smoothly. The behavior of $M_{u,d,s}$ as a function of angular velocity Ω is very similar to the case of finite quark chemical potential [38, 39], but the first order phase transition will occur at $\mu_c \simeq 0.33$ GeV at finite density.

At a higher temperature, T = 150 MeV, FIG.2 reveals that chiral phase transition will be crossover which occurs around $\Omega_c \sim 0.4$ GeV. The phase transition takes place in a smaller angular velocity. Furthermore, It is noticed that the mass decreases slowly before the phase transition. Above all, the rotational effect on the dynamical quark mass is similar to that of the chemical potential μ_q .




FIG. 3. ϕ meson mass as a function of angular velocity at temperature T = 150 MeV and $\mu = 0$ MeV.




FIG. 4. The deviation of ϕ meson mass as a function of angular velocity at temperature T = 150 MeV.

B. Mass spectra of ϕ and ρ meson under rotation

After we obtain the dynamical quark masses in different temperatures and angular velocities, we can apply this result in Eq.(18) and obtain corresponding meson masses.

At zero temperature and low temperatures, similar to ρ meson, ϕ meson mass with different spin components $s_z = 0, \pm 1$ also show mass splitting effect with $M_{\phi}(\Omega) = M_{\phi}(0) - s_z \Omega$.

At T = 150 MeV and $\mu = 0$ MeV, FIG.3 shows ϕ meson mass with different spin components $s_z = 0, \pm 1$ as a function of angular velocity. The mass of $s_z = 0$ component for ϕ meson almost remain unchanged with the angular velocity. The mass of $s_z = -1$ component of ϕ meson grows almost linearly with the angular velocity. It implies that ϕ meson will be less possible to stay at $s_z = -1$ state. In contrast, the $s_z = +1$ component ϕ meson mass decreases almost linearly with the angular velocity. $s_z = +1$ component will be a preferred state under the rotation. Above all, the nearly linear mass splitting behavior of ϕ meson at T = 150 MeV can be summarized in the following expression:

$$M_{\phi}(\Omega) \sim M_{\phi}(0) - s_z \Omega \tag{42}$$

In FIG.4, we reveal the deviation of ϕ meson mass from




FIG. 5. The spectral function $\xi(\omega)$ for ϕ meson with different spin component as a function of the frequency ω under different angular velocities $\Omega = 0, 0.1, 0.20.3$ GeV at T = 150MeV and $\mu = 0$.

Eq.(42) at T = 150 MeV. The deviation is caused by the inhibition of chiral condensate. The dashed line shows the deviation of $s_z = 0$ component, which is caused by quark mass descending in finite temperature and angular velocity. ϕ meson is an almost pure $s\bar{s}$ state [34], so its mass is influenced by the quark mass obviously. It is seen that the deviation of $s_z = \pm 1$ components states are larger than that of the $s_z = 0$ component state.

FIG.5 and FIG .6 reveal rotational effect on the spectra function $\xi(\omega)$ for ϕ meson and ρ meson with different spin components as a function of the frequency ω respectively. The blue lines stand for spectral functions at $\Omega = 0$ without rotation, while the red, orange and gray dashed lines stand for spectral functions under finite rotation in the case of $\Omega = 0.1, 0.2$ and 0.3 GeV, respectively. In the zero rotation case, ϕ mesons with different spin states have shared the same spectral function which is constituted by a delta function part and a continuum part. The




FIG. 6. The spectral function $\xi(\omega)$ for ρ meson with different spin component as a function of the frequency ω under under different angular velocities $\Omega = 0, 0.1, 0.20.3$ GeV at T = 150 MeV and $\mu = 0$.

location of the delta function indicates the pole mass.

For $s_z = 0$ component of ϕ meson, the rotational effect is less remarkable than the other two cases. So, the scale has been amplified and we only present the $\xi(\omega)$ in the range of energy ω from 0.85 GeV to 1.20 GeV. It is found that spectral functions are shifted to the left slightly. And the peaks of the continuum parts are enhanced significantly.

For $s_z = +1$ component of ϕ meson, the rotational effect will shift the spectral function to the left side, and the rotation will change the height or the shape of the continuum part of the spectral function as well. For the $s_z = -1$ component, the spectral function is shifted to the right side correspondingly.

A similar analysis can be applied for the vector meson ρ , since we have assumed $M_u = M_d$. We can obtain the spectral function by substituting M_s for M_u/M_d . For ϕ mesons, a bound state is labeled by mass M_{ϕ} which is




FIG. 7. Spin alignment ρ_{00} for vector meson ϕ as a function of angular velocity at temperature T = 150 MeV.

very close to the $2M_s$. However, ρ mesons are dissociated at the temperature T = 150 MeV. So, in FIG.(6), a spectral function only has a continuum part and appears as a single peak. The top panel in FIG.(6) shows the spectral functions of ρ mesons with spin components $s_z = 0$. Different colored lines stand for different strengths of angular velocities ranging from $\Omega = 0, 0.1, 0.2$ and 0.3 GeV. Rotational effects are reflected in two aspects: the heights of the peaks are suppressed and the widths are broadened by the angular velocities. It can be understood that mesons tend to be less bounded in a rotating medium. The location of the peaks is almost unchanged in the case of $s_z = 0$. However, in the case of $s_z = +1$, the locations of resonance peaks are shifted to the left side by rotation. Similarly, in the case of $s_z = -1$, mass spectra are shifted to the right side by rotation. Above all, on the shape of spectral functions, the rotation effects are similar.

C. Spin alignment of vector meson ϕ and ρ

In FIG.7, we show the deviation of spin alignment ρ_{00} from 1/3 for ϕ meson as a function of angular velocity at a finite temperature T=150 MeV. In the case of rotation, ρ_{00} is always smaller than 1/3, and the deviation will become more significant as the angular velocity grows. Furthermore, resonance states will have less contribution for spin alignment and the deviation between the bounded state and the total result is even negligible. In FIG. 8, we compare the spin alignment ρ_{00} from 1/3 for ϕ meson with ρ meson, and it is found the difference is quite small.

Here, we mention the result from the quark coalescence model [40]:

$$\rho_{00}^{\phi}(\Omega) = \frac{1}{3} - \frac{1}{9}(\beta\Omega)^2, \qquad (43)$$

where $\beta = 1/T$ is the inverse of temperature. If we fit our result with polynomial functions, numerical results give $\rho_{00}^{\phi} \sim 1/3 - 3.1\Omega^2$, and the absolute value of the coefficient is smaller than $-\frac{1}{9}\beta^2 = 4.94$.




FIG. 8. Spin alignment ρ_{00} for resonance states of vector meson ρ and ϕ as a function of angular velocity at temperature T = 150 MeV.

Compared with the spin alignment under an external magnetic field and rotation, the deviation $\rho_{00} - 1/3$ is positive under the magnetic field while it is negative in the presence of rotation. It is natural to be understood from quark dynamics that the spin of a particle tends to align along the direction of angular momentum due to the spin-orbital coupling. For $s_z = +1$ component, the vector ϕ, ρ mesons masses are suppressed in the rotating medium. As a consequence, vector mesons are more possible to occupy the $s_z = +1$ state and less possible to occupy $s_z = 0$ state. So, $\rho_{00} - 1/3$ is negative in the rotating medium. On the contrary, $\rho_{00} - 1/3$ of ϕ meson is positive under the magnetic field [11]. Actually, vector meson masses in the magnetic field are charge dependent. The ϕ meson is a neutral particle, its property under the magnetic field can be extended from the result of neutral ρ^0 meson mass spectra under the magnetic field [41]. Under the magnetic field, neutral ϕ meson with $s_z = \pm 1$ will have a larger mass than ϕ meson with $s_z = 0$. So, ϕ mesons are more possible to occupy the $s_z = 0$ state in presence of the magnetic field, which naturally explains $\rho_{00} - 1/3$ for ϕ meson is positive under the magnetic field.

Similarly, our theoretical method can be applied to other species of vector mesons, such as ρ and K^{*0} . The difference is the dynamical mass of constituent quark u, dand s in the rotating medium. In Eq.(18), only one specie of quark propagator exists in the one loop polarization function. Since we have applied the assumption $M_u =$ M_d , spin alignment of ρ meson is demonstrated in FIG.8. In the rotating medium with a temperature of 150 MeV, ρ mesons are resonance states. So, we compare it with the resonance states of ϕ mesons in FIG.8. The tendency of the deviation $\rho_{00} - 1/3$ is still close to the quadratic polynomial.

It is worth to remind that those results are calculated for vector mesons that stay at rest in a rotating medium, i.e. at $\vec{q} = 0$. In the $\vec{q} \neq 0$ case, the calculation will be more complicated, and it is still a puzzle to switch the physical quantities in the rotating frame into the counterparts in the lab frame. Above all, the contribution from the rotating medium is significant, although the spin alignment of vector mesons is affected by a combination of many factors.

IV. CONCLUSION AND DISCUSSION

In this work, we investigate the spin alignment of vector mesons ϕ and ρ induced by rotation. By applying a three flavor NJL model with a vector interaction channel, we obtain the dynamical quark mass under rotation. The curves of $M_f(\Omega)$ is similar to $M_f(\mu)$. For the *s* quark, the first order phase transition occurs at a critical angular velocity Ω_c , and M_s decreases smoothly after the phase transition, which is similar to the quark mass behavior at a finite chemical potential.

After substituting the dynamical quark mass, the mass spectra of vector mesons can be obtained through the quark-antiquark polarization function. The rotating angular velocity induces mass splitting of spin components for vector ϕ, ρ mesons $M_{\phi,\rho}(\Omega) \simeq M_{\phi,\rho}(\Omega = 0) - s_z \Omega$. This behavior contributes to the spin alignment of vector mesons ϕ, ρ in an equilibrium medium. In a rotating medium, ρ_{00} of vector mesons has a negative deviation from 1/3 which implies a spin alignment phenomenon, which can be easily understood from quark dynamics that the spin of a particle tends to align along the direction of angular momentum due to the spin-orbital coupling. For $s_z = +1$ component, the vector mesons masses are suppressed in the rotating medium. As a consequence, vector mesons are more possible to occupy the $s_z = +1$ state and less possible to occupy $s_z = 0$ state. So, $\rho_{00} - 1/3$ is negative in the rotating medium.

On the contrary, the deviation $\rho_{00} - 1/3$ is positive under the magnetic field, which can also be easily understood from quark dynamics. Under the magnetic field, $s_z = \pm 1$ components of ϕ , ρ meson will have a larger mass than that of the $s_z = 0$ component of ϕ , ρ mesons. So, ϕ , ρ mesons are more possibly to occupy the $s_z = 0$ state in the presence of the magnetic field, which naturally explains positive $\rho_{00} - 1/3$ for ϕ meson under magnetic field.

Based on a dynamical quark model, in the next step, we are able to calculate the spin alignment ρ_{00} for the transverse momentum dependent case. Furthermore, we can apply it for vector meson K^{*0} in the future. In this series of studies, we have studied the vector meson ϕ and ρ . In this case, quark loops only contain s and \bar{s} quarks, and light quarks are assumed to have the same value of mass, i.e. $M_u = M_d$. For vector meson K^{*0} , the mass difference of quarks is expected to explain the different measurement between vector meson ϕ and K^{*0} in the experiment. Our work is an attempt at this target in a rotating medium.

ACKNOWLEDGMENTS

We thank the helpful discussion with Li Yan, Anping Huang, Xinli Sheng and Kun Xu. This work is supported in part by the National Natural Science Foundation of China (NSFC) Grant Nos. 12235016, 12221005, the Strategic Priority Research Program of Chinese Academy of Sciences under Grant Nos XDB34030000, the startup funding from University of Chinese Academy of Sciences(UCAS), and the Fundamental Research Funds for the Central Universities.

Appendix A: Quark propagator in a rotating medium

To obtain the quark propagator in a rotating and dense medium, we adopt the method from Vladimir A. Miransky and Igor A. Shovkovy[16]. This derivation has considered the chemical potential μ and the rotation term $\Omega \cdot \hat{J}_z$. Accord to an alternative definition, the quark propagator is given by:

$$S\left(\tilde{r},\tilde{r}'\right) = i\left\langle \tilde{r} \left| \left[\left(i\partial_t + \mu + \Omega \cdot \hat{J}_z \right) \gamma^0 - \vec{\pi} \cdot \vec{\gamma} - M_f \right]^{-1} \right| \tilde{r}' \right\rangle,\tag{A1}$$

where $\vec{\pi}$ is the canonical momentum and $\vec{\gamma}$ is the Dirac matrix. Their expressions depend on the coordinates of the position \tilde{r} . At this moment, we treat them as abstract operators and the propagator can be rewritten by:

$$S\left(\tilde{r},\tilde{r}'\right) = i\left\langle \tilde{r} \left| \left[\left(i\partial_t + \mu + \Omega \cdot \hat{J}_z \right) \gamma^0 - \vec{\pi} \cdot \vec{\gamma} + M_f \right] \left[\left(i\partial_t + \mu + \Omega \cdot \hat{J}_z \right) \gamma^0 - \vec{\pi} \cdot \vec{\gamma} + M_f \right]^{-1} \right] \right\rangle$$

$$\left[\left(i\partial_t + \mu + \Omega \cdot \hat{J}_z \right) \gamma^0 - \vec{\pi} \cdot \vec{\gamma} - M_f \right]^{-1} \left| \tilde{r}' \right\rangle$$

$$= i\left\langle \tilde{r} \left| \left[\left(\partial_t + \mu + \omega \cdot \hat{J}_z \right) \gamma^0 - \vec{\pi} \cdot \vec{\gamma} + M_f \right] \left[\left(i\partial_t + \mu + \Omega \cdot \hat{J}_z \right)^2 - \vec{\pi}^2 - M_f^2 \right]^{-1} \right| \tilde{r}' \right\rangle$$
(A2)

Due to the translation invariance in t- and z- direction, we can perform the Fourier transformation on the quark propagator as follows:

$$S(E, k_z; \mathbf{r}_{\perp}, \mathbf{r}'_{\perp}) = \int dt dz e^{iE(t-t') - ik_z(z-z')} S(\tilde{r}, \tilde{r}').$$
(A3)

Here, $\mathbf{r}_{\perp} = (r, \theta)$ is the position in cylindrical coordinates and E is the energy. The propagator can be expressed as follows:

$$S(E, k_{z}; \mathbf{r}_{\perp}, \mathbf{r}'_{\perp}) = i \left[\left(E + \mu + \Omega \cdot \hat{J}_{z(\mathbf{r}_{\perp})} \right) \gamma^{0} - \vec{\pi}_{\mathbf{r}_{\perp}} \cdot \vec{\gamma}_{\perp} - k_{z} \gamma^{3} + M_{f} \right] \\ \cdot \left\langle \mathbf{r}_{\perp} \left| \left[\left(E + \mu + \Omega \cdot \hat{J}_{z} \right)^{2} - (k_{z})^{2} - \vec{\pi}_{\perp}^{2} - M_{f}^{2} \right]^{-1} \right| \mathbf{r}'_{\perp} \right\rangle$$
(A4)

where $\vec{\pi}_{\mathbf{r}_{\perp}}$ and $\hat{J}_{z(r_{\perp})}$ are the canonical momentum and the angular momentum operator in cylindrical coordinate space, respectively. According to Ref.[12], the operators $\vec{\pi}_{\perp}^2$ and \hat{L}_z commute with each other and have a common eigenstate $|nk_t\rangle$, which the explicit form in coordinate space is given by Eq.(9) and (10). Here, we present several useful equations:

where \hat{L}_z is the operator of orbital angular momentum. As a result, the right hand side of Eq.(A4) can be evaluated as follows:

$$\left\langle \mathbf{r}_{\perp} \left| \left[\left(E + \mu + \Omega \cdot \hat{J}_{z} \right)^{2} - (k_{z})^{2} - \vec{\pi}_{\perp}^{2} - M_{f}^{2} \right]^{-1} \right| \mathbf{r}_{\perp}' \right\rangle$$

$$= \sum_{n} \int k_{t} dk_{t} \left\langle \mathbf{r}_{\perp} \left| \left[\left(E + \mu + \Omega \cdot \hat{J}_{z} \right)^{2} - (k_{z})^{2} - \vec{\pi}_{\perp}^{2} - M_{f}^{2} \right]^{-1} \right| nk_{t} \right\rangle \langle nk_{t} \mid \mathbf{r}_{\perp}' \rangle$$

$$= \sum_{n} \int k_{t} dk_{t} \left\langle \mathbf{r}_{\perp} \left| \left[\left(E + \mu + \Omega \cdot \hat{L}_{z} \right)^{2} + 2(E + \mu)\Omega \cdot S_{z} + \Omega^{2} (S_{z})^{2} - k_{z}^{2} - \vec{\pi}_{\perp}^{2} - M_{f}^{2} \right]^{-1} \right| nk_{t} \right\rangle \langle nk_{t} \mid \mathbf{r}_{\perp}' \rangle$$

$$= \sum_{n} \int_{0}^{+\infty} k_{t} dk_{t} J_{n} (k_{t}r) J_{n} (k_{t}r') e^{in(\theta - \theta')} \left[(E + \mu + \Omega n)^{2} + 2(E + \mu)\Omega S_{z} + \Omega^{2} \cdot \frac{1}{4} - k_{z}^{2} - k_{t}^{2} - M_{f}^{2} \right]^{-1}$$

$$(A6)$$

where $S_z = \frac{i}{2}\gamma^1\gamma^2$ is the spin angular momentum term. And we have insert the completeness condition in the second line of Eq.(A6). Now, it is easy to obtain Eq.(11) by means of projection operator $\mathcal{P}_{\pm} = \frac{1}{2} (1 \pm i \gamma^1 \gamma^2)$. By inserting $I_4 = \mathcal{P}_- + \mathcal{P}_+$, the summation in Eq.(A6) can be replaced by:

$$\sum_{n} \frac{J_{n}(k_{t}r) J_{n}(k_{t}r') e^{in(\theta-\theta')} \mathcal{P}_{+} + J_{n+1}(k_{t}r) J_{n+1}(k_{t}r') e^{i(n+1)(\theta-\theta')} \mathcal{P}_{-}}{\left[E + \mu + \left(n + \frac{1}{2}\right)\Omega\right]^{2} - k_{z}^{2} - k_{t}^{2} - M_{f}^{2}}.$$
(A7)

Finally, we can calculate Eq.(A4) by acting the operators $\hat{J}_{z(\mathbf{r}_{\perp})}\gamma^{0}, \vec{\pi}_{\mathbf{r}_{\perp}} \cdot \vec{r}_{\perp}$ and $k_{z}\gamma^{3}$ on Eq.(A7).

Appendix B: Spectral functions

The explicit form and the derivation of spectral functions can be found in Ref.[19, 32]. In our previous work, we have utilized the properties of Bessel functions, i.e. $J_n(0) = 0$ for $n \neq 0$ and $J_0(0) = 1$. As a consequence, the infinite summation is reduced to finite terms. Then, Eq.(18) can be evaluated at zero and finite temperature. For saving space in this manuscript, we merely present one component of polarization functions. For example, the imaginary part of 00-component is [19]:

$$\operatorname{Im} \Pi^{00}(\omega, \vec{q}) = -\frac{\pi}{2} N_f N_c \sum_{\eta=\pm 1} \int \frac{d^3 \vec{p}}{(2\pi)^3} \frac{1}{E_p E_k} \left\{ \left[E_p E_k + \vec{p} \cdot \vec{k} + M_f^2 \right] \left[f \left(E_p - \mu - \frac{\eta \Omega}{2} \right) + f \left(E_p + \mu - \frac{\eta \Omega}{2} \right) \right] \right\}$$

$$\times \left[\delta \left(\omega + E_p - E_k \right) - \delta \left(\omega - E_p + E_k \right) \right] + \left[E_p E_k - \vec{p} \cdot \vec{k} - M_f^2 \right] \delta \left(\omega - E_p - E_k \right) \right]$$

$$\times \left[1 - f \left(E_p - \mu - \frac{\eta \Omega}{2} \right) - f \left(E_p + \mu - \frac{\eta \Omega}{2} \right) \right] \right\}.$$
(B1)

where k = p + q and $E_k = \sqrt{\vec{k}^2 + M_f^2}$. Here, N_f and N_c are the flavor number and color number in the quark

loop, and f(x) is Fermi-Dirac distribution functions with a finite temperature T. Other components can be obtained similarly.

- I. Arsene *et al.* (BRAHMS), Quark gluon plasma and color glass condensate at RHIC? The Perspective from the BRAHMS experiment, Nucl. Phys. A **757**, 1 (2005), arXiv:nucl-ex/0410020.
- [2] F. Becattini, F. Piccinini, and J. Rizzo, Angular momentum conservation in heavy ion collisions at very high energy, Phys. Rev. C 77, 024906 (2008), arXiv:0711.1253 [nucl-th].
- Y. Jiang, Z.-W. Lin, and J. Liao, Rotating quark-gluon plasma in relativistic heavy ion collisions, Phys. Rev. C 94, 044910 (2016), [Erratum: Phys.Rev.C 95, 049904 (2017)], arXiv:1602.06580 [hep-ph].
- [4] W.-T. Deng and X.-G. Huang, Vorticity in Heavy-Ion Collisions, Phys. Rev. C 93, 064907 (2016), arXiv:1603.06117 [nucl-th].
- [5] D. E. Kharzeev, L. D. McLerran, and H. J. Warringa, The Effects of topological charge change in heavy ion collisions: 'Event by event P and CP violation', Nucl. Phys. A 803, 227 (2008), arXiv:0711.0950 [hep-ph].
- [6] V. Skokov, A. Y. Illarionov, and V. Toneev, Estimate of the magnetic field strength in heavy-ion collisions, Int. J. Mod. Phys. A 24, 5925 (2009), arXiv:0907.1396 [nucl-th].
- [7] W.-T. Deng and X.-G. Huang, Event-by-event generation of electromagnetic fields in heavy-ion collisions, Phys. Rev. C 85, 044907 (2012), arXiv:1201.5108 [nucl-th].
- [8] S. Acharya *et al.* (ALICE), Evidence of Spin-Orbital Angular Momentum Interactions in Relativistic Heavy-Ion Collisions, Phys. Rev. Lett. **125**, 012301 (2020), arXiv:1910.14408 [nucl-ex].
- [9] M. S. Abdallah *et al.* (STAR), Pattern of global spin alignment of ϕ and K^{*0} mesons in heavy-ion collisions 10.1038/s41586-022-05557-5 (2022), arXiv:2204.02302 [hep-ph].
- [10] X.-L. Sheng, L. Oliva, and Q. Wang, What can we learn from the global spin alignment of ϕ mesons in heavy-ion collisions?, Phys. Rev. D **101**, 096005 (2020), [Erratum: Phys.Rev.D 105, 099903 (2022)], arXiv:1910.13684 [nucl-th].
- [11] X.-L. Sheng, S.-Y. Yang, Y.-L. Zou, and D. Hou, Mass splitting and spin alignment for φ mesons in a magnetic field in NJL model, (2022), arXiv:2209.01872 [nucl-th].
- [12] Y. Jiang and J. Liao, Pairing Phase Transitions of Matter under Rotation, Phys. Rev. Lett. **117**, 192302 (2016), arXiv:1606.03808 [hep-ph].
- [13] S. P. Klevansky and R. H. Lemmer, Chiral symmetry restoration in the Nambu-Jona-Lasinio model with a constant electromagnetic field, Phys. Rev. D 39, 3478 (1989).
- [14] K. G. Klimenko, Three-dimensional Gross-Neveu model at nonzero temperature and in an external magnetic field, Z. Phys. C 54, 323 (1992).
- [15] V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, Dimensional reduction and catalysis of dynamical symmetry breaking by a magnetic field, Nucl. Phys. B 462, 249 (1996), arXiv:hep-ph/9509320.
- [16] V. A. Miransky and I. A. Shovkovy, Quantum field theory in a magnetic field: From quantum chromodynamics

to graphene and Dirac semimetals, Phys. Rept. 576, 1 (2015), arXiv:1503.00732 [hep-ph].

- [17] H.-L. Chen, K. Fukushima, X.-G. Huang, and K. Mameda, Analogy between rotation and density for Dirac fermions in a magnetic field, Phys. Rev. D 93, 104052 (2016), arXiv:1512.08974 [hep-ph].
- [18] H.-L. Chen, X.-G. Huang, and J. Liao, QCD phase structure under rotation, Lect. Notes Phys. 987, 349 (2021), arXiv:2108.00586 [hep-ph].
- [19] M. Wei, C. A. Islam, and M. Huang, Production rate and ellipticity of lepton pairs from a rotating hot and dense QCD medium, Phys. Rev. D 105, 054014 (2022), arXiv:2111.05192 [hep-ph].
- [20] X. Wang, M. Wei, Z. Li, and M. Huang, Quark matter under rotation in the NJL model with vector interaction, Phys. Rev. D 99, 016018 (2019), arXiv:1808.01931 [hepph].
- [21] S. Ebihara, K. Fukushima, and K. Mameda, Boundary effects and gapped dispersion in rotating fermionic matter, Phys. Lett. B 764, 94 (2017), arXiv:1608.00336 [hep-ph].
- [22] M. N. Chernodub and S. Gongyo, Effects of rotation and boundaries on chiral symmetry breaking of relativistic fermions, Phys. Rev. D 95, 096006 (2017), arXiv:1702.08266 [hep-th].
- [23] M. N. Chernodub, Inhomogeneous confining-deconfining phases in rotating plasmas, Phys. Rev. D 103, 054027 (2021), arXiv:2012.04924 [hep-ph].
- [24] Y. Fujimoto, K. Fukushima, and Y. Hidaka, Deconfining Phase Boundary of Rapidly Rotating Hot and Dense Matter and Analysis of Moment of Inertia, Phys. Lett. B 816, 136184 (2021), arXiv:2101.09173 [hep-ph].
- [25] L. Wang, Y. Jiang, L. He, and P. Zhuang, Local suppression and enhancement of the pairing condensate under rotation, Phys. Rev. C 100, 034902 (2019), arXiv:1901.00804 [nucl-th].
- [26] Y. Chen, D. Li, and M. Huang, Inhomogeneous chiral condensation under rotation in the holographic QCD, Phys. Rev. D 106, 106002 (2022), arXiv:2208.05668 [hepph].
- [27] M. N. Chernodub and S. Gongyo, Interacting fermions in rotation: chiral symmetry restoration, moment of inertia and thermodynamics, JHEP 01, 136, arXiv:1611.02598 [hep-th].
- [28] Z. Zhang, C. Shi, X.-T. He, X. Luo, and H.-S. Zong, Chiral phase transition inside a rotating cylinder within the Nambu–Jona-Lasinio model, Phys. Rev. D 102, 114023 (2020), arXiv:2012.01017 [hep-ph].
- [29] Z. Zhang, C. Shi, X. Luo, and H.-S. Zong, Chiral phase transition in a rotating sphere, Phys. Rev. D 101, 074036 (2020), arXiv:2003.03765 [nucl-th].
- [30] S. Chen, J. Zhao, and P. Zhuang, Charmonium transition in electromagnetic and rotational fields, Phys. Rev. C 103, L031902 (2021), arXiv:2005.08473 [nucl-th].
- [31] M. He, R. J. Fries, and R. Rapp, Heavy-Quark Diffusion and Hadronization in Quark-Gluon Plasma, Phys. Rev. C 86, 014903 (2012), arXiv:1106.6006 [nucl-th].

- [32] M. Wei, Y. Jiang, and M. Huang, Mass splitting of vector mesons and spontaneous spin polarization under rotation *, Chin. Phys. C 46, 024102 (2022), arXiv:2011.10987 [hep-ph].
- [33] A. Vilenkin, QUANTUM FIELD THEORY AT FINITE TEMPERATURE IN A ROTATING SYSTEM, Phys. Rev. D 21, 2260 (1980).
- [34] S. P. Klevansky, The Nambu-Jona-Lasinio model of quantum chromodynamics, Rev. Mod. Phys. 64, 649 (1992).
- [35] K. Fukushima, Phase diagrams in the three-flavor Nambu-Jona-Lasinio model with the Polyakov loop, Phys. Rev. D 77, 114028 (2008), [Erratum: Phys.Rev.D 78, 039902 (2008)], arXiv:0803.3318 [hep-ph].
- [36] J. I. Kapusta and C. Gale, *Finite-Temperature Field Theory: Principles and Applications*, 2nd ed., Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2006).
- [37] Z.-T. Liang and X.-N. Wang, Spin alignment of vector mesons in non-central A+A collisions, Phys. Lett. B 629,

20 (2005), arXiv:nucl-th/0411101.

- [38] P. Rehberg, S. P. Klevansky, and J. Hufner, Hadronization in the SU(3) Nambu-Jona-Lasinio model, Phys. Rev. C 53, 410 (1996), arXiv:hep-ph/9506436.
- [39] M. Buballa, NJL model analysis of quark matter at large density, Phys. Rept. 407, 205 (2005), arXiv:hepph/0402234.
- [40] Y.-G. Yang, R.-H. Fang, Q. Wang, and X.-N. Wang, Quark coalescence model for polarized vector mesons and baryons, Phys. Rev. C 97, 034917 (2018), arXiv:1711.06008 [nucl-th].
- [41] K. Xu, S. Shi, H. Zhang, D. Hou, J. Liao, and M. Huang, Extracting the magnitude of magnetic field at freeze-out in heavy-ion collisions, Phys. Lett. B 809, 135706 (2020), arXiv:2004.05362 [hep-ph].
- [42] X.-L. Sheng, Q. Wang, and X.-N. Wang, Improved quark coalescence model for spin alignment and polarization of hadrons, Phys. Rev. D 102, 056013 (2020), arXiv:2007.05106 [nucl-th].