

On the Relations between Fermion Masses and Isospin Couplings in the Microscopic Model

Bodo Lampe

II. Institut für theoretische Physik der Universität Hamburg
Luruper Chaussee 149, 22761 Hamburg, Germany

Abstract

Quark and lepton masses and mixings are considered in the framework of the microscopic model. The most general ansatz for the interactions among tetrons leads to a Hamiltonian H_T involving Dzyaloshinskii-Moriya (DM), Heisenberg and torsional isospin forces. Diagonalization of the Hamiltonian provides for 24 eigenvalues which are identified as the quark and lepton masses. While the masses of the third and second family arise from DM and Heisenberg type of isospin interactions, light family masses are related to torsional interactions among tetrons. Neutrino masses turn out to be special in that they are given in terms of tiny isospin non-conserving DM, Heisenberg and torsional couplings.

The approach not only leads to masses, but also allows to calculate the quark and lepton eigenstates, an issue, which is important for the determination of the CKM and PMNS mixing matrices. Compact expressions for the eigenfunctions of H_T are given. The almost exact isospin conservation of the system dictates the form of the lepton states and makes them independent of all the couplings in H_T . Much in contrast, there is a strong dependence of the quark states on the coupling strengths, and a promising hierarchy between the quark family mixings shows up.

I. Introduction

Our universe according to the microscopic model[1] is a 3-dimensional elastic substrate expanding within some higher dimensional space. The elastic substrate is built from tiny invisible constituents, called tetrons, with bond length about the Planck length and binding energy the Planck energy. Tetrons transform under the fundamental spinor representation of $SO(6,1)$. This representation is 8-dimensional and sometimes called the octonion representation[2, 3].

Details of the approach provide a powerful unified picture for particle physics and cosmology. All physical properties in the universe can be derived from properties of the tetrons. This philosophy is applied here to the Standard Model mass and mixing parameters which are shown to be determined by the interactions among tetrons.

The 24 known quarks and leptons arise as eigenmode excitations of a tetrahedral fiber structure, which is made up from 4 tetrons and extends into 3 extra ‘internal’ dimensions. While the laws of gravity are due to the elastic properties of the tetron bonds[4], particle physics interactions take place within the internal fibers, with the characteristic internal energy being the Fermi scale. All ordinary matter quarks and leptons are constructed as quasiparticle excitations of this internal fiber structure. Since the quasiparticles fulfill Lorentz covariant wave equations, they perceive the universe as a 3+1 dimensional spacetime continuum.

More in detail, the ground state of our universe looks like illustrated in Fig. 1. In this figure the tetrahedrons (=‘fibers’) extend into the 3 extra dimensions. The picture is a little misleading because in the tetron model physical space and the extra (‘internal’) dimensions are assumed to be completely orthogonal. This means the whole game has to be played within a larger, at least 6 dimensional space, 3 physical dimensions and 3 internal ones¹.

Each tetrahedron in Fig. 1 is made up from 4 tetrons, depicted as dots. With respect to the decomposition of $SO(6,1) \rightarrow SO(3,1) \times SO(3)$ into the (3+1)-dimensional base space and the 3-dimensional internal space, a tetron Ψ possesses spin $\frac{1}{2}$ and isospin $\frac{1}{2}$. This means it can rotate both in physical space and in the extra di-

¹There are indications that the system actually lives in 7+1 dimensions instead of 6+1; this however does not play a role in the calculations presented on the following pages.

mensions, and corresponds to the fact that Ψ decomposes into an isospin doublet $\Psi = (U, D)$ of two ordinary $SO(3,1)$ Dirac fields U and D .

$$8 \rightarrow (1, 2, 2) + (2, 1, 2) = ((1, 2) + (2, 1), 2) \quad (1)$$




Figure 1: The global ground state of the universe after the electroweak symmetry breaking has occurred, considered at Planck scale distances. The big black arrow represents 3-dimensional physical space. Before the symmetry breaking the isospin vectors are directed randomly, thus exhibiting a local $SU(2)$ symmetry, but once the temperature drops below the Fermi scale Λ_F , they become ordered into a repetitive tetrahedral structure, thereby spontaneously breaking the initial $SU(2)$. Note that the SM Higgs vev is related to the length of the aligned isospin vectors. Quarks and leptons glide on this background as quasiparticle excitations. The background has the properties of a Lorentz ether and is thereby not in conflict with Michelson-Morley type of experiments.

Why this tetrahedral structure? It is needed in order to explain the observed quark and lepton spectrum, which means to get exactly 24 excitation states with the correct multiplet structure². In fact, the tetrahedral symmetry is rather uniquely determined by this condition[7]. As shown below, under reasonable assumptions on the tetron dynamics, the numerical mass values of quarks and leptons can be correctly reproduced.

The arrows in Fig. 1 denote the isospins, i.e. internal spin vectors of the tetrons.

²The quark triplets are triplets under tetrahedral transformations at this point. For the question how to interpret them as $SU(3)$ color triplets one may consult Appendix B.

More precisely, each arrow stands for two(!) vectors $\langle \vec{Q}_L \rangle = \langle \vec{Q}_R \rangle$ where[8]

$$\vec{Q}_L = \frac{1}{4}\Psi^\dagger(1 - \gamma_5)\vec{\tau}\Psi \quad \vec{Q}_R = \frac{1}{4}\Psi^\dagger(1 + \gamma_5)\vec{\tau}\Psi \quad (2)$$

and $\langle \rangle$ denotes the ground state/vacuum expectation values (i.e. after the SSB). In other words, the ground state values $\langle \vec{Q}_{Li} \rangle$ and $\langle \vec{Q}_{Ri} \rangle$ are assumed to be equal on each tetrahedral site $i=1,2,3,4$ and given by one of the arrows in Fig. 1. $\vec{\tau}$ are the internal spin Pauli matrices.

According to (1) the tetron representation 8 contains both particle and antiparticle degrees of freedom. \vec{Q}_L and \vec{Q}_R cover 6 of its 8 dof³. Furthermore, \vec{Q}_L and \vec{Q}_R are particularly useful to handle because quantum mechanically they commute with each other[8]. As turns out, the interactions of these internal spins play an essential role for particle physics and for electroweak symmetry breaking.

Due to the pseudovector property of the isospin vectors their tetrahedral symmetry group actually is a Shubnikov group[9]. This means, while the coordinate symmetry is S_4 , the arrangement of isospin vectors respects the tetrahedral Shubnikov symmetry[10]

$$G_4 := A_4 + CPT(S_4 - A_4) \quad (3)$$

where $A_4(S_4)$ is the (full) tetrahedral symmetry group and CPT the usual CPT operation except that P is the parity transformation in physical space only. Since the elements of $S_4 - A_4$ contain an implicit factor of internal parity, the symmetry (3) certifies CPT invariance of the local ground state in the full of R^{6+1} .

Note that in the situation depicted in Fig. 1 the $SU(2)$ symmetry breaking has already occurred, because the isospins are aligned between all the tetrahedrons. Before the symmetry breaking, which means above a certain temperature, isospins are directed randomly, corresponding to a local $SU(2) \times U_1$ symmetry⁴, but when the universe cools down, there is a phase transition, and the isospins freeze into the aligned structure, breaking the $SU(2)$ symmetry to the discrete ‘family group’

³The remaining 2 dof correspond to the ‘densities’ $\Psi^\dagger\Psi$ and $\Psi^\dagger\gamma_5\Psi$ whose fluctuations actually are dark matter candidates[1]. Other dark matter candidates in the model arise from the fact that its Higgs sector corresponds to an inert 2HDM[7, 17].

⁴Weak parity violation, vulgo the appearance of index L in $SU(2)_L$, arises from the chirality of the isospin tetrahedrons. This, as well as the Z- γ mixing, is discussed in detail in [1].

G_4 . And the important point to note is, this temperature can be identified with the Fermi scale[7]. Moreover, the remaining symmetry $G_4 \times U(1)_{em}$ is valid down to the lowest energies.

As elaborated in the following sections, the mathematical treatment of the excitations arising from the isospin interactions (7) and (27) is similar to that of magnons in ordinary magnetism. However, the physics is quite different, because in contrast to magnons the isospin excitations are pointlike, i.e. they can exist within one point of physical space, because they are vibrations of the isospin vectors of the tetrons within one internal tetrahedron. Note, that these internal vibrations are spin- $\frac{1}{2}$ because they inherit their fermion nature from the fermion property of the vibrating tetrons in their 3-dimensional physical ‘base space’.

Similar to magnons, the vibrations can move in physical space[9] by hopping from one tetrahedron to another (particle picture) or propagating as quasiparticle waves through physical space (wave picture). Thus, although it can exist at one point of physical space, when one tries to exactly measure its location, for example by scattering with another particle, the excitation will start to move on physical space, and this movement will follow a wave equation which naturally has an uncertainty in it according to Schwarz’ inequality. Planck’s constant enters this uncertainty because the whole process is taking place on a discrete system with Planck length ‘lattice constant’ and Planck energy ‘response energy’.

The second part of the article deals with the mixing of families and the question to what extent it can be deduced from tetron ideas. Since as much as 8 of the 19 free SM parameters arise from those mixings, there have been many attempts to reduce this freedom by BSM ideas. That is the reason why in the literature a lot of suggestions for relations among the CKM resp PMNS matrix elements can be found, e.g. [23, 24, 25, 26], mostly on the basis of assumptions on additional discrete symmetries, from which such relations then are derived.

II. Quark and Lepton Masses from the Interactions of Isospins

The SM SSB being realized by an alignment of the tetron isospins, it is not surprising that the masses of quarks and leptons, and thus the SM Yukawa couplings are determined by the interactions among those isospins. The simplest interaction

Hamiltonian between isospin vectors of 2 tetrons i and j looks like

$$H = -J \vec{Q}_i \vec{Q}_j \quad (4)$$

So it has the form of a Heisenberg interaction - but for isospins, not for spins. The coupling J may be called an ‘isomagnetic exchange coupling’. Note that the language of magnetism often is used in this paper, although interactions of isospins and not of spins are considered. Note further that isospin is not an abstract symmetry here, but corresponds to real rotations in the 3 extra dimensions.

In reality, the Hamiltonian H is more complicated than (4), for several reasons:

- There are inner- and inter-tetrahedral interactions of isospins, i.e. within the same and with a neighboring tetrahedron. The inner ones must have an energy minimum at the tetrahedral angle $\theta = \theta_{tet} = \arccos(-\frac{1}{3})$, while the inter ones correspond to a minimum at the collinear configuration $\theta = 0$, cf. Fig. 1.
- The appearance of antitetron degrees of freedom should be accounted for by using interactions both of \vec{Q}_L and \vec{Q}_R defined in (2) instead of \vec{Q} in (4). The Heisenberg Hamiltonian for the interaction between 2 tetrons i and j then reads:

$$H_H = -J_{LL} \vec{Q}_{Li} \vec{Q}_{Lj} - J_{LR} \vec{Q}_{Li} \vec{Q}_{Rj} - J_{RR} \vec{Q}_{Ri} \vec{Q}_{Rj} \quad (5)$$

As shown later in Sect. IV, the three couplings J_{LL} , J_{LR} and J_{RR} can be roughly associated to the masses of the second family fermions, m_c , m_μ and m_s , respectively.

- In addition to the Heisenberg Hamiltonian (5) Dzyaloshinskii interactions[11] are to be considered. They will be shown to give the dominant mass contributions to the heavy family. As well known, the form of the DM couplings \vec{D}_{ab} in (27) is restricted by the ground state symmetry through the so-called Moriya rules[12]. Applying these rules to the given tetrahedral structure, the DM Hamiltonian can be shown to have the form (29).

- Heisenberg and DM terms do not contribute at all to the masses m_e , m_u and m_d of the first family. Therefore, small torsional interactions are introduced in Sect. V. They are characterized by the exerting torques $dQ_{L,R}/dt$ being proportional to the isospins $Q_{L,R}$ themselves, cf. Eq. (42).

- The masses of the neutrinos are yet another story. While the interactions discussed so far are isospin conserving and leave the neutrinos massless, neutrino masses can arise only from isospin violation. Generation of these masses will be discussed in

Sect. VI, and a physical explanation for the origin of the isospin violation will be given.

The DM-couplings K_{LL} , K_{LR} and K_{RR} introduced in (29) are much larger than both Heisenberg and torsional interactions and essentially determine the masses m_τ , m_b and m_t of the third family particles. K_{LL} will be shown to be particularly large. It gives the dominant contribution to the top mass and is the main source for the parallel arrangement of isospins and the SU(2) SSB.

All the types of interaction mentioned above contribute to the angular dependence of the energy of 2 tetron isospins at angle θ which is basically of the form

$$E = A + B \cos(\theta) + C \cos^2(\theta) \quad (6)$$

where A , B and C are determined by the Heisenberg, torsional and DM coupling strengths. (For example, the Heisenberg coupling J_{LL} in (5) concerns $\theta_{LL} = \angle(\vec{Q}_{Li}, \vec{Q}_{Lj})$ and gives a contribution to B_{LL} only.) Altogether, they fix the *relative* directions of the ground state isospins at the energy minimum, both locally and globally in the way depicted in Fig. 1 - whereas the *absolute* arrangement of the tetrahedrons is spontaneous.

Furthermore, they give rise to fermionic excitations which are to be interpreted as quarks and leptons. Masses can then be calculated using the Hamiltonians discussed above. Indeed, 24 eigen energies arise from the tetrahedral configuration by diagonalizing equations for the isospin torque which are generically of the form

$$\frac{d\vec{Q}}{dt} = i[H, \vec{Q}] \quad (7)$$

While the masses correspond to the eigenvalues, CKM and PMNS mixings can be deduced from the eigenvectors. This point will be discussed in Sects. VII ff.

More in detail, the quarks and leptons are vibrations δ of the isospin vectors \vec{Q}_{Li} and \vec{Q}_{Ri} of the tetrons i at sites $i = 1, 2, 3, 4$, i.e. fluctuations of the ground state values within one tetrahedron.

$$\vec{Q}_{Li} = \langle \vec{Q}_{Li} \rangle + \vec{\delta}_{Li} \quad \vec{Q}_{Ri} = \langle \vec{Q}_{Ri} \rangle + \vec{\delta}_{Ri} \quad (8)$$

where

$$\langle \vec{Q}_{Li} \rangle = \frac{1}{4} \langle \Psi^\dagger (1 - \gamma_5) \vec{\tau} \Psi \rangle \quad \langle \vec{Q}_{Ri} \rangle = \frac{1}{4} \langle \Psi^\dagger (1 + \gamma_5) \vec{\tau} \Psi \rangle \quad (9)$$

are the ground state radial isospin vectors of a tetrahedron in Fig. 1 assumed to be pointing outward

$$\langle \vec{Q}_{Li} \rangle = \langle \vec{Q}_{Ri} \rangle = \vec{e}_r \quad (10)$$

In the present model, gauge and Higgs bosons are constructed as excitations of a single pair of a tetron and an antitetron belonging to 2 neighboring tetrahedrons⁵. As excitations of single pairs they do not feel their tetrahedral environment, which means they transform under isospin but do not appear in multiplets of the tetrahedral group.

One may rightfully ask whether the Higgs mass and in general the parameters of the SM Higgs potential can be determined from similar considerations as the fermion masses. I do not have a complete answer to this question, but will draw below an analogy with a ferromagnet. Actually, this is a very meaningful analogy because in the microscopic model the ‘isomagnetic’ alignment between isospin vectors of neighboring tetrahedrons plays an important role for the SSB of SU(2).

Within the SM the mass of the Higgs particle is given by

$$m_H = \sqrt{2}|\mu| \quad (11)$$

where μ is defined through the potential of the Higgs doublet ϕ

$$V(\phi) = -\mu^2 \phi^\dagger \phi + \lambda(\phi^\dagger \phi)^2 \quad (12)$$

It is instructive to compare this to the free energy of a ferromagnet given in terms of its magnetization M

$$F = U - TS = \alpha \vec{M}^2 + \beta \vec{M}^4 \quad (13)$$

Here α gets a negative contribution from the Heisenberg interaction energy U between 2 spins and a positive contribution from the temperature/entropy term. The transition energy is then obtained from the condition $\alpha = 0$ as $kT_C = 2U$, i.e. the transition energy is directly related to the interaction between 2 spins.

⁵Note that these excitations are bosons because they involve tensor products of 2 tetron fields over 2 points in physical space.

This may be compared to the present model: for the tetrahedrons to align as in Figure 1, pairs of isospin vectors from neighboring tetrahedrons must align. As described after (36), the second term in (36) is a Heisenberg type of ‘ferromagnetic’ coupling with strength $\frac{2}{3}K_{LL} + J_{LL}$ (where $K_{LL} \gg J_{LL}$). This means the coupling K_{LL} is the dominant driving source for the SSB alignment, and at the same time according to (34) the coupling K_{LL} gives the dominant contribution to the top mass. One is thus led to the conclusion that m_H and m_t are roughly of the same order of magnitude and, more specifically, that the Higgs mass should be about two thirds of the top mass.

III. Physical Origin of the Isospin Interactions

This chapter is devoted to the question how the Heisenberg, DM and torsional interactions introduced in the last section can be understood from a more fundamental interaction among tetrons.

First of all, the interested reader should remember that Heisenberg used the Heitler-London results for the hydrogen molecule to understand the phenomenon of ferromagnetism. Heisenberg showed[13] that ferromagnetism is a quantum effect arising from the Pauli principle, more precisely, from the large exchange energies due to the overlap of the antisymmetrized electron wave functions.

The situation here is in principle similar - but in practice somewhat more complicated, because one deals with 6 dimensions with 2 types of rotations: spin and isospin.

In the non-relativistic limit $SO(6, 1) \rightarrow SO(6)$ the tetron representation 8 of $SO(6,1)$ reduces to

$$SO(6, 1) \rightarrow SO(6) \tag{14}$$

$$8 \rightarrow 4 + \bar{4} \tag{15}$$

where 4 is the spinor representation of $SO(6)$ and $\bar{4}$ its complex conjugate. Since the universal covering of $SO(6)$ is given by $SU(4)$, the 4-representation actually is the fundamental representation of $SU(4)$. This representation contains the spin $(\pm\frac{1}{2})$ and isospin $(\pm\frac{1}{2})$ of the tetron, while the $\bar{4}$ -representation corresponds to the antitetron degrees of freedom.

Within a non-relativistic quantum mechanics the binding energy between a tetron and an (anti)tetron should generally be calculable from the expectation value

$$E_F = \int d^6x_i d^6x_j \Phi_F^*(x_i, x_j) U_F(|x_i - x_j|) \Phi_F(x_i, x_j) \quad (16)$$

of a non-relativistic potential U_F , where Φ is the complete wavefunction for the tetron-(anti)tetron system and F denotes its combined list of quantum numbers, i.e. spin, isospin, orbital angular momentum etc.

Φ_F may be approximated by a sum of products of two 1-tetron wave functions concentrated at the two tetrahedral sites x_i and x_j . Antisymmetrization of this sum will lead E_F to consist of two terms, the classical ‘direct’ integral D_F and the quantum mechanical exchange contribution J_F .

$$E_F = D_F + J_F \quad (17)$$

While D_F determines the elastic binding among tetrons and thus the gravitational properties of the substrate, the exchange integral J_F can be used to understand the isospin interactions and thus the phenomena of particle physics. Actually, as seen below, J_F is directly related to the ‘isomagnetic’ Heisenberg and DM couplings J and K defined in the last section.

If one assumes the single tetron wave functions to be fairly localized at their tetrahedral sites, there is a hierarchy $|J_F| \ll |D_F|$. This is different from ordinary 3-dimensional ferromagnetism and is even enhanced by the 12-dimensional integration in (16), through which any overlap contribution becomes strongly suppressed as compared to a direct one. In the extreme case of delta functions, D_F reflects the form of the potential, while J_F vanishes. In the general case, D_F will still be much larger than J_F . For example, assuming the single tetron wave function to fall off by a factor of 10 at half the distance between the 2 sites i and j , J_F will be smaller than D_F roughly by a factor of 10^{-12} . This, en passant, is the way the hierarchy between the Planck scale and the Fermi scale can be understood within the tetron approach. The item has been discussed more thoroughly in [4].

One may ask how the potential U_F transforms under $SU(4)$. Since the energy must be a singlet, one has to have

$$(4 + \bar{4}) \times R_U \times (4 + \bar{4}) = 1 + \dots \quad (18)$$

where R_U is the representation under which U_F transforms. Since $4 \times \bar{4} = 1 + 15$ and $4 \times 4 = 6 + 10$ and $15 \times 15 = 1 + \dots$ and $6 \times 6 = 1 + \dots$ [2], it follows that U_F is either a scalar U_1 , an adjoint $U_{15}^a \lambda^a$, $a=1, \dots, 15$, or a vector $U_6^i e^i$, $i=1, \dots, 6$, where λ^a are the generators of $SU(4)$ and e^i are vectors which span 6-dimensional space. U_1 and U_{15} describe interactions among a tetron and an antitetron and U_6 is a tetron-tetron interaction.

In the present context, where the tetrahedrons are completely orthogonal to physical space, spin and isospin essentially decouple from each other, and the above analysis may be strongly simplified, in the following way: instead of (15) one may consider

$$SO(6,1) \rightarrow SO(3)_{spin} \times SO(3)_{isospin} \quad (19)$$

$$8 \rightarrow (1, 2) + (1, 2) + (2, 1) + (2, 1) \quad (20)$$

Since the ordinary spin of the tetrons (i.e. the spin in physical space) is irrelevant for the internal interactions, it is enough to look for $SO(3)_{isospin}$ singlets in $2 \times R \times 2 = 1 + \dots$, which implies $R = 1$ or $R = 3$, i.e. only an isospin singlet or a triplet potential V_1 or V_3 are allowed for the isospin interactions among tetrons.

V_1 and V_3 may be considered as part of the above $SO(6)$ potentials U_1 and U_{15} and induced by them within the $SO(3)_{isospin}$ fibers. Alternatively, V_1 and V_3 can also be shown to arise in the relativistic framework, i.e. sticking to the original octonion representation 8 of $SO(6,1)$ instead of using (15). Namely, a relativistic potential W_7 is allowed that transforms as 7 under $SO(6,1)$, and the product[2]

$$8 \times 7 \times 8 = 1 + 7 + 7 + 21 + 21 + 27 + 35 + 35 + 105 + 189 \quad (21)$$

contains a singlet.

W_7 may well be a gauge potential and the basis for the fundamental tetron interaction. Furthermore, V_1 and V_3 are part of W_7 due to

$$SO(6,1) \rightarrow SO(3)_{spin} \times SO(3)_{isospin} \quad (22)$$

$$7 \rightarrow (1, 1) + (1, 3) + (3, 1) \quad (23)$$

where V_1 transforms as (1,1) and

$$V_3 = V_3^a \tau^a \quad (24)$$

as the isospin triplet (1,3)⁶.

While the Heisenberg interaction $\sim Q_i^a Q_j^a$ is associated to the singlet potential V_1 in the usual way[13], DM terms $\sim \epsilon^{abc} Q_i^b Q_j^c$ in (27) arise from the V_3 contributions. This can be shown by inserting the completeness relation for Pauli matrices

$$\delta_{sv}\delta_{ut} = 2\tau_{st}^a \tau_{uv}^a + \frac{1}{2}\delta_{st}\delta_{uv} \quad (25)$$

into the V_3 -exchange integral and afterwards noting that the factor of τ^a in (24) can be merged with one of the factors τ in (25) via

$$\tau^a \tau^b = i\epsilon^{abc} \tau^c + \delta^{ab} \quad (26)$$

The ϵ tensor part in (26) then directly yields the 'antisymmetric exchange' (=DM) contribution (27).

IV. Dzyaloshinskii Masses for the Heavy Family; Heisenberg Masses for the Second Family

My presentation of the mass calculations begins with the Dzyaloshinskii-Moriya (DM) coupling, firstly because it is the dominant isospin interaction and secondly it gives masses only to the third family, i.e. to top, bottom and τ , while leaving all other quarks and leptons massless.

Among all the fermion masses the top quark mass is by far the largest and is of the order of the Fermi scale. As turns out, this is no accident, but has to do with the largeness of the relevant DM coupling.

In the simplest version the isospin DM interaction[1, 11] is

$$H_{DM} = -K \sum_{i \neq j=1}^4 \vec{D}_{ij} (\vec{Q}_i \times \vec{Q}_j) \quad (27)$$

⁶In contrast to the suggestion in [7] the SM photon should *not* be assumed to be part of the gauge field W_7 . As explained before, the photon as well as all the other SM gauge bosons are excitations of tetron-antitetron bonds of neighboring tetrahedrons. Nevertheless, they transform under $SO(3) \times SO(3)$, the weak bosons, for example, as (3,3), i.e. they are spin 1 and isospin 1 particles.

to be compared to the Heisenberg interaction (4). The form of the vectors \vec{D}_{ij} is dictated by the tetrahedral symmetry to be [12]

$$\vec{D}_{ij} = \vec{Q}_i \times \vec{Q}_j \quad (28)$$

As explained before, interactions among \vec{Q}_L and \vec{Q}_R have to be considered in order to cover all degrees of freedom. The complete DM Hamiltonian then reads

$$\begin{aligned} H_D = & -K_{LL} \sum_{i \neq j=1}^4 (\vec{Q}_{Li} \times \vec{Q}_{Lj})^2 - K_{LR} \sum_{i \neq j=1}^4 (\vec{Q}_{Li} \times \vec{Q}_{Rj})^2 \\ & - K_{RR} \sum_{i \neq j=1}^4 (\vec{Q}_{Ri} \times \vec{Q}_{Rj})^2 \end{aligned} \quad (29)$$

with DM couplings (= V_3 exchange integrals) K_{LL} , K_{LR} and K_{RR} .

It is convenient to already include at this point the Heisenberg terms

$$H_H = -J_{LL} \sum_{i \neq j=1}^4 \vec{Q}_{Li} \vec{Q}_{Lj} - J_{LR} \sum_{i \neq j=1}^4 \vec{Q}_{Li} \vec{Q}_{Rj} - J_{RR} \sum_{i \neq j=1}^4 \vec{Q}_{Ri} \vec{Q}_{Rj} \quad (30)$$

with V_1 exchange couplings J_{LL} , J_{LR} and J_{RR} . They are smaller than the DM interactions and turn out to give masses both to the second and third family (but not to the first one).

Phenomenologically, the Heisenberg couplings J are typically smaller than 1 GeV, while the DM couplings K are larger than 1 GeV. Altogether, Heisenberg and DM terms provide the most general isotropic and isospin conserving interactions within the internal space. Apart from that there will only be tiny torsional interactions responsible for the mass of the first family and still smaller isospin violating interactions giving masses to the neutrinos (to be discussed in Sects. V and VI).

The masses m of the corresponding excitations δ defined in (8) arise from the exponents in the vibrations

$$\delta \sim \exp(imt) = \exp(iXt) \quad (31)$$

where X stands for the appropriate linear combination of the isospin couplings J and K introduced in (29) and (30). The combinations X will be obtained from the

torque equations (7), using the angular momentum commutation relations for the isospin vectors[8]

$$[Q_{Ri}^a, Q_{Rj}^b] = i\delta_{ij}\epsilon^{abc}Q_{Ri}^c \quad [Q_{Li}^a, Q_{Lj}^b] = i\delta_{ij}\epsilon^{abc}Q_{Li}^c \quad [Q_{Ri}^a, Q_{Lj}^b] = 0 \quad (32)$$

where $i, j = 1, 2, 3, 4$ count the 4 tetrahedral edges and $a, b, c = 1, 2, 3$ the 3 internal directions(=extra dimensions).

It may be stressed that I have noch undertaken to calculate the couplings J and K in terms of the 12-dimensional V_1 and V_3 exchange integrals as defined in (17) and (16). What is done here, is to use the J and K as free parameters and calculate the masses of the excitations in terms of these couplings. This is the usual approach in magnetic theories, where it often turns out that calculation of integrals like (16) are plagued with large and uncertain corrections. Keeping the couplings as free parameters usually is more rewarding for physical applications.

When carrying out the calculation, care must be taken concerning the unique choice of a quantization axis \vec{Q}_0 [21], because this is the condition under which (32) holds. One may choose one of the tetrahedral edges, e.g.

$$\vec{Q}_0 \equiv \langle \vec{Q}_1 \rangle = \frac{1}{\sqrt{3}}(-1, -1, -1) \quad (33)$$

to define the axis of quantization and then has to rotate the other isospins to this system.

The 24 first order differential equations for dQ/dt arising from H_H and H_D are rather lengthy. In linear approximation they read

$$\begin{aligned} \frac{d\vec{\delta}_{Li}}{dt} &= 2K_{LL}\{\vec{Q}_0 \times \vec{\Delta}_{LLi} + i[-\vec{\Delta}_{LLi} + (\vec{\Delta}_{LLi} \cdot \vec{Q}_0) \vec{Q}_0]\} \\ &+ 2K_{LR}\{\vec{Q}_0 \times \vec{\Delta}_{LRi} + i[-\vec{\Delta}_{LRi} + (\vec{\Delta}_{LRi} \cdot \vec{Q}_0) \vec{Q}_0]\} \\ &+ J_{LL}(\vec{Q}_0 \times \vec{\Delta}_{LLi}) + J_{LR}(\vec{Q}_0 \times \vec{\Delta}_{LLi}) \end{aligned} \quad (34)$$

$$\begin{aligned} \frac{d\vec{\delta}_{Ri}}{dt} &= 2K_{RR}\{\vec{Q}_0 \times \vec{\Delta}_{RRi} + i[-\vec{\Delta}_{RRi} + (\vec{\Delta}_{RRi} \cdot \vec{Q}_0) \vec{Q}_0]\} \\ &+ 2K_{LR}\{\vec{Q}_0 \times \vec{\Delta}_{RLi} + i[-\vec{\Delta}_{RLi} + (\vec{\Delta}_{RLi} \cdot \vec{Q}_0) \vec{Q}_0]\} \\ &+ J_{RR}(\vec{Q}_0 \times \vec{\Delta}_{RRi}) + J_{LR}(\vec{Q}_0 \times \vec{\Delta}_{RLi}) \end{aligned} \quad (35)$$

In these equations $\vec{\delta}_{Li} = \vec{Q}_{Li} - \langle \vec{Q}_{Li} \rangle$ and $\vec{\delta}_{Ri} = \vec{Q}_{Ri} - \langle \vec{Q}_{Ri} \rangle$, $a = 1, 2, 3, 4$, denote the isospin vibrations and the Δ 's are certain linear combinations of them which

will play an important role in discussing isospin conservation in Sect. VI:

$$\begin{aligned}
\vec{\Delta}_{LLi} &= -3\vec{\delta}_{Li} + \sum_{j \neq i} \vec{\delta}_{Lj} \\
\vec{\Delta}_{LRi} &= -3\vec{\delta}_{Li} + \sum_{j \neq i} \vec{\delta}_{Rj} \\
\vec{\Delta}_{RLi} &= -3\vec{\delta}_{Ri} + \sum_{j \neq i} \vec{\delta}_{Lj} \\
\vec{\Delta}_{RRi} &= -3\vec{\delta}_{Ri} + \sum_{j \neq i} \vec{\delta}_{Rj}
\end{aligned} \tag{36}$$

Eqs. (34) and (35) are the basis of the Mathematica program included in Appendix A and correspond to a 24×24 eigenvalue problem which - after the SSB - leads to 6 singlet and 6 triplet states of the Shubnikov group (3), the latter ones each consisting of 3 degenerate eigenstates (corresponding to three colors, cf. Appendix B).

After diagonalization one obtains the following results: the first family excitations are still massless at this point, but will get masses from the torsional interactions to be discussed in the next section. The DM exchange coupling K_{LL} is consistently of the order of the transition energy Λ_F and the DM and Heisenberg couplings can be accommodated to reproduce the third and second family masses.

Namely, assuming the DM couplings K to dominate over the Heisenberg couplings J , one can prove the following approximate relations

$$\begin{aligned}
m_t &= 4K_{LL} + O(J) & m_\tau &= \frac{3}{2}K_{LR} + O(J) & m_b &= 4K_{RR} + O(J) \\
m_c &= J_{LL} & m_\mu &= \frac{3}{2}J_{LR} & m_s &= J_{RR}
\end{aligned} \tag{37}$$

In this approximation, the masses of quarks and leptons arise from different isospin interaction terms in (29) and (30), each mass associated essentially to one of the interactions.

Because of the DM dominance one may say that a single tetrahedron of isospin vectors is a ‘DM isomagnet’.

Due to the tetrahedral ‘star’ configuration of the 4 isospin vectors pointing outward, it may also be called a ‘frustrated’ isomagnet[22] based on isospin interactions with ‘antiferromagnetic’ couplings.

There is, however, a different interpretation arising from (29) and (30), where one attains attraction among isospins instead of frustration, and furthermore both inner- and inter-tetrahedral interactions turn out to be of order Λ_F . Namely, there is a Hamiltonian for the interaction between 2 isospins \vec{Q}_i and \vec{Q}_j with minimum energy at the tetrahedral angle $\theta_{tet} = \arccos(-\frac{1}{3})$, thus stabilizing the tetrahedral 'star' arrangement. As compared to (4) and (27) this Hamiltonian has the form

$$H \sim \sum_{i \neq j=1}^4 \vec{Q}_i \vec{Q}_j - \frac{3}{2} \sum_{i \neq j=1}^4 (\vec{Q}_i \times \vec{Q}_j)^2 \quad (38)$$

Since the Heisenberg term is $\sim \cos(\theta)$ and the DM-term involves $\sin(\theta)$, their linear combination (38) can be shown to have a minimum at θ_{tet} . One can then rewrite the top (K_{LL}) and charm (J_{LL}) mass part of the Hamiltonian $H_H + H_D$ eqs. (29) and (30) as a sum of 2 contributions

$$\frac{2}{3}K_{LL} \left[\sum_{i \neq j=1}^4 \vec{Q}_{Li} \vec{Q}_{Lj} - \frac{3}{2} \sum_{i \neq j=1}^4 (\vec{Q}_{Li} \times \vec{Q}_{Lj})^2 \right] - \left(\frac{2}{3}K_{LL} + J_{LL} \right) \sum_{i \neq j=1}^4 \vec{Q}_{Li} \vec{Q}_{Lj} \quad (39)$$

where the first term is assumed to arise from the inner tetrahedral interactions, and the second from the inter ones. Both the inner and inter contributions now are of order $\Lambda_F (\sim K_{LL})$, the inner having a minimum at θ_{tet} thus stabilizing any tetrahedron of isospins, and the inter with coupling $J := \frac{2}{3}K_{LL} + J_{LL}$ being a 'ferromagnetic' Heisenberg interaction which supports the alignment of any 2 neighboring tetrahedrons of isospins.

V. Isospin Conserving Torsion and the Masses of the First Family

In the previous sections it was shown how the heaviness of the third family is related to large DM couplings. Afterwards masses of the quarks and leptons of the second family were obtained from Heisenberg exchange. In this section it will be seen how the small masses of the first family can be obtained from isospin conserving torsional interactions.

It turns out that torsional interactions give contributions to the masses of all families. However, since they are assumed to be small, the 2 heavy families remain dominated by DM and Heisenberg couplings, as given in (37).

The structure of torsional interactions is quite simple. They correspond to a generalization of Hooke's law to rotations, where instead of an exerting force which is proportional to the stretch x there is an exerting torque which is proportional to the stretch angle φ .

$$I \frac{d^2 \varphi}{dt^2} = -C_T^2 \varphi \quad (40)$$

with some constant C_T . The energy of the system is given by

$$E_T = \frac{1}{2} I \left(\frac{d\varphi}{dt} \right)^2 + \frac{1}{2} C_T^2 \varphi^2 \quad (41)$$

with I the moment of inertia.

By differentiation one can see that the second order differential equation (40) is equivalent to $d\varphi/dt = iC_T\varphi$ and thus to the first order equation

$$\frac{dQ}{dt} = iC_T Q \quad (42)$$

where $Q = Id\varphi/dt$ is the angular momentum and dQ/dt the torque.

In the present context (42) is more suitable than (40), because it can be immediately added to the system of differential equations for the \vec{Q}_{Li} and \vec{Q}_{Ri} which was obtained in (34) and (35) for the DM and Heisenberg interactions. Using the notation introduced in (36) one has

$$\frac{d\vec{\delta}_{Li}}{dt} = iC_{LL}\vec{\Delta}_{LLi} + iC_{LR}\vec{\Delta}_{LRi} \quad (43)$$

$$\frac{d\vec{\delta}_{Ri}}{dt} = iC_{LR}\vec{\Delta}_{RLi} + iC_{RR}\vec{\Delta}_{RRi} \quad (44)$$

where the couplings C_{LL} , C_{LR} and C_{RR} generalize C_T to \vec{Q}_L and \vec{Q}_R .

In the formulation (44) care has been taken to maintain isospin conservation as defined in (52). This requirement leads to the appearance of the linear combinations Δ given in (36).

Since (43) and (44) give the only mass contributions to the first family, the C -couplings can be chosen to accommodate the mass of the up quark, down quark and electron, respectively. Namely, one arrives at the mass formulas

$$m_e = 6C_{LR} \quad (45)$$

$$m_u = -2C_{LL} + 3C_{LR} + 2C_{RR} - W_C \quad (46)$$

$$m_d = -2C_{LL} + 3C_{LR} + 2C_{RR} + W_C \quad (47)$$

where

$$W_C := \sqrt{4(C_{LL} + C_{RR})^2 + C_{LR}^2} \quad (48)$$

Then, using the phenomenological values

$$m_e = 0.51 \text{ MeV} \quad m_u = 1.7 \text{ MeV} \quad m_d = 4.7 \text{ MeV} \quad (49)$$

one obtains

$$C_{LR} = 0.085 \text{ MeV} \quad C_{LL} = 1.13 \text{ MeV} \quad C_{RR} = 0.49 \text{ MeV} \quad (50)$$

VI. Neutrino Masses and Isospin Nonconservation

In discussions of neutrino masses there is always the question whether they are of Dirac or Majorana type. Within the tetron model, neutrinos have the same spacetime properties as the other quarks and leptons, because all isospin excitations inherit their $SO(3,1)$ transformation properties from the underlying octonion representation of $SO(6,1)$ - which is Dirac.

This means, neutrinos are special only because of their small masses. In the tetron model small neutrino masses arise in the following way: among the 24 isospin excitations, which are the quarks and leptons, there are always 3 G_4 -singlet modes which are approximately massless. This has to do with the conservation of total isospin. The 3 masses are suppressed because they correspond to the vibrations of the 3 components of the total internal angular momentum vector within one tetrahedron

$$\vec{\Sigma} := \sum_{i=1}^4 (\vec{Q}_{Li} + \vec{Q}_{Ri}) = \sum_{i=1}^4 \vec{Q}_i = \frac{1}{2} \sum_{i=1}^4 \Psi_i^\dagger \vec{r} \Psi_i \quad (51)$$

Whenever this quantity is conserved

$$d\vec{\Sigma}/dt = 0 \quad (52)$$

the neutrino masses will strictly vanish. In fact, the combinations of Heisenberg, DM and torsional interactions (29), (30), (43) and (44) considered so far, conserve total isospin. They fulfill (52) and give no contribution to the neutrino masses. A

signal for the conservation of isospin is the appearance in all those equations of the linear combinations

$$\vec{\Delta}_i = -3\vec{\delta}_i + \sum_{j \neq i} \vec{\delta}_j \quad (53)$$

The Δ_i enter $d\vec{\Sigma}$ in the form of the sum $\sum_i \vec{\Delta}_i$ - and this sum trivially vanishes.

Nonvanishing contributions to the neutrino masses will be derived below in a systematic and comprehensive way. In order to enlighten the procedure, first consider as a simple example an isospin conserving torque of the form

$$\frac{d\vec{Q}_1}{dt} \sim (\vec{Q}_2 - \vec{Q}_1) + (\vec{Q}_3 - \vec{Q}_1) + (\vec{Q}_4 - \vec{Q}_1) = \vec{\Delta}_1 \quad (54)$$

and compare it with an isospin violating one

$$\frac{d\vec{Q}_1}{dt} = iN_T(\vec{Q}_1 - \vec{Q}_0) = N_T\vec{\delta}_1 \quad (55)$$

with some tiny new coupling N_T and $\vec{Q}_0 = \langle \vec{Q}_1 \rangle$ denoting the ground state value of \vec{Q}_1 . Similarly $dQ_j/dt = iN_T(\vec{Q}_j - \langle \vec{Q}_j \rangle)$ for $j = 2, 3, 4$.

What is the physical meaning of such an isospin violating contribution? After all, (55) does not exhibit any interaction of \vec{Q}_1 with the other $\vec{Q}_{2,3,4}$. It is an isospin non conserving reset torque towards \vec{Q}_0 and effects a mysterious steady gain or loss of isospin, which certainly needs understanding.

In my opinion there is only one plausible explanation: in order that isospin does not disappear into nirvana, the most straightforward assumption is the existence of some kind of nucleus sitting at the center of each tetrahedron and to which isospin can be transferred, at least in tiny doses. There may be other explanations, but I find this one particularly appealing, because one may speculate that the nuclei are responsible for an additional stabilization of the substrate's skeleton structure in Fig. 1.

As seen below, in addition to giving neutrino masses, the coupling N_T also enters all the other quark and lepton mass formulas. Therefore, there is *always* this tiny exchange of isospins with the nucleus, whenever a tetrahedron of isospins gets excited to a quark or a lepton.

With contributions (55) alone, all 3 neutrinos ν_e , ν_μ and ν_τ get the same mass of order N_T . To obtain different masses it is instructive to remember how the

different masses for the 3 families were obtained in the case of the other quarks and leptons, namely by use of isospin-preserving Heisenberg, torsional and DM interactions. Analogously, one may construct isospin violating DM and Heisenberg interactions by replacing $\Delta \rightarrow \delta$ in (34) and (35). One obtains

$$\begin{aligned} \frac{d\vec{Q}_{Li}}{dt} &= iN_T(\vec{Q}_{Li} - \vec{Q}_0) + N_H(\vec{Q}_{Li} \times \vec{Q}_0) \\ &\quad + 2N_D\{-\vec{Q}_{Li} \times \vec{Q}_0 + i(-(\vec{Q}_{Li} - \vec{Q}_0) + ((\vec{Q}_{Li} - \vec{Q}_0)\vec{Q}_0)\vec{Q}_0)\} \\ &= iN_T\vec{\delta}_{Li} + N_H(\vec{\delta}_{Li} \times \vec{Q}_0) + 2N_D\{-\vec{\delta}_{Li} \times \vec{Q}_0 + i(-\vec{\delta}_{Li} + ((\vec{\delta}_{Li}\vec{Q}_0)\vec{Q}_0))\} \end{aligned} \quad (56)$$

and similarly for \vec{Q}_{Ri} . This procedure leads to different masses for the 3 neutrino mass eigenstates ν_1 , ν_2 and ν_3 of the following form

$$m(\nu_1) = 4N_T \quad m(\nu_2) = N_T + N_H \quad m(\nu_3) = N_H + 4N_D - 4N_T \quad (57)$$

As mentioned before, all other quarks and leptons get similar contributions to their masses from N_T , N_H and N_D . However, since the isospin violating couplings N are assumed to be tiny ($\leq 1\text{eV}$), they can be neglected in the mass formulas which were presented in the preceding sections.

One may accommodate (57) to the results from neutrino oscillation experiments. Consider first the case of the so called ‘normal mass hierarchy’ $m(\nu_1) < m(\nu_2) \ll m(\nu_3)$ where

$$m(\nu_1)/\text{eV} = 0.001 \quad m(\nu_2)/\text{eV} = 0.0087 \quad m(\nu_3)/\text{eV} = 0.048 \quad (58)$$

Lacking experimental informations on $m(\nu_1)$ I have guessed here a value of 0.001 eV. In the normal hierarchy limit $m(\nu_1) \ll m(\nu_2) \ll m(\nu_3)$ one sees that $m(\nu_1)$ is a measure of the torsional coupling N_T , $m(\nu_2)$ measures the strength N_H of the Heisenberg coupling and $m(\nu_3)$ of the DM coupling N_D . The situation is thus similar as for the other quarks and leptons, where the heavy family mass is dominated by DM interactions, the second family by Heisenberg and the light family by torsional couplings.

In the case of the so called ‘inverted hierarchy’ one has

$$m(\nu_1)/\text{eV} \approx m(\nu_2)/\text{eV} = 0.0245 \quad m(\nu_3)/\text{eV} = 0.001 \quad (59)$$

where this time the assumption is made on the (unknown) mass $m(\nu_3)$. Trying to accommodate (59) with (57) one obtains $N_H \approx 0$. At the same time a small $m(\nu_1)$

leads to $N_D \approx N_T$, i.e. an accidental compensation between torsional und DM contribution is needed to occur.

In summary, it was found in this section, that the masses $m(\nu_1)$, $m(\nu_2)$ and $m(\nu_3)$ are a measure of the strength of the isospin-violating torsional, Heisenberg and DM interactions, respectively. This happens in a similar way, as the masses of the first, second and third family of quarks and charged leptons are determined by the strength of the isospin-conserving torsional, Heisenberg and DM interactions, cf. the discussion at the beginning of Sect. V.

VII. Lepton Eigenstates

In the previous sections the focus of discussion has been laid on the eigenvalues of the system, i.e. on quark and lepton masses. In the course of the calculations the transition from isospin to mass eigenstates has been carried out via an appropriate diagonalization process and has led to numerical values for the quark and lepton masses.

Actually, the dynamic equations for the isospin vectors allow to calculate the eigenfunctions as well. Namely, the Mathematica program presented in Appendix A gives the physical mass eigenstates in terms of the isospin eigenstates, provided one simply changes the command 'eigenvalues' to 'eigensystem' in the last line of the code.

While the masses correspond to the eigenvalues, CKM and PMNS mixings can be inferred from the eigenvectors. Details of this deduction are presented in Sect. IX. Here I will concentrate on explicitly representing the quark and lepton mass states in terms of the isospin eigenstate vectors. To that end, the following definitions are used:

$$|\vec{S}\rangle := \vec{\delta}_L \qquad |\vec{T}\rangle := \vec{\delta}_R \qquad (60)$$

Dirac's notation with bra and ket states is applied to make the mechanism more transparent. The index $i = 1 - 4$ counting the tetrahedral sites is left out for reasons discussed below.

The quantities (60) are orthonormal vector states and can be used to write down

the equations for the neutrino mass eigenstates, as obtained from Appendix A:

$$\begin{aligned}
|\nu_{e,m}\rangle &= \frac{1}{\sqrt{6}}[(|S_x\rangle + |T_x\rangle) + (|S_y\rangle + |T_y\rangle) + (|S_z\rangle + |T_z\rangle)] \\
|\nu_{\mu,m}\rangle &= \frac{1}{\sqrt{6}}[(|S_x\rangle + |T_x\rangle) + \omega(|S_y\rangle + |T_y\rangle) + \bar{\omega}(|S_z\rangle + |T_z\rangle)] \\
|\nu_{\tau,m}\rangle &= \frac{1}{\sqrt{6}}[(|S_x\rangle + |T_x\rangle) + \bar{\omega}(|S_y\rangle + |T_y\rangle) + \omega(|S_z\rangle + |T_z\rangle)]
\end{aligned} \tag{61}$$

The corresponding result for the charged leptons is

$$\begin{aligned}
|e_m\rangle &= \frac{1}{\sqrt{6}}[(|T_x\rangle - |S_x\rangle) + (|T_y\rangle - |S_y\rangle) + (|T_z\rangle - |S_z\rangle)] \\
|\mu_m\rangle &= \frac{1}{\sqrt{6}}[(|T_x\rangle - |S_x\rangle) + \omega(|T_y\rangle - |S_y\rangle) + \bar{\omega}(|T_z\rangle - |S_z\rangle)] \\
|\tau_m\rangle &= \frac{1}{\sqrt{6}}[(|T_x\rangle - |S_x\rangle) + \bar{\omega}(|T_y\rangle - |S_y\rangle) + \omega(|T_z\rangle - |S_z\rangle)]
\end{aligned} \tag{62}$$

Reference is made to the isospin of only 1 of the 4 tetrons within a tetrahedron, because the contributions from the other 3 tetrons to the eigenstates are identical, and for simplicity not included.

The appearance of the complex numbers

$$\omega = -\frac{1 - i\sqrt{3}}{2} \qquad \bar{\omega} = -\frac{1 + i\sqrt{3}}{2} \tag{63}$$

corresponding to rotations by 120 and 240 degrees are an effect of the underlying tetrahedral symmetry. They turn the expressions (61) and (62) into G_4 -symmetry adapted functions.

The lepton mass states actually can be brought to the much more compact form

$$\begin{aligned}
\begin{bmatrix} |\nu_{em}\rangle \\ |\nu_{\mu m}\rangle \\ |\nu_{\tau m}\rangle \end{bmatrix} &= Z \begin{bmatrix} |V_x\rangle \\ |V_y\rangle \\ |V_z\rangle \end{bmatrix} & \qquad \begin{bmatrix} |e_m\rangle \\ |\mu_m\rangle \\ |\tau_m\rangle \end{bmatrix} &= Z \begin{bmatrix} |A_x\rangle \\ |A_y\rangle \\ |A_z\rangle \end{bmatrix}
\end{aligned} \tag{64}$$

by using the definitions

$$|\vec{V}\rangle = \frac{1}{\sqrt{2}}(|\vec{S}\rangle + |\vec{T}\rangle) \qquad |\vec{A}\rangle = \frac{1}{\sqrt{2}}(|\vec{T}\rangle - |\vec{S}\rangle) \tag{65}$$

and the Z_3 Fourier transform matrices

$$Z = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega & \bar{\omega} \\ 1 & \bar{\omega} & \omega \end{bmatrix} \qquad Z^\dagger = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \bar{\omega} & \omega \\ 1 & \omega & \bar{\omega} \end{bmatrix} \tag{66}$$

Note in passing $|\vec{V}\rangle$ and $|\vec{A}\rangle$ describe fluctuations of the isospin vectors and axial vectors $\Psi^\dagger \vec{\tau} \Psi$ and $\Psi^\dagger \vec{\tau} \gamma_5 \Psi$ around their ground state values.

My calculations show that the eigenfunctions (61), (62) and (64) are stable against variations of all the isospin couplings introduced in the last chapters. As shown in Sect. IX, this implies that the neutrino mixing matrix does not depend on any fermion mass values and leads to a stable and unambiguous prediction for the PMNS matrix.

VIII. Quark Eigenstates

The Mathematica output from Appendix A for the quark mass states in terms of the isospin eigenstates (60) at first sight looks rather cumbersome, but can be simplified for several reasons. First of all, on grounds of symmetry it is not necessary to write down the full 24×24 output. As discussed in the last section, for the case of leptons all 4 tetrons I, II, III and IV on the tetrahedron contribute in the same way, i.e. the structure of the eigenstate is always of the form of a sum I+II+III+IV, so that for the presentation it sufficed to write down the contribution from tetron I. Similarly, the quark states have a recurring form $3 \times \text{I-II-III-IV}$ (for one color, and $3 \times \text{II-I-III-IV}$ and $3 \times \text{III-I-II-IV}$ for the other two). Knowing this, it is enough to present the contribution of one of the tetrons to one of the colors⁷.

A complication arises from the fact that Mathematica cannot distinguish between degenerate eigenstates. Therefore, in order to determine for each quark flavor the state of a definite color (e.g. the one of the form $3 \times \text{I-II-III-IV}$) I had to introduce artificially a tiny color breaking coupling in the program Appendix A. My choice was an additional contribution $0.00001 * \text{del1u}$ and $0.00001 * \text{eel1u}$ to the terms zx5 and zx1 for the isospins of tetron I. Using this trick the program in Appendix A

⁷The quark triplets are triplets under the Shubnikov group initially. For the question how to interpret them as SU(3) color triplets one may consult the appendix in [6].

finally leads to the mass eigenstates for the up-type quarks

$$\begin{aligned}
u_m &= \frac{1}{\sqrt{3}\sqrt{1+\epsilon_1^2}} [(|S_x\rangle + \epsilon_1 |T_x\rangle) + (|S_y\rangle + \epsilon_1 |T_y\rangle) + (|S_z\rangle + \epsilon_1 |T_z\rangle)] \\
c_m &= \frac{1}{\sqrt{3}\sqrt{1+\epsilon_2^2}} [(|S_x\rangle + \epsilon_2 |T_x\rangle) + \omega(|S_y\rangle + \epsilon_2 |T_y\rangle) + \bar{\omega}(|S_z\rangle + \epsilon_2 |T_z\rangle)] \\
t_m &= \frac{1}{\sqrt{3}\sqrt{1+\epsilon_3^2}} [(|S_x\rangle + \epsilon_3 |T_x\rangle) + \bar{\omega}(|S_y\rangle + \epsilon_3 |T_y\rangle) + \omega(|S_z\rangle + \epsilon_3 |T_z\rangle)] \quad (67)
\end{aligned}$$

and for the down quarks

$$\begin{aligned}
d_m &= \frac{1}{\sqrt{3}\sqrt{1+\epsilon_1^2}} [(|T_x\rangle - \epsilon_1 |S_x\rangle) + (|T_y\rangle - \epsilon_1 |S_y\rangle) + (|T_z\rangle - \epsilon_1 |S_z\rangle)] \\
s_m &= \frac{1}{\sqrt{3}\sqrt{1+\epsilon_2^2}} [(|T_x\rangle - \epsilon_2 |S_x\rangle) + \omega(|T_y\rangle - \epsilon_2 |S_y\rangle) + \bar{\omega}(|T_z\rangle - \epsilon_2 |S_z\rangle)] \\
b_m &= \frac{1}{\sqrt{3}\sqrt{1+\epsilon_3^2}} [(|T_x\rangle - \epsilon_3 |S_x\rangle) + \bar{\omega}(|T_y\rangle - \epsilon_3 |S_y\rangle) + \omega(|T_z\rangle - \epsilon_3 |S_z\rangle)] \quad (68)
\end{aligned}$$

As discussed before, reference is made to the isospins \vec{S} and \vec{T} of tetron I only⁸.

Three coefficients $\epsilon_{1,2,3}$ appear in these equations. They depend on the DM, HH and torsional isospin couplings introduced in Sects. IV, V and VI and can be calculated within the model (numbers will be given below). Since there is a one-to-one relation between these couplings and the Yukawa couplings and fermion masses, the ϵ_i may be considered to depend on the quark (and lepton) masses. Qualitatively, this dependence is such, that variation of the i -th family masses modifies ϵ_i only, and hardly the other ϵ_j . Furthermore, ϵ_i goes up when increasing the charged lepton mass of family i , while it goes down with increasing quark masses of family i . It is worthwhile to stress that in any case there is an appreciable dependence on the LR-couplings, which determine the charged lepton masses.

In the case of the light family ϵ_1 as well as the fermion masses are determined solely by the isospin conserving torsional couplings (43), (44) and (50). Masses are given in (47) and the ϵ_1 parameter is

$$\epsilon_1 = \frac{8C_{LL}^3 - 3C_{LL}C_{LR}^2 + 8C_{LL}^2C_{RR} - 5C_{LR}^2C_{RR} + (4C_{LL}^2 + 2C_{LR}^2)W_C}{C_{LR}[-2C_{LL}^2 + 2C_{LR}^2 - 2C_{RR}^2 + (C_{LL} + C_{RR})W_C]} \quad (69)$$

⁸Formally, the lepton eigenfunctions (61) and (62) are recovered by choosing $\epsilon_3 = \epsilon_2 = \epsilon_1 = 1$. It should be stressed, however, that this is only formally true, because the quark states (67) and (68) are defined in a different space than the lepton states; see the above discussion on eigenfunctions I+II+III+IV for leptons and 3×I-II-III-IV for quarks.

With a little algebra one can rewrite this formula so that ϵ_1 depends only on the masses $M_{U1} = m_u$, $M_{D1} = m_d$ and $M_{L1} = m_e$ of the up quark, the down quark and the electron

$$\epsilon_1 = \frac{2f_+(f_+ + f_-)^2 + f_0^2(f_- - 4f_+) + [2f_0^2 - (f_+ + f_-)^2]\sqrt{4f_+^2 + f_0^2}}{f_0[f_- \sqrt{4f_+^2 + f_0^2} + 2f_0^2 - f_+^2 - f_-^2]} \quad (70)$$

where I have introduced the abbreviations

$$f_+ = \frac{1}{4}\sqrt{(M_{U1} + M_{D1})^2 - \frac{2}{3}M_{L1}^2} \quad f_- = \frac{1}{4}(M_{U1} - M_{D1} + M_{L1}) \quad f_0 = \frac{M_{L1}}{6} \quad (71)$$

This complicated result can be approximated to very good precision by

$$\epsilon_1 = \frac{1}{6} \frac{M_{L1}}{M_{U1} + M_{D1}} \quad (72)$$

Interestingly, it turns out that the corresponding results for ϵ_2 and ϵ_3 are formally identical to (72), except that one has to replace the values for the up, down and electron mass by the corresponding mass values of the second and the third family.

$$\epsilon_i = \frac{1}{6} \frac{M_{Li}}{M_{Ui} + M_{Di}} \quad (73)$$

In other words, one obtains the result for the eigenstates irrespective of what kind of coupling (DM, Heisenberg or torsion) is considered.

Another interesting point is that there is the dependence of (73) on the lepton masses. Since the ϵ_i enter the CKM matrix, a lepton mass dependence appears in the CKM elements, cf. (101) and (106).

Using the plain quark and lepton mass values given by the particle data group[28] the formula (73) yields

$$\epsilon_1 = 0.0140 \quad \epsilon_2 = 0.0128 \quad \epsilon_3 = 0.00171 \quad (74)$$

It will be argued later in (103) that instead of the low energy values (74) ϵ_i -values near the Planck scale should better be considered. They show a more pronounced hierarchy $\epsilon_1 \gg \epsilon_2 \gg \epsilon_3$ than (74).

Instead of $\epsilon_{1,2,3}$ one may introduce three angles $\alpha_{1,2,3}$ to describe the rotations between S and T states implicit in the eigenfunctions (67) and (68), with $\alpha_i := \arctan(\epsilon_i)$. The α_i turn out to be related to the angles appearing in the standard parametrization of the CKM matrix.

IX. Quark and Lepton Mixing Matrices

Experiments show that there is a mixing between the flavor and mass eigenstates of the 3 neutrino species, and this can be described by a unitary matrix, the PMNS neutrino mixing matrix[32, 33]. The experimentally relevant quantities are the absolute values of the matrix elements, which describe the amount of admixture of the flavor into mass eigenstates, and the leptonic Jarlskog invariant which describes any possible CP violation in the leptonic sector.

Since the discovery of neutrino oscillations, many models of neutrino mass and mixings have been constructed. The most straightforward approach is to incorporate Dirac neutrino masses into the Standard Model by introducing three right-handed neutrinos coupled to a Higgs field analogously to the quarks and charged leptons - but unfortunately, within the SM the values of the mixing parameters cannot be predicted.

Leading symmetric Approximation

In a first step a leading order result for the mixing matrix will be derived which is

$$\begin{aligned}
 V_{PMNS} &= \exp \left\{ \frac{i}{\sqrt{3}} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 1 & -1 \\ 0 & -1 & -1 \end{bmatrix} \right\} \\
 &= \begin{bmatrix} 0.8467 - i0.0300 & -0.1489 + i0.4861 & 0.1532 - i0.00051 \\ -0.1489 - i0.4861 & 0.5446 + i0.4568 & -0.00433 - i0.4858 \\ 0.1532 - i0.00051 & -0.00433 - i0.4858 & 0.6892 - i0.5153 \end{bmatrix} \quad (75)
 \end{aligned}$$

while an improved formula will be given later in (93).

The leading order expression (75) is a complex, symmetric and unitary matrix, and the absolute values of the matrix elements can be calculated numerically and compared to measurements

$$\begin{bmatrix} 0.843 & 0.510 & 0.153 \\ 0.510 & 0.711 & 0.486 \\ 0.153 & 0.486 & 0.861 \end{bmatrix} \quad vs. \quad \begin{bmatrix} 0.80 - 0.85 & 0.51 - 0.58 & 0.142 - 0.155 \\ 0.23 - 0.51 & 0.46 - 0.69 & 0.63 - 0.78 \\ 0.25 - 0.53 & 0.47 - 0.70 & 0.61 - 0.76 \end{bmatrix} \quad (76)$$

By inspection one concludes that the agreement is reasonable but not optimal, with the 23 entry being the most critical. The first row, which is best measured, is

also best fitting. Concerning the other rows, the experimental results in (76) are non-symmetric, though with very large errors. It will be described later, in connection with (93) and (94), how (75) can be improved by additional non-symmetric contributions so that complete agreement within the errors is obtained.

A prediction for the leptonic Jarlskog invariant[27] can be calculated from (75) as

$$J_{PMNS} = \Im(V_{e1}V_{\mu 2}\bar{V}_{e2}\bar{V}_{\mu 1}) = -0.0106 \quad (77)$$

This value is large as compared to the Jarlskog parameter of the CKM matrix[28]. J_{PMNS} has not been measured so far, although there are experimental indications that leptonic CP violation is indeed rather large[29].

Motivation and Proof of (75)

My calculations show that the eigenfunctions (61), (62) and (64) are stable against variations of all the isospin couplings one may use in the Hamiltonian H in (7). In consequence, the neutrino mixing matrix does not depend on any fermion mass values. This implies a stable and unambiguous prediction for the PMNS matrix and is in contrast to the CKM matrix in the quark sector, where a mass dependence shows up.

As well known, the defining equation for the PMNS matrix is

$$\left[\langle \nu_{ew} | \quad \langle \nu_{\mu w} | \quad \langle \nu_{\tau w} | \right] W_{\mu}^+ \begin{bmatrix} |e_w\rangle \\ |\mu_w\rangle \\ |\tau_w\rangle \end{bmatrix} = \left[\langle \nu_{em} | \quad \langle \nu_{\mu m} | \quad \langle \nu_{\tau m} | \right] W_{\mu}^+ V_{PMNS} \begin{bmatrix} |e_m\rangle \\ |\mu_m\rangle \\ |\tau_m\rangle \end{bmatrix} \quad (78)$$

where the index w denotes weak interaction eigenstates, and it is understood that we talk about left handed fields only. The mixing matrix is formally given by

$$V_{PMNS} = V_N V_L^{\dagger} = \begin{bmatrix} V_{1e} & V_{1\mu} & V_{1\tau} \\ V_{2e} & V_{2\mu} & V_{2\tau} \\ V_{3e} & V_{3\mu} & V_{3\tau} \end{bmatrix} \quad (79)$$

where

$$V_N = \begin{bmatrix} \langle \nu_{em} | \\ \langle \nu_{\mu m} | \\ \langle \nu_{\tau m} | \end{bmatrix} \left[|\nu_{ew}\rangle \quad |\nu_{\mu w}\rangle \quad |\nu_{\tau w}\rangle \right] \quad V_L^{\dagger} = \begin{bmatrix} \langle e_w | \\ \langle \mu_w | \\ \langle \tau_w | \end{bmatrix} \left[|e_m\rangle \quad |\mu_m\rangle \quad |\tau_m\rangle \right] \quad (80)$$

Replacing the mass eigenstates by the isospin excitations according to (64) one obtains

$$V_{PMNS} = Z \left\{ \begin{array}{c} \langle V_x | \\ \langle V_y | \\ \langle V_z | \end{array} \right. \left[\begin{array}{ccc} |\nu_{ew}\rangle & |\nu_{\mu w}\rangle & |\nu_{\tau w}\rangle \end{array} \right] \left. \begin{array}{c} \langle e_w | \\ \langle \mu_w | \\ \langle \tau_w | \end{array} \right. \left[\begin{array}{ccc} |A_x\rangle & |A_y\rangle & |A_z\rangle \end{array} \right] \Big\} Z^\dagger \quad (81)$$

By inspection one sees that (81) exactly compensates all the matrix transformations in (78) and (64) so as to maintain lepton universality and keep the weak current diagonal in the weak eigenstates.

The brace in (81) comprises a matrix of expectation values of the form

$$Y := \begin{array}{c} \langle V_x | \\ \langle V_y | \\ \langle V_z | \end{array} \mathcal{O} \left[\begin{array}{ccc} |A_x\rangle & |A_y\rangle & |A_z\rangle \end{array} \right] \quad (82)$$

where the inner product

$$\mathcal{O} := \left[\begin{array}{ccc} |\nu_{ew}\rangle & |\nu_{\mu w}\rangle & |\nu_{\tau w}\rangle \end{array} \right] \begin{array}{c} \langle e_w | \\ \langle \mu_w | \\ \langle \tau_w | \end{array} \quad (83)$$

is a dyadic 1-dimensional operator which acts between the complex 3-dimensional spaces of charged lepton ($\sim \vec{S} - \vec{T}$) and antineutrino ($\sim \vec{S} + \vec{T}$) states. One may say that it contains all information about what the charged W-boson does to the lepton fields: it changes isospin, mixes families and so on. Weak SU(2) and tetrahedral symmetry force \mathcal{O} to have the form

$$\begin{aligned} \mathcal{O} = & |S_x\rangle \langle T_x| + |S_y\rangle \langle T_y| + |S_z\rangle \langle T_z| - |T_x\rangle \langle S_x| - |T_y\rangle \langle S_y| - |T_z\rangle \langle S_z| \\ & + \frac{i}{\sqrt{3}} [|S_y\rangle \langle S_z| + |S_z\rangle \langle S_y| - |T_y\rangle \langle T_z| - |T_z\rangle \langle T_y|] \\ & + \frac{i}{\sqrt{3}} [\omega |S_x\rangle \langle S_y| + \bar{\omega} |S_y\rangle \langle S_x| - \omega |T_x\rangle \langle T_y| - \bar{\omega} |T_y\rangle \langle T_x|] \\ & + \frac{i}{\sqrt{3}} [\bar{\omega} |S_x\rangle \langle S_z| + \omega |S_z\rangle \langle S_x| - \bar{\omega} |T_x\rangle \langle T_z| - \omega |T_z\rangle \langle T_x|] \end{aligned} \quad (84)$$

In order to derive (84) one has to note that SU(2) invariance allows the appearance of dot products and triple products only. The coefficients of these products are then dictated by the tetrahedral symmetry of the isospin vectors. For example, to derive

the triple product coefficients one should remember that the W^+ -boson is defined in the 3 internal dimensions in an analogous manner as a plus circularly polarized wave in 3 spatial dimensions, namely by means of an ‘internal polarization’ vector $\vec{e}_+ = (\vec{e}_1 + i\vec{e}_2)/\sqrt{2}$ which is perpendicular to the axis of quantization, in this case given by $\sim (1, 1, 1)$.

$$\vec{e}_1 = \frac{1}{\sqrt{2}}(0, 1, -1) \quad \vec{e}_2 = \frac{1}{\sqrt{6}}(-2, 1, 1) \quad (85)$$

Introducing the vector

$$\vec{\Omega} = \frac{1}{\sqrt{3}}(1, \omega, \bar{\omega}) \quad (86)$$

allowed contributions to \mathcal{O} are of the triple product form

$$\begin{aligned} \varepsilon_{ijk} \frac{1}{\sqrt{2}}(\vec{e}_1 + i\vec{e}_2)_i |Q_j\rangle \langle Q'_k| &= -\frac{i}{\sqrt{3}} \vec{\Omega}(\vec{Q} \times \vec{Q}') = -\frac{i}{\sqrt{3}} [|Q'_y\rangle \langle Q_z| - |Q'_z\rangle \langle Q_y| \\ &\quad -\omega(|Q'_x\rangle \langle Q_z| - |Q'_z\rangle \langle Q_x|) + \bar{\omega}(|Q'_x\rangle \langle Q_y| - |Q'_y\rangle \langle Q_x|)] \end{aligned} \quad (87)$$

for the ket and bra states belonging to any 2 internal angular momenta Q and Q' . These contributions are anti-hermitean, and care must be taken in the definition of the complex triple product when using complex conjugation in the determination of \mathcal{O} .

Note that \mathcal{O} as given in (84) is universal in the sense that it depends only on properties of the Ψ field, and therefore will appear in identical form within the quark sector and the calculation of the CKM matrix. This fact reflects the quark lepton universality of the W-boson interactions.

Inserting (84) into (82) one obtains

$$Y = \begin{bmatrix} \langle V_x | \\ \langle V_y | \\ \langle V_z | \end{bmatrix} \mathcal{O} \begin{bmatrix} |A_x\rangle & |A_y\rangle & |A_z\rangle \end{bmatrix} = I + X \quad (88)$$

i.e. a sum of a hermitean part (the unit matrix I) and an anti-hermitean matrix

$$X = -\frac{i}{\sqrt{3}} \begin{bmatrix} 0 & \bar{\omega} & \omega \\ \omega & 0 & 1 \\ \bar{\omega} & 1 & 0 \end{bmatrix} \quad (89)$$

The invariant structure which gives the unit matrix in (88) is the dot product, while the invariant structure belonging to the anti-hermitean contribution X is the triple product. The unit matrix corresponds to no mixing at all, so the origin of a non-trivial PMNS matrix is to be found solely in the triple product terms (87).

Since the result (88) is not unitary, an exponentiation suggests itself which gives a unitary PMNS matrix of the form

$$\begin{aligned}
V_{PMNS} &= Ze^X Z^\dagger = e^{ZXZ^\dagger} \\
&= \frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega & \bar{\omega} \\ 1 & \bar{\omega} & \omega \end{bmatrix} \exp \left\{ \frac{-i}{\sqrt{3}} \begin{bmatrix} 0 & \bar{\omega} & \omega \\ \omega & 0 & 1 \\ \bar{\omega} & 1 & 0 \end{bmatrix} \right\} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \bar{\omega} & \omega \\ 1 & \omega & \bar{\omega} \end{bmatrix} \\
&= \begin{bmatrix} 0.8467 - i0.0300 & -0.1489 + i0.4861 & 0.1532 - i0.00051 \\ -0.1489 - i0.4861 & 0.5446 + i0.4568 & -0.00433 - i0.4858 \\ 0.1532 - i0.00051 & -0.00433 - i0.4858 & 0.6892 - i0.5153 \end{bmatrix} \quad (90)
\end{aligned}$$

identical to what was claimed in (75).

Improved Formula for the PMNS Matrix

So far only dot product and triple product terms (87) have been considered as contributing to the operator (84) and the PMNS result. Actually, there is a third kind of term that needs consideration. Using $\vec{\Omega}^2 = 0$ and the same normalization as in (87) it is of the form

$$-(\vec{\Omega} \times \vec{Q}) (\vec{\Omega} \times \vec{Q}') = (\vec{\Omega} \vec{Q}) (\vec{\Omega} \vec{Q}') \quad (91)$$

As shown in the previous sections, quark and lepton masses are related to torsional, Heisenberg and Dzyaloshinskii isospin interactions of the fundamental tetron Ψ field. Furthermore, these three types of interactions completely fix the structure of the model.

This fact is reflected in the contributions to the operator \mathcal{O} : while the dot products and triple products appearing in (84) parallel the torsional and Heisenberg interactions, (91) corresponds to the Dzyaloshinskii Hamiltonian. Working out the products $|Q_i\rangle \langle Q'_j|$ arising from (91), it leads to an additional contribution to (84)

which can be comprised by a matrix

$$D := \frac{1}{6} \begin{bmatrix} 1 & \omega & \bar{\omega} \\ \omega & \bar{\omega} & 1 \\ \bar{\omega} & 1 & \omega \end{bmatrix} - h.c. \quad (92)$$

The role of D for (91) is analogous to that of X for the triple product term. Combining the X and D contributions an improved formula for the PMNS matrix is obtained

$$V_{PMNS} = \exp \left\{ \frac{1}{2} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix} \right\} \exp \left\{ \frac{i}{\sqrt{3}} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 1 & -1 \\ 0 & -1 & -1 \end{bmatrix} \right\} \quad (93)$$

This represents a complex and unitary matrix whose absolute value matrix $|V_{PMNS}|$ is not symmetric, in contrast to (75). Its elements are given by

$$\begin{bmatrix} 0.847 & 0.510 & 0.153 \\ 0.430 & 0.505 & 0.748 \\ 0.311 & 0.698 & 0.645 \end{bmatrix} \quad vs. \quad \begin{bmatrix} 0.80 - 0.85 & 0.51 - 0.58 & 0.142 - 0.155 \\ 0.23 - 0.51 & 0.46 - 0.69 & 0.63 - 0.78 \\ 0.25 - 0.53 & 0.47 - 0.70 & 0.61 - 0.76 \end{bmatrix} \quad (94)$$

and fit the phenomenological numbers to within one standard error.

The value of the leptonic Jarlskog invariant now is

$$J_{PMNS} = 0.0118 \quad (95)$$

Thus, while the improvement (93) only moderately corrects the absolute values (76), it strongly modifies the prediction (77) for J_{PMNS} (changing even the sign!). This is because in contrast to the absolute values the Jarlskog invariant is dominated by higher orders of the exponential expansion.

Outlook to Quark Mixing

Mixing in the quark sector has been known since the time of Cabibbo[31]. Although the mixing percentages are smaller, it is much better measured than in the lepton sector. On the other hand, concerning theory, the predictions for the CKM mixing elements in the present model are somewhat more difficult to obtain, though parts of the arguments for leptons can be taken over to the quark sector. The idea is again that the mixing matrix counterbalances the deviation of the mass eigenstates

from the weak eigenstates in such a way that the charged current effectively acts diagonal on the isospin operators (60). The main complication is the appearance of mass dependent factors in the quark eigenstates, see below.

Within the Standard Model, quark masses arise from Yukawa interactions involving the Higgs field, left-handed⁹ quark doublets, and right-handed down- or up-type quark singlets. The Yukawa couplings are given as complex 3×3 matrices, and the mass terms are obtained when the Higgs field acquires a vacuum expectation value v . The physical(=mass) states are attained by diagonalizing the Yukawa matrices Y^U and Y^D using 4 unitary matrices $V_{L,R}^{U,D}$ to obtain diagonal mass matrices for up- and down-type quarks

$$\frac{v}{\sqrt{2}} Y^U = V_L^{U\dagger} M_{diag}^U V_R^U \quad \frac{v}{\sqrt{2}} Y^D = V_L^{D\dagger} M_{diag}^D V_R^D \quad (96)$$

These equations show how the SM Yukawa couplings can be built from the mass eigenvalues and the matrices $V_{L,R}^{U,D}$. One may compare this to the present model where

-the entries of M_{diag}^U and M_{diag}^D are obtained as excitation energies of the isospin Hamiltonian.

-the matrices $V_L^{U,D}$ can be expressed in terms of the isospin states $|\vec{S}\rangle$ and $|\vec{T}\rangle$ as will be seen in the explicit representation (98) below.

As a result of the diagonalization (96), the left-handed weak charged-current interactions couple to the physical quarks with couplings involving the CKM matrix which is defined as

$$\begin{aligned} V_{CKM} &= V_U V_D^\dagger = \begin{bmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{bmatrix} \\ &= \begin{bmatrix} \langle u_m | u_w \rangle & \langle u_m | c_w \rangle & \langle u_m | t_w \rangle \\ \langle c_m | u_w \rangle & \langle c_m | c_w \rangle & \langle c_m | t_w \rangle \\ \langle t_m | u_w \rangle & \langle t_m | c_w \rangle & \langle t_m | t_w \rangle \end{bmatrix} \begin{bmatrix} \langle d_w | d_m \rangle & \langle d_w | s_m \rangle & \langle d_w | b_m \rangle \\ \langle s_w | d_m \rangle & \langle s_w | s_m \rangle & \langle s_w | b_m \rangle \\ \langle b_w | d_m \rangle & \langle b_w | s_m \rangle & \langle b_w | b_m \rangle \end{bmatrix} \quad (97) \end{aligned}$$

Here I have used Dirac's ket and bra notation as described in the last section. m denotes mass eigenstates (the physical states) and w weak interaction eigenstates.

⁹The appearance in the microscopic model of chiral fermions and left handed weak interactions has been discussed detailedly in previous work. In essence it is due to the handedness of the tetrahedrons formed by the isospin vectors in isospin space.

Knowing the quark mass eigenstates (64) and (65) one may write down the CKM matrix in an analogous fashion as the PMNS matrix (78) for leptons

$$V_{CKM} = \left\{ RZ \begin{bmatrix} \langle S_x | \\ \langle S_y | \\ \langle S_z | \end{bmatrix} + REZ \begin{bmatrix} \langle T_x | \\ \langle T_y | \\ \langle T_z | \end{bmatrix} \right\} \left[|u_w\rangle \quad |c_w\rangle \quad |t_w\rangle \right] \times \\ \begin{bmatrix} \langle d_w | \\ \langle s_w | \\ \langle b_w | \end{bmatrix} \left\{ \left[|T_x\rangle \quad |T_y\rangle \quad |T_z\rangle \right] Z^\dagger R - \left[|S_x\rangle \quad |S_y\rangle \quad |S_z\rangle \right] Z^\dagger ER \right\} \quad (98)$$

where the first line is the matrix V_L^U and the second line is $V_L^{D\dagger}$ and the matrices

$$E := \begin{bmatrix} \epsilon_1 & 0 & 0 \\ 0 & \epsilon_2 & 0 \\ 0 & 0 & \epsilon_3 \end{bmatrix} \quad R := \begin{bmatrix} \frac{1}{\sqrt{1+\epsilon_1^2}} & 0 & 0 \\ 0 & \frac{1}{\sqrt{1+\epsilon_2^2}} & 0 \\ 0 & 0 & \frac{1}{\sqrt{1+\epsilon_3^2}} \end{bmatrix} \quad (99)$$

have been introduced.

Just as in the case of leptons (83) there is a 1-dimensional dyadic transformation

$$\mathcal{O} = \left[|u_w\rangle \quad |c_w\rangle \quad |t_w\rangle \right] \begin{bmatrix} \langle d_w | \\ \langle s_w | \\ \langle b_w | \end{bmatrix} \quad (100)$$

which operates between the 3-dimensional spaces of up- and down-type quark states. Due to quark-lepton universality, when expressed in terms of operators \vec{S} and \vec{T} , the operator \mathcal{O} for quarks must be identical to what was used for leptons in (84).

Restricting, for a moment, on the dot and triple product contributions (84) as input, one may then calculate V_{CKM} given in (98) to be

$$V_{CKM} = I + RZXZ^\dagger ER + REZZXZ^\dagger R \rightarrow \exp\{RZXZ^\dagger ER + REZZXZ^\dagger R\} \quad (101)$$

where I is the 3×3 unit matrix arising from the dot product terms in (84). The other terms in (101) are the anti-hermitian contributions from the triple product in (87) and (84). They replace the expression ZXZ^\dagger in (90) for leptons.

Just as in the case of leptons one may improve on this result by including the contributions from (91), in order to obtain the desired non-symmetric contributions

to $|V_{CKM}|$. The improved formula for the CKM matrix reads

$$V_{CKM} = \exp\{2[RZDZ^\dagger ER - REZD^\dagger Z^\dagger R]\} \exp\{RZXZ^\dagger ER + REZXZ^\dagger R\} \quad (102)$$

In contrast to X in (89) the matrix D in (92) is not anti-hermitian. This fact has been accounted for in the first exponential factor.

Eq. (102) allows to evaluate $|V_{CKM}|$ using appropriate values for the fermion masses entering (73). It must be noted, however, that the low energy values (74) of the ϵ_i are not useful in this context. Instead one should use running masses near the Planck scale, because the dynamics generates fermion masses originally at Planck scale distances¹⁰. Unfortunately, the predictions for running masses are not very precise because higher order contributions become appreciable at very large scales. Nevertheless, I am using results from the literature[30, 38] to determine the ϵ_i at high scales.

$$\epsilon_1 = 0.35 \quad \epsilon_2 = 0.070 \quad \epsilon_3 = 0.0040 \quad (103)$$

unfortunately with a large theoretical error[38], whose magnitude even is hard to estimate. The numbers are for a 2HDM (2 Higgs doublet model) which is known to be the low-energy limit of the microscopic model[7]. They exhibit a family hierarchy which will be seen to induce a corresponding hierarchy in the mixing of the quark families. Actually, as discussed in earlier work[5], this is to be expected within the present model due to the large top mass which forces the up- and down-type mass eigenstates to be approximately $\sim \vec{S}$ and $\sim \vec{T}$, respectively, in (67) and (68), much unlike the lepton states which are $\sim \vec{S} \pm \vec{T}$ according to (64).

Just as masses, CKM matrix elements are running, i.e. dependent on the scale paramter $t = \ln \frac{E}{\mu}$ where E is the relevant energy scale and μ the renormalization scale. The running of the absolute values of the CKM matrix elements has been discussed for the 2HDM in [38]. It turns out to be remarkably simple, at least in

¹⁰A GUT scale is not present in the model. There is only the Fermi scale, defined as the interaction energy of the isospin vectors, and the Planck scale, defined as the binding energy of the fields Ψ [7].

leading order, because it can be given in terms of one universal function $h(t)$.

$$|V_{CKM}(t)| \approx \begin{bmatrix} |V_{ud}(0)| & |V_{us}(0)| & \frac{|V_{ub}(0)|}{h(t)} \\ |V_{cd}(0)| & |V_{cs}(0)| & \frac{|V_{cb}(0)|}{h(t)} \\ \frac{|V_{td}(0)|}{h(t)} & \frac{|V_{ts}(0)|}{h(t)} & |V_{tb}(0)| \end{bmatrix} \quad (104)$$

For the Jarlskog invariant one has

$$J_{CKM}(t) \approx \frac{J_{CKM}(0)}{h^2(t)} \quad (105)$$

In the 2HDM case $h(t)$ is a moderately varying function. According to [38] it increases by about 20% when going from GeV to Planck scale energies.

Using (102) and (103) I have calculated the CKM elements at high energies and then extrapolated them back to GeV energies according to (104). I obtain the matrix $|V_{CKM}|$ of absolute values

$$\begin{bmatrix} 0.974 & 0.224 & 0.0035 \\ 0.224 & 0.973 & 0.044 \\ 0.0080 & 0.043 & 0.9991 \end{bmatrix} \text{ vs. } \begin{bmatrix} 0.9734 - 0.9740 & 0.2235 - 0.2251 & 0.00362 - 0.00402 \\ 0.217 - 0.225 & 0.969 - 0.981 & 0.0394 - 0.0422 \\ 0.0083 - 0.0088 & 0.0404 - 0.0424 & 0.985 - 1.043 \end{bmatrix} \quad (106)$$

The numbers look reasonable, as compared to the phenomenological values, and show the correct hierarchy and orders of magnitude. However, the theoretical uncertainty from the scale evolution is large and difficult to estimate, in particular concerning quark mass values near the Planck scale. For example, ϵ_1 accommodates the Cabbibo angle correctly, whereas the ‘23’-matrix elements $|V_{ts}|$ and $|V_{cb}|$ tendentially come out too large, while the ‘13’-elements $|V_{ub}|$ and $|V_{td}|$ are typically too small. These deviations may seem being just 2σ effects, but as stressed before the theoretical error from the quark mass evolution is extremely difficult to handle.

Similarly, concerning the Jarlskog invariant one obtains $J_{CKM} = 0.000027$, a bit small when compared to the observed value $J_{CKM} = (3.00 + 0.15 - 0.09) \times 10^{-5}$.

In conclusion, explicit analytic and numerical results for the mixing matrices have been presented in this work. Of particular interest are the prediction for the PMNS matrix (93) and the fermion mass dependence of the CKM matrix as given by (102). Actually, (102) is universal in that it embraces (i) the case of no mixing ($\epsilon_1 = \epsilon_2 = \epsilon_3 = 0$), (ii) the CKM prediction obtained with ϵ_i -values (103) and (iii)




Figure 2: Transition between the CKM and the PMNS limit of the matrix elements $|V_{12}|$ and $|V_{13}|$ and the Jarlskog invariant (from left to right) as a function of the parameter α defined in the main text. For example, $|V_{12}|$ starts with its CKM value 0.224 at $\alpha = 0$ and grows towards the PMNS value at $\alpha = 1$.

the PMNS formula which formally is given using $\epsilon_1 = \epsilon_2 = \epsilon_3 = 1$. To make this visible, I have drawn in Fig. 1 the ‘12’ (i.e. Cabibbo) and the ‘13’ matrix element and the Jarlskog invariant as a function of a parameter α . α is introduced to avoid drawing the full ϵ_i -dependence of the matrix elements and defined in such a way that it vanishes in the CKM case and takes the value of 1 in the PMNS limit. More precisely, one has

$$\epsilon_1 = 0.35 + 0.65 \alpha \quad \epsilon_2 = 0.07 + 0.93 \alpha \quad \epsilon_3 = 0.004 + 0.996 \alpha \quad (107)$$

IX. Summary and Discussion

This work has shown in detail how the observed spectrum of quarks and leptons can be related to isospin interactions among tetrons. After clarifying the connection between SM Yukawa couplings and the isomagnetic couplings, the magnitudes of the latter were adapted to the observed mass values. Furthermore, in Sect. III it was explained how these coupling parameters themselves can be calculated from exchange integrals involving the fundamental scalar and triplet potentials V_1 and V_3 among tetrons. The resulting optimized predictions for the masses can be found at the end of the Mathematica program in Appendix A. Numbers are understood in GeV.

As turns out, Dzyaloshinskii-Moriya couplings are the largest, while Heisenberg interaction terms are smaller. It is a feature of the DM interaction to give masses only to the third family. In particular the top mass is the only excitation with mass of order Λ_F , because it corresponds to a minimum energy of the tetrahedral

isospin Hamiltonian (38). This is linked to the SSB of the ordered isospins, i.e. to how the aligned tetrahedral 'stars' are oriented collectively in internal space, thus breaking weak isospin SU(2) symmetry. The ordering takes place below the transition temperature Λ_F , while isospins are distributed randomly (and thus SU(2) symmetric) at temperatures above the Fermi scale.

All other quark and lepton masses naturally turn out to be much smaller than m_t . For example, the Heisenberg interactions characteristically give equal contributions to the masses of the second and third family, keeping the first family massless. The first family then obtains its masses from still smaller torsional interactions, as explained in Sect. V.

As a byproduct of the calculations, solutions within the microscopic model to several outstanding classical problems of particle physics have appeared:

-The hierarchy problem of why the Fermi scale is so small as compared to the Planck scale. In the microscopic model this is due to the smallness of exchange integrals as compared to direct ones, see the discussion after (17) in Sect. III.

-The tinytness of neutrino masses arises from the conservation of tetron isospin. Isospin violating interactions have been introduced in Sect. VI in order to accommodate reasonable neutrino mass values, and their physical origin has been clarified. Note that isospin is not an abstract concept here but corresponds to real rotations in the 3 extra dimensions.

Concerning the observed quark and lepton mixing a detailed analysis will follow in [46]. As basis for such an examination in the present work all quark and lepton states of the 3 families have been listed as eigenfunctions of the tetron isospin Hamiltonian. In the microscopic model the internal dynamics of quarks and leptons is intertwined, and therefore it is not surprising that the quark states not only depend on the quark but also on the lepton masses. Details of these and other dependencies were discussed in Sect. VIII.

Neutrino eigenstates are given as vibrations $\vec{S} + \vec{T} = \delta\vec{Q}_L + \delta\vec{Q}_R$, i.e. of the total isospin vector, and the 3 charged leptons as vibrations $\vec{S} - \vec{T} = \delta\vec{Q}_L - \delta\vec{Q}_R$. As shown in [46] this corresponds to large mixing in the lepton sector and large values of the PMNS matrix elements[28]. On a qualitative level this should be no surprise in view of the discussion of isospin conservation in Sect. VI, because isospin conservation

explains the appearance of sums $\vec{S} + \vec{T}$ for neutrinos and - for reasons of orthogonality - of differences $\vec{S} - \vec{T}$ for the charged leptons. Neutrino mass eigenstates are thus 'far away' from the isospin states \vec{S} and \vec{T} , and the resulting PMNS mixing matrix will be 'far away' from the unit matrix. This is much in contrast to the case of quarks where the mass eigenstates are small deviations $\vec{S} + \epsilon\vec{T}$ and $-\epsilon\vec{S} + \vec{T}$ from the states \vec{S} and \vec{T} , with small numbers ϵ_i that measure the mixing contribution from the i -th family and show a hierarchy $\epsilon_3 \ll \epsilon_2 \ll \epsilon_1 \ll 1$ as needed to understand the observed hierarchy in the CKM matrix[46].

Finally, the question: are there limitations of the approach? Certainly yes. The calculations presented are leading order with respect to many types of corrections: -For one, a linear approximation has been used throughout for the fluctuations δ describing the quark and lepton states. Probably there will be important higher order corrections, for example effects from the heavy quarks on the light families. More concretely, next-to-leading effects from the large DM-couplings may overwhelm the tiny contributions from torsional interactions on the first family.

-Secondly, the calculations in this work include only *intra*-tetrahedral interactions of tetron isospins, i.e. interactions within one tetrahedron. Although an attempt has been made in connection with (39), *inter*-tetrahedral interactions may not have been fully taken into account. In other words, besides the top quark contributions (39) there may be other inter-tetrahedral effects from the lighter fermions.

It is an interesting question whether there are phenomenological predictions which distinguish the present model from other BSM theories and from the SM. First of all, it must be noted that since the model has been constructed to have a low energy limit which is as close to the SM as possible, many of its BSM effects are suppressed by negative powers of the Planck mass. However, I can see 2 major areas which go beyond the standard predictions:

-the parameter-free formula (98) for the PMNS matrix. In contrast to the CKM result (94) which strongly depends on running quark mass values chosen, its theoretical error is expected to be very small. As future neutrino experiments become more and more accurate, discrepancies to (98) would shake the validity of the microscopic model.

-the other area is cosmology. In previous work it was pointed out that the accelerated expansion of the universe obtained in the present model differs from the cosmologi-

cal standard model (CSM), at least in very large time scales, and that an improved observation of the dark energy effect may be able to discriminate between the 2 models. These considerations were published before the so called ‘Hubble tension’ (saying that measurements at different time scales provide contradictory values for the cosmological constant) was found, and it will be interesting to see whether the present model can contribute to its solution.

Appendix A: Mathematica Program to calculate the Quark and Lepton Masses and Eigenstates

The following code allows to calculate quark and lepton masses and eigenstates, given the isospin couplings as defined in the main text. The resulting masses can be found at the bottom line of the program (in GeV).

The program’s output for the eigenstates is not included in the code, but presented in (61), (62), (67) and (68).

```

s10:={-1, -1, -1}/√3
del1u:={d1x, d1y, d1z} * ef
del2u:={d2x, -d2y, -d2z} * ef
del3u:={-d3x, d3y, -d3z} * ef
del4u:={-d4x, -d4y, d4z} * ef

t10:= + s10
eel1u:={e1x, e1y, e1z} * ef
eel2u:={e2x, -e2y, -e2z} * ef
eel3u:={-e3x, e3y, -e3z} * ef
eel4u:={-e4x, -e4y, e4z} * ef

dd1:=del2u + del3u + del4u - 3 * del1u
dd2:=del1u + del3u + del4u - 3 * del2u
dd3:=del1u + del2u + del4u - 3 * del3u
dd4:=del1u + del2u + del3u - 3 * del4u

```

ed1:=eel2u + eel3u + eel4u - 3 * del1u
ed2:=eel1u + eel3u + eel4u - 3 * del2u
ed3:=eel1u + eel2u + eel4u - 3 * del3u
ed4:=eel1u + eel2u + eel3u - 3 * del4u

del1:=del2u + del3u + del4u - 3 * eel1u
de2:=del1u + del3u + del4u - 3 * eel2u
de3:=del1u + del2u + del4u - 3 * eel3u
de4:=del1u + del2u + del3u - 3 * eel4u

ee1:=eel2u + eel3u + eel4u - 3 * eel1u
ee2:=eel1u + eel3u + eel4u - 3 * eel2u
ee3:=eel1u + eel2u + eel4u - 3 * eel3u
ee4:=eel1u + eel2u + eel3u - 3 * eel4u

vdd1:= - 2 * dd1 + 2 * dd1.s10 * s10
vdd2:= - 2 * dd2 + 2 * dd2.s10 * s10
vdd3:= - 2 * dd3 + 2 * dd3.s10 * s10
vdd4:= - 2 * dd4 + 2 * dd4.s10 * s10

ved1:= - 2 * ed1 + 2 * ed1.s10 * s10
ved2:= - 2 * ed2 + 2 * ed2.s10 * s10
ved3:= - 2 * ed3 + 2 * ed3.s10 * s10
ved4:= - 2 * ed4 + 2 * ed4.s10 * s10

vde1:= - 2 * de1 + 2 * de1.s10 * s10
vde2:= - 2 * de2 + 2 * de2.s10 * s10
vde3:= - 2 * de3 + 2 * de3.s10 * s10
vde4:= - 2 * de4 + 2 * de4.s10 * s10

vee1:= - 2 * ee1 + 2 * ee1.s10 * s10
vee2:= - 2 * ee2 + 2 * ee2.s10 * s10
vee3:= - 2 * ee3 + 2 * ee3.s10 * s10

$$\text{vee4} := -2 * \text{ee4} + 2 * \text{ee4.s10} * \text{s10}$$

$$\text{ss} := -10.700000000000000000$$

$$\text{st} := -0.077000000000000000$$

$$\text{tt} := -0.220000000000000000$$

$$\text{jss} := 0.320000000000000000$$

$$\text{jtt} := 0.010200000000000000$$

$$\text{jst} := 0.017500000000000000$$

$$\text{ff} := 0.000490000000000000$$

$$\text{gg} := 0.001130000000000000$$

$$\text{fg} := 0.000085000000000000$$

$$\text{ne} := -0.00000000000103000$$

$$\text{nm} := -0.00000000000790000$$

$$\text{nt} := 0.00000000001350000$$

$$\text{ndd1} := -2 * \text{del1u} + 2 * \text{del1u.s10} * \text{s10}$$

$$\text{ndd2} := -2 * \text{del2u} + 2 * \text{del2u.s10} * \text{s10}$$

$$\text{ndd3} := -2 * \text{del3u} + 2 * \text{del3u.s10} * \text{s10}$$

$$\text{ndd4} := -2 * \text{del4u} + 2 * \text{del4u.s10} * \text{s10}$$

$$\text{nee1} := -2 * \text{eel1u} + 2 * \text{eel1u.s10} * \text{s10}$$

$$\text{nee2} := -2 * \text{eel2u} + 2 * \text{eel2u.s10} * \text{s10}$$

$$\text{nee3} := -2 * \text{eel3u} + 2 * \text{eel3u.s10} * \text{s10}$$

$$\text{nee4} := -2 * \text{eel4u} + 2 * \text{eel4u.s10} * \text{s10}$$

$$\text{zx1} :=$$

$$\text{Coefficient}[\text{ss} * (2 * \text{Cross}[\text{s10}, \text{dd1}] + i * \text{vdd1}) +$$

$$\text{nt} * (2 * \text{Cross}[\text{s10}, \text{del1u}] + i * \text{ndd1})$$

$$+ \text{st} * (2 * \text{Cross}[\text{s10}, \text{ed1}] + i * \text{ved1})$$

$$+ \text{jss} * \text{Cross}[\text{s10}, \text{dd1}] + \text{jst} * \text{Cross}[\text{s10}, \text{ed1}] + \text{nm} * \text{Cross}[\text{s10}, \text{del1u}]$$

$$+ i * \text{ff} * \text{dd1} + i * \text{fg} * \text{ed1} + i * \text{ne} * \text{del1u}, \text{ef}, 1]$$

$$\text{zx2} :=$$

$$\text{Coefficient}[\text{ss} * (2 * \text{Cross}[\text{s10}, \text{dd2}] + i * \text{vdd2}) +$$

$$\begin{aligned}
& nt * (2 * \text{Cross}[s10, del2u] + i * ndd2) \\
& + st * (2 * \text{Cross}[s10, ed2] + i * ved2) \\
& + jss * \text{Cross}[s10, dd2] + jst * \text{Cross}[s10, ed2] + nm * \text{Cross}[s10, del2u] \\
& + i * ff * dd2 + i * fg * ed2 + i * ne * del2u, ef, 1] \\
& zx3:= \\
& \text{Coefficient}[ss * (2 * \text{Cross}[s10, dd3] + i * vdd3)+ \\
& nt * (2 * \text{Cross}[s10, del3u] + i * ndd3) \\
& + st * (2 * \text{Cross}[s10, ed3] + i * ved3) \\
& + jss * \text{Cross}[s10, dd3] + jst * \text{Cross}[s10, ed3] + nm * \text{Cross}[s10, del3u] \\
& + i * ff * dd3 + i * fg * ed3 + i * ne * del3u, ef, 1] \\
& zx4:= \\
& \text{Coefficient}[ss * (2 * \text{Cross}[s10, dd4] + i * vdd4)+ \\
& nt * (2 * \text{Cross}[s10, del4u] + i * ndd4) \\
& + st * (2 * \text{Cross}[s10, ed4] + i * ved4) \\
& + jss * \text{Cross}[s10, dd4] + jst * \text{Cross}[s10, ed4] + nm * \text{Cross}[s10, del4u] \\
& + i * ff * dd4 + i * fg * ed4 + i * ne * del4u, ef, 1] \\
& \\
& zx5:=\text{Coefficient}[st * (2 * \text{Cross}[s10, de1] + i * vde1) \\
& + tt * (2 * \text{Cross}[s10, ee1] + i * vee1) + nt * (2 * \text{Cross}[s10, eel1u] + i * nee1) \\
& + jst * \text{Cross}[s10, de1] + jtt * \text{Cross}[s10, ee1] + nm * \text{Cross}[s10, eel1u] \\
& + i * gg * ee1 + i * fg * de1 + i * ne * eel1u, ef, 1] \\
& zx6:=\text{Coefficient}[st * (2 * \text{Cross}[s10, de2] + i * vde2) \\
& + tt * (2 * \text{Cross}[s10, ee2] + i * vee2) + nt * (2 * \text{Cross}[s10, eel2u] + i * nee2) \\
& + jst * \text{Cross}[s10, de2] + jtt * \text{Cross}[s10, ee2] + nm * \text{Cross}[s10, eel2u] \\
& + i * gg * ee2 + i * fg * de2 + i * ne * eel2u, ef, 1] \\
& zx7:=\text{Coefficient}[st * (2 * \text{Cross}[s10, de3] + i * vde3) \\
& + tt * (2 * \text{Cross}[s10, ee3] + i * vee3) + nt * (2 * \text{Cross}[s10, eel3u] + i * nee3) \\
& + jst * \text{Cross}[s10, de3] + jtt * \text{Cross}[s10, ee3] + nm * \text{Cross}[s10, eel3u] \\
& + i * gg * ee3 + i * fg * de3 + i * ne * eel3u, ef, 1] \\
& zx8:=\text{Coefficient}[st * (2 * \text{Cross}[s10, de4] + i * vde4) \\
& + tt * (2 * \text{Cross}[s10, ee4] + i * vee4) + nt * (2 * \text{Cross}[s10, eel4u] + i * nee4) \\
& + jst * \text{Cross}[s10, de4] + jtt * \text{Cross}[s10, ee4] + nm * \text{Cross}[s10, eel4u] \\
& + i * gg * ee4 + i * fg * de4 + i * ne * eel4u, ef, 1]
\end{aligned}$$

S535:=Flatten[i{zx1, zx2, zx3, zx4, zx5, zx6, zx7, zx8}]

Eigenvalues[

{

Coefficient[S535, d1x, 1],

Coefficient[S535, d1y, 1],

Coefficient[S535, d1z, 1],

Coefficient[S535, d2x, 1],

–Coefficient[S535, d2y, 1],

–Coefficient[S535, d2z, 1],

–Coefficient[S535, d3x, 1],

Coefficient[S535, d3y, 1],

–Coefficient[S535, d3z, 1],

–Coefficient[S535, d4x, 1],

–Coefficient[S535, d4y, 1],

Coefficient[S535, d4z, 1],

Coefficient[S535, e1x, 1],

Coefficient[S535, e1y, 1],

Coefficient[S535, e1z, 1],

Coefficient[S535, e2x, 1],

–Coefficient[S535, e2y, 1],

–Coefficient[S535, e2z, 1],

–Coefficient[S535, e3x, 1],

Coefficient[S535, e3y, 1],

–Coefficient[S535, e3z, 1],

–Coefficient[S535, e4x, 1],

–Coefficient[S535, e4y, 1],

Coefficient[S535, e4z, 1]

}

]

{170.794, 170.794, 170.794, 4.35497, 4.35497, 4.35497, 1.74351,

1.33497, 1.33497, 1.33497, 0.10551, 0.097825, 0.097825, 0.097825,

0.00477782, 0.00477782, 0.00477782, 0.00221218, 0.00221218,
0.00221218, 0.00051, $4.7123 * 10^{\wedge} - 11$, $8.92766 * 10^{\wedge} - 12$, $1.02624 * 10^{\wedge} - 12$

Appendix B: How to include the Strong Interaction in the Tetron Scheme

As shown in a series of papers[1, 4, 5, 6, 7] the properties of gravity and of electroweak processes can be reduced to properties of tetrons. The present calculations are an example for this statement proving explicitly that the fermion mass spectrum and family mixings can be correctly obtained within the microscopic model.

In these calculations each quark flavor appears as a triplet of the Shubnikov group G_4 , eq. (3), with 3 degenerate masses. But how does the strong interaction, where quarks usually are considered as SU(3) triplets, fit into the tetron picture?

The dominant features of the strong interaction are the linear attractive potential at low energies and asymptotic freedom at high energies. In the tetron model the strong interaction is related to disturbances by the triplet isospin excitations(=quarks) of the local ground state which is formed by a single tetrahedron of isospin vectors. As triplet states of G_4 , quarks disturb the ground state's isomagnetism, whereas leptons are G_4 -singlets, i.e. 'isomagnetically' neutral. They do not disturb the ground state and can exist freely, not taking part in the strong interaction.

As discussed in [1], the isomagnetic ground state energy of 2 neighboring tetrahedrons is roughly $E_{QCD} \approx 1$ GeV corresponding to a characteristic length scale $L_{QCD} \approx 10^{-15}$ m. The linear potential between two G_4 -triplets, vulgo a quark Q and an antiquark \bar{Q} , then arises as follows: Since the inter-tetrahedral exchange energy $j = E_{QCD}$ is relatively small, its physical effects have a much longer range L_{QCD} than the weak interactions which are induced by the inner-tetrahedral exchange energies $J = O(100)$ GeV. The triplet excitations corresponding to the two quarks are characterized by small vibrations $\vec{\delta}_Q$ and $\vec{\delta}_{\bar{Q}}$ of the isospin vectors (2). When the distance between the two excitations becomes larger than L_{QCD} , an additional pair $\vec{\delta}_q$ and $\vec{\delta}_{\bar{q}}$ is excited on intermediate tetrahedrons in order to reduce the original 'isomagnetic' suspense between $\vec{\delta}_Q$ and $\vec{\delta}_{\bar{Q}}$. The associated cost in energy is proportional to the number of $q\bar{q}$ pairs created, and the potential V between Q and \bar{Q}

therefore increases linearly with distance:

$$V = F|x| \qquad F \approx -j\langle\vec{\delta}_Q\vec{\delta}_{\bar{Q}}\rangle/L_Q \qquad (108)$$

where $\langle\vec{\delta}_Q\vec{\delta}_{\bar{Q}}\rangle$ is the isospin correlation between the sites, on which the isospins vibrate, and L_{QCD} the length where all this becomes relevant. The confinement energy is hence proportional to the original ‘ferromagnetic’ exchange energy j induced on the disturbances Q and \bar{Q} . The ratio x/L_{QCD} is the number of times, an additional pair of excitations has to be created from the ‘sea’.

In the tetron model quarks are disturbances $\vec{\delta}_Q$ of the isospin vectors (2). Due to isospin interactions like (4) not only the ground state vectors but also the disturbances tend to align. This tendency of the triplet excitations gives rise to a ‘mass gap’ $\langle\vec{\delta}_Q\vec{\delta}_{\bar{Q}}\rangle \neq 0$ which signals a phase transition in the form of the usual breakdown of chiral symmetry due to the strong interactions.

In summary, a single quark Q increases the energy of the system in its neighborhood L_{QCD} not only by its flavor-dependent mass(=excitation energy) but by an additional energy necessary to ‘pick up a $q\bar{q}$ pair from the sea’. This energy is flavor independent, because it does not depend on the flavor Q , which flavors q are excited. The flavors q correspond to an average of the light quarks u , d and s . So when a Q and a \bar{Q} are torn apart, at some distance $x \approx L_{QCD}$ a light $q\bar{q}$ pair is formed, because otherwise the single quark Q could not endure the disturbance of the ground state. In the end a sort of string appears obtained by $Q\bar{q}q\bar{q}'\dots\bar{Q}$ pairs. Any time a new $q\bar{q}$ pair is created, energy is to be taken from the environment, so the associated cost in energy is proportional to the number of $q\bar{q}$ pairs and the potential between quark and antiquark therefore increases linearly with distance as indicated in (108).

Readers familiar with the strong interaction, will recognize that one is led this way to the classic ideas of the quark model. For example, using the linear potential (108) masses of mesons and baryons can be estimated just as in the quark model. Since mesons and baryons are G_4 -singlets, the ‘isomagnetic’ disturbances induced by quarks get neutralized in these bound states, i.e. mesons and baryons do not disturb the ground state of a single tetrahedron.

The role of the length L_{QCD} , where the creation of a light quark-antiquark is enforced, is the same as in the Standard Model. At distances above L_{QCD} one has

confinement, while below L_{QCD} the strong force diminishes. Virtual bound quark-antiquark pairs are formed which as gluons mediate a strong interaction of the original $Q\bar{Q}$ pair which effectively can be described by the QCD Lagrangian with its local $SU(3)$ gauge symmetry. As well known, this interaction dies out when the energies involved go to infinity, i.e. one has asymptotic freedom.

References

- [1] see the review B. Lampe, arXiv:1505.03477 [hep-ph] (2015).
- [2] R. Slansky, Phys. Rept. 79 (1981) 1.
- [3] G. Ross, Grand Unified Theories, Cummings Publishing Company, 1984.
- [4] B. Lampe, Progr. of Phys. 69 (2021) 2000072.
- [5] B. Lampe, Int. J. Mod. Phys. A30 (2015) 1550025.
- [6] B. Lampe in Proc. of the MG16 Meeting on General Relativity, 5–10 July 2021, editors R. Ruffini and G. Vereshchagin, <https://doi.org/10.1142/13149> (2023) 999.
- [7] B. Lampe, Int. J. Mod. Phys. A30 (2015) 1550026.
- [8] S. Oneda and K. Terasaki, Prog. Theor. Phys. Suppl. 82 (1985) 1.
- [9] R.M. White, Quantum Theory of Magnetism, Springer Verlag, ISBN-10 3-540-65116-0.
- [10] A.P. Cracknell, Progr. Theor. Phys. 35 (1966) 196.
- [11] I. Dzyaloshinskii, J. Phys. Chem. Sol. 4 (1958) 241.
- [12] T. Moriya, Phys. Rev. 120 (1960) 91.
- [13] W. Heisenberg, Zeitschrift f. Phys. 38 (1926) 411.
- [14] J. W. Moffat, Int. J. Mod, Phys. D2, 351 (1993).

- [15] J. D. Barrow, Phys. Rev. Lett. 82 (1999) 884.
- [16] J. D. Bekenstein, J. D. Phys.Rev. D25 (1982) 1527.
- [17] A. Belyaev et al., Phys. Rev. D97 (2018) 035011.
- [18] H. Fritzsche and J. Sola, Adv. High Energy Phys. 2014 (2014) 361587.
- [19] J. Magueijo, Rept. Prog. Phys. 66 (2003) 2025.
- [20] J. K. Webb et al. Phys.Rev.Lett. 87 (2001) 091301.
- [21] M. Elhadj, B. Canals, R.S. Borrell and C. Lacroix, Phys. Rev. B71 (2005) 094420.
- [22] W. Zheng et al., Phys. Rev. Lett. 96 (2006) 057201.
- [23] H. Ishimori et. al., Prog. Theor. Phys. Suppl. 183 (2010) 1.
- [24] M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, JHEP 1411, 052 (2014).
- [25] G. Altarelli et al., JHEP 21 (2012).
- [26] H. Fritzsche, Mod. Phys. Lett. A36 (2021) 2150097.
- [27] C. Jarlskog, Z. Phys. C29 (1985) 491.
- [28] Particle Data Group, Prog. Theor. Exp. Phys. (2022) 083C01.
- [29] A. R. Ellis, K. J. Kelly and S. W. Li, Phys. Rev. D 102 (2020) 115027 .
- [30] K. Bora, arXiv:1206.5909 [hep-ph] (2012).
- [31] N. Cabibbo, Phys. Rev. Lett. 10 (1963) 531.
- [32] Z. Maki, M. Nakagawa and S. Sakata, Progr.Theor. Phys. 28 (1962) 870.
- [33] C. Giganti, S. Lavignac and M. Zito, Progr. Part. Nucl.Phys 98 (2018) 1.
- [34] J. A. Belinchon et al., Int. J. Mod. Phys. D12 (2003) 1113.
- [35] J. C. Carvalho et al. Phys. Rev. D46 (1992) 2404.

- [36] T. Chiba, Prog. Theor. Phys. 126 (2011) 993.
- [37] X. Calmet et al., Phys. Lett. B540 (2002) 173.
- [38] W. Juarez et al., Phys. Rev. D66 (2002) 116007.
- [39] C. L. Gardner, Phys. Rev. D68 (2003) 043513.
- [40] D. F. Mota et al. Phys.Lett. B581 (2004) 141.
- [41] D. Parkinson et al. Phys. Lett. B578 (2004) 235.
- [42] E. J. Copeland, Phys. Rev. D69 (2004) 023501.
- [43] A.S. Borovik-Romanov and H. Grimmer, International Tables for Crystallography D (2006) 105.
- [44] G. C. Branco, L. Lavoura and J. P. Silva, CP Violation, Oxford University Press, 1999.
- [45] Wolfram Research, Inc., Mathematica, Version 13.2, Champaign, IL (2022).
- [46] B. lampe, arXiv:2308.08498 [hep-ph].