On the Work of Cartan and Münzner on Isoparametric Hypersurfaces

Thomas E. Cecil

November 8, 2024

Abstract

A hypersurface M^n in a real space form \mathbf{R}^{n+1} , S^{n+1} , or H^{n+1} is isoparametric if it has constant principal curvatures. This paper is a survey of the fundamental work of Cartan and Münzner on the theory of isoparametric hypersurfaces in real space forms, in particular, spheres. This work is contained in four papers of Cartan [3]–[6] published during the period 1938–1940, and two papers of Münzner [44]–[45] that were published in preprint form in the early 1970's, and as journal articles in 1980–1981. These papers of Cartan and Münzner have been the foundation of the extensive field of isoparametric hypersurfaces, and they have all been recently translated into English by the author. The paper concludes with a brief survey of the recently completed classification of isoparametric hypersurfaces in spheres in Section 4.

1 Introduction

A hypersurface M^n in a real space form \mathbf{R}^{n+1} , S^{n+1} , or H^{n+1} is isoparametric if it has constant principal curvatures. This paper is a survey of the fundamental work of Cartan and Münzner on the theory of isoparametric hypersurfaces in real space forms. During the period 1938–1940, Cartan [3]–[6] published four papers that classified isoparametric hypersurfaces in \mathbf{R}^{n+1} and H^{n+1} , and laid the groundwork for the theory of isoparametric hypersurfaces in the sphere S^{n+1} , especially in the case where the principal curvatures of the hypersurface all have the same multiplicity. Approximately thirty years later in the early 1970's, Münzner [44]–[45] published two preprints which extended some of Cartan's main results to the arbitrary case where there are no assumptions on the multiplicities of the principal curvatures. Münzner then proved his major result that the number g of distinct principal curvatures of an isoparametric hypersurface in a sphere satisfies the restriction g = 1, 2, 3, 4 or 6. Münzner's articles were later published as journal articles in 1980–1981.

These papers of Cartan and Münzner have proven to be the foundation for the extensive theory of isoparametric hypersurfaces, which has been developed by many researchers over the years. The classification of isoparametric hypersurfaces in spheres has been recently completed (see Chi [20]), and this paper concludes with a brief survey of the classification results in Section 4.

The author has recently published English translations of all six of these papers Cartan and Münzner at CrossWorks, the publishing site for the College of the Holy Cross. References for these translations are listed together with the references for the papers of Cartan and Münzner in the bibliography.

In general, this paper follows the treatment of the theory of isoparametric hypersurfaces in the book of Cecil and Ryan [13, pp. 85–137], and some passages are taken directly from that book. Several proofs of theorems are omitted in this paper, but they are included in the book [13], and detailed references are given for the proofs.

The author wishes to acknowledge valuable collaborations over many years on the theory of isoparametric hypersurfaces with Quo-Shin Chi, Gary Jensen, and Patrick Ryan.

2 Cartan's Work

2.1 Families of Isoparametric Hypersurfaces

We begin with some preliminary notation and definitions, which generally follow those in the book [13]. By a real space form of dimension n, we mean a complete, connected, simply connected manifold $\widetilde{M}^n(c)$ with constant sectional curvature c. If c = 0, then $\widetilde{M}^n(c)$ is the *n*-dimensional Euclidean space \mathbb{R}^n ; if c = 1, then $\widetilde{M}^n(c)$ is the unit sphere $S^n \subset \mathbb{R}^{n+1}$; and if c = -1, then $\widetilde{M}^n(c)$ is the *n*-dimensional real hyperbolic space H^n (see, for example, [38, Vol. I, pp. 204–209]). For any value of c > 0, this subject is basically the same as for the case c = 1, and for any value of c < 0, it is very similar to the case c = -1. So we will restrict our attention to the cases c = 0, 1, -1.

The original definition of a family of isoparametric hypersurfaces in a real space form \widetilde{M}^{n+1} was formulated in terms of the level sets of an isoparametric function, as we now describe. Let $F: \widetilde{M}^{n+1} \to \mathbf{R}$ be a nonconstant smooth function. The classical Beltrami differential parameters of F are defined by

$$\Delta_1 F = |\text{grad } F|^2, \quad \Delta_2 F = \Delta F \text{ (Laplacian of } F),$$
(1)

where grad F denotes the gradient vector field of F.

The function F is said to be *isoparametric* if there exist smooth functions ϕ_1 and ϕ_2 from **R** to **R** such that

$$\Delta_1 F = \phi_1(F), \quad \Delta_2 F = \phi_2(F). \tag{2}$$

That is, both of the Beltrami differential parameters are constant on each level set of F. This is the origin of the term *isoparametric*. The collection of level sets of an isoparametric function is called an *isoparametric family* of sets in \widetilde{M}^{n+1} .

An isoparametric family in \mathbb{R}^{n+1} consists of either parallel planes, concentric spheres, or coaxial spherical cylinders, and their focal sets. This was first shown for n = 2 by Somigliana [56] (see also B. Segre [52] and Levi-Civita [40]), and for arbitrary n by B. Segre [53].

In the late 1930's, shortly after the publication of the papers of Levi-Civita and Segre, Cartan [3]–[6] began a study of isoparametric families in arbitrary real space forms. Cartan began his first paper [3] with an extract of a letter that he had recently written to B. Segre. Cartan states that he can extend Segre's results for isoparametric families in \mathbb{R}^{n+1} to isoparametric families in hyperbolic space H^{n+1} , and that he has made progress in the study of isoparametric families in the sphere S^{n+1} . Cartan notes that he employs a different technique than the one that Segre used. Cartan uses the method of moving frames and the theory of parallel hypersurfaces, which we will briefly review here.

2.2 Parallel Hypersurfaces

Let $f: M^n \to \widetilde{M}^{n+1}(c)$ be an oriented hypersurface with field of unit normals ξ . For $x \in M^n$, let $T_x M^n$ denote the tangent space to M^n at x. For any vector X in the tangent space $T_x M^n$, we have the fundamental equation

$$\widetilde{\nabla}_{f_*(X)}\xi = -f_*(A_\xi X),\tag{3}$$

where $\widetilde{\nabla}$ is the Levi-Civita connection in \widetilde{M} , f_* is the differential of f, and A_{ξ} is the *shape operator* determined by the normal vector field ξ .

The shape operator A_{ξ} is a symmetric tensor of type (1, 1) on M^n . An eigenvalue λ of A_{ξ} is called a *principal curvature* of A_{ξ} , and a corresponding eigenvector is called a *principal vector*. We often write A instead of A_{ξ} , in the case where the choice of field of unit normals ξ has been specified.

The parallel hypersurface to $f(M^n)$ at signed distance $t \in \mathbf{R}$ is the map $f_t: M^n \to \widetilde{M}^{n+1}(c)$ such that for each $x \in M^n$, the point $f_t(x)$ is obtained by traveling a signed distance t along the geodesic in $\widetilde{M}^{n+1}(c)$ with initial point f(x) and initial tangent vector $\xi(x)$. For $\widetilde{M}^{n+1}(c) = \mathbf{R}^{n+1}$, the formula for f_t is

$$f_t(x) = f(x) + t \xi(x),$$
 (4)

and for $\widetilde{M}^{n+1}(c) = S^{n+1}$, the formula for f_t is

$$f_t(x) = \cos t \ f(x) + \sin t \ \xi(x). \tag{5}$$

There is a similar formula in hyperbolic space H^{n+1} (see, for example, [10]).

Locally, for sufficiently small values of t, the map f_t is also an immersed hypersurface. However, the map f_t may develop singularities at focal points of the original hypersurface $f(M^n)$. Specifically, a point $p = f_t(x)$ is called a focal point of (M^n, x) of multiplicity m > 0 if the differential $(f_t)_*$ has nullity m at x.

In the cases $\widetilde{M}^{n+1}(c) = \mathbf{R}^{n+1}$, respectively S^{n+1} , the point $p = f_t(x)$ is a focal point of (M^n, x) of multiplicity m > 0 if and only if 1/t, respectively $\cot t$, is a principal curvature of M^n of multiplicity m at x. Thus, in the case $\widetilde{M}^{n+1}(c) = S^{n+1}$, each principal curvature λ gives rise to two antipodal focal points along the normal geodesic, since $\lambda = \cot t = \cot(t + \pi)$ (see, for example, [13, p. 11]). There is a similar formula for focal points in the ambient space H^{n+1} (see, for example, [10] or [13, p. 11]).

Now let $f: M^n \to \widetilde{M}^{n+1}(c)$ be an oriented hypersurface with constant principal curvatures. One can show from the formulas for the principal curvatures of a parallel hypersurface that if f has constant principal curvatures, then each immersed hypersurface f_t also has constant principal curvatures (see, for example, [13, pp. 17–18] or Subsection 3.1 of this paper).

Since the principal curvatures are constant on M^n , the focal points along the normal geodesic to $f(M^n)$ at f(x) occur for the same values of t, independent of the choice of the point $x \in M^n$. In the case of \mathbf{R}^{n+1} , for example, this means that if μ is a nonzero constant principal curvature of multiplicity m > 0 of M^n , then the map $f_{1/\mu}$ has constant rank n - m on M^n , and the set $f_{1/\mu}(M^n)$ is an (n - m)-dimensional submanifold of \mathbf{R}^{n+1} , called a *focal* submanifold of $f(M^n)$. Similar results hold for the other ambient space forms (see, for example, [13, pp. 18–32]).

2.3 Isoparametric and Parallel Hypersurfaces

In Section 2 of his first paper, Cartan [3] showed that an isoparametric family, defined as the collection of level sets of an isoparametric function F, is equivalent to a family of parallel hypersurfaces, each of which has constant principal curvatures, together with their focal submanifolds.

We now discuss the main steps in this argument. We will omit most of the proofs, and we refer the reader to [13, pp. 86–90] for detailed proofs of these steps, which are contained in Theorems 2.1 - 2.5 below.

Let $F: \widetilde{M}^{n+1} \to \mathbf{R}$ be a nonconstant smooth function defined on a real space form \widetilde{M}^{n+1} . Suppose that grad F does not vanish on the level set $M = F^{-1}(0)$. Then M is a smooth hypersurface in \widetilde{M}^{n+1} , and the shape operator A of M satisfies the equation

$$\langle AX, Y \rangle = -\frac{H_F(X, Y)}{|\text{grad } F|},$$
(6)

where X and Y are tangent vectors to M, and H_F is the Hessian of the function F.

Then one can show by a calculation, that the mean curvature h of the level hypersurface M is given by the following theorem.

Theorem 2.1. The mean curvature h of the level hypersurface M is given by

$$h = \frac{1}{n\rho^2} (\langle \text{grad } F, \text{ grad } \rho \rangle - \rho \Delta F), \qquad (7)$$

where $\rho = |\text{grad } F|$.

The following theorem of Cartan is important in showing that the level sets of an isoparametric function have constant principal curvatures.

Theorem 2.2. If $F : \widetilde{M}^{n+1} \to \mathbf{R}$ is an isoparametric function on a real space form, then each level hypersurface of F has constant mean curvature.

The vector field $\xi = \text{grad } F/|\text{grad } F|$ is defined on the open subset of \widetilde{M}^{n+1} on which grad F is nonzero. A direct calculation then yields the following theorem that the integral curves of ξ are geodesics in \widetilde{M}^{n+1} .

Theorem 2.3. Let $F : \widetilde{M}^{n+1} \to \mathbf{R}$ be a function for which |grad F| is a function of F. Then on the subset of \widetilde{M}^{n+1} where |grad F| is nonzero, the integral curves of the vector field $\xi = \text{grad } F/|\text{grad } F|$ are geodesics in \widetilde{M}^{n+1} .

Theorem 2.3 shows that a family of level hypersurfaces of an isoparametric function is a family of parallel hypersurfaces in \widetilde{M}^{n+1} , modulo reparametrization to take into account the possibility that |grad F| is not identically equal to one. The next step is to show that each of these level hypersurfaces has constant principal curvatures. This follows from Theorem 2.2 and the next theorem of Cartan.

Theorem 2.4. Let $f_t : M \to \widetilde{M}^{n+1}$, $-\varepsilon < t < \varepsilon$, be a family of parallel hypersurfaces in a real space form. Then f_0M has constant principal curvatures if and only if each f_tM has constant mean curvature.

As a consequence of Theorems 2.2–2.4, we obtain the following theorem of Cartan.

Theorem 2.5. If $F : \widetilde{M}^{n+1} \to \mathbf{R}$ is an isoparametric function on a real space form, then each level hypersurface of F has constant principal curvatures.

Conversely, let $f_t : M \to \widetilde{M}^{n+1}$, $-\varepsilon < t < \varepsilon$, be a family of parallel hypersurfaces such that f_0 has constant principal curvatures. We see that this is an isoparametric family of hypersurfaces as follows.

By the well known formulas for the principal curvatures of a parallel hypersurface in a real space form (see, for example, [13, pp. 17–18] and Subsection 3.1 of this paper), each $f_t M$ has constant principal curvatures, and thus each $f_t M$ has constant mean curvature. Then the function F defined by F(x) = t, if $x \in f_t M$, is a smooth function defined on an open subset of \widetilde{M}^{n+1} with the property that grad $F = \xi$ is a unit length vector field such that $\widetilde{\nabla}_{\xi} \xi = 0$.

Furthermore, since the function $\rho = |\text{grad } F| = 1$ is constant, we see from Theorem 2.1 that the constancy of the mean curvature h on each level hypersurface $f_t M$ implies that the Laplacian of F is also constant on each level set $f_t M$, and therefore F is an isoparametric function, as defined by equation (2). Thus, this family of parallel hypersurfaces is a family of isoparametric hypersurfaces in the original sense, that is, it is the family of level sets of an isoparametric function.

In summary, the analytic definition of an isoparametric family of hypersurfaces in terms of level sets of an isoparametric function on a real space form \widetilde{M}^{n+1} is equivalent to the geometric definition of an isoparametric family as a family of parallel hypersurfaces to a hypersurface with constant principal curvatures.

Note that this is not true if \widetilde{M}^{n+1} is only assumed to be a Riemannian manifold, as can be seen by examples in complex projective space due to Q.-M. Wang [73] (see also Thorbergsson [72] and [13, pp. 526–530]).

We will therefore say that a connected hypersurface M^n immersed in a real space form \widetilde{M}^{n+1} is an *isoparametric hypersurface* if it has constant principal curvatures.

2.4 Cartan's Formula

In Section 3 of his first paper on isoparametric hypersurfaces, Cartan [3] derived what he called a "fundamental formula," which is the basis for many of his results on the subject.

The formula involves the distinct principal curvatures $\lambda_1, \ldots, \lambda_g$, and their respective multiplicities m_1, \ldots, m_g , of an isoparametric hypersurface $f: M^n \to \widetilde{M}^{n+1}(c)$ in a space form of constant sectional curvature c. If the number g of distinct principal curvatures is greater than one, Cartan showed that for each $i, 1 \leq i \leq g$, the following equation holds,

$$\sum_{j \neq i} m_j \frac{c + \lambda_i \lambda_j}{\lambda_i - \lambda_j} = 0.$$
(8)

Cartan's original formulation of the formula was given in a more complicated form, but he gave the formulation in equation (8) in his third paper [5, p. 1484]. Equation (8) is now known as "Cartan's formula." Another proof of this formula is given in [13, pp. 91–96].

2.5 Isoparametric Hypersurfaces in \mathbb{R}^{n+1}

Using his formula, Cartan was able to completely classify isoparametric hypersurfaces in Euclidean space \mathbf{R}^{n+1} (c = 0) and in hyperbolic space H^{n+1}

(c = -1), and to make significant progress on the theory of isoparametric hypersurfaces in the sphere S^{n+1} (c = 1), as we will discuss below.

We first consider an isoparametric hypersurface in Euclidean space. The classification is a local theorem, so we consider the hypersurface to be embedded in \mathbf{R}^{n+1} . Here $S^k(r)$ denotes a k-dimensional sphere of radius r in a totally geodesic $\mathbf{R}^{k+1} \subset \mathbf{R}^{n+1}$.

Theorem 2.6. Let $M^n \subset \mathbf{R}^{n+1}$ be a connected isoparametric hypersurface. Then M^n is an open subset of a flat hyperplane, a metric hypersphere, or a spherical cylinder $S^k(r) \times \mathbf{R}^{n-k}$.

Proof. If the number g of distinct principal curvatures of M^n is one, then M^n is totally umbilic, and it is well known that M^n is an open subset of a hyperplane or hypersphere in \mathbf{R}^{n+1} (see, for example, [57, Vol. 4, p. 110]).

If $g \geq 2$, then by taking an appropriate choice of unit normal field ξ , one can assume that at least one of the principal curvatures is positive. If λ_i is the smallest positive principal curvature, then each term $\lambda_i \lambda_j / (\lambda_i - \lambda_j)$ in Cartan's formula (8) with c = 0 is non-positive, and thus it equals zero. Hence, there can be at most two distinct principal curvatures, and if there are two, then one of them equals zero.

If g = 2, then by an appropriate choice of the unit normal field ξ , we can arrange that the nonzero principal curvature is positive. Thus, suppose that the principal curvatures are $\lambda_1 > 0$ with multiplicity $m_1 = k$, and $\lambda_2 = 0$ with multiplicity $m_2 = n - k$. Then for $t = 1/\lambda_1$, the focal submanifold $V = f_t(M^n)$ has dimension n - k, and it is totally geodesic in \mathbb{R}^{n+1} , since the formulas for the shape operator of a parallel map f_t [13, pp. 17-18] show that for every unit normal η to V at every point p of V, the shape operator A_η has one distinct principal curvature given by

$$\frac{\lambda_2}{(1-t\lambda_2)} = 0. \tag{9}$$

Thus, V is contained in a totally geodesic submanifold $\mathbf{R}^{n-k} \subset \mathbf{R}^{n+1}$, and M^n is an open subset of a tube of radius $1/\lambda_1$ over \mathbf{R}^{n-k} . Such a tube is a spherical cylinder $S^k(r) \times \mathbf{R}^{n-k}$, where $S^k(r)$ is a k-dimensional sphere of radius $r = 1/\lambda_1$ in a totally geodesic $\mathbf{R}^{k+1} \subset \mathbf{R}^{n+1}$ orthogonal to \mathbf{R}^{n-k} . \Box

2.6 Isoparametric Hypersurfaces in H^{n+1}

Next we consider an isoparametric hypersurface $M^n \subset H^{n+1}$ in hyperbolic space (c = -1). As we will see in the proof of Theorem 2.7 below, Cartan used his formula again to show that the number g of distinct principal curvatures must be less than or equal to 2. The rest of the proof then follows in a way similar to the Euclidean case above. Again this is a local result, so we consider the hypersurface to be embedded.

Theorem 2.7. Let $M^n \subset \mathbf{H}^{n+1}$ be a connected isoparametric hypersurface. Then M^n is an open subset of a totally geodesic hyperplane, an equidistant hypersurface, a horosphere, a metric hypersphere, or a tube over a totally geodesic submanifold of codimension greater than one in H^{n+1} .

Proof. Let g be the number of distinct principal curvatures of M^n . If g = 1, then M^n is totally umbilic, and it is an open subset of a totally geodesic hyperplane, an equidistant hypersurface, a horosphere, or a metric hypersphere in H^{n+1} (see, for example, [57, Vol. 4, p. 114]).

If $g \geq 2$, then by an appropriate choice of the unit normal field ξ , we can arrange that at least one of the principal curvatures is positive. Then there exists a positive principal curvature λ_i such that no principal curvature lies between λ_i and $1/\lambda_i$. (In fact, λ_i is either the largest principal curvature between 0 and 1, or the smallest principal curvature greater than or equal to one.)

For this λ_i , each term $(-1 + \lambda_i \lambda_j)/(\lambda_i - \lambda_j)$ in Cartan's formula (8) with c = -1 is negative unless $\lambda_j = 1/\lambda_i$, in which case the term is zero. Therefore, there can be at most two distinct principal curvatures, and if there are two, they are reciprocals of each other.

In that case, suppose that the two principal curvatures are $\lambda_1 = \coth \theta$ with multiplicity $m_1 = k$, and $\lambda_2 = 1/\lambda_1 = \tanh \theta$ with multiplicity $m_2 = n-k$. If we take $t = \theta$, then $V = f_t(M^n)$ is a focal submanifold of dimension n-k, and it is totally geodesic in H^{n+1} by the formulas for the shape operator of a parallel map f_t [13, pp. 17-18].

Thus, this focal submanifold V is contained in a totally geodesic submanifold $H^{n-k} \subset H^{n+1}$, and $f(M^n)$ is an open subset of a tube of radius $t = \theta$ over H^{n-k} . Such a tube is standard Riemannian product $S^k(c_1) \times H^{n-k}(c_2)$ in hyperbolic space H^{n+1} , where $c_1 = 1/\sinh^2 \theta$ and $c_2 = -1/\cosh^2 \theta$ are the constant sectional curvatures of the sphere $S^k(c_1)$ and the hyperbolic space $H^{n-k}(c_2)$, respectively (see Ryan [50]).

2.7 Isoparametric Hypersurfaces in S^{n+1}

In the sphere S^{n+1} , however, Cartan's formula does not lead to the conclusion that $g \leq 2$. In fact, Cartan produced examples with g = 1, 2, 3 or 4 distinct principal curvatures. Moreover, he classified isoparametric hypersurfaces $M^n \subset S^{n+1}$ with $g \leq 3$ in [4] (see Section 2.9).

In the case g = 1, the hypersurface M^n is totally umbilic, and it is well known that M^n is an open subset of a great or small hypersphere in S^{n+1} (see [57, Vol. 4, p. 112]).

If g = 2, then M^n is an open subset of a standard product of two spheres,

$$S^{p}(r) \times S^{q}(s) \subset S^{n+1}(1) \subset \mathbf{R}^{p+1} \times \mathbf{R}^{q+1} = \mathbf{R}^{n+2}, \quad r^{2} + s^{2} = 1,$$
 (10)

where n = p + q, and r > 0, s > 0. The proof of this result is similar to the proofs of Theorems 2.6 and 2.7 above, i.e., one shows that a focal submanifold is totally geodesic, and M^n is a tube of constant radius over it. This is true for both focal submanifolds in this case (see, for example, [13, pp. 110–111]).

The case of g = 3 distinct principal curvatures is much more difficult, and Cartan's thorough treatment of that case in [4] is a highlight of his work on isoparametric hypersurfaces.

Specifically, Cartan [4] showed that in the case g = 3, all the principal curvatures must have the same multiplicity m, which must be one of the values m = 1, 2, 4 or 8. Furthermore, M^n must be an open subset of a tube of constant radius over a standard embedding of a projective plane \mathbf{FP}^2 into S^{3m+1} , where \mathbf{F} is the division algebra \mathbf{R} , \mathbf{C} , \mathbf{H} (quaternions), \mathbf{O} (Cayley numbers), for m = 1, 2, 4, 8, respectively.

Thus, up to congruence, there is only one such family of isoparametric hypersurfaces for each value of m. For each of these hypersurfaces, the focal set of M^n consists of two antipodal standard embeddings of \mathbf{FP}^2 , and M^n is a tube of constant radius over each focal submanifold. We will describe Cartan's work in the case g = 3 in more detail in Section 2.9 below.

2.8 The case where all multiplicities are equal

Cartan [3, pp. 186–187] gave a method for determining the values of the principal curvatures of an isoparametric hypersurface M^n , if one knows the value of g and the multiplicities m_1, \ldots, m_g , as in equation (17). This is method is based on an analytic calculation involving Cartan's formula (8).

In the case where all of the principal curvatures have the same multiplicity m = n/g, Cartan's method [5, pp. 34–35] yields the formula for the principal curvatures given in Theorem 2.8 below.

This formula implies that for any point $x \in M$, there are 2g focal points of (M, x) along the normal geodesic to M through x, and they are evenly distributed at intervals of length π/g .

Theorem 2.8 (Cartan). Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface with g principal curvatures $\lambda_i = \cot \theta_i, \ 0 < \theta_1 < \cdots < \theta_g < \pi$, all having the same multiplicity m = n/g. Then

$$\theta_i = \theta_1 + (i-1)\frac{\pi}{g}, \quad 1 \le i \le g.$$
(11)

For any point $x \in M$, there are 2g focal points of (M, x) along the normal geodesic to M through x, and they are evenly distributed at intervals of length π/g .

Remark 2.1. Münzner [44] later showed that formula (11) is valid for all isoparametric hypersurfaces in S^{n+1} , even those where the multiplicities are not all the same. Münzner's method is different than the method of Cartan. Münzner's proof is based on the fact that the set of focal points along a normal geodesic circle to $M \subset S^{n+1}$ is invariant under the dihedral group D_g of order 2g that acts on the normal circle and is generated by reflections in the focal points. Münzner's proof also yields the important result that the multiplicities satisfy $m_i = m_{i+2}$ (subscripts mod g). We will give a detailed proof of Münzner's result in Theorem 3.3 in Subsection 3.2.2 below.

Using a lengthy calculation involving equation (11), Cartan [4, pp. 364-367] proved the important result that any isoparametric family with g distinct principal curvatures all having the same multiplicity is algebraic in sense of Theorem 2.9 below.

Recall that a function $F : \mathbf{R}^{n+2} \to \mathbf{R}$ is homogeneous of degree g if $F(tx) = t^g F(x)$, for all $t \in \mathbf{R}$ and $x \in \mathbf{R}^{n+2}$.

Theorem 2.9 (Cartan). Let $M \subset S^{n+1} \subset \mathbf{R}^{n+2}$ be a connected isoparametric hypersurface with g principal curvatures $\lambda_i = \cot \theta_i$, $0 < \theta_1 < \cdots < \theta_g < \pi$, all having the same multiplicity m. Then M is an open subset of a level set of the restriction to S^{n+1} of a homogeneous polynomial F on \mathbf{R}^{n+2} of degree g satisfying the differential equations,

$$\Delta_1 F = |\text{grad } F|^2 = g^2 r^{2g-2} = g^2 (x_1^2 + x_2^2 + \dots + x_{n+2}^2)^{g-1}, \qquad (12)$$

$$\Delta_2 F = \Delta F = 0 \ (F \ is \ harmonic), \tag{13}$$

where r = |x|.

Cartan's Theorem 2.9 was a forerunner of Münzner's general result (see Theorem 3.5 below) that every isoparametric hypersurface with g distinct principal curvatures is algebraic (regardless of the multiplicities of the principal curvatures), and its defining homogeneous polynomial F of degree gsatisfies certain conditions on $\Delta_1 F$ and $\Delta_2 F$, which generalize the conditions that Cartan found in Theorem 2.9.

2.9 Hypersurfaces With Three Principal Curvatures

The classification of isoparametric hypersurfaces with g = 3 distinct principal curvatures in [4] is certainly one of the highlights of Cartan's work on isoparametric hypersurfaces. In this paper, Cartan gives a complete classification of isoparametric hypersurfaces in S^{n+1} with g = 3 principal curvatures, and he describes each example in great detail.

Let $M^n \subset S^{n+1} \subset \mathbf{R}^{n+2}$ be a connected, oriented hypersurface with field of unit normals ξ having g = 3 distinct principal curvatures at every point. The first step in Cartan's classification theorem is to prove the following theorem [4, pp. 359–360].

Theorem 2.10. Let $M^n \subset S^{n+1}$ be a connected isoparametric hypersurface with g = 3 principal curvatures at each point. Then all of the principal curvatures have the same multiplicity m.

Cartan proves this using the method of moving frames and conditions on the structural formulas that follow from the isoparametric condition and the assumption that g = 3, and we refer the reader to Cartan's paper [4, pp. 359–360] for the proof.

Next, since all of the multiplicities are equal, Cartan's Theorem 2.9 [4, pp. 364-367] above implies that M^n is an open subset of a level set of the restriction to S^{n+1} of a harmonic homogeneous polynomial F on \mathbb{R}^{n+2} of degree g = 3 satisfying the differential equations stated in Theorem 2.9.

Cartan's approach to classifying isoparametric hypersurfaces with g = 3 is to try to directly determine all homogeneous harmonic polynomials F of degree g = 3 that satisfy the differential equation from Theorem 2.9,

$$\Delta_1 F = |\text{grad } F|^2 = g^2 r^{2g-2} = g^2 (x_1^2 + x_2^2 + \dots + x_{n+2}^2)^{g-1}.$$
(14)

After a long calculation, Cartan shows that the algebraic determination of such a harmonic polynomial requires the possibility of solving the algebraic problem:

Problem 1: Represent the product

$$(u_1^2 + u_2^2 + \dots + u_m^2)(v_1^2 + v_2^2 + \dots + v_m^2)$$

of two sums of m squares by the sum of the squares of m bilinear combinations of the u_i and the v_j .

By a theorem of Hurwitz [35] on normed linear algebras, this is only possible if m = 1, 2, 4, or 8.

Cartan [4, pp. 340–341] relates the solution of Problem 1 above to the theory of Riemannian spaces admitting an isogonal absolute parallelism, and he uses results from a joint paper of himself and J. A. Schouten [7] on that topic to determine the possibilities for F.

Ultimately, Cartan [4] shows that the required harmonic homogeneous polynomial F of degree 3 on \mathbb{R}^{3m+2} must be of the form F(x, y, X, Y, Z) given by

$$x^{3} - 3xy^{2} + \frac{3}{2}x(X\overline{X} + Y\overline{Y} - 2Z\overline{Z}) + \frac{3\sqrt{3}}{2}y(X\overline{X} - Y\overline{Y}) + \frac{3\sqrt{3}}{2}(XYZ + \overline{ZYX}).$$
(15)

In this formula, x and y are real parameters, while X, Y, Z are coordinates in the division algebra $\mathbf{F} = \mathbf{R}, \mathbf{C}, \mathbf{H}$ (quaternions), **O** (Cayley numbers), for m = 1, 2, 4, 8, respectively.

Note that the sum $XYZ + \overline{ZYX}$ is twice the real part of the product XYZ. In the case of the Cayley numbers, multiplication is not associative, but the real part of XYZ is the same whether one interprets the product as (XY)Z or X(YZ).

The isoparametric hypersurfaces in the family are the level sets M_t in S^{3m+1} determined by the equation $F = \cos 3t$, $0 < t < \pi/3$, where F is the polynomial in equation (15). The focal submanifolds are obtained by taking t = 0 and $t = \pi/3$. These focal submanifolds are a pair of antipodal standard embeddings of the projective plane \mathbf{FP}^2 , for the appropriate division algebra \mathbf{F} listed above (see, for example, [13, pp. 74–78]). In the case of $\mathbf{F} = \mathbf{R}$, these focal submanifolds are standard Veronese surfaces in $S^4 \subset \mathbf{R}^5$.

For the cases $\mathbf{F} = \mathbf{R}, \mathbf{C}, \mathbf{H}$, Cartan gave a specific parametrization of the focal submanifold M_0 defined by the condition F = 1 as follows:

$$X = \sqrt{3}v\overline{w}, \quad Y = \sqrt{3}w\overline{u}, \quad Z = \sqrt{3}u\overline{v}, \quad (16)$$
$$x = \frac{\sqrt{3}}{2}(|u|^2 - |v|^2), \quad y = |w|^2 - \frac{|u|^2 + |v|^2}{2},$$

where u, v, w are in **F**, and $|u|^2 + |v|^2 + |w|^2 = 1$. This map is invariant under the equivalence relation

$$(u, v, w) \sim (u\lambda, v\lambda, w\lambda), \quad \lambda \in \mathbf{F}, \ |\lambda| = 1.$$

Thus, it is well-defined on \mathbf{FP}^2 , and it is easily shown to be injective on \mathbf{FP}^2 . Therefore, it is an embedding of \mathbf{FP}^2 into S^{3m+1} .

Cartan states that he does not know of an analogous representation of the focal variety in the case m = 8. In that case, if u, v, w are taken to be Cayley numbers, then the formulas (16) are not well defined on the Cayley projective plane. For example, the product $v\overline{w}$ is not preserved, in general, if we replace v by $v\lambda$, and w by $w\lambda$, where λ is a unit Cayley number.

Note that even though there is no parametrization corresponding to (16), the Cayley projective plane can be described (see, for example, Kuiper [39], Freudenthal [28]) as the submanifold

$$V = \{ A \in M_{3 \times 3}(\mathbf{O}) \mid \overline{A}^T = A = A^2, \text{ trace } A = 1 \},\$$

where $M_{3\times 3}(\mathbf{O})$ is the space of 3×3 matrices of Cayley numbers. This submanifold V lies in a sphere S^{25} in a 26-dimensional real subspace of $M_{3\times 3}(\mathbf{O})$.

In his paper, Cartan [4, pp. 342–358] writes a separate section for each of the cases m = 1, 2, 4, 8, that is, $\mathbf{F} = \mathbf{R}, \mathbf{C}, \mathbf{H}, \mathbf{O}$. He gives extensive details in each case, describing many remarkable properties, especially in the case m = 8. In particular, he shows that in each case the isoparametric hypersurfaces and the two focal submanifolds are homogeneous, that is, they are orbits of points in S^{n+1} under the action of a closed subgroup of SO(n+2).

Cartan's results show that up to congruence, there is only one isoparametric family of hypersurfaces with g = 3 principal curvatures for each value of m. This classification is closely related to various characterizations of the standard embeddings of \mathbf{FP}^2 . (See Ewert [23], Little [41], and Knarr-Kramer [37].)

For alternative proofs of Cartan's classification of isoparametric hypersurfaces with g = 3 principal curvatures, see the papers of Knarr and Kramer [37], and Console and Olmos [21]. In a related paper, Sanchez [51] studied Cartan's isoparametric hypersurfaces from an algebraic point of view. (See also the paper of Giunta and Sanchez [29].)

2.10 Hypersurfaces With Four Principal Curvatures

In his fourth paper on the subject, Cartan [6] continues his study of isoparametric hypersurfaces with the property that all the principal curvatures have the same multiplicity m. In that paper, Cartan produces a family of isoparametric hypersurfaces with g = 4 principal curvatures of multiplicity m = 1in S^5 , and a family with g = 4 principal curvatures of multiplicity m = 2 in S^9 .

In both cases, Cartan describes the geometry and topology of the hypersurfaces and the two focal submanifolds in great detail, pointing out several notable properties that they have. (See also the papers of Nomizu [46]–[47] and the book [13, pp.155–159] for more detail on the example with m = 1.) Cartan also proves that the hypersurfaces and the focal submanifolds are homogeneous in both cases, m = 1 and m = 2.

It is worth noting that Cartan's example with four principal curvatures of multiplicity m = 1 is the lowest dimensional example of an isoparametric hypersurface of FKM-type constructed by Ferus, Karcher and Münzner [27] using representations of Clifford algebras. Furthermore, Cartan's example with four principal curvatures of multiplicity m = 2 is not of FKM-type. Its principal curvatures do not have the correct multiplicities for a hypersurface of FKM-type (see [27]).

Cartan says that he can show by a very long calculation (which he does not give) that for an isoparametric hypersurface with g = 4 principal curvatures having the same multiplicity m, the only possibilities are m = 1 and m = 2. This was shown to be true later by Grove and Halperin [30]. Furthermore, up to congruence, Cartan's examples given in [6] are the only possibilities for m = 1 and m = 2. This was shown to be true by Takagi [60] in the case m = 1, and by Ozeki and Takeuchi [48]–[49] in the case m = 2.

2.11 Cartan's Questions

Cartan's third paper [5] is basically a survey of the results that he obtained in the other three papers. As we have noted, all of the examples that Cartan found are homogeneous, each being an orbit of a point under an appropriate closed subgroup of SO(n + 2). Based on his results and the properties of his examples, Cartan asked the following three questions at the end of his survey paper [5]. All of his questions were answered in the 1970's, as we will describe below.

- 1. For each positive integer g, does there exist an isoparametric family with g distinct principal curvatures of the same multiplicity?
- 2. Does there exist an isoparametric family of hypersurfaces with more than three distinct principal curvatures such that the principal curvatures do not all have the same multiplicity?
- 3. Does every isoparametric family of hypersurfaces admit a transitive group of isometries?

In the early 1970's, Nomizu [46]–[47] wrote two papers describing the highlights of Cartan's work. He also generalized Cartan's example with four principal curvatures of multiplicity one to produce examples with four principal curvatures having multiplicities $m_1 = m_3 = m$, and $m_2 = m_4 = 1$, for any positive integer m. This answered Cartan's Question 2 in the affirmative.

In 1972, Takagi and Takahashi [61] gave a complete classification of all homogeneous isoparametric hypersurfaces in S^{n+1} based on the work of Hsiang and Lawson [33]. Takagi and Takahashi [61, p.480] showed that each homogeneous isoparametric hypersurface in S^{n+1} is a principal orbit of the isotropy representation of a Riemannian symmetric space of rank 2, and they gave a complete list of examples. This list contains examples with g = 6 principal curvatures as well as those with g = 1, 2, 3, 4 principal curvatures. In some cases with g = 4, the principal curvatures do not all have the same multiplicity, so this also provided an affirmative answer to Cartan's Question 2.

At about the same time as the papers of Nomizu and Takagi-Takahashi, Münzner [44]–[45] published two preprints that greatly extended Cartan's work and have served as the basis for much of the research in the field since that time. The preprints were eventually published as journal articles [44]– [45] in 1980–1981, and they are the basis of Section 3 of this paper below.

One of Münzner's primary results is that the number g of distinct principal curvatures of an isoparametric hypersurface in a sphere must be equal to 1, 2, 3, 4 or 6, and thus the answer to Cartan's Question 1 is negative.

Finally, the answer to Cartan's Question 3 is also negative, as was first shown by the construction of some families of inhomogeneous isoparametric hypersurfaces with g = 4 principal curvatures by Ozeki and Takeuchi [48] in 1975. Their construction was then generalized in 1981 to yield even more inhomogeneous examples by Ferus, Karcher and Münzner [27].

Remark 2.2 (Isoparametric submanifolds of codimension greater than one). There is also an extensive theory of isoparametric submanifolds of codimension greater than one in the sphere, due primarily to Carter and West [8]–[9], West [74], Terng [65]–[69], and Hsiang, Palais and Terng [34]. (See also Harle [32] and Strübing [59].)

Terng [65] formulated the definition as follows: a connected, complete submanifold V in a real space form \widetilde{M}^{n+1} is said to be *isoparametric* if it has flat normal bundle and if for any parallel section of the unit normal bundle $\eta: V \to B^n$, the principal curvatures of A_η are constant.

After considerable development of the theory, Thorbergsson [71] showed that a compact, irreducible isoparametric submanifold M substantially embedded in S^{n+1} with codimension greater than one is homogeneous. Thus, M is a principal orbit of an isotropy representation of a symmetric space (also called *s*-representations), as in the codimension one case. Orbits of isotropy representations of symmetric spaces are also known as *generalized flag manifolds* or *R*-spaces. (See Bott-Samelson [2] and Takeuchi-Kobayashi [62]).

3 Münzner's Work

In this section, we cover the work of Münzner in his two papers [44]–[45]. We will give detailed proofs of several of the main results. Our treatment is based primarily on Münzner's first paper [44] and on the book of Cecil and Ryan [13, pp. 102–137]. Some passages in this section are taken directly from that book.

3.1 Principal Curvatures of Parallel Hypersurfaces

We now begin our treatment of Münzner's theory, which is contained in Subsections 3.1–3.4 of this paper. We start by deriving the well-known formulas for the principal curvatures of a parallel hypersurface to a given hypersurface. We include this for the sake of completeness, since these formulas are important in the development of the theory.

For the following local calculations, we consider a connected, oriented isoparametric hypersurface $M \subset S^{n+1} \subset \mathbf{R}^{n+2}$ with field of unit normals ξ . Assume that M has g distinct principal constant curvatures at each point, which we label by,

$$\lambda_i = \cot \theta_i, \ 0 < \theta_i < \pi, \ 1 \le i \le g, \tag{17}$$

where the θ_i form an increasing sequence, and λ_i has multiplicity m_i on M.

We denote the corresponding principal distribution by,

$$T_i(x) = \{ X \in T_x M \mid AX = \lambda_i X \}, \tag{18}$$

where A is the shape operator determined by the field of unit normals ξ .

Using the Codazzi equations in the case of multiplicity $m_i > 1$, and by the theory of ordinary equations in the case $m_i = 1$, one can show that each T_i is integrable, i.e., it is a foliation of M with leaves of dimension m_i . Furthermore, the leaves of the principal foliation T_i corresponding to λ_i are open subsets of m_i -dimensional metric spheres in S^{n+1} , and the space of leaves M/T_i is an $(n - m_i)$ -dimensional manifold with the quotient topology (see [13, pp. 18–32] for proofs).

We consider the parallel hypersurface $f_t: M \to S^{n+1}$ defined by

$$f_t(x) = \cos t \ x + \sin t \ \xi(x), \tag{19}$$

that is, $f_t(x)$ is the point in S^{n+1} at an oriented distance t along the normal geodesic in S^{n+1} to M through the point x. Note that f_0 is the original embedding f defined by f(x) = x, and we suppress the mention of f in equation (19).

In the following calculations, we show that f_t is an immersion at x if and only if $\cot t$ is not a principal curvature of M at x. In that case, we then find the principal curvatures of f_t in terms of the principal curvatures of the original embedding f.

Let $X \in T_x M$. Then differentiating equation (19) in the direction X, we get

$$(f_t)_*X = \cos t \ X + \sin t \ D_X \xi = \cos t \ X - \sin t \ AX = (\cos t \ I - \sin t \ A)X,$$

where on the right side we are identifying X with its Euclidean parallel translate at $f_t(x)$. If $X \in T_i(x)$, this yields,

$$(f_t)_* X = (\cos t - \sin t \cot \theta_i) \ X = \frac{\sin(\theta_i - t)}{\sin \theta_i} X.$$
(20)

Since $T_x M$ is the direct sum of the principal spaces $T_i(x)$, we see that $(f_t)_*$ is injective on $T_x M$, unless $t = \theta_i \pmod{\pi}$ for some *i*, that is, unless $f_t(x)$ is a focal point of (M, x).

Thus, we see that a parallel hypersurface $f_t M$ is an immersed hypersurface if $t \neq \theta_i \pmod{\pi}$ for any *i*. In that case, we want to find the principal curvatures of $f_t M$.

Theorem 3.1. Let $M \subset S^{n+1}$ be an oriented isoparametric hypersurface and let $\lambda = \cot \theta$ be a principal curvature of multiplicity m on M. Suppose that f_t is an immersion of M, and let x be a point in M. Then the parallel hypersurface $f_t M$ has a principal curvature $\tilde{\lambda} = \cot(\theta - t)$ at $f_t(x)$ having the same multiplicity m and (up to parallel translation in \mathbf{R}^{n+2}) the same principal space $T_{\lambda}(x)$ as λ at x.

Proof. Let ξ be the field of unit normals on M, and denote the value ξ at a point $y \in M$ by ξ_y . Then we can easily compute that the vector

$$\tilde{\xi}_y = -\sin t \ y + \cos t \ \xi_y,\tag{21}$$

when translated to $\tilde{y} = f_t(y)$, is a unit normal to the hypersurface $f_t M$ at the point \tilde{y} . We want to find the shape operator A_t determined by this field of unit normals $\tilde{\xi}$ on $f_t M$. Let $X \in T_{\lambda}(x)$. Since $\langle X, \xi \rangle = 0$, we have

$$D_X \xi = \widetilde{\nabla}_X \xi - \langle X, \xi \rangle \ x = \widetilde{\nabla}_X \xi = -AX = -\lambda X = -\cot\theta \ X, \tag{22}$$

where D is the Euclidean covariant derivative on \mathbb{R}^{n+2} , and $\widetilde{\nabla}$ is the induced Levi-Civita connection on S^{n+1} . By definition, the shape operator A_t is given by

$$(f_t)_*(A_tX) = -\widetilde{\nabla}_{(f_t)_*X}\widetilde{\xi} = -D_{(f_t)_*X}\widetilde{\xi},$$
(23)

since $\langle (f_t)_*X, \tilde{\xi} \rangle = 0$. To compute this, let x_u be a curve in M with initial point $x_0 = x$ and initial tangent vector $\overrightarrow{x_0} = X$. Then we have using equation (22),

$$D_{(f_t)_*X}\tilde{\xi} = \frac{d}{du}(\tilde{\xi}_{x_u})|_{u=0} = -\sin t \ X - \cos t \cot \theta \ X$$
(24)
$$= \frac{-\cos(\theta - t)}{\sin \theta} X,$$

where we again identify X on the right with its Euclidean parallel translate at $f_t(x)$. If we compare this with equation (20) and use equation (23), we see that

$$A_t X = \cot(\theta - t)X,\tag{25}$$

for $X \in T_{\lambda}(x)$, and so $\tilde{\lambda} = \cot(\theta - t)$ is a principal curvature of A_t with the same principal space $T_{\lambda}(x)$ and same multiplicity m as λ at x.

As a consequence of Theorem 3.1, we get the following corollary regarding the principal curvatures of a family of parallel isoparametric hypersurfaces.

Corollary 3.1. Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface having g distinct principal curvatures $\lambda_i = \cot \theta_i$, $1 \leq i \leq g$, with respective multiplicities m_i . If t is any real number not congruent to any $\theta_i \pmod{\pi}$, then the map f_t immerses M as an isoparametric hypersurface with principal curvatures $\tilde{\lambda}_i = \cot(\theta_i - t)$, $1 \leq i \leq g$, with the same multiplicities m_i . Furthermore, for each i, the principal foliation corresponding to $\tilde{\lambda}_i$ is the same as the principal foliation T_i corresponding to λ_i on M.

Remark 3.1. It follows from Münzner's theory that if M is an isoparametric hypersurface embedded in S^{n+1} , then each parallel isoparametric hypersurface $f_t M$ is also embedded in S^{n+1} and not just immersed. This is because M and its parallel hypersurfaces are level sets of the restriction to S^{n+1} of a certain polynomial function on \mathbf{R}^{n+2} , as will be discussed later in Theorem 3.5.

3.2 Focal Submanifolds

The geometry of the focal submanifolds is a crucial element in the theory of isoparametric hypersurfaces. In this subsection, we obtain some important basic results about isoparametric hypersurfaces and their focal submanifolds due to Münzner [44].

We use the notation introduced in Subsection 3.1. So we let $M^n \subset S^{n+1}$ be a connected, oriented isoparametric hypersurface with field of unit normals ξ having g distinct constant principal curvatures,

$$\lambda_i = \cot \theta_i, \ 0 < \theta_i < \pi, \ 1 \le i \le g, \tag{26}$$

where the θ_i form an increasing sequence. Denote the multiplicity of λ_i by m_i .

As noted earlier, the leaves of the principal foliation T_i corresponding to λ_i are open subsets of m_i -dimensional metric spheres in S^{n+1} , and the space of leaves M/T_i is an $(n - m_i)$ -dimensional manifold with the quotient topology (see [13, pp. 18–32] for proofs).

We consider the map f_t defined as in equation (19) by

$$f_t(x) = \cos t \ x + \sin t \ \xi(x),$$

and we use the formulas developed in Subsection 3.1 for making computations.

If $t = \theta_i$, then the map f_t has constant rank $n - m_i$ on M, and it factors through an immersion $\psi_i : M/T_i \to S^{n+1}$, defined on the space of leaves M/T_i , i.e., $f_t = \psi_i \circ \pi$, where π is the projection from M onto M/T_i . Thus, f_t is a focal map, and we denote image of ψ_i by V_i .

We now want to find the principal curvatures of this focal submanifold V_i . This is similar to the calculation for parallel hypersurfaces in Subsection 3.1, although we must make some adjustments because V_i has codimension greater than one.

Let $x \in M$. Then we have the orthogonal decomposition of the tangent space $T_x M$ as

$$T_x M = T_i(x) \oplus T_i^{\perp}(x), \tag{27}$$

where $T_i^{\perp}(x)$ is the direct sum of the spaces $T_j(x)$, for $j \neq i$. By equation (20) in Subsection 3.1, the map $(f_t)_* = 0$ on $T_i(x)$, and $(f_t)_*$ is injective on $T_i^{\perp}(x)$, for $t = \theta_i$. We consider a map $h : M \to S^{n+1}$ given by

$$h(x) = -\sin t \ x + \cos t \ \xi(x). \tag{28}$$

This is basically the same function that we considered in equation (21) for the field $\tilde{\xi}$ of unit normals to $f_t M$, in the case where f_t is an immersion.

In the case $t = \theta_i$, we see that the inner product $\langle f_t(x), h(x) \rangle = 0$, and so the vector h(x) is tangent to the sphere S^{n+1} at the point $p = f_t(x)$. Furthermore, $\langle h(x), X \rangle = 0$ for all $X \in T_i^{\perp}(x)$, and so by equation (20), the vector h(x) is normal to the focal submanifold V_i at the point p.

We can use the map h to find the shape operator determined by a normal vector to V_i as follows. Let p be an arbitrary point in V_i . Then the set $C = f_t^{-1}(p)$ is an open subset of an m_i -sphere in S^{n+1} . For each $x \in C$, the vector h(x) is a unit normal to the focal submanifold V_i at p. Thus, the restriction of h to C is a map from C into the m_i -sphere $S_p^{\perp}V_i$ of unit normal vectors to V_i at p. At a point $x \in C$, the tangent space T_xC equals $T_i(x)$. Since $t = \theta_i$, we compute for $x \in C$ and a nonzero vector $X \in T_i(x)$ that

$$h_*(X) = -\sin tX + \cos t(-AX) = -\sin \theta_i X + \cos \theta_i (-\cot \theta_i) X$$
(29)
$$= \frac{-1}{\sin \theta_i} X \neq 0.$$

Thus, h_* has full rank m_i on C, and so h is a local diffeomorphism of open subsets of m_i -spheres. This enables us to prove the following important result due to Münzner [44].

Theorem 3.2. Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface, and let $V_i = f_t M$ for $t = \theta_i$ be a focal submanifold of M. Let η be a unit normal vector to V_i at a point $p \in V_i$, and suppose that $\eta = h(x)$ for some $x \in f_t^{-1}(p)$. Then the shape operator A_η of V_i is given in terms of its principal vectors by

$$A_{\eta}X = \cot(\theta_j - \theta_i)X, \text{ for } X \in T_j(x), \ j \neq i.$$
(30)

(As before we are identifying $T_j(x)$ with its Euclidean parallel translate at p.)

Proof. Let $\eta = h(x)$ for some $x \in C = f_t^{-1}(p)$ for $t = \theta_i$. The same calculation used in proving Theorem 3.1 is valid here, and it leads to equation (25), which we write as in equation (30),

$$A_{\eta}X = \cot(\theta_j - \theta_i)X$$
, for $X \in T_j(x), \ j \neq i$.

Corollary 3.2. Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface, and let $V_i = f_t M$, for $t = \theta_i$, be a focal submanifold of M. Then for every unit normal vector η at every point $p \in V_i$, the shape operator A_{η} has principal curvatures $\cot(\theta_j - \theta_i)$ with multiplicities m_j , for $j \neq i$, $1 \leq j \leq g$. Proof. By Theorem 3.2, the corollary holds on the open subset h(C) of the m_i -sphere $S_p^{\perp}V_i$ of unit normal vectors to V_i at p. Consider the characteristic polynomial $P_u(\eta) = \det(A_\eta - uI)$ as a function of η on the normal space $T_p^{\perp}V_i$. Since A_η is linear in η , we have for each fixed $u \in \mathbf{R}$ that the function $P_u(\eta)$ is a polynomial of degree $n - m_i$ on the vector space $T_p^{\perp}V_i$. Thus, the restriction of $P_u(\eta)$ to the sphere $S_p^{\perp}V_i$ is an analytic function of η . Then since $P_u(\eta)$ is constant on the open subset h(C) of $S_p^{\perp}V_i$, it is constant on all of $S_p^{\perp}V_i$.

3.2.1 Minimality of the focal submanifolds

Münzner also obtained the following consequence of Corollary 3.2. This result was obtained independently with a different proof by Nomizu [46].

Corollary 3.3. Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface. Then each focal submanifold V_i of M is a minimal submanifold in S^{n+1} .

Proof. Let η be a unit normal vector to a focal submanifold V_i of M. Then $-\eta$ is also a unit normal vector to V_i . By Corollary 3.2, the shape operators A_η and $A_{-\eta}$ have the same eigenvalues with the same multiplicities. So

trace
$$A_{-n} = \text{trace } A_n$$
.

On the other hand, trace $A_{-\eta} = -$ trace A_{η} , since $A_{-\eta} = -A_{\eta}$. Thus, we have trace $A_{\eta} = -$ trace A_{η} , and so trace $A_{\eta} = 0$. Since this is true for all unit normal vectors η , we conclude that V_i is a minimal submanifold in S^{n+1} .

As a consequence of Theorem 3.2, we can give a proof of Cartan's formula (see equation (31) below) for isoparametric hypersurfaces in S^{n+1} . The proof shows that Cartan's formula is essentially equivalent to the minimality of the focal submanifolds.

Corollary 3.4. (Cartan's formula) Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface with g principal curvatures

 $\lambda_i = \cot \theta_i, \quad 0 < \theta_1 < \dots < \theta_g < \pi,$

with respective multiplicities m_i . Then for each $i, 1 \leq i \leq g$, Cartan's formula holds, that is,

$$\sum_{j \neq i} m_j \frac{1 + \lambda_i \lambda_j}{\lambda_i - \lambda_j} = 0.$$
(31)

Proof. We will show that for each i and for any unit normal η to the focal submanifold V_i , the left side of equation (31) equals trace A_{η} , which equals zero by Corollary 3.3. Represent the principal curvatures of M as $\lambda_i = \cot \theta_i$, $0 < \theta_1 < \cdots < \theta_g < \pi$, with respective multiplicities m_i . Let V_i be the focal submanifold corresponding to λ_i . By Corollary 3.2, we have

$$0 = \operatorname{trace} A_{\eta} = \sum_{j \neq i} m_j \cot(\theta_j - \theta_i) = \sum_{j \neq i} m_j \frac{1 + \cot \theta_i \cot \theta_j}{\cot \theta_i - \cot \theta_j} \quad (32)$$
$$= \sum_{j \neq i} m_j \frac{1 + \lambda_i \lambda_j}{\lambda_i - \lambda_j}.$$

3.2.2 Formula for the principal curvatures

Münzner [44, p. 61] showed that the principal curvatures of an isoparametric hypersurface in S^{n+1} have a very specific form, given in equation (33) in Theorem 3.3 below. Cartan [3, pp. 186–187] obtained this same formula in the case where the principal curvatures all have the same multiplicity (see Theorem 2.8), but he did not have the result in the general case.

Moreover, Münzner's Theorem 3.3 also contains the result that in the general case, the multiplicities of the principal curvatures must satisfy the relation $m_i = m_{i+2}$ (subscripts mod g).

Theorem 3.3 is a key element in Münzner's extension to the general case of Cartan's Theorem 2.9 concerning the algebraic nature of isoparametric hypersurfaces (see Theorem 3.5 below). Our treatment of the proof of Münzner's Theorem 3.3 is taken from [13, pp. 108–110].

Theorem 3.3 (Münzner). Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface with g principal curvatures $\lambda_i = \cot \theta_i, \ 0 < \theta_1 < \cdots < \theta_g < \pi$, with respective multiplicities m_i . Then

$$\theta_i = \theta_1 + (i-1)\frac{\pi}{g}, \quad 1 \le i \le g, \tag{33}$$

and the multiplicities satisfy $m_i = m_{i+2}$ (subscripts mod g). For any point $x \in M$, there are 2g focal points of (M, x) along the normal geodesic to M through x, and they are evenly distributed at intervals of length π/g .

Proof. If g = 1, then the theorem is trivially true, so we now consider g = 2. Let V_1 be the focal submanifold determined by the map f_t for $t = \theta_1$. By Corollary 3.2, the principal curvature $\cot(\theta_2 - \theta_1)$ of the shape operator A_η is the same for every choice of unit normal η at every point $p \in V_1$. Since $A_{-\eta} = -A_\eta$, this says that

$$\cot(\theta_2 - \theta_1) = -\cot(\theta_2 - \theta_1).$$

Thus, $\cot(\theta_2 - \theta_1) = 0$, so $\theta_2 - \theta_1 = \pi/2$ as desired. In the case g = 2, there is no restriction on the multiplicities.

Next we consider the case $g \geq 3$. For a fixed value of $i, 1 \leq i \leq g$, let V_i be the focal submanifold determined by the map f_t for $t = \theta_i$. By Corollary 3.2, the set

$$\{\cot(\theta_j - \theta_i) \mid j \neq i\}$$

of principal curvatures of the shape operator A_{η} is the same for every choice of unit normal η at every point $p \in V_i$. Since $A_{-\eta} = -A_{\eta}$, this says that the two sets

$$\{\cot(\theta_j - \theta_i) \mid j \neq i\}$$
 and $\{-\cot(\theta_j - \theta_i) \mid j \neq i\}$

are the same. In the case $2 \leq i \leq g-1$, the largest principal curvature of A_{η} is $\cot(\theta_{i+1}-\theta_i)$ with multiplicity m_{i+1} , while the largest principal curvature of $A_{-\eta}$ is $\cot(\theta_i - \theta_{i-1})$ with multiplicity m_{i-1} . Since these two largest principal curvatures and their respective multiplicities are equal, we conclude that

$$\theta_{i+1} - \theta_i = \theta_i - \theta_{i-1}, \quad m_{i+1} = m_{i-1}, \quad 2 \le i \le g - 1.$$
 (34)

If i = 1, the largest principal curvature of A_{η} is $\cot(\theta_2 - \theta_1)$ with multiplicity m_2 , and the largest principal curvature of $A_{-\eta}$ is

$$\cot(\theta_1 - \theta_g) = \cot(\theta_1 - (\theta_g - \pi)),$$

with multiplicity m_g , and we have

$$\theta_2 - \theta_1 = \theta_1 - (\theta_g - \pi), \quad m_2 = m_g. \tag{35}$$

If we let $\theta_2 - \theta_1 = \delta$, then equation (34) implies that $\theta_g - \theta_1 = (g-1)\delta$, while equation (35) implies that $\theta_g - \theta_1 = \pi - \delta$. Combining these two equations, we get that $g\delta = \pi$, and thus $\delta = \pi/g$. From this we get the formula in equation (33) for θ_i . The formula for the multiplicities in the theorem follows from equations (34) and (35). If x is any point of M, then each principal curvature $\cot \theta_i$ of M gives rise to a pair of antipodal focal points along the normal geodesic to M through x. Thus, there are 2g focal points of (M, x) along this normal geodesic, and they are evenly distributed at intervals of length π/g by equation (33).

Remark 3.2 (Isoparametric submanifolds and their Coxeter groups). It follows from Theorem 3.3 that the set of focal points along a normal circle to $M \subset S^{n+1}$ is invariant under the dihedral group D_g of order 2g that acts on the normal circle and is generated by reflections in the focal points. This is a fundamental idea that generalizes to isoparametric submanifolds of higher codimension in the sphere (see Remark 2.2). Specifically, for an isoparametric submanifold M^n of codimension k > 1 in S^{n+1} , Carter and West [8] (in the case k = 2) and Terng [65] for arbitrary k > 1 found a Coxeter group (finite group generated by reflections) that acts in a way similar to this dihedral group in the codimension one case. This Coxeter group is important in the overall development of the theory in the case of isoparametric submanifolds of higher codimension (see Terng [65] and Thorbergsson [72]).

Since the multiplicities in Theorem 3.3 satisfy $m_i = m_{i+2}$ (subscripts mod g), we have the following immediate corollary.

Corollary 3.5. Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface with g distinct principal curvatures. If g is odd, then all of the principal curvatures have the same multiplicity. If g is even, then there are at most two distinct multiplicities.

3.3 Cartan-Münzner Polynomials

Our goal in this subsection is to outline the proof of Münzner's generalization of Cartan's Theorem 2.9 concerning the algebraic nature of isoparametric hypersurfaces in S^{n+1} . Here we follow the treatment in the book [13, pp. 111–130] closely.

As noted at the beginning of Section 2, the original definition of an isoparametric family of hypersurfaces in a real space form \widetilde{M}^{n+1} was formulated in terms of the level sets of an isoparametric function, where a smooth function $V: \widetilde{M}^{n+1} \to \mathbf{R}$ is called isoparametric if both of the classical Beltrami differential parameters,

$$\Delta_1 V = |\text{grad } V|^2, \quad \Delta_2 V = \Delta V \text{ (Laplacian } V), \tag{36}$$

are functions of V itself, and are therefore constant on the level sets of V in \widetilde{M}^{n+1} .

In the case of an isoparametric hypersurface M in the unit sphere S^{n+1} in \mathbf{R}^{n+2} with g distinct principal curvatures, Münzner [44] showed that the corresponding isoparametric function $V: S^{n+1} \to \mathbf{R}$ is the restriction to S^{n+1} of a homogeneous polynomial $F: \mathbf{R}^{n+2} \to \mathbf{R}$ of degree g satisfying certain differential equations. Thus, it is useful to relate the Beltrami differential parameters of a homogeneous function F on \mathbf{R}^{n+2} to those of its restriction V to S^{n+1} .

Recall that a function $F : \mathbf{R}^{n+2} \to \mathbf{R}$ is homogeneous of degree g if $F(tx) = t^g F(x)$, for all $t \in \mathbf{R}$ and $x \in \mathbf{R}^{n+2}$. By Euler's Theorem, we know that for any $x \in \mathbf{R}^{n+2}$,

$$\langle \operatorname{grad}^E F, x \rangle = gF(x).$$
 (37)

Here the superscript E is used to denote the Euclidean gradient of F. The gradient of the restriction V of F to S^{n+1} will be denoted by $\operatorname{grad}^S F$. Similarly, we will denote the respective Laplacians by $\Delta^E F$ and $\Delta^S F$.

The following theorem relates the various differential operators for a homogeneous function F of degree g, and it is useful in proving Theorem 3.5. (See [13, pp. 112–113] for a proof.)

Theorem 3.4. Let $F : \mathbb{R}^{n+2} \to \mathbb{R}$ be a homogeneous function of degree g. Then

- (a) $|\text{grad}^{S}F|^{2} = |\text{grad}^{E}F|^{2} g^{2}F^{2}$,
- (b) $\Delta^S F = \Delta^E F g(g-1)F g(n+1)F$.

Münzner's generalization of Cartan's Theorem 2.9 is the following.

Theorem 3.5 (Münzner). Let $M \subset S^{n+1} \subset \mathbf{R}^{n+2}$ be a connected isoparametric hypersurface with g principal curvatures $\lambda_i = \cot \theta_i, 0 < \theta_1 < \cdots < \theta_g < \pi$, with respective multiplicities m_i . Then M is an open subset of a level set of the restriction to S^{n+1} of a homogeneous polynomial F on \mathbf{R}^{n+2} of degree g satisfying the differential equations,

$$|\text{grad}^E F|^2 = g^2 r^{2g-2},$$
 (38)

$$\Delta^E F = cr^{g-2},\tag{39}$$

where r = |x|, and $c = g^2(m_2 - m_1)/2$.

Remark 3.3. Recall from Corollary 3.5 that there are at most two distinct multiplicities m_1, m_2 , and the multiplicities satisfy $m_{i+2} = m_i$ (subscripts mod g).

Münzner [44, p. 65] called F the Cartan polynomial of M, and now F is usually referred to as the Cartan-Münzner polynomial of M. Equations (38)–(39) are called the Cartan-Münzner differential equations. By Theorem 3.4 the restriction V of F to S^{n+1} satisfies the differential equations,

$$|\text{grad}^S V|^2 = g^2 (1 - V^2),$$
 (40)

$$\Delta^S V = c - g(n+g)V, \tag{41}$$

where $c = g^2(m_2 - m_1)/2$. Thus V is an isoparametric function in the sense of Cartan, since both $|\text{grad}^S V|^2$ and $\Delta^S V$ are functions of V itself. (Here we are using the superscript S in the notations $\text{grad}^S V$ and $\Delta^S V$ to emphasize that V is a function defined on S^{n+1} .)

We now describe Münzner's [44, pp. 62–65] construction of this polynomial F in detail. This is basically the same approach used by Cartan [4] in proving Theorem 2.9 under the more restrictive assumption that all of the principal curvatures have the same multiplicity.

Note that Cartan [4, pp. 364–365] proved that if the principal curvatures all have the same multiplicity m, then the polynomial F must be harmonic. This agrees with Münzner's condition (39) in the case where all of the multiplicities are equal, since then the number $c = g^2(m_2 - m_1)/2$ in equation (39) equals zero, and so F is harmonic.

Let $M \subset S^{n+1}$ be a connected, oriented isoparametric hypersurface with g distinct principal curvatures $\lambda_i = \cot \theta_i$, $0 < \theta_1 < \cdots < \theta_g < \pi$, with respective multiplicities m_i . The normal bundle NM of M in S^{n+1} is trivial and is therefore diffeomorphic to $M \times \mathbf{R}$. Thus we can consider the normal exponential map $E: M \times \mathbf{R} \to S^{n+1}$ defined by

$$E(x,t) = f_t(x) = \cos t \ x + \sin t \ \xi(x),$$
 (42)

where ξ is the field of unit normals to M in S^{n+1} .

By the standard calculation for the location of focal points (see, for example, [13, pp. 11–14]), we know that the differential of E has rank n + 1at $(x,t) \in M \times \mathbf{R}$, unless $\cot t$ is a principal curvature of M at x. Thus, for any noncritical point (x,t) of E, there is an open neighborhood U of (x,t) in $M \times \mathbf{R}$ on which E restricts to a diffeomorphism onto an open subset $\widetilde{U} = E(U)$ in S^{n+1} . We define a function $\tau : \widetilde{U} \to \mathbf{R}$ by

$$\tau(p) = \theta_1 - \pi_2(E^{-1}(p)), \tag{43}$$

where π_2 is projection onto the second coordinate. That is, if p = E(x, t), then

$$\tau(p) = \theta_1 - t. \tag{44}$$

Then we define a function $V: \widetilde{U} \to \mathbf{R}$ by,

$$V(p) = \cos(g\tau(p)). \tag{45}$$

Clearly, τ and V are constant on each parallel hypersurface $M_t = f_t(M)$ in \widetilde{U} .

The number $\tau(p)$ is the oriented distance from p to the first focal point along the normal geodesic to the parallel hypersurface M_t through p. Thus, if we begin the construction with a parallel hypersurface near M rather than with M itself, we get the same functions τ and V. If we begin the construction with the opposite field of unit normals $-\xi$ instead of ξ , then we obtain the function -V instead of V.

We next extend V to a homogeneous function of degree g on the cone in \mathbf{R}^{n+2} over \widetilde{U} by the formula

$$F(rp) = r^g \cos(g(\tau(p))), \ p \in \widetilde{U}, \ r > 0.$$

$$(46)$$

The first step in the proof of Theorem 3.5 is to show that the function F in equation (46) satisfies the Cartan-Münzner differential equations (38)–(39). One then completes the proof of Theorem 3.5 by showing that F is the restriction to the cone over \tilde{U} of a homogeneous polynomial of degree g.

These two steps involve lengthy calculations based on the formula for the principal curvatures of an isoparametric hypersurface given in Theorem 3.3. We refer the reader to [44, pp. 62–67] or [13, pp. 115–125] for the details of the proof, which we will omit here.

We now want to make a few remarks concerning some important consequences of Theorem 3.5.

Remark 3.4 (Consequences of Theorem 3.5). From equation (40), we see that the range of the restriction V of F to S^{n+1} is contained in the closed interval [-1, 1], since the left side of the equation is nonnegative. We can

see that the range of V is all of the interval [-1, 1] as follows. Since V is not constant on S^{n+1} , it has distinct maximum and minimum values on S^{n+1} . By equation (40) these maximum and minimum values are 1 and -1, respectively, since grad^SV is nonzero at any point where V is not equal to ± 1 . For any s in the open interval (-1, 1), the level set $V^{-1}(s)$ is a compact hypersurface, since grad^SV is never zero on $V^{-1}(s)$. Münzner also proves that each level set of V is connected, and therefore, the original connected isoparametric hypersurface M is contained in a unique compact, connected isoparametric hypersurface.

For $s = \pm 1$, grad^SV is identically equal to zero on $V^{-1}(s)$, and the sets $M_+ = M_1 = V^{-1}(1)$ and $M_- = M_{-1} = V^{-1}(-1)$ are submanifolds of codimension greater than one in S^{n+1} . Münzner showed that M_+ and $M_$ are connected (see Theorem 3.6), and that they are the focal submanifolds of any isoparametric hypersurface $V^{-1}(s)$, -1 < s < 1, in the family of isoparametric hypersurfaces. Thus, there are only two focal submanifolds regardless of the number g of distinct principal curvatures. By Theorem 3.3, there are 2g focal points evenly distributed along each normal geodesic to the family $\{V^{-1}(s)\}$ of isoparametric hypersurfaces. Münzner proves in part (a) of Theorem 3.7 below that these focal points lie alternately on the two focal submanifolds M_+ and M_- .

Suppose we consider the isoparametric hypersurface $V^{-1}(0)$. From equation (45), we see that the function τ equals $\pi/2g$ on $V^{-1}(0)$. The function τ is the distance from a point x in $V^{-1}(0)$ to the first focal point along the normal geodesic through x. By Theorem 3.3, this means that the largest principal curvature of $V^{-1}(0)$ is $\cot(\pi/2g)$, and the principal curvatures of $V^{-1}(0)$ are given by $\cot \theta_i$, where

$$\theta_i = \frac{\pi}{2g} + \frac{(i-1)}{g}\pi, \quad 1 \le i \le g, \tag{47}$$

with multiplicities satisfying $m_{i+2} = m_i$ (subscripts mod g). This simple form form for θ_i is helpful in the calculations involved in the proof of Theorem 3.5.

3.4 Global Structure Theorems

In this section, we discuss several results concerning the global structure of an isoparametric family of hypersurfaces in S^{n+1} that stem from Münzner's construction of the Cartan-Münzner polynomials, as discussed in the previous subsection (see [44, pp. 67–69]).

These structure theorems ultimately lead to Münzner's main structure theorem which states that a compact, connected isoparametric hypersurface M divides the sphere S^{n+1} into two ball bundles over the two focal submanifolds M_+ and M_- , which lie on different sides of M in S^{n+1} .

From this theorem, Münzner ultimately uses methods from algebraic topology to derive his primary result that the number g of distinct principal curvatures of an isoparametric hypersurface in S^{n+1} must be 1, 2, 3, 4 or 6.

Let $F : \mathbf{R}^{n+2} \to \mathbf{R}$ be the Cartan-Münzner polynomial of degree g constructed from a connected isoparametric hypersurface in $S^{n+1} \subset \mathbf{R}^{n+2}$ with g distinct principal curvatures as in Theorem 3.5, and let V denote the restriction of F to S^{n+1} . As noted in Remark 3.4, the range of the function Vis the closed interval [-1, 1], and each level set $M_t = V^{-1}(t), -1 < t < 1$, is an isoparametric hypersurface in S^{n+1} .

For the sake of definiteness, we let $M = M_0 = V^{-1}(0)$ be the isoparametric hypersurface discussed in Remark 3.4. We denote the two (possibly equal) multiplicities of M by $m_+ = m_1$ and $m_- = m_{-1}$.

The first goal is to show that M and all of the parallel hypersurfaces M_t , -1 < t < 1, as well as the two focal submanifolds, are connected. We omit the proof here and refer the reader to Münzner's paper [44] or to [13, pp. 130–137].

Theorem 3.6 (Connectedness of the level sets of F). Let $F : \mathbb{R}^{n+2} \to \mathbb{R}$ be a Cartan-Münzner polynomial of degree g and V its restriction to S^{n+1} . Then each isoparametric hypersurface

$$M_t = V^{-1}(t), \ -1 < t < 1,$$

is connected. Moreover, $M_+ = V^{-1}(1)$ and $M_- = V^{-1}(-1)$ are the focal submanifolds, and they are also connected.

Remark 3.5. A consequence of Theorem 3.6 is that any connected isoparametric hypersurface M lies in a unique compact, connected isoparametric hypersurface of the form $V^{-1}(t)$, -1 < t < 1, where V is the restriction to S^{n+1} of the Cartan-Münzner polynomial of M. For the rest of Section 3, we will assume that each isoparametric hypersurface and each focal submanifold is compact and connected.

The next step is Münzner's important structure theorem which states that a compact, connected isoparametric hypersurface M divides the sphere S^{n+1} into two ball bundles over the two focal submanifolds M_+ and M_- , which lie on different sides of M in S^{n+1} . The precise wording of the theorem is as follows. (See Münzner [44, pp. 67–69] or [13, pp. 132–133] for a proof.)

Theorem 3.7. Let $k = \pm 1$, and let Z be a normal vector to the focal submanifold M_k in S^{n+1} . Let $\exp : NM_k \to S^{n+1}$ denote the normal exponential map for M_k . Then

- (a) $V(\exp Z) = k \cos(g|Z|).$
- (b) Let $B_k = \{q \in S^{n+1} \mid kV(q) \ge 0\}$, and let $(B^{\perp}M_k, S^{\perp}M_k)$ be the bounded unit ball bundle in NM_k . Then

$$\psi_k : (B^\perp M_k, S^\perp M_k) \to (B_k, M),$$

where $M = V^{-1}(0)$ and $\psi_k(Z) = \exp(\frac{\pi}{2g}Z)$, is a diffeomorphism of manifolds with boundary.

As a consequence of the fact that the set of normal geodesics to each focal submanifold M_k , $k = \pm 1$, is the same as the set of normal geodesics to each of the parallel isoparametric hypersurfaces, we immediately obtain the following corollary.

Corollary 3.6. Let M_k , $k = \pm 1$, be a focal submanifold of an isoparametric hypersurface M. Then the focal set of M_k is the same as the focal set of M, *i.e.*, it is $M_k \cup M_{-k}$.

3.4.1 Münzner's restriction on the number of principal curvatures

Münzner's major result is that the number g of distinct principal curvatures of an isoparametric hypersurface M in S^{n+1} is 1, 2, 3, 4 or 6. This is a lengthy and delicate computation involving the cohomology rings of the hypersurface M and its two focal submanifolds M_+ and M_- . The structure of these rings is determined by the basic topological fact that a compact, connected isoparametric hypersurface $M \subset S^{n+1}$ divides S^{n+1} into two ball bundles over the two focal submanifolds, as in Theorem 3.7 (b).

Theorem 3.8 below does not assume that M is isoparametric, but only that it divides the sphere into two ball bundles. This is important, since the theorem can be applied to more general settings, in particular, to the case of a compact, connected proper Dupin hypersurface embedded in S^{n+1} , as was shown by Thorbergsson [70] (see also [13, pp. 140–143]). **Remark 3.6** (Proper Dupin hypersurfaces). Recall that an oriented hypersurface hypersurface M in a real space form \widetilde{M}^{n+1} is called a *proper Dupin* hypersurface if the number g of distinct principal curvatures is constant on M, and if each continuous principal curvature function on M is constant along each leaf of its corresponding principal foliation.

Using methods of algebraic topology, Münzner [45] proved the theorem below, and we refer the reader to Münzner's paper for the proof.

Theorem 3.8. Let M be a compact, connected hypersurface in S^{n+1} which divides S^{n+1} into two ball bundles over submanifolds M_+ and M_- . Then the number $\alpha = (1/2) \dim H^*(M, R)$ can only assume the values 1, 2, 3, 4 and 6. (The ring R of coefficients is \mathbf{Z} if both M_+ and M_- are orientable, and \mathbf{Z}_2 otherwise.)

Münzner then proved Theorem 3.9 below regarding the cohomology of an isoparametric hypersurface and its focal submanifolds. Since all of the parallel hypersurfaces $M_t = V^{-1}(t)$ are diffeomorphic, it is sufficient to consider the case $M = V^{-1}(0)$. In that case, M has two focal submanifolds $M_1 = V^{-1}(1)$ of dimension $n - m_1$ and $M_{-1} = V^{-1}(-1)$ of dimension $n - m_{-1}$, where m_1 and m_{-1} are the two (possibly equal) multiplicities of the principal curvatures of M. Then by Theorem 3.7, the sets,

$$B_1 = \{ q \in S^{n+1} \mid V(q) \ge 0 \}, \quad B_{-1} = \{ q \in S^{n+1} \mid V(q) \le 0 \},$$

are $(m_k + 1)$ -ball bundles over the focal submanifolds M_k , for k = 1, -1, respectively.

The dimension n of M is equal to $g(m_1 + m_{-1})/2$, and so $g = 2n/\mu$, where $\mu = m_1 + m_{-1}$. Thus, an isoparametric hypersurface M satisfies the hypothesis of the following theorem of Münzner [45]. The proof is done using algebraic topology, and we will omit it here. We refer the reader to Münzner's original proof [45] (see also [13, pp. 134–136]).

Theorem 3.9. Let M be a compact, connected hypersurface in S^{n+1} such that:

- (a) S^{n+1} is divided into two manifolds (B_1, M) and (B_{-1}, M) with boundary along M.
- (b) For k = ±1, the manifold B_k has the structure of a differentiable ball bundle over a compact, connected manifold M_k of dimension n - m_k.

Let the ring R of coefficients be Z if both M_1 and M_{-1} are orientable, and Z_2 otherwise. Let $\mu = m_1 + m_{-1}$. Then $\alpha = 2n/\mu$ is an integer, and for $k = \pm 1$,

$$H^{q}(M_{k}) = \begin{cases} R & \text{for } q \equiv 0 \pmod{\mu}, \ 0 \leq q < n, \\ R & \text{for } q \equiv m_{-k} \pmod{\mu}, \ 0 \leq q < n, \\ 0 & \text{otherwise.} \end{cases}$$

Further,

$$H^{q}(M) = \begin{cases} R \text{ for } q = 0, n, \\ H^{q}(M_{1}) \oplus H^{q}(M_{-1}), \text{ for } 1 \le q \le n - 1. \end{cases}$$

For a compact, connected isoparametric hypersurface $M \subset S^{n+1}$ with g distinct principal curvatures, we have

dim
$$M = n = g(m_1 + m_{-1})/2 = g\mu/2.$$

Thus, $\alpha = 2n/\mu = g$. By Theorem 3.9, we see that α is also equal to $\dim_R H^*(M, R)/2$. Hence by Münzner's Theorem 3.8, the number $g = \alpha$ can only assume the values 1, 2, 3, 4 or 6, and we have Münzner's [45] major theorem.

Theorem 3.10. Let $M \subset S^{n+1}$ be a connected isoparametric hypersurface with g distinct principal curvatures. Then g is 1, 2, 3, 4 or 6.

Note that we do not have to assume that M is compact in the theorem, because any connected isoparametric hypersurface is contained in a unique compact, connected isoparametric hypersurface to which the arguments above can be applied (see Remark 3.5). We also note that there exist isoparametric hypersurfaces for each of the values of g in the theorem, as mentioned in Subsection 2.11.

Remark 3.7 (Crystallographic groups). A consequence of Münzner's Theorem 3.10 is that the dihedral group D_g associated to M (see Remark 3.2) is crystallographic (see L.C. Grove and C.T. Benson [31, pp. 21–22]). A direct proof that D_g must be crystallographic could possibly give a simpler proof of Theorem 3.10 (see also K. Grove and S. Halperin [30, pp. 437–438]).

3.4.2 Multiplicities of the principal curvatures

As discussed in Subsection 2.9, Cartan [4] classified isoparametric hypersurfaces with $g \leq 3$ principal curvatures. In the cases g = 4 and g = 6, many results concerning the possible multiplicities of the principal curvatures have been obtained from the topological situation described in Theorem 3.9, i.e., that a compact, connected isoparametric hypersurface M in S^{n+1} divides S^{n+1} into two ball bundles over its two focal submanifolds.

In the case of g = 4 principal curvatures, several mathematicians, including Münzner [44]–[45], Abresch [1], Grove and Halperin [30], Tang [63]–[64] and Fang [24]–[25], found restrictions on the multiplicities (m_1, m_2) . This series of results culminated with the paper of Stolz [58], who used techniques from homotopy theory to prove Theorem 3.11 below. This theorem of Stolz was ultimately an important part in the proof of the classification of isoparametric hypersurfaces with g = 4 principal curvatures, which will be discussed in Section 4.

Theorem 3.11 (Stolz). The multiplicities (m_1, m_2) of the principal curvatures of an isoparametric hypersurface $M \subset S^{n+1}$ with g = 4 principal curvatures are the same as those in the examples due to Ferus, Karcher and Münzner [27] or the two homogeneous examples with $(m_1, m_2) = (2, 2)$ or $(m_1, m_2) = (4, 5)$ that are not of FKM-type.

Remark 3.8 (Application to proper Dupin hypersurfaces). Stolz proved Theorem 3.11 under more general assumption that M is a compact, connected proper Dupin hypersurface with four principal curvatures embedded in S^{n+1} . Such a result is possible because Thorbergsson [70] had shown earlier that a compact, connected proper Dupin hypersurface $M \subset S^{n+1}$ separates S^{n+1} into two ball bundles over the first focal submanifolds on either side of M, as in the case of a compact, connected isoparametric hypersurface (see also [13, p. 143]).

In the case of g = 6 principal curvatures, Münzner [45] showed that all of the principal curvatures have the same multiplicity m, and Abresch [1] showed that m equals 1 or 2. Thus we have:

Theorem 3.12. For an isoparametric hypersurface with g = 6 principal curvatures, all the principal curvatures have the same multiplicity m, and m equals 1 or 2.

Later Grove and Halperin [30] showed that this restriction on the multiplicities also holds for compact, connected proper Dupin hypersurfaces with g = 6 principal curvatures embedded in S^{n+1} .

4 Classification Results

Due to the work of many mathematicians, isoparametric hypersurfaces in S^{n+1} have been completely classified, and we will briefly state the classification results now. As noted above, Münzner proved that the number g of distinct principal curvatures must be 1, 2, 3, 4 or 6. We now summarize the classification results for each value of g. Many of these have been discussed already.

If g = 1, then the isoparametric hypersurface M is totally umbilic, and it must be an open subset of a great or small sphere. If g = 2, Cartan [3] showed that M must be an open subset of a standard product of two spheres,

$$S^{k}(r) \times S^{n-k-1}(s) \subset S^{n}, \quad r^{2} + s^{2} = 1.$$

In the case g = 3, Cartan [4] showed that all the principal curvatures must have the same multiplicity m = 1, 2, 4 or 8, and the isoparametric hypersurface must be an open subset of a tube of constant radius over a standard embedding of a projective plane \mathbf{FP}^2 into S^{3m+1} , where \mathbf{F} is the division algebra \mathbf{R} , \mathbf{C} , \mathbf{H} (quaternions), \mathbf{O} (Cayley numbers), for m = 1, 2, 4, 8, respectively. Thus, up to congruence, there is only one such family for each value of m. (See Subsection 2.9.)

In the case of an isoparametric hypersurface with g = 4 four principal curvatures, Münzner (Theorem 3.3) proved that the principal curvatures can have at most two distinct multiplicities m_1, m_2 . Next Ferus, Karcher and Münzner [27] used representations of Clifford algebras to construct for any positive integer m_1 an infinite series of isoparametric hypersurfaces with four principal curvatures having respective multiplicities (m_1, m_2) , where m_2 is nondecreasing and unbounded in each series. These examples are now known as isoparametric hypersurfaces of *FKM-type*, and they are also described in [13, pp. 162–180]. This construction of Ferus, Karcher and Münzner was a generalization of an earlier construction due to Ozeki and Takeuchi [48].

Stolz [58] (see Theorem 3.11) next proved that the multiplicities (m_1, m_2) of the principal curvatures of an isoparametric hypersurface with g = 4 principal curvatures must be the same as those of the hypersurfaces of FKM-type,

or else (2, 2) or (4, 5), which are the multiplicities for certain homogeneous examples that are not of FKM-type (see the classification of homogeneous isoparametric hypersurfaces of Takagi and Takahashi [61]).

Cecil, Chi and Jensen [12] then showed that if the multiplicities of an isoparametric hypersurface with four principal curvatures satisfy the condition $m_2 \geq 2m_1 - 1$, then the hypersurface is of FKM-type. (A different proof of this result, using isoparametric triple systems, was given later by Immervoll [36].)

Taken together with known results of Takagi [60] for $m_1 = 1$, and Ozeki and Takeuchi [48] for $m_1 = 2$, this result of Cecil, Chi and Jensen handled all possible pairs of multiplicities except for four cases, the FKM pairs (3, 4), (6, 9) and (7, 8), and the homogeneous pair (4, 5).

In a series of recent papers, Chi [14]-[19] completed the classification of isoparametric hypersurfaces with four principal curvatures. Specifically, Chi showed that in the cases of multiplicities (3, 4), (6, 9) and (7, 8), the isoparametric hypersurface must be of FKM-type, and in the case (4, 5), it must be homogeneous.

The final conclusion is that an isoparametric hypersurface with g = 4 principal curvatures must either be of FKM-type, or else a homogeneous isoparametric hypersurface with multiplicities (2, 2) or (4, 5).

In the case of an isoparametric hypersurface with g = 6 principal curvatures, Münzner [44]–[45] showed that all of the principal curvatures must have the same multiplicity m, and Abresch [1] showed that m must equal 1 or 2. By the classification of homogeneous isoparametric hypersurfaces due to Takagi and Takahashi [61], there is up to congruence only one homogeneous family in each case, m = 1 or m = 2.

These homogeneous examples have been shown to be the only isoparametric hypersurfaces in the case g = 6 by Dorfmeister and Neher [22] in the case of multiplicity m = 1, and by Miyaoka [42]–[43] in the case m = 2 (see also the papers of Siffert [54]–[55]).

For general surveys on isoparametric hypersurfaces in spheres, see the papers of Thorbergsson [72], Cecil [11], and Chi [20].

References

[1] U. Abresch, Isoparametric hypersurfaces with four or six distinct principal curvatures, Math. Ann. 264 (1983), 283–302.

- R. Bott and H. Samelson, Applications of the theory of Morse to symmetric spaces, Amer. J. Math. 80 (1958), 964–1029. Corrections in 83 (1961), 207–208.
- [3] E. Cartan, Familles de surfaces isoparamétriques dans les espaces à courbure constante, Annali di Mat. 17 (1938), 177–191 (see also an English translation by T.E. Cecil, https://crossworks.holycross.edu/math_fac_scholarship/17).
- [4] E. Cartan, Sur des familles remarquables d'hypersurfaces isoparamétriques dans les espaces sphériques, Math. Z. 45 (1939), 335–367 (see also an English translation by T.E. Cecil, https://crossworks.holycross.edu/math_fac_scholarship/18).
- [5] E. Cartan, Sur quelque familles remarquables d'hypersurfaces, C.R. Congrès Math. Liège, 1939, 30–41 (see also an English translation by T.E. Cecil, https://crossworks.holycross.edu/math_fac_scholarship/19).
- [6] E. Cartan, Sur des familles d'hypersurfaces isoparamétriques des espaces sphériques à 5 et à 9 dimensions, Revista Univ. Tucuman, Serie A, 1 (1940), 5–22 (see also an English translation by T.E. Cecil, https://crossworks.holycross.edu/math_fac_scholarship/20).
- [7] E. Cartan and J.A. Schouten, On the riemannian Geometries admitting an absolute parallelism, Proc. Akad. Amsterdam 29 (1926), 933–946.
- [8] S. Carter and A. West, *Isoparametric systems and transnormality*, Proc. London Math. Soc. 51 (1985), 520–542.
- [9] S. Carter and A. West, *Isoparametric and totally focal submanifolds*, Proc. London Math. Soc. 60 (1990), 609–624.
- [10] T.E. Cecil, A characterization of metric spheres in hyperbolic space by Morse theory, Tôhoku Math. J. 26 (1974), 341–351.
- T. Cecil, Isoparametric and Dupin hypersurfaces, SIGMA 4 (2008), 062, 28 pages, 2008.
- [12] T. Cecil, Q.-S. Chi and G. Jensen, Isoparametric hypersurfaces with four principal curvatures, Ann. Math. 166 (2007), 1–76.

- [13] T. Cecil and P. Ryan, Geometry of Hypersurfaces, Springer, New York, 2015.
- [14] Q.-S. Chi, Isoparametric hypersurfaces with four principal curvatures revisited, Nagoya Math. J. 193 (2009), 129–154.
- [15] Q.-S. Chi, A note on the paper "Isoparametric hypersurfaces with four principal curvatures", Hongyou Wu Memorial Volume, Pacific Journal of Applied Mathematics 3 (2011), 127–134.
- [16] Q.-S. Chi, Isoparametric hypersurfaces with four principal curvatures II, Nagoya Math. J. 204 (2011), 1–18.
- [17] Q.-S. Chi, A new look at Condition A, Osaka J. Math. 49 (2012), 133– 166.
- [18] Q.-S. Chi, Isoparametric hypersurfaces with four principal curvatures III, J. Differential Geom. 94 (2013), 469–504.
- [19] Q.-S. Chi, Isoparametric hypersurfaces with four principal curvatures IV, J. Differential Geom. 115 (2020), 225–301.
- [20] Q.-S. Chi, The isoparametric story, a heritage of Élie Cartan, Proceedings of the International Consortium of Chinese Mathematicians, 2018, 197–260, International Press, 2021.
- [21] S. Console and C. Olmos, Clifford systems, algebraically constant second fundamental form and isoparametric hypersurfaces, Manuscripta Math. 97 (1998), 335–342.
- [22] J. Dorfmeister and E. Neher, *Isoparametric hypersurfaces, case* g = 6, m = 1, Communications in Algebra 13 (1985), 2299–2368.
- [23] H. Ewert, Uber isoparametrische Hyperflächen in Räumen konstanter Krümmung, Diplomarbeit, Universität Köln (1994).
- [24] F. Fang, Multiplicities of principal curvatures of isoparametric hypersurfaces, preprint, Max Planck Institut f
 ür Mathematik, 1996.
- [25] F. Fang, On the topology of isoparametric hypersurfaces with four distinct principal curvatures, Proc. Amer. Math. Soc. 127 (1999), 259–264.

- [26] F. Fang, Topology of Dupin hypersurfaces with six principal curvatures, Math. Z. 231(1999), 533–555.
- [27] D. Ferus, H. Karcher and H.-F. Münzner, *Cliffordalgebren und neue isoparametrische Hyperflächen*, Math. Z. 177 (1981), 479–502 (see also an English translation by T.E. Cecil, arXiv:1112.2780v1 [mathDG] 13 Dec 2011).
- [28] H. Freudenthal, Oktaven, Ausnahmegruppen und Oktavengeometrie, Geom. Dedicata 19 (1985), 7–63.
- [29] A. Giunta and C. Sanchez, Projective spaces in the algebraic sets of planar normal sections of homogeneous isoparametric submanifolds, Rev. Un. Mat. Argentina 55 (2014), 139–154.
- [30] K. Grove and S. Halperin, Dupin hypersurfaces, group actions, and the double mapping cylinder, J. Differential Geom. 26 (1987), 429–459.
- [31] L.C. Grove and C.T. Benson, *Finite Reflection Groups*, second edition, Graduate Texts in Mathematics 99, Springer, New York, 1985.
- [32] C. Harle, Isoparametric families of submanifolds, Bol. Soc. Brasil Mat. 13 (1982), 35–48.
- [33] W.-Y. Hsiang and H.B. Lawson, Jr., Minimal submanifolds of low cohomogeneity, J. Differential Geom. 5 (1971), 1–38.
- [34] W.-Y. Hsiang, R. Palais and C.-L. Terng, The topology of isoparametric submanifolds, J. Differential Geom. 27 (1988), 423–460.
- [35] A. Hurwitz, Über die Komposition der quadratischen Formen, Math. Ann. 88 (1923), 1–25.
- [36] S. Immervoll, On the classification of isoparametric hypersurfaces with four distinct principal curvatures in spheres, Ann. Math. 168 (2008), 1011–1024.
- [37] N. Knarr and L. Kramer, Projective planes and isoparametric hypersurfaces, Geom. Dedicata 58 (1995), 193–202.
- [38] S. Kobayashi and K. Nomizu, Foundations of Differential Geometry, Vols. I, II, Wiley–Interscience, New York, 1963, 1969.

- [39] N. Kuiper, Tight embeddings and maps. Submanifolds of geometrical class three in Eⁿ, The Chern Symposium 1979 (Proc. Internat. Sympos., Berkeley, Calif., 1979), 97–145, Springer, Berlin–New York, 1980.
- [40] T. Levi-Civita, Famiglie di superficie isoparametrische nell'ordinario spacio euclideo, Atti. Accad. naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. 26 (1937), 355–362.
- [41] J.A. Little, Manifolds with planar geodesics, J. Differential Geom. 11 (1976), 265–285.
- [42] R. Miyaoka, Isoparametric hypersurfaces with (g,m) = (6,2), Ann. Math. 177 (2013), 53–110.
- [43] R. Miyaoka, Errata on isoparametric hypersurfaces with (g, m) = (6, 2), Ann. Math. 183 (2016), 1057–1071.
- [44] H.-F. Münzner, Isoparametrische Hyperflächen in Sphären I, Math. Ann. 251 (1980), 57–71 (see also an English translation by T.E. Cecil, https://crossworks.holycross.edu/math_fac_scholarship/14).
- [45] H.-F. Münzner, Isoparametrische Hyperflächen in Sphären II, Math. Ann. 256 (1981), 215–232 (see also an English translation by T.E. Cecil, https://crossworks.holycross.edu/math_fac_scholarship/13).
- [46] K. Nomizu, Some results in E. Cartan's theory of isoparametric families of hypersurfaces, Bull. Amer. Math. Soc. 79 (1973), 1184–1188.
- [47] K. Nomizu, Élie Cartan's work on isoparametric families of hypersurfaces, Proc. Sympos. Pure Math. 27, Part 1, 191–200, Amer. Math. Soc., Providence, RI, 1975.
- [48] H. Ozeki and M. Takeuchi, On some types of isoparametric hypersurfaces in spheres I, Tôhoku Math. J. 27 (1975), 515–559.
- [49] H. Ozeki and M. Takeuchi, On some types of isoparametric hypersurfaces in spheres II, Tôhoku Math. J. 28 (1976), 7–55.
- [50] P.J. Ryan, Hypersurfaces with parallel Ricci tensor, Osaka J. Math. 8 (1971), 251–259.

- [51] C. Sanchez, Triality and the normal sections of Cartan's isoparametric hypersurfaces, Rev. Un. Mat. Argentina 52 (2011), 73–88.
- [52] B. Segre, Una proprietá caratteristica de tre sistemi ∞^1 di superficie, Atti Acc. Sc. Torino 59 (1924), 666–671.
- [53] B. Segre, Famiglie di ipersuperficie isoparametrische negli spazi euclidei ad un qualunque numero di demesioni, Atti. Accad. naz, Lincei Rend. Cl. Sci. Fis. Mat. Natur. 27 (1938), 203–207.
- [54] A. Siffert, Classification of isoparametric hypersurfaces in spheres with (g,m) = (6,1), Proc. Amer. Math. Soc. 144 (2016), 2217–2230.
- [55] A. Siffert, A new structural approach to isoparametric hypersurfaces in spheres, Ann. Global Anal. Geom. 52 (2017), 425–456.
- [56] C. Somigliana, Sulle relazione fra il principio di Huygens e l'ottica geometrica, Atti Acc. Sci. Torino 54 (1918–1919), 974–979. Also in: Memorie Scelte, 434–439.
- [57] M. Spivak, A Comprehensive Introduction to Differential Geometry, Vols. 1–5, 3rd edition, Publish or Perish, Houston, Texas, 1999.
- [58] S. Stolz, Multiplicities of Dupin hypersurfaces, Invent. Math. 138 (1999), 253–279.
- [59] W. Strübing, Isoparametric submanifolds, Geom. Dedicata 20 (1986), 367–387.
- [60] R. Takagi, A class of hypersurfaces with constant principal curvatures in a sphere, J. Differential Geom. 11 (1976), 225–233.
- [61] R. Takagi and T. Takahashi, On the principal curvatures of homogeneous hypersurfaces in a sphere, Differential Geometry in honor of K. Yano, 469–481, Kinokuniya, Tokyo, 1972.
- [62] M. Takeuchi and S. Kobayashi, *Minimal imbeddings of R-spaces*, J. Differential Geom. 2 (1968), 203–215.
- [63] Z.-Z. Tang, Isoparametric hypersurfaces with four distinct principal curvatures, Chinese Sci. Bull. 36 (1991), 1237–1240.

- [64] Z.-Z. Tang, Multiplicities of equifocal hypersurfaces in symmetric spaces, Asian J. Math. 2 (1998), 181–214.
- [65] C.-L. Terng, Isoparametric submanifolds and their Coxeter groups, J. Differential Geom. 21 (1985), 79–107.
- [66] C.-L. Terng, Convexity theorem for isoparametric submanifolds, Invent. Math. 85 (1986), 487–492.
- [67] C.-L. Terng, Submanifolds with flat normal bundle, Math. Ann. 277 (1987), 95–111.
- [68] C.-L. Terng, Proper Fredholm submanifolds of Hilbert space, J. Differential Geom. 29 (1989), 9–47.
- [69] C.-L. Terng, Recent progress in submanifold geometry, Proc. Sympos. Pure Math. 54, Part 1, 439-484, Amer. Math. Soc., Providence, RI, 1993.
- [70] G. Thorbergsson, Dupin hypersurfaces, Bull. London Math. Soc. 15 (1983), 493–498.
- [71] G. Thorbergsson, Isoparametric foliations and their buildings, Ann. Math. 133 (1991), 429–446.
- [72] G. Thorbergsson, A survey on isoparametric hypersurfaces and their generalizations, Handbook of Differential Geometry, Vol. I, 963–995, North-Holland, Amsterdam, 2000.
- [73] Q.-M. Wang, Isoparametric hypersurfaces in complex projective spaces, pp. 1509–1523, Symposium on Differential Geometry and Differential Equations (Beijing, 1980), vol. 3, Science Press, Beijing, 1982.
- [74] A. West, *Isoparametric systems*, Geometry and Topology of Submanifolds I, 222–230, World Scientific, River Edge, NJ, 1989.

Department of Mathematics and Computer Science College of the Holy Cross Worcester, MA 01610, U.S.A. email: tcecil@holycross.edu