

Compatibility between e^+e^- and τ decay data in the di-pion channel and implications for a_{μ}^{SM} and CVC tests

Alejandro Miranda^{*a*,*}

^aInstitut de Física d'Altes Energies (IFAE) and The Barcelona Institute of Science and Technology (BIST), Campus UAB, 08193 Bellaterra (Barcelona), Spain.

E-mail: jmiranda@ifae.es

We have revisited the isospin-breaking corrections relating $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$ and $\Gamma(\tau^- \rightarrow \pi^-\pi^0 \nu_{\tau})$ [1]. We confirm that the associated uncertainty is under control, so that tau data can also be used to predict accurately the leading hadronic contribution to the muon anomalous magnetic moment and precision conserved vector current tests can be carried out.

10th International Conference on Quarks and Nuclear Physics (QNP2024) 8-12 July, 2024 Barcelona, Spain

*Speaker

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).



1. Introduction:

The measurement of the muon anomalous magnetic moment, $a_{\mu} = (g_{\mu} - 2)/2$ has achieved an extraordinary precision, thanks to the (perfectly compatible) BNL [2] and FNAL [3, 4] results, leading to the combination

$$a_{\mu}^{\exp} = 0.00116592059 \pm 0.0000000022 \,. \tag{1}$$

On the contrary, the corresponding Standard Model (SM) prediction is not clear at the moment, due to the uncertainties on the hadronic vacuum polarization piece and, specifically, on its 2π part. For the Muon g-2 Theory Initiative White Paper [5], this contribution was dominated by the combination of the KLOE [6] and BaBar [7] results, which were in tension (but still at the limit of compatibility, so they could be combined). The White Paper a_{μ} prediction is 5.1 σ smaller than the experimental average, Eq. (1). Although it was not used for the White Paper, the BMW lattice collaboration obtained a result for a_{μ}^{HVP} [8] that is only 1.1 σ smaller than a_{μ}^{exp} . Its recent improvement combining the lattice simulation with data [9] achieved accord within 0.9 σ with a_{μ}^{exp} . In 2023 the situation with e^+e^- data became puzzling, because the new CMD-3 measurement [10, 11] disagrees so much with KLOE, that they cannot be combined, a discrepancy which is not yet understood.

Given this conundrum, we recall that we have advocated since [12] (and emphasized and updated in [1, 13]) that two-pion tau decay data should be used again ¹ to obtain the corresponding contribution to a_{μ}^{HVP} , given the fact that:

- 1. All measurements, by the ALEPH [23], Belle [24], CLEO [25] and OPAL [26] collaborations are consistent, within errors.
- 2. The uncertainty related to the isospin breaking (IB) corrections required is small enough to make this procedure competitive.

Here we revisit the IB corrections relating e^+e^- and τ data, with a particular focus on those arising from the different neutral (electromagnetic) and charged (weak) form factors.

2. IB corrections

In the data-driven approach, at lowest order (LO), the HVP contribution to a_{μ} is [27]

$$a_{\mu}^{\text{HVP, LO}} = \frac{1}{4\pi^3} \int_{m_{\pi^0}}^{\infty} ds \, K(s) \, \sigma_{e^+ e^- \to \text{hadrons}(+\gamma)}^0(s), \tag{2}$$

where $\sigma_{e^+e^- \to \text{hadrons}(+\gamma)}^0(s)$ is the bare hadronic cross-section at hadrons invariant mass \sqrt{s} with vacuum polarization (VP) effects removed [28] and K(s) is a smooth kernel enhancing the low-energy effects [29].

¹The usefulness of τ data for this purpose was put forward by Alemany, Davier and Höcker in Ref. [14], see also e.g. Refs. [15–22].

Alternatively, it can also be obtained from ²

$$\sigma_{\pi\pi(\gamma)}^{0}\Big|_{\text{CVC}} = \left[\frac{K_{\sigma}(s)}{K_{\Gamma}(s)}\frac{d\Gamma_{\pi\pi[\gamma]}}{ds}\right] \times \frac{R_{\text{IB}}(s)}{S_{\text{EW}}},\tag{3}$$

where the ratio of K functions depends on G_F , V_{ud} , α and m_{τ} , and the measured spectrum of the di-pion tau decays ³, $\frac{d\Gamma_{\pi\pi[\gamma]}}{ds}$, needs to be corrected for the short-distance universal electroweak radiative corrections, S_{EW} [34], and the isospin-breaking (IB) factor

$$R_{\rm IB}(s) = \frac{\rm FSR(s)}{G_{\rm EM}(s)} \frac{\beta_{\pi^+\pi^-}^3(s)}{\beta_{\pi^+\pi^0}^3(s)} \left| \frac{F_V(s)}{f_+(s)} \right|^2.$$
(4)

 $R_{\rm IB}(s)$ depends on two factors which are straightforward: final-state radiative corrections (FSR) [35] and the kinematical factor depending on the ratio of β functions. However, it also depends on two corrections that are challenging. The first of these corresponds to the long-distance QED virtual plus real photon corrections, encoded in the $G_{\rm EM}$ function, which has been the focus of Refs. [12, 15, 16, 36, 37] ⁴. Here we concentrate on the second of such corrections, given by the ratio of the neutral F_V and charged f_+ pion form factors, where one of the leading IB corrections, the $\rho - \omega$ mixing entering F_V , takes place. In this factor, also the mass and (partial) width differences of the neutral and charged $\rho \to \pi\pi(\gamma)$ channels we rely on the results of Ref. [47].

In this contribution we will focus on two observables which are sensitive to IB. We will consider the IB corrections to the $\pi\pi$ contribution to $a_{\mu}^{\text{HVP, LO}}$ obtained using tau data,

$$\Delta a_{\mu}^{\text{HVP, LO}}[\pi\pi, \tau] = \frac{1}{4\pi^3} \int_{4m_{\pi}^2}^{m_{\tau}^2} ds \, K(s) \left[\frac{K_{\sigma}(s)}{K_{\Gamma}(s)} \frac{d\Gamma_{\pi\pi[\gamma]}}{ds} \right] \left(\frac{R_{\text{IB}}(s)}{S_{\text{EW}}} - 1 \right),\tag{5}$$

and also the modification to the di-pion tau decay branching ratio obtained from the e^+e^- measurement,

$$\Delta \mathcal{B}_{\pi\pi^0}^{\text{CVC}} = \mathcal{B}_e \int_{4m_\pi^2}^{m_\tau^2} ds \,\sigma_{\pi^+\pi^-(\gamma)}(s) \mathcal{N}(s) \left(\frac{S_{\text{EW}}}{R_{\text{IB}}(s)} - 1\right),\tag{6}$$

where \mathcal{B}_e is the electronic tau decay branching ratio and $\mathcal{N}(s)$ depends on α , V_{ud} and m_{τ} .

3. Form factor parametrizations

We considered different descriptions of the electromagnetic, F_V , and weak, f_+ , pion form factors (see Ref. [1] for details). As typically done by the experimental collaborations, we employed the Gounaris-Sakurai (GS) [48] and Kühn-Santamaría (KS) [49] form factors. We also used the Guerrero-Pich (GP) [50] parametrization. However, since it is limited to the ρ (770) resonance, we considered its extension including the ρ' and ρ'' resonances, along the lines of Refs. [51, 52], to

²CVC stands for conserved vector current, which relates the $e^+e^- \rightarrow \pi^+\pi^-$ and $\tau^- \rightarrow \pi^-\pi^0\nu_{\tau}$ in absence of IB corrections.

³New physics could affect the tau decays and not the e^+e^- cross-section into two pions [30–33], nevertheless.

⁴We stick here to the ' $O(p^4)$ ' result of Ref. [12], that uses Resonance Chiral Theory [38, 39], which has been successfully employed in computing different contributions to a_{μ} [13, 16, 40–46]. Noteworthy, this result is consistent with the vector meson dominance one employed by the Orsay group since [19].

construct the input phase shift ⁵ of an unsubtracted dispersive form factor, where we accounted for $\rho - \omega - \phi$ mixing (we verified that complex mixing coefficients were needed) and inelastic effects (captured by a conformal polynomial), following Ref. [53].

4. Results

We have computed the IB corrections in Eqs. (5) and (6), according to the different form factor parametrizations: GS, KS, GP, Seed and Dispersive. Both the fits to data and the analyticity tests (see Ref. [1]) work best (and better when KLOE is excluded) for GS and Dispersive. Taking this into account, we will take the latter as our reference result and add linearly a systematic uncertainty coming from its difference with GS in our final results.

In Table 1 we collect the different IB corrections to the CVC di-pion tau decay branching ratio, obtained using the different form factors considered, and we split them according to their type. Focusing on GS and Dispersive, there is good agreement in all contributions (but for the individual effect of the ρ mass difference and the $\rho - \omega$ interference, which are compensated in their sum, however) and the final results agree nicely.

Source	$\Delta \mathcal{B}_{\pi\pi}^{\rm CVC} (10^{-2})$						
	GS	KS	GP	Seed	Dispersive		
					p_{4-1}		
$S_{\rm EW}$			+0.57(1)				
$G_{ m EM}$			$-0.09(^{3}_{1})$				
FSR			-0.19(2)				
$m_{\pi^{\pm}} - m_{\pi^0}$ effect on σ			+0.20				
$m_{\pi^{\pm}} - m_{\pi^0}$ effect on Γ_{ρ}	-0.21	-0.22	-0.22	-0.23	-0.20		
$m_{K^{\pm}} - m_{K^0}$ effect on Γ_{ρ}	-	-	-0.02	-0.03	+0.01		
$m_{ ho^{\pm}} - m_{ ho^0}$	+0.08(8)	+0.09(8)	-0.02(2)	-0.02(2)	$-0.02(^{2}_{1})$		
$\rho - \omega$ interference	$-0.08(0)({}^{12}_{0})$	$-0.09(0)({}^{11}_0)$	$-0.09(0)\binom{16}{1}$	$-0.06(0)\binom{10}{0}$	$-0.01(0)\binom{6}{4}$		
$\rho - \phi$ interference	$-0.00(0)(^{1}_{0})$	-0.00(0)(0)	-	-0.01(0)(0)	-0.01(0)(1)		
$\pi\pi\gamma$, electromagnetic decays	+0.34(3)	+0.37(4)	+0.34(4)	+0.34(4)	+0.37(4)		
TOTAL	$+0.62(^{15}_{9})$	$+0.64(^{15}_{9})$	$+0.48(^{17}_{5})$	$+0.48(^{12}_{5})$	$+0.63\binom{8}{6}\binom{0}{3}$		

Table 1: IB contributions to BR($\tau^- \rightarrow \pi^- \pi^0 \nu_{\tau}$) according to the different form factor inputs. The uncertainties are mostly of systematic origin and are specified in Ref. [1]. In the last entry, we take as an additional uncertainty (last shown) the difference between our preferred option for the conformal polynomial (p_{4-1}) and the other dispersive results that we considered [1].

In Table 2 we display the different IB corrections to the di-pion contribution to $a_{\mu}^{\text{HVP, LO}}$ using the diverse FF considered. Concentrating on the GS and dispersive results, we made analogous observations as with Table 1, highlighting again the good consistency of our reference results.

In Fig. 1 we show the IB corrections to the di-pion tau decay branching ratio and contribution to $a_{\mu}^{\text{HVP, LO}}$, using the inputs in Tables 1 and 2, where we also compare to the results in Ref. [19] (in blue).

Finally, in Fig. 2 we compare the measured di-pion tau decay branching fractions and the predictions from the $e^+e^- \rightarrow \pi^+\pi^-$ spectral functions, applying the IB corrections given in Table

⁵We also used the form factor giving this phase, which we labelled 'Seed'.

Compatibility between e^+e^- and τ decay data in the di-pion channel and implications for a_{μ}^{SM} and CVC tests Alejandro Miranda

Source	$\Delta a_{\mu}^{ m had,LO}[\pi\pi,\tau] (10^{-10})$							
	GS	KS	GP	Seed	Dispersive			
					p_{4-1}			
$S_{\rm EW}$			-11.96(0.15)					
G_{EM}			$-1.71(^{0.61}_{1.48})$					
FSR			+4.56(0.46)					
$m_{\pi^{\pm}} - m_{\pi^0}$ effect on σ			-7.47					
$m_{\pi^{\pm}} - m_{\pi^0}$ effect on Γ	+3.74	+4.12	+4.07	+4.13	+3.58			
$m_{K^{\pm}} - m_{K^0}$ effect on Γ	-	-	+0.37	+0.36	-0.22			
$m_{\rho^{\pm}} - m_{\rho^0}$	$+0.10(^{0.18}_{0.09})$	$-0.04(^{0.06}_{0.00})$	$+1.87(^{1.75}_{1.68})$	$+1.86(^{1.75}_{1.68})$	$+1.85(^{1.69}_{1.66})$			
ρ – ω interference	$+3.84(0.08)(^{0.23}_{2.17})$	$+4.00(0.08)(^{0.25}_{1.96})$	$+4.33(0.07)\binom{0.33}{2.83}$	$+3.57(0.07)(^{0.18}_{1.77})$	$+2.72(0.08)(^{0.65}_{1.15})$			
$\rho - \phi$ interference	$+0.09(0.03)(^{0.06}_{0.10})$	$+0.03(0.03)({}^{0.03}_{0.05})$	-	$+0.13(0.03)(^{0.05}_{0.07})$	$+0.12(0.02)(^{0.14}_{0.08})$			
ππγ	-6.09(0.67)	-6.68(0.74)	-6.19(0.68)	-6.21(0.68)	-6.62(0.73)			
TOTAL	$-14.90(\frac{1.08}{2.76})$	$-15.15(^{1.11}_{2.61})$	$-12.13(^{2.06}_{3.70})$	$-12.74(^{2.04}_{2.97})$	$-15.15(^{2.11}_{2.65})(^{0.24}_{0.09})$			

Table 2: Contributions to $a_{\mu}^{had,LO}[\pi\pi, \tau](10^{-10})$ from the isospin-breaking (IB) corrections according to the different form factor inputs. The uncertainties are mostly systematic, and are discussed in Ref. [1]. In the last entry, we take as an additional uncertainty (last shown) the difference between p_{4-1} and the other dispersive results that we considered.



Figure 1: IB corrections in the ratio of the form factors $|F_V(s)/f_+(s)|$ to $a_\mu^{\text{HVP, LO}}$ and $\mathcal{B}_{\pi\pi}^{\text{CVC}}$.

1. We remark the very good consistency between all results using the CMD-3 data and updated IB corrections (even more for our preferred results, GS and dispersive, and those of Ref. [22]).

5. Conclusions

We have revisited the IB corrections relating the e^+e^- and τ decay di-pion observables, particularly focusing on the one coming from the ratio of the electromagnetic and weak form factors. We have considered the popular parametrizations Gounaris-Sakurai, Kühn-Santamaría, Guerrero-Pich (and its extension with excited isovector mesons) and a dispersive form factor, favoring the first and last one through fits to data and analyticity tests (detailed in Ref. [1]). For our main observables of interest, we find $\Delta \mathcal{B}_{\pi\pi}^{\text{CVC}} = (+0.63^{+0.09}_{-0.08}) \times 10^{-2}$, $\Delta a_{\mu}^{\text{had,LO}[\pi\pi,\tau]} = (-15.15^{+2.37}_{-2.90}) \times 10^{-10}$, corresponding to $\Delta a_{\mu} = a_{\mu}^{\text{exp}} - a_{\mu}^{\text{SM}} = (14.8^{+5.1}_{-5.4}) \times 10^{-10}$, a 2.7 σ difference, with agrees nicely with Refs. [19, 22]. Our analysis confirms the reliability of these IB corrections, supporting the use of tau data in the updated SM prediction of a_{μ} .



Figure 2: Comparison between the measured branching fractions for $\tau^- \rightarrow \pi^- \pi^0 v_{\tau}$ and the prediction from the $e^+e^- \rightarrow \pi^+\pi^-$ spectral functions, applying the isospin-breaking corrections given in Table 1. Our reference result comes from the dispersive evaluation, and we add linearly to the final theory uncertainty (second error shown, preceded by the statistical one) its difference with the GS value, that also complies well with analyticity.

Acknowledgments

We gladly acknowledge the QNP24 organizing committees. We thank financial support through the Ministerio de Ciencia e Innovación under Grant No. PID2020–112965 GB-I00, by the Departament de Recerca i Universitats from Generalitat de Catalunya to the Grup de Recerca "Grup de Física Teòrica UAB/IFAE" (Codi: 2021 SGR 00649) and partial Conahcyt support, from project CB2023-2024-3226.

References

- [1] G. L. Castro, A. Miranda and P. Roig, "Isospin breaking corrections in 2π production in tau decays and e^+e^- annihilation: consequences for the muon g - 2 and CVC tests," [arXiv:2411.07696 [hep-ph]].
- [2] G. W. Bennett *et al.* [Muon g-2], "Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL," Phys. Rev. D 73 (2006), 072003, doi:10.1103/PhysRevD.73.072003, [arXiv:hep-ex/0602035 [hep-ex]].
- [3] B. Abi et al. [Muon g-2], "Measurement of the Positive Muon Anomalous Magnetic Moment to

0.46 ppm," Phys. Rev. Lett. **126** (2021) no.14, 141801, doi:10.1103/PhysRevLett.126.141801, [arXiv:2104.03281 [hep-ex]].

- [4] D. P. Aguillard et al. [Muon g-2], "Measurement of the Positive Muon Anomalous Magnetic Moment to 0.20 ppm," Phys. Rev. Lett. 131 (2023) no.16, 161802, doi:10.1103/PhysRevLett.131.161802, [arXiv:2308.06230 [hep-ex]].
- [5] T. Aoyama, et al. "The anomalous magnetic moment of the muon in the Standard Model," Phys. Rept. 887 (2020), 1-166, doi:10.1016/j.physrep.2020.07.006, [arXiv:2006.04822 [hep-ph]].
- [6] A. Anastasi *et al.* [KLOE-2], "Combination of KLOE $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma(\gamma))$ measurements and determination of $a_{\mu}^{\pi^+\pi^-}$ in the energy range 0.10 < s < 0.95 GeV²," JHEP **03** (2018), 173, doi:10.1007/JHEP03(2018)173, [arXiv:1711.03085 [hep-ex]].
- [7] J. P. Lees *et al.* [BaBar], "Precise Measurement of the $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ Cross Section with the Initial-State Radiation Method at BABAR," Phys. Rev. D **86** (2012), 032013, doi:10.1103/PhysRevD.86.032013, [arXiv:1205.2228 [hep-ex]].
- [8] S. Borsanyi, et al. "Leading hadronic contribution to the muon magnetic moment from lattice QCD," Nature 593 (2021) no.7857, 51-55 doi:10.1038/s41586-021-03418-1 [arXiv:2002.12347 [hep-lat]].
- [9] A. Boccaletti, et al. "High precision calculation of the hadronic vacuum polarisation contribution to the muon anomaly," [arXiv:2407.10913 [hep-lat]].
- [10] F. V. Ignatov *et al.* [CMD-3], "Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section from threshold to 1.2 GeV with the CMD-3 detector," Phys. Rev. D 109 (2024) no.11, 112002 doi:10.1103/PhysRevD.109.112002 [arXiv:2302.08834 [hep-ex]].
- [11] F. V. Ignatov et al. [CMD-3], "Measurement of the Pion Form Factor with CMD-3 Detector and its Implication to the Hadronic Contribution to Muon (g-2)," Phys. Rev. Lett. 132 (2024) no.23, 231903, doi:10.1103/PhysRevLett.132.231903, [arXiv:2309.12910 [hep-ex]].
- [12] J. A. Miranda and P. Roig, "New τ-based evaluation of the hadronic contribution to the vacuum polarization piece of the muon anomalous magnetic moment," Phys. Rev. D 102 (2020), 114017, doi:10.1103/PhysRevD.102.114017, [arXiv:2007.11019 [hep-ph]].
- [13] P. Masjuan, A. Miranda and P. Roig, "τ data-driven evaluation of Euclidean windows for the hadronic vacuum polarization," Phys. Lett. B 850 (2024), 138492, doi:10.1016/j.physletb.2024.138492, [arXiv:2305.20005 [hep-ph]].
- [14] R. Alemany, M. Davier and A. Höcker, "*Improved determination of the hadronic contribution to the muon (g-2) and to alpha (M(z)) using new data from hadronic tau decays*," Eur. Phys. J. C 2 (1998), 123-135, doi:10.1007/s100520050127 [arXiv:hep-ph/9703220 [hep-ph]].
- [15] V. Cirigliano, G. Ecker and H. Neufeld, "Isospin violation and the magnetic moment of the muon," Phys. Lett. B 513 (2001), 361-370, doi:10.1016/S0370-2693(01)00764-X, [arXiv:hepph/0104267 [hep-ph]].

- [16] V. Cirigliano, G. Ecker and H. Neufeld, "*Radiative tau decay and the magnetic moment of the muon*," JHEP **08** (2002), 002, doi:10.1088/1126-6708/2002/08/002, [arXiv:hep-ph/0207310 [hep-ph]].
- [17] M. Davier, S. Eidelman, A. Hocker and Z. Zhang, "Confronting spectral functions from e⁺e⁻ annihilation and tau decays: Consequences for the muon magnetic moment," Eur. Phys. J. C 27 (2003), 497-521 doi:10.1140/epjc/s2003-01136-2 [arXiv:hep-ph/0208177 [hep-ph]].
- [18] M. Davier, S. Eidelman, A. Hocker and Z. Zhang, "Updated estimate of the muon magnetic moment using revised results from e⁺e⁻ annihilation," Eur. Phys. J. C **31** (2003), 503-510 doi:10.1140/epjc/s2003-01362-6 [arXiv:hep-ph/0308213 [hep-ph]].
- [19] M. Davier, A. Hoecker, G. López Castro, B. Malaescu, X. H. Mo, G. Toledo Sánchez, P. Wang, C. Z. Yuan and Z. Zhang, "The Discrepancy Between tau and e+e- Spectral Functions Revisited and the Consequences for the Muon Magnetic Anomaly," Eur. Phys. J. C 66 (2010), 127-136, doi:10.1140/epjc/s10052-009-1219-4, [arXiv:0906.5443 [hep-ph]].
- M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, "*Reevaluation of the Hadronic Contributions to the Muon g-2 and to alpha(M_Z)*," Eur. Phys. J. C 71 (2011), 1515, [erratum: Eur. Phys. J. C 72 (2012), 1874], doi:10.1140/epjc/s10052-012-1874-8, [arXiv:1010.4180 [hep-ph]].
- [21] M. Davier, A. Höcker, B. Malaescu, C. Z. Yuan and Z. Zhang, "Update of the ALEPH nonstrange spectral functions from hadronic τ decays," Eur. Phys. J. C **74** (2014) no.3, 2803, doi:10.1140/epjc/s10052-014-2803-9, [arXiv:1312.1501 [hep-ex]].
- [22] M. Davier, A. Hoecker, A. M. Lutz, B. Malaescu and Z. Zhang, "Tensions in e⁺e⁻ → π⁺π⁻(γ) measurements: the new landscape of data-driven hadronic vacuum polarization predictions for the muon g 2," Eur. Phys. J. C 84 (2024) no.7, 721 doi:10.1140/epjc/s10052-024-12964-7 [arXiv:2312.02053 [hep-ph]].
- [23] S. Schael et al. [ALEPH], "Branching ratios and spectral functions of tau decays: Final ALEPH measurements and physics implications," Phys. Rept. 421 (2005), 191-284, doi:10.1016/j.physrep.2005.06.007, [arXiv:hep-ex/0506072 [hep-ex]].
- [24] M. Fujikawa *et al.* [Belle], "*High-Statistics Study of the tau- to pi- pi0 nu(tau) Decay*," Phys. Rev. D 78 (2008), 072006, doi:10.1103/PhysRevD.78.072006, [arXiv:0805.3773 [hep-ex]].
- [25] S. Anderson *et al.* [CLEO], "*Hadronic structure in the decay tau- to pi- pi0 neutrino(tau)*," Phys. Rev. D 61 (2000), 112002, doi:10.1103/PhysRevD.61.112002, [arXiv:hep-ex/9910046 [hep-ex]].
- [26] K. Ackerstaff et al. [OPAL], "Measurement of the strong coupling constant alpha(s) and the vector and axial vector spectral functions in hadronic tau decays," Eur. Phys. J. C 7 (1999), 571-593, doi:10.1007/s100529901061, [arXiv:hep-ex/9808019 [hep-ex]].
- [27] M. Gourdin and E. De Rafael, "*Hadronic contributions to the muon g-2 factor*," Nucl. Phys. B 10 (1969), 667-674, doi:10.1016/0550-3213(69)90333-2.

- [28] S. Eidelman and F. Jegerlehner, "Hadronic contributions to g-2 of the leptons and to the effective fine structure constant $\alpha(M_Z^2)$," Z. Phys. C **67** (1995), 585-602 doi:10.1007/BF01553984 [arXiv:hep-ph/9502298 [hep-ph]].
- [29] S. J. Brodsky and E. De Rafael, "SUGGESTED BOSON LEPTON PAIR COUPLINGS AND THE ANOMALOUS MAGNETIC MOMENT OF THE MUON," Phys. Rev. 168 (1968), 1620-1622 doi:10.1103/PhysRev.168.1620
- [30] J. A. Miranda and P. Roig, "*Effective-field theory analysis of the* $\tau^- \rightarrow \pi^- \pi^0 v_\tau$ *decays*," JHEP **11** (2018), 038, doi:10.1007/JHEP11(2018)038, [arXiv:1806.09547 [hep-ph]].
- [31] V. Cirigliano, A. Falkowski, M. González-Alonso and A. Rodríguez-Sánchez, "Hadronic τ Decays as New Physics Probes in the LHC Era," Phys. Rev. Lett. 122 (2019) no.22, 221801, doi:10.1103/PhysRevLett.122.221801, [arXiv:1809.01161 [hep-ph]].
- [32] S. Gonzàlez-Solís, A. Miranda, J. Rendón and P. Roig, "Exclusive hadronic tau decays as probes of non-SM interactions," Phys. Lett. B 804 (2020), 135371, doi:10.1016/j.physletb.2020.135371, [arXiv:1912.08725 [hep-ph]].
- [33] V. Cirigliano, D. Díaz-Calderón, A. Falkowski, M. González-Alonso and A. Rodríguez-Sánchez, "Semileptonic tau decays beyond the Standard Model," JHEP 04 (2022), 152, doi:10.1007/JHEP04(2022)152, [arXiv:2112.02087 [hep-ph]].
- [34] W. J. Marciano and A. Sirlin, "Electroweak Radiative Corrections to tau Decay," Phys. Rev. Lett. 61 (1988), 1815-1818, doi:10.1103/PhysRevLett.61.1815.
- [35] M. Drees and K. i. Hikasa, "Scalar top production in e+ e- annihilation," Phys. Lett. B 252 (1990), 127-134 doi:10.1016/0370-2693(90)91094-R
- [36] F. Flores-Báez, A. Flores-Tlalpa, G. López Castro and G. Toledo Sánchez, "Long-distance radiative corrections to the di-pion tau lepton decay," Phys. Rev. D 74 (2006), 071301, doi:10.1103/PhysRevD.74.071301, [arXiv:hep-ph/0608084 [hep-ph]].
- [37] A. Flores-Tlalpa, F. Flores-Báez, G. López Castro and G. Toledo Sánchez, "Model-dependent radiative corrections to tau to pi- pi0 nu revisited," Nucl. Phys. B Proc. Suppl. 169 (2007), 250-254, doi:10.1016/j.nuclphysbps.2007.03.011, [arXiv:hep-ph/0611226 [hep-ph]].
- [38] G. Ecker, J. Gasser, A. Pich and E. de Rafael, "The Role of Resonances in Chiral Perturbation Theory," Nucl. Phys. B 321 (1989), 311-342, doi:10.1016/0550-3213(89)90346-5.
- [39] G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, "*Chiral Lagrangians for Massive Spin 1 Fields*," Phys. Lett. B 223 (1989), 425-432, doi:10.1016/0370-2693(89)91627-4.
- [40] P. Roig, A. Guevara and G. López Castro, "*VV'P form factors in resonance chiral theory and the* $\pi \eta \eta'$ *light-by-light contribution to the muon* g 2," Phys. Rev. D **89** (2014) no.7, 073016, doi:10.1103/PhysRevD.89.073016, [arXiv:1401.4099 [hep-ph]].

- [41] A. Guevara, P. Roig and J. J. Sanz-Cillero, "Pseudoscalar pole light-by-light contributions to the muon (g - 2) in Resonance Chiral Theory," JHEP 06 (2018), 160, doi:10.1007/JHEP06(2018)160, [arXiv:1803.08099 [hep-ph]].
- [42] P. Roig and P. Sánchez-Puertas, "Axial-vector exchange contribution to the hadronic light-bylight piece of the muon anomalous magnetic moment," Phys. Rev. D 101 (2020) no.7, 074019, doi:10.1103/PhysRevD.101.074019, [arXiv:1910.02881 [hep-ph]].
- [43] W. Qin, L. Y. Dai and J. Portolés, "Two and three pseudoscalar production in e⁺e⁻ annihilation and their contributions to (g-2)_μ," JHEP **03** (2021), 092, doi:10.1007/JHEP03(2021)092, [arXiv:2011.09618 [hep-ph]].
- [44] S. J. Wang, Z. Fang and L. Y. Dai, "Two body final states production in electronpositron annihilation and their contributions to (g-2)_μ," JHEP 07 (2023), 037, doi:10.1007/JHEP07(2023)037, [arXiv:2302.08859 [hep-ph]].
- [45] B. H. Qin, W. Qin and L. Y. Dai, "*Electron-positron annihilation into* $K\bar{K}\pi$ and their contributions to $(g 2)_{\mu}$," [arXiv:2403.14294 [hep-ph]].
- [46] E. J. Estrada, S. Gonzàlez-Solís, A. Guevara and P. Roig, "Improved π^0 , η , η' transition form factors in resonance chiral theory and their a_{μ}^{HLbL} contribution," [arXiv:2409.10503 [hep-ph]].
- [47] F. V. Flores-Baez, G. L. Castro and G. Toledo Sánchez, "The Width difference of rho vector mesons," Phys. Rev. D 76 (2007), 096010 doi:10.1103/PhysRevD.76.096010 [arXiv:0708.3256 [hep-ph]].
- [48] G. J. Gounaris and J. J. Sakurai, "*Finite width corrections to the vector meson dominance prediction for* $\rho \rightarrow e^+e^-$," Phys. Rev. Lett. **21** (1968), 244-247 doi:10.1103/PhysRevLett.21.244
- [49] J. H. Kühn and A. Santamaría, "Tau decays to pions," Z. Phys. C 48 (1990), 445-452 doi:10.1007/BF01572024
- [50] F. Guerrero and A. Pich, "Effective field theory description of the pion form-factor," Phys. Lett. B 412 (1997), 382-388 doi:10.1016/S0370-2693(97)01070-8 [arXiv:hep-ph/9707347 [hep-ph]].
- [51] D. Gómez Dumm and P. Roig, "Dispersive representation of the pion vector form factor in $\tau \rightarrow \pi \pi \nu_{\tau}$ decays," Eur. Phys. J. C **73** (2013) no.8, 2528 doi:10.1140/epjc/s10052-013-2528-1 [arXiv:1301.6973 [hep-ph]].
- [52] S. Gonzàlez-Solís and P. Roig, "A dispersive analysis of the pion vector form factor and $\tau^- \rightarrow K^- K_S v_\tau$ decay," Eur. Phys. J. C **79** (2019) no.5, 436 doi:10.1140/epjc/s10052-019-6943-9 [arXiv:1902.02273 [hep-ph]].
- [53] G. Colangelo, M. Hoferichter and P. Stoffer, "Two-pion contribution to hadronic vacuum polarization," JHEP 02 (2019), 006 doi:10.1007/JHEP02(2019)006 [arXiv:1810.00007 [hepph]].