W Hya: molecular inventory by ISO-SWS *

K. Justtanont¹, T. de Jong^{2,3}, A.G.G.M. Tielens^{4,5}, H. Feuchtgruber⁶, and L.B.F.M. Waters³

- ¹ Stockholm Observatory, AlbaNova, Dept. of Astronomy, SE-106 91 Stockholm, Sweden
- ² SRON-Utrecht, Sorbonnelaan 2, NL-3584 CA Utrecht, The Netherlands
- ³ University of Amsterdam, Kruislaan 403, NL-1098 SJ Amsterdam, The Netherlands
- ⁴ Kapteyn Institute, P.O. Box 800, NL-9700 AV Groningen, The Netherlands
- ⁵ SRON-Groningen, P.O. Box 800, NL-9700 AV Groningen, The Netherlands
- ⁶ Max-Planck Institut fuer extraterrestrische Physik, Giessenbachstrasse, D-85740 Garching, Germany

Received; accepted

Abstract. Infrared spectroscopy is a powerful tool to probe the inventory of solid state and molecular species in circumstellar ejecta. Here we analyse the infrared spectrum of the Asymptotic Giant Branch star W Hya, obtained by the Short and Long Wavelength Spectrometers on board of the Infrared Satellite Observatory. These spectra show evidence for the presence of amorphous silicates, aluminum oxide, and magnesium-iron oxide grains. We have modelled the spectral energy distribution using laboratory measured optical properties of these compounds and derive a total dust mass loss rate of $3 \times 10^{-10} \ \mathrm{M_{\odot} \ yr^{-1}}$. We find no satisfactory fit to the $13\mu\mathrm{m}$ dust emission feature and the identification of its carrier is still an open issue. We have also modelled the molecular absorption bands due to H₂O, OH, CO, CO₂, SiO, and SO₂ and estimated the excitation temperatures for different bands which range from 300 to 3000K. It is clear that different molecules giving rise to these absorption bands originate from different gas layers. We present and analyse high resolution Fabry-Perot spectra of the three CO₂ bands in the $15\mu\mathrm{m}$ region. In these data, the bands are resolved into individual Q-lines in emission, which allows the direct determination of the excitation temperature and column density of the emitting gas. This reveals the presence of a warm ($\simeq 450$ K) extended layer of CO₂, somewhere between the photosphere and the dust formation zone. The gas in this layer is cooler than the 1000K CO₂ gas responsible for the low-resolution absorption bands at 4.25 and 15μ m. The rotational and vibrational excitation temperatures derived from the individual Q-branch lines of CO_2 are different (~ 450 K and 150K, respectively) so that the CO_2 level population is not in LTE.

Key words. Stars: circumstellar matter – Stars: evolution – Stars: individual : W Hya – Stars: late-type – Stars: mass-loss – Infrared: stars

1. Introduction

One of the characteristics of Asymptotic Giant Branch (AGB) stars is their mass loss. As a star evolves up the AGB, its mass loss rate increases. Just before it evolves off to become a planetary nebula, it sheds most of its mass in a short time, i.e., superwind. Signatures of mass loss can be studied in the infrared by tracing the dust (e.g., Bedijn 1987, Justtanont & Tielens 1992), and in the submillimetre by using molecular rotational lines such as CO (e.g., Knapp & Morris 1985, Groenewegen 1994, Skinner et al. 1999).

An example of an early AGB star is W Hya. It has a relatively low mass loss rate as deduced from the

Send offprint requests to: K. Justtanont

weak CO millimeter observations (Wannier & Sahai 1986; Bujarrabal et al. 1989; Cernicharo et al. 1997). It is classified as a semiregular variable (SRa) from its light curve. The visual magnitude ranges from 6 to 10 over a period of 382 days (Lebzelter et al. 2000). It exhibits SiO (e.g., Schwartz et al. 1982; Clark et al. 1985), H₂O (e.g., Spencer et al. 1979; Lane et al. 1987) and OH main line masers (e.g., Szymczak et al. 1998; Chapman et al. 1994). It has a weak silicate dust emission at $10\mu m$ along with a smaller emission feature at $13\mu m$. The carrier of the latter is still being debated. W Hya is one of the first AGB stars for which H₂O rotational emission was reported after the launch of the Infrared Space Observatory (ISO, Kessler et al. 1996). Neufeld et al. (1996) and Barlow et al. (1996) reported that these lines were detected in Short and Long Wavelength Spectrometers (SWS and LWS), respectively. Both authors presented models accounting for the emission. Recently, Zubko & Zuckerman (2000) also reported a model predicting H₂O line fluxes observed by ISO. Also,

^{*} Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands and the United Kingdom) with the participation of ISAS and NASA.

W Hya has been detected in the ortho ground state line by SWAS at 557GHz (Harwit & Bergin 2002) In our paper, we mainly discuss the SWS spectra taken during the ISO mission. We also present the SWS Fabry-Perot spectrum of W Hya centered on CO_2 bands. This band has been reported by Ryde (1999) and Justtanont et al. (1998).

2. Observations

Spectroscopic observations of W Hya were taken aboard ISO using both the SWS (de Graauw et al. 1996) and LWS (Clegg et al. 1996). The SWS data were taken using the standard AOT1 (full scan from $2.38-45.2\mu m$) speed 3, giving an average resolution of 510. We also obtained the spectrum using the full grating resolution of 2000 between $12-16\mu m$ (AOT6) in order to study the CO₂ bands. The data were reduced using the data reduction package provided by the SWS calibration team based at SRON-Groningen. We were able to observe the three CO₂ bands using the SWS Fabry-Perot (FP) centering on the three Q-branches in order to resolve individual transitions. Care was taken to deglitch the spectrum due to cosmic ray hits on the detectors at the time of observation. As the data were taken in small wavelength intervals, each subband spectrum had to be stitched together. The final spectra were rebinned to a resolution of 4 10⁴, with an oversampling of 6. Also the flux level for each wavelength band has been normalised to the AOT6 flux.

The LWS data were taken from the archive (PI: M. Barlow). The reduction was done using the ISAP data reduction package provided by IPAC.

3. Dust and gas mass loss rates

In this section, we present a model fit to the infrared spectral energy distribution (SED) in order to derive the dust mass loss rate. This is then used to calculate the rate of gas which is driven by the dust to the observed terminal velocity by momentum transfer between gas and dust. At the same time, the temperature structure of the circumstellar envelope is calculated. The main gas heating is by the dust drag while cooling is done through adiabatic expansion of the gas and radiative cooling via rotational molecular line emission. We also include vibrational cooling by H₂ (Goldreich & Scoville 1976, Justtanont et al. 1994). Hence the dust and gas mass loss rates are derived in a self-consistent manner. The issue of mass loss rates is readdressed here as there is a wide range of reported rates from 310^{-5} (Neufeld et al. 1996) to 5.210^{-8} M_{\odot} yr⁻¹ (Wannier & Sahai 1986); see also Zubko & Elitzur (2000) for a recent reassessment of this controversy.

3.1. The dusty envelope

To derive the dust mass loss rate, we have fitted the SED of W Hya using ISO SWS and LWS data, together with photometry from ground based observations (Wilson et al. 1972) and the IRAS point source fluxes. We adjusted

the flux levels for the LWS spectrum to coincide with the IRAS 60μ m flux which also joins the SWS spectrum very well. There are broad features present in the LWS spectrum which are due to the internal reflection within the instrument at 54 and 110μ m. Here, we calculate the radiative transfer through the dust shell, using a code based on Haisch (1979) which takes into account thermal emission and reemission and multiple scattering. It also allows for a grain size distribution which we assumed to be the same as the interstellar grain sizes (Mathis et al. 1977, hereafter MRN). We also employ multiple grain species.

The input parameters are listed in Table 1. The distance is taken from the Hipparcos catalogue (ESA 1997). From the study by Cami (2002), the emission features between $10\text{-}20\mu\text{m}$ are a combination of amorphous silicates giving rise to the $10\mu\text{m}$ emission, compact Al_2O_3 which gives a feature at $11\mu\text{m}$, spinel (MgAl₂O₄) is responsible for the $13\mu\text{m}$ feature (Posch et al. 1999), and the $19\mu\text{m}$ emission which is attributed to Mg_{0.1}Fe_{0.9}O. In our attempt to fit these dust features, we used silicate dust properties employed by Justtanont & Tielens (1992). For other compounds, we extracted the optical constants from the University of Jena database and calculate the extinction coefficients using the Mie theory.

The fit to the SED for a three-dust component model is reasonably good for the complete SWS and LWS data coverage (Fig. 1). The fit is very satisfactory for all observed emission features, providing support for the presence of amorphous silicates, Al₂O₃ and MgFeO, except for the 13μ m feature. A model with spinel which produces a fit to the $13\mu m$ emission feature predicts an additional feature at $16\mu m$, not seen in the observation (Fig. 2). Hence, while spinel provides a good fit to the 13μ m band, this identification is suspect. Further laboratory studies on related materials are required to test this suggestion. Onaka et al. (1989) suggested aluminum oxide as a carrier for the $13\mu m$ dust emission and following that, Kozasa & Sogawa (1998) were able to explain the $13\mu m$ feature using crystalline aluminum oxide core-silicate mantle grains. However, the optical constants used for aluminum oxide were coarse (1 μ m spacing, Toon et al. 1976) and the interpolation of them close to the resonance could result in a misleading profile. A possible carrier proposed by Speck (2000) is SiO_2 which has features at 9.5, 13 and $20\mu m$ while Begemann et al. (1997) suggested a silicate-based material. For the moment, the issue of the dust species responsible for the observed $13\mu m$ feature is still unresolved.

At a distance of 115pc, we derive a total dust mass loss rate of $3\,10^{-10}~\rm M_\odot$ yr⁻¹ for the three species of silicates, ${\rm Al_2O_3}$ and MgFeO, assuming a constant outflow velocity of 6 km s⁻¹ (Bujarrabal et al. 1989). We note here that we assume that the dust and gas velocity is the same as the fitting of the SED allows us to derive $\dot{M}_d/v_{\rm dust}$. We will discuss the effect of the dust drift velocity in Sect. 3.2. Our input values are very similar to those used by Zubko & Elitzur (2000) but we derive a dust mass loss rate which is lower by a factor of ten. We investigated whether this could be due to the difference in the dust properties used.




Fig. 1. A fit to the spectral energy distribution of W Hya (solid line) with a three-dust component of amorphous silicates, Al_2O_3 and MgFeO. Open circles are from simbad (V magnitudes) and ground based observations (Wilson et al. 1972) while filled squares are IRAS PSC and crosses are SWS and LWS data. Note that the model does not account for the 13μ m emission feature.

Zubko & Elitzur used the silicate optical constants derived for AGN dust (Laor & Draine 1993), which has its infrared characteristics based upon astronomical silicates (Draine & Lee 1984). We have also analysed the observed spectra using dust parameters for the astronomical silicates and for ${\rm Mg_{0.5}Fe_{0.5}SiO_3}$ from the University of Jena database. The derived dust mass loss rates which fit the observed strength of the $10\mu{\rm m}$ dust feature agree with our value. Hence the difference in the derived dust mass loss rate is not due to the differences in the optical properties used. Rather, Zubko & Elitzur (2000) did not attempt to fit the

observed $10\mu\mathrm{m}$ feature in detail, but using only ground based-photometry and the ISO-LWS spectrum. We emphasize that the high mass loss rate of Zubko & Elitzur (2000) gives too strong 10 and $20\mu\mathrm{m}$ silicate emission as shown in Fig. 2. We find that extending the calculation to include the dust envelope size observed by IRAS at 60 and $100\mu\mathrm{m}$ (Hawkins 1990) does not affect the far-IR flux as the density is far too low in the outer part to affect the SED. For our case, we fitted the silicate $10\mu\mathrm{m}$ feature and the other dust species (except for the $13\mu\mathrm{m}$ emission feature) which givess a more reliable mass loss estimate




Fig. 2. The plot of a close-up of the dust features. The long dashed line represent a model with Zubko & Elitzur (2000) parameters. The heavy solid line is the three dust model (Fig. 1) while the heavy dashed line is a model which includes spinel. The contribution from individual dust component is shown as dotted lines. From top to bottom: spinel, amorphous silicate, Al_2O_3 , and $Mg_{0.1}Fe_{0.9}O$.

as dust mass loss rate is proportional to the strength of the $10\mu \rm m$ silicate feature. However, since we did not fit the $13\mu \rm m$ feature, our estimate of the mass loss rate should be viewed as a lower limit, pending the correct identification of the feature.

If we assume that the elemental abundance of the outflow is that of the Sun, our estimated dust mass loss rate for each dust component indicates that aluminum condenses out at a higher rate than the silicate, magnesium and iron: the latter three having abundances an order of magnitude higher than that of Al (Anders & Grevesse 1989). The amount of the silicate dust is the same as the aluminum dust while that of MgFeO dust is five times less (Table 1). Thermodynamic consideration of the dust condensation sequence starts with gas phase condensation of Al₂O₃, leading finally to Mg- and Fe-incorporation into silicate grains (Tielens 1990). Alternatively, Al and Si condense out at the same time but as Al has a high affinity for oxygen, aluminum oxide will be formed at a higher rate than silicate (Stencel et al. 1990) thereby giving rise to an $11\mu m$ feature which is as strong as the silicate feature even though the abundance of Al is much lower than that of Si. In either case, the high fraction of Al in dust as compared to Si, reflects a freeze out of the condensation process well before all the silicon has been consumed (e.g., well before magnesium-iron silicates condense out). W Hya is prototypical for a set of O-rich stars which have $10-20\mu m$ spectra which are not dominated by the wellknown broad (magnesium/iron) silicate emission features but rather by a set of oxide features (Cami 2002). We surmise that this difference reflects a steep density gradient in the dust condensation layer in these sources as compared to typical Mira variables. The presence of such a steep density gradient is also indicated by the asymmetric light curve of these types of stars.

3.2. The molecular envelope

Using the total dust mass loss rate obtained by fitting the SED which drives the gas to the terminal velocity, we obtain a dust-to-gas mass ratio of $8.7\,10^{-3}$, giving a gas mass loss rate of $3.5\,10^{-8}$ M_{\odot} yr⁻¹ (Model 1) from solving the equation of motion, assuming that the gas and dust are fully momentum coupled.

$$v\frac{dv}{dr} = (\Gamma - 1)\frac{GM_*}{r^2} \tag{1}$$

where v is the gas velocity, G is the gravitational constant, M_* is the mass of the star, and Γ is the ratio of the radiation pressure force on dust grains to the gravitational force. Following Justtanont et al. (1994), Γ can be written to include the dust size distribution, $n_d(a)$, and $Q(a, \lambda)$, the radiation pressure efficiency for a grain size, a, at wavelength λ as

$$\Gamma = \int \int \frac{\sigma_d Q(a,\lambda) L(\lambda)}{4\pi c G M_*} \frac{n_d(a)}{\rho} d\lambda \ da \eqno(2)$$

where σ_d is the geometric cross section of the dust, $L(\lambda)$ is the luminosity at the wavelength λ , ρ is the gas density (see Justtanont et al. (1994) for full details). We use radiation pressure for purely silicate dust in the calculation of the equation of motion. As a matter of fact, we find that the dust velocity is much greater than that of the gas due to the drift velocity in stars with such a low mass loss rate.

$$v_{\text{drift}}^{2}(a) = \int \frac{Q(a,\lambda)L(\lambda)v}{\dot{M}c}da$$
 (3)

The drift velocity is calculated for each grain size and then weighted by the size distribution of $a^{-3.5}$. The ratio for the drift velocity for the largest and smallest grains $(0.25 - 0.005 \mu \text{m})$ is 8.1. In our calculation, the heating is dominated by the largest grains (heating $\propto v_d^3$). This results in a high heating rate, therefore the calculated gas kinetic temperature is very high. In the fitting of the SED, we derived an M/v_{dust} in which we can ignore the drift velocity. However, if we take the drift velocity into account when calculating the dust mass loss rate and then solve for both the equations of motion and energy balance in a self consistent way, we derive a mass loss rate of $1.5 \, 10^{-7}$ $M_{\odot} \text{ yr}^{-1}$ (Model 2) and the calculated CO line fluxes are higher than those observed for the low J transitions (see Table 2). From running a number of models with various different densities and temperatures, we find that the line flux ratio of the CO 2-1/1-0 is a sensitive measure of the assumed mass loss rate while the temperature plays a

Table 1. Input parameters for modelling the dust and gas mass loss rates in W Hya

Stellar mass	$1~{ m M}_{\odot}$
Stellar effective temperature	2500 K
Stellar radius	$410^{13}~{\rm cm}$
Outflow velocity	6 km s^{-1}
Distance	115 pc
Dust condensation radius	$210^{14}~{\rm cm}$
Dust mass loss rates	$1.5 10^{-10} \mathrm{M}_{\odot} \mathrm{yr}^{-1}$ - silicates
	$1.310^{-10}~{\rm M}_{\odot}~{\rm yr}^{-1}$ - ${\rm Al_2O_3}$
	$2.510^{-11}~{\rm M}_{\odot}~{\rm yr}^{-1}$ - MgFeO
Dust-to-gas mass ratio	8.710^{-3}
Gas mass loss rate	$3.510^{-8}~{\rm M}_{\odot}~{\rm yr}^{-1}$

minor role For the observed ratio of 14.7, an upper limit for the mass loss rate is $8\,10^{-8}~{\rm M_{\odot}~yr^{-1}}$ (Model 3, with the temperature derived from solving the energy balance equation). This model gives the best description of the observed line fluxes from low to higher transitions. As a test, we calculated CO line fluxes using this mass loss rate and a simple temperature law $T \propto (r/r_i)^{-\alpha}$ where α ranges from 0.5 to 0.8. The resulting line fluxes are insensitive to the change of temperature at this low mass loss rate $(\leq 10\%)$. This mass loss rate is within a factor of two of our detailed calculation and puts a stringent upper limit on the mass loss rate for W Hya. We also calculated the expected CO line fluxes using parameters from Zubko & Elitzur (2000), including their temperature structure. The calculated line fluxes are higher than those by Barlow et al. (1996), reflecting the higher mass loss rate.

We calculate the peak main beam antenna temperatures for Model 1 to be 0.1K for CO J=1-0 observed with the IRAM telescope (cf. 0.3K from Bujarrabal et al. 1989) and 0.7K for J=2-1 observed with a clear double-peaked profile (cf. a flat-top profile with a peak temperature of 2.3K from Cernicharo et al. (1997)), with a canonical value of the CO abundance of 310^{-4} . Our calculated gas kinetic temperature, obtained from solving the energy balance, in the outer part of the shell is very high due to the large drift velocity, hence the populations of low-lying levels are excited to higher levels. This, coupled with a low density, results in very low antenna temperatures for J=1-0 and 2-1 lines. In order to reconcile these with the observations, we need to increase the mass loss rate to $8 \times 10^{-8} \,\mathrm{M}_{\odot} \,\mathrm{yr}^{-1}$ (Model 3). We also calculated expected line fluxes using input parameters from Barlow et al. (1996) and their temperature, and compared them to the IRAM results plus J=16-15 and J=17-16 in their ISO-LWS spectrum. The results of these calculations are summarized in Table 2. Our model predicts line fluxes which are lower than the observed values, which is due to the low derived mass loss rate for lower transitions while Barlow et al.'s parameters predict too high line fluxes by a factor of ten. However, we find that the high rotational lines are insensitive to the input mass loss rate. The fact that there is a combination of different types of dust, not just silicates, may lead to dif-

Table 2. A comparison of between the observed lines and those from the models using parameters by Barlow et al. (1996) and those derived in this paper. We list the available antenna temperatures $(T_{\rm MB})$, line intensities (I) and line fluxes (F) from various sources.

Transition	Observed	Barlow	Model 1	Model 2	Model 3
		et al.			
$T_{MB}(1-0)$	0.3K	4.7	0.06	0.4	0.1
I (1-0)	1.5^{a}	55.0	0.8	4.6	1.57
$T_{\rm MB}(2-1)$	2.3K	20.9	0.7	5.7	1.9
I (2-1)	22.0^{b}	229.6	7.5	51.4	19.3
F(16-15)	$3.3E-20^{c}$	8.0E-20	1.6E-20	5.9E-20	2.9E-20
F(17-16)	$2.3E-20^{c}$	8.6E-20	1.7E-20	6.2E-20	3.1E-20

- ^a CO line intensity (K km s⁻¹) from Bujarrabal et al. (1989)
- b CO line intensity (K km s⁻¹) Cernicharo et al. (1997)
- c ISO : Barlow et al. (1996) flux in W $\rm cm^{-2}$

ferent radiation pressure. A change in this parameter will affect the drift velocity and hence the heating rate and the gas kinetic temperature. Another factor which affects the CO line fluxes is the adopted turbulent velocity (Kemper et al. 2003). We assume a constant value of 1 km s⁻¹ in our calculation but in reality this could vary within the circumstellar envelope. An important issue which we touched upon in the previous section is the dust drift velocity. We calculated the drift velocity of $\sim 15 \text{ km s}^{-1}$ for the averaged radiation pressure for the grains ($\langle Q(a,\lambda) \rangle =$ 0.0057). However, the largest grains with a high drift velocity may no longer be momentum-coupled to the gas $(v_{\rm drift} \sim 100 \ {\rm km \ s^{-1}!})$. Hence these grains will not partake in the heating of the gas, and hence the temperature structure of the envelope should be steeper. Further progress in this area has to await a more detailed radiative transfer calculation that includes a realistic model for the heating and cooling.

Our value of the gas mass loss rate agrees within a factor of two to three with those from Wannier & Sahai (1986) and Olofsson et al. (2002) after scaling for the distance and velocity assumed here. However, our value falls short by an order of magnitude compared with that from Barlow et al. (1996) who calculated radiative transfer for H₂O lines in the LWS wavelength range and from Neufeld et al. (1996) who modelled H₂O lines from the SWS. The gas mass loss rate derived recently by modelling H₂O emission lines of W Hya by Zubko & Elitzur (2000) is slightly higher than that from Barlow et al. (1996). In an attempt to reconcile these differences, we can say that our assumed CO abundance of 3×10^{-4} may be too high for this object. But we need to reduce this by a factor of 30 to bring the derived mass loss rate in line with those from the water lines. However, the column density of the CO seen in absorption in SWS seems to contradict the very low abundance (Table 3). The abundance of water used in these calculations may be too low. By increasing the water abundance and lowering the mass loss rate in those cases, or by increasing our mass loss rate and lowering

the CO abundance in our case, a convergence in mass loss rate can be found between these approaches. Also, it is difficult to get the kinematic information from ISO LWS observations of H₂O lines because they are not resolved. Although SWAS and Odin observations of the 557GHz line should be resolved, currently the signal-to-noise ratio is not good enough to be confident about the line profile. In the future it will be possible to study water lines in more detail using the heterodyne instrument (HIFI) aboard the Herschel Telescope due to be launched in 2007 which has a resolution of 1 km s^{-1} or better. Equally important, the collisional cross sections between ortho- and para-H₂O and ortho- and para-H₂ for high temperature extending to high quantum numbers will be available as part of the study project in conjunction with HIFI. In the case where the lines are subthermally excited, the input collisional rates have a strong influence in the calculation of radiative transfer.

4. Modelling absorption bands

The near-IR spectrum of W Hya shows several absorption bands due to various molecules, such as CO, H₂O. In order to see these bands clearly, we divided the spectrum by our fit to the SED and normalised it at 4μ m. We have modelled these absorption bands assuming thermodynamic equilibrium. The molecular data used are taken from Le Floch (1991) for the CO molecule, Schwenke (1998, private communication) for OH and H₂O molecules and Hitran (Rothman et al., 1998) for others. The absorption line profiles are convolved with a Voigt profile with the appropriate full width at half maximum for direct comparison with the observation (see Helmich 1996). From our SED fit to the spectrum, we can define the stellar continuum satisfactorily up to $\lambda < 9\mu m$. By fitting the observed continuum-divided spectrum with our models, we can derive an excitation temperature (T_x) for each molecule, along with the column density, N, assuming a Doppler parameter, b_D . In this paper, we try fitting the different molecular bands independently and derive the best set of parameters for each band. A more sophisticated approach has been made by Cami (2002) using a chi-square technique.

We fitted, in all, six different molecules from $2.3\text{-}9\mu\text{m}$. The input parameters are listed in Table 3, while the fits are shown in Figs. 3 and 4. From the temperatures, we can deduce that these molecules originate from different layers in the circumstellar shell. We see bandheads of CO Δv =2 from v=5-3 up to v=8-6 which indicates a very hot gas. This is reflected in the input parameter for CO which has a temperature of 3000K. The column density for the band is high, 10^{21} cm⁻² also indicating that CO bands originate from a hot, dense region very close to the stellar photosphere. We believe that the gas is possibly excited by shocks since the effective temperature of the star is below the excitation temperature which gives rise to the observed v=8-6 band. This is also consistent with




Fig. 3. ISO SWS observation in band 1 (dotted line) compared with the model absorption of CO Δv =2, H₂O stretching modes and OH absorption bands (solid line). Each absorption component is also depicted separately.

the inferred Doppler width of 9 km $\rm s^{-1}$ used to model the CO absorption band.




Fig. 4. ISO SWS observation in band 2 (dotted line) compared with the model absorption of CO_2 , H_2O bending mode, SO_2 stretching mode and SiO v=1-0 band (solid line). Each absorption component is individually shown.

Table 3. Parameters for the spectral absorption calculation. For each molecular band, we also indicate the wavelength coverage.

	λ	$T_{\mathbf{x}}$	N	b_D
	$(\mu \mathrm{m})$	(K)	$({\rm cm}^{-2})$	$(\mathrm{km}\ \mathrm{s}^{-1})$
$CO \Delta v = 2$	2.3-2.6	3 000	110^{21}	9
$H_2O \nu_1 + \nu_3$	2.3 - 4.0	1000	510^{20}	7
ОН	2.7 - 4.0	1500	710^{17}	8
CO_2 ν_3	4.0 - 4.6	1000	810^{17}	7
CO v=1-0	4.2 - 5.5	2000	510^{18}	8
$H_2O \nu_2$	4.0 - 9.0	300	510^{18}	3
SO_2	7.1 - 7.7	500	210^{17}	3
SiO $v=1-0$	7.2 - 9.0	1000	110^{19}	7
CO_2 ν_2	14.8 - 15.0	1000	110^{17}	7

We also see the stretching modes of $\rm H_2O$ molecules at $3\mu\rm m$. We modelled these bands assuming an ortho to para ratio of 3. We are able to reproduce the absorption band relatively well with a gas temperature of 1000K for the $\rm H_2O$ stretching modes. However, the band is narrower than the observed one, especially in the red wing. The water band overlaps the OH ro-vibrational bands ($\Pi_{3/2}1-\Pi_{3/2}0$ and $\Pi_{1/2}1-\Pi_{1/2}0$) which cover from 2.9 to $4\mu\rm m$. The highly excited lines of OH also indicate that they come from a warm region, close to the star where $\rm H_2O$ is probably photodissociated by shocks. We also modelled the $\rm CO_2$ stretching mode at $4.25\mu\rm m$. The band seems to come from the same region as the others. The width of the band is consistent with a gas of 1000K.

The fit to the bending mode of H_2O at $6\mu m$ is, however, not consistent with the observed data. In view of the various ratios of adjacent lines, we estimate that the excitation temperature is only 300K for a column density of $5\,10^{18}$ cm⁻². The observed absorption is much deeper than calculated in the blue wing where the CO fundamental band is present. The $4.6\mu m$ CO R-branch blends with the CO₂ absorption wing while its P-branch blends with H₂O. At the resolution of the observations, individual absorption lines are not resolved. We roughly estimate the excitation temperature of the CO fundamental absorption to be $\sim 2\,000\,\mathrm{K}$ and put an upper limit for the column density of 10^{19} cm⁻². From very high resolution studies of CO lines in Mira variables (Hinkle et al. 1982), different bands come from different regions, with the fundamental band coming from a cool region while the overtone bands come from a hotter region, close to or from the pulsating photosphere. We can also fit the SiO band at $8\mu m$ with a warm gas of 1000K, possibly from the same layer as the H₂O stretching modes. This is consistent with emission coming from gas interior to the dust forming region where SiO is not bound in dust grains. There is evidence for the weak SO_2 absorption band at 7.3 μ m. This band was discovered in a few O-rich AGB stars by Yamamura et al. (1999). We also see $CO_2 \nu_2$ absorption band at 15 μ m which we will discuss in more detail in the next section.

Table 4. Line fluxes of rotational water lines detected in the ISO-SWS wavelength range.

wavelength	line flux	transition
$(\mu \mathrm{m})$	$(\mathrm{W}~\mathrm{cm}^{-2})$	
32.98	1.9E-18	$6_{61} - 5_{50}, 6_{60} - 5_{51}$
35.45	5.3E-19	$5_{33} - 4_{04}, 7_{43} - 6_{34}$
35.92	4.3E-19	$6_{52} - 5_{41}, 6_{51} - 5_{42}$
39.35	1.6E-18	$5_{50} - 4_{41}, 5_{51} - 4_{40}, 6_{42} - 5_{33}$
40.31	4.8E-19	$6_{43} - 5_{32}$
40.71	1.4E-18	$4_{32} - 3_{03}, 6_{33} - 5_{24}$
43.91	6.6E-19	$5_{41} - 4_{32}$
44.14	6.1E-19	$5_{42} - 4_{31}$
44.68	4.4E-19	$8_{36} - 7_{25}$
45.10	6.8E-19	$5_{23} - 4_{14}$

Although the overall fit to each band is good, there is still some residue which we are not currently able to account for. There is an absorption band just longward of the $4.25\mu \text{m CO}_2$ band. It is likely that the hotter component of water from the stretching modes contributes between 4.5- 6μ m. We compared our results to that from Cami (2002; Table 6.2) and found that although our derived excitation temperatures agree within a factor of two, the column densities are very different (up to two magnitudes difference) in the cases of OH and SiO, while others agree reasonably well. This may lie in the fact that we adopt different Doppler broadening for different regions, i.e., the hotter region has a higher thermal broadening than the cooler one, while Cami (2002) assumed a constant velocity of 3 km s⁻¹ throughout. As it turns out, different combinations of column density and Doppler width can give similar resulting absorption. The depth of the absorption is very sensitive to the adopted Doppler velocity used, more so than the column density.

5. Molecular emission lines

Both Barlow et al. (1996) and Neufeld et al. (1996) reported emission due to $\rm H_2O$ rotational lines from LWS and SWS FP mode observations, respectively. From the full SWS scans, we detected 10 more $\rm H_2O$ rotational lines in the region between 30-45 μ m. We estimated the line fluxes by assuming a gaussian profile and these are reported in Table 4. Some lines are a blend of the orthoand para-water as the wavelengths of their transitions are too close to be resolved by the SWS scan. Barlow et al. (1996) also noted some CO lines present in their LWS spectrum. Surprisingly, only two CO lines have been detected in LWS (J=16-15 and J=17-16, see Sect. 3). In this section, we will limit our discussion mainly to new results on $\rm CO_2$.

5.1. The SWS grating spectrum of CO₂

We detected CO_2 in W Hya between $13-16\mu m$ from the full scan SWS AOT1. We see these CO_2 bands in all of






Fig. 6. SWS-FP observations of CO₂ in W Hya with the wavelength corrected for the heliocentric velocity. The top panels show the partly resolved Q-branches of the 13.87 and 16.18 μ m bands and the bottom panel shows the 14.97 μ m. band. Flux units are 10⁻¹¹ W m⁻² μ m⁻¹. Vertical lines indicate positions of Q lines. The first few Q lines are not resolved.

the stars which exhibit the so called $13\mu m$ dust feature in our guaranteed time program (PI. de Jong). From this we proposed that objects with the $13\mu m$ dust have CO_2 associated with it (see Justtanont et al. 1998, Fig. 1). On this basis, we obtained further observing time to scan objects with this dust feature at the highest grating resolution (AOT6). All the objects observed in the supplementary program confirm our idea, i.e., all stars with the $13\mu m$ dust show CO_2 bands. However, the reverse is not always the case, e.g., one of the stars in our program, o Ceti, has

no $13\mu\mathrm{m}$ dust but shows CO_2 emission and IRC+10011 also shows CO_2 while the $10\mu\mathrm{m}$ silicate is self-absorbed (Markwick & Millar 2000). Cami et al. (2000) show a detailed calculation of CO_2 emission bands in EP Aqr, one of the stars in our program. If the association between the $13\mu\mathrm{m}$ dust feature and CO_2 gas is real then CO_2 is very prevalent because Sloan et al. (1996) argue that up to 50% of O-rich semiregular variables show such a dust feature. The formation of CO_2 is thought to be via $\mathrm{CO}+\mathrm{OH}$ (Nercessian et al. 1989; Willacy & Millar 1997) where OH





is the product of either photodissociation or of shock destruction of ${\rm H_2O}$ molecules.

In W Hya, the transitions $10^{0}0-01^{1}0$ at $13.87\mu m$ and $02^{0}0-01^{1}0$ at $16.18\mu m$ are in emission while $01^{1}0-00^{0}0$ at $14.97\mu m$ is in absorption. We modelled the absorption band which is shifted from the normal position as seen in emission in other stars. We attribute this shift towards a shorter wavelength to the importance of highly excited bands. To excite these bands (e.g., 02^20-01^10 , 03^30-02^20 , 04^40-03^30), we need the gas temperature to be ~ 1000 K (Fig. 5). These bands have their Q-branches at successively shorter wavelengths than the $01^{1}0-00^{0}0$ band. By including these bands in our calculation, we are able to match the peak absorption at $14.957\mu m$ as observed. One thing to note from our observations of objects with CO₂ is that only stars with the $14.97\mu m$ absorption show the corresponding $4.25\mu m$ absorption. Those which show the $14.97\mu m$ in emission lack the stretching mode counterpart (for the full sample, see Cami (2002)). This probably indicates that the radiative pumping by the $4.25\mu m$ band plays a very minor role, if at all, in populating the levels which result in the 13-16 μ m emission.

5.2. The Fabry-Perot observation of CO₂ bands

We subsequently obtained high resolution spectra of these CO_2 bands using the SWS-FP which resolved the individual Q branches (Fig. 6). The small dip at $13.877\mu m$ in the band 10^00-01^10 in this figure is an instrumental effect due to the end of the scan of FP wavelength setting. The re-




Fig. 7. The rotation diagram of CO_2 . Crosses are from the 01^10-00^00 , triangles represent the 02^00-01^10 and squares represent 10^00-01^10 . The solid lines are the best fit for these bands.

solved Q-lines are listed in Table 5 where the wavelength, the energy of the upper level, the measured line flux and the identification are given. The flux errors given represent the estimated uncertainties in the continuum and do not take into account possible calibration errors so that they should be taken as lower limits.

It is quite remarkable though that we can discern emission of the Q-branch lines in the depth of the $14.97\mu m$ band while the absorption blends into a single band. As discussed in Sect. 5.1, this absorption is formed in a hot $(1\,000\mathrm{K})$ gas layer in the outer photosphere and lines may be smeared out by turbulence $(b_D=7\,\mathrm{km~s^{-1}},\mathrm{see~Table~3})$ and/or optical depth effects. The first of the individual Q-line to be resolved in emission is Q10. The close spacing between Q2-Q8 means that even at this resolution we cannot separate them from each other. We conclude that the emission lines are produced in a gas layer which is more extended and lying farther out in the circumstellar shell than the hot layer which is responsible for the absorption band at $14.95\text{-}14.98\mu\text{m}$, in line with earlier suggestions by Justtanont et al. (1998) and Cami et al. (2000).

We have Fabry-Perot observations of individual Q-branch lines for three bands of CO_2 (see Fig.6). In the fundamental band we detected lines up to Q42 and in the other bands up to Q24. We can use the observed flux of the individual line in each band to construct a rotation diagram from which we can derive a rotational excitation temperature $\mathrm{T}_{\mathrm{rot}}$ characterizing the population distribution within the upper vibrational level of each band, using

Table 5. A list of the CO₂ emission lines detected from **Table 6.** Quantities derived from the rotation diagram the FP observations.

wavelength	\mathbf{E}_{upper}	line flux	transition
$(\mu \mathrm{m})$	(K)	$(10^{-19} \mathrm{W} \ \mathrm{cm}^{-2})$	
13.87564	2059.0606	$2.6 {\pm} 0.5$	$10^{0}0-01^{1}0 \text{ Q}10$
13.87658	2084.8831	1.0 ± 0.5	$10^{0}0-01^{1}0 \text{ Q}12$
13.87768	2115.1954	2.2 ± 0.2	$10^{0}0-01^{1}0 \text{ Q}14$
13.87894	2149.9974	2.7 ± 0.2	$10^{0}0-01^{1}0 \text{ Q}16$
13.88036	2189.2882	1.9 ± 0.2	$10^{0}0-01^{1}0 \text{ Q}18$
13.88194	2233.0677	5.3 ± 1.9	$10^{0}0-01^{1}0 \text{ Q}20$
13.88367	2281.3351	3.1 ± 1.5	$10^{0}0\text{-}01^{1}0 \text{ Q}22$
13.88556	2334.0896	4.3 ± 1.1	$10^{0}0\text{-}01^{1}0 \text{ Q}24$
13.88760	2391.3305	$1.6 {\pm} 0.5$	$10^{0}0\text{-}01^{1}0 \text{ Q}26$
14.94229	1976.2415	1.6 ± 0.3	$01^{1}0\text{-}00^{0}0 \text{ Q}42$
14.94609	1882.9064	$2.8 {\pm} 0.5$	$01^{1}0-00^{0}0 \text{ Q}40$
14.94972	1794.0594	2.1 ± 0.3	$01^10-00^00 \text{ Q}38$
14.95317	1709.7018	1.5 ± 0.3	$01^10-00^00 \text{ Q}36$
14.95643	1629.8354	0.9 ± 0.4	$01^{1}0\text{-}00^{0}0 \text{ Q}34$
14.95952	1554.4611	1.0 ± 0.3	$01^{1}0\text{-}00^{0}0 \text{ Q}32$
14.96243	1483.5803	1.9 ± 0.5	$01^{1}0\text{-}00^{0}0 \text{ Q}30$
14.96515	1417.1943	$2.1 {\pm} 0.4$	$01^10\text{-}00^00 \text{ Q}28$
14.96769	1355.3043	2.1 ± 0.5	$01^{1}0\text{-}00^{0}0 \text{ Q}26$
14.97005	1297.9109	1.2 ± 0.4	$01^{1}0\text{-}00^{0}0 \text{ Q}24$
14.97223	1245.0155	1.2 ± 0.4	$01^{1}0-00^{0}0 \text{ Q}22$
14.97422	1196.6187	1.3 ± 0.5	$01^{1}0-00^{0}0 \text{ Q}20$
14.97603	1152.7216	$2.1 {\pm} 0.7$	$01^10-00^00 \text{ Q}18$
14.97765	1113.3246	$2.6 {\pm} 0.7$	$01^{1}0-00^{0}0 \text{ Q}16$
14.97909	1078.4285	3.8 ± 0.3	$01^{1}0-00^{0}0 \text{ Q}14$
14.98034	1048.0339	$2.2 {\pm} 0.5$	$01^{1}0-00^{0}0 \text{ Q}12$
14.98141	1022.1414	2.7 ± 0.7	$01^{1}0-00^{0}0 \text{ Q}10$
16.18366	1937.0742	4.2 ± 0.2	$02^{0}0-01^{1}0 \text{ Q}12$
16.18476	1967.4081	2.2 ± 0.2	$02^{0}0-01^{1}0 \text{ Q}14$
16.18603	2002.2345	1.7 ± 0.2	$02^{0}0-01^{1}0 \text{ Q}16$
16.18747	2041.5522	$2.5 {\pm} 0.2$	$02^{0}0-01^{1}0 \text{ Q}18$
16.18908	2085.3610	6.0 ± 2.0	$02^{0}0-01^{1}0 \text{ Q}20$
16.19087	2133.6598	$4.1 {\pm} 0.5$	$02^{0}0-01^{1}0 \text{ Q}22$
16.19283	2186.4477	$5.6 {\pm} 0.1$	$02^{0}0-01^{1}0 \text{ Q}24$

the relation

$$E_j/kT_{\rm rot} = ln(N_v/\Phi(T_{\rm rot})d^2) - ln(4\pi F_{ji}/A_{ji}h\nu_{ji}g_j)$$

$$\equiv ln(N_v/\Phi(T_{\rm rot})d^2) - Y_j$$
 (4)

where E_j and g_j are the energy and statistical weight of the upper state, F_{ji} and ν_{ji} are the line flux and frequency of the transition, A_{ji} is the spontaneous transition probability, N_v is the total number of molecules in the upper vibrational level v, $\Phi(T_{\rm rot})$ is the partition function, and d is the distance to W Hya. Fig. 7 shows the rotation diagram of all the resolved Q-lines. There is considerable scatter in the data points due to uncertainties in the calibration (e.g., joining of different pieces of the spectrum within each band) and in the continuum. The offset between the data points in the 01^{10} level and in the 02^{00} and $10^{0}0$ levels is due to differences in the number of molecules in these levels.

In Table 6, we collect a number of quantities derived from the fits in Fig. 7. The derived rotation temperatures are 435 K for the 01¹0 level and 460 K for the other two

Level	T _{rot} (K)	$\Phi(T_{\rm rot})$	Y_0	N_v
$10^{0}0$	460	4.73	-7.57	3.010^{38}
$02^{0}0$	460	6.56	-7.50	4.210^{38}
$01^{1}0$	435	47.2	-3.20	2.410^{41}

levels combined. The partition functions for each vibrational level are computed assuming that the levels are populated in thermal equilibrium at the rotational temperature. The quantities Y_0 are the intercepts of the fits in Fig. 7 at $E_j = 0$. Putting $E_j = 0$ and $Y_j = Y_0$ in Eq. (3) we derive values of N_v , the total number of molecules in each vibrational level, listed in column 5 of Table 6. These numbers are a lower limit as we assumed that the lines are optically thin.

From these, we then find the vibrational excitation temperatures T_{vib} (10⁰0-01¹0) = 150 K and T_{vib} $(02^{0}0-01^{1}0) = 170 \text{ K}$, using the relation $T_{\rm vib} = 1.44$ $E_{ij}/\ln(N_j/N_i)$ with $E(10^00-01^10) = 720 \text{ cm}^{-1}$ and $E(02^00-01^10) = 720 \text{ cm}^{-1}$ $01^{1}0) = 617 \text{ cm}^{-1}$ and taking into account that the number of rotational levels in level 01¹0 is twice as large (J = even and odd) as in levels $10^{0}0$ and $02^{0}0$ (J = even only). Assuming that the relative number of molecules in the ground vibrational state 00⁰0 and the first excited vibrational state 01^{10} is also characterized by $T_{\rm vib} \sim 160~{\rm K}$ we find a total number of CO_2 molecules of 5×10^{43} .

Clearly the temperature characterizing the distribution of rotational levels in the vibrational states of CO₂ $(\sim 450 \text{ K})$ is different from the temperature characterizing the distribution of molecules over the vibrational states $(\sim 160~{\rm K})$ so that the ${\rm CO_2}$ molecules are close to, but not in thermal equilibrium. This non-LTE behaviour can be understood on the basis of the symmetry of the CO₂ molecule which does not allow pure rotational radiative transitions and the magnitudes of the cross sections for (de-)excitation of CO_2 by collisions with H_2 , which are about three orders of magnitude larger for pure rotational transitions than for ro-vibrational transitions (Banks & Clary 1987). It can be shown by more detailed non-LTE multi-level line transfer calculations (de Jong, in preparation) that at a few stellar radii from the star and at H₂ densities above $\sim 10^8 \ \mathrm{cm}^{-3}$ the rotational levels are more strongly coupled to the gas kinetic temperature ($\sim 400 \text{ K}$) while the population of the vibrational levels is dominated by radiation trapping in the ro-vibrational lines (the ones that we observe) resulting in a vibrational temperature somewhat below the kinetic temperature.

We suggest that CO₂ absorption and emission originate in two different regions of an extended dynamical atmosphere: one close to the star in the outer photosphere where the temperature is 1000K, giving rise to the absorption seen at 4.25 and 14.95 μ m, and the other farther out at few stellar radii, where the density is still high enough to thermally distribute the pure rotational levels at ~ 450 K. Assuming that the cooler CO_2 layer is located at ~ 3 stellar radii we find, from the total number of molecules observed a column density $N(CO_2) \sim 10^{15}$ two orders of magnitude smaller than that derived for the hot CO_2 gas (see Table 3) and also several orders of magnitude smaller than found for the warm gas in the star EP Aqr by Cami et al. (2000)

As proposed earlier by Tsuji et al. (1997), Yamamura et al. (1999), and Cami et al. (2000), we believe that the emission line gas detected here is part of the warm gas layer located between the stellar photosphere and the dust condensation zone. This gas is probably related to the blown-up atmosphere observed by Reid & Menten (1990) and found by Höfner et al. (1998) in their calculations of the dynamical atmospheres of pulsating stars. How CO₂ molecules are formed in this gas is unclear but they may well be a left-over from the preceding photosphere phase. According to Höfner et al. (1998), this gas is expected to have turbulent velocities of order 1 km s^{-1} . Eventually a little further out (at ~ 5 stellar radii) the dust is formed which then accelerates the flow to 5-10 km $\rm s^{-1}$. The CO₂ is probably quickly photodissociated in the expanding envelope. We note here that the line widths of the emission are much narrower than those of the absorption, suggesting that the emission comes from a cooler (since we do not see higher overtone bands in emission), much less turbulent layer than that which produces the absorption. Whether the hot and cool layers of CO₂ are manifestations of the same gas needs to be investigated further in a detailed model for CO_2 excitation.

6. Summary

By fitting the SED of W Hya from the near- to far-IR, we derive a dust mass loss rate of $310^{-10} \text{ M}_{\odot} \text{ yr}^{-1}$ using a combination of three different dust species of amorphous silicate, Al₂O₃ and MgFeO. We also tried adding a fourth component of spinel to account for the $13\mu m$ dust feature but in order to fit it, we also expect to see another strong peak at $16\mu m$ which is not present in the observed spectrum. From this, we conclude that spinel cannot be the carrier of the $13\mu m$ dust emission feature. Hence, the debate on the carrier of the $13\mu m$ feature is still open. With the estimated value of the our dust mass loss rate, we get a gas mass loss rate range of $(3.5-8) \, 10^{-8} \, \mathrm{M}_{\odot} \, \mathrm{yr}^{-1}$ from simple consideration of momentum dust driven wind. Our derived mass loss rate agrees well with those previously derived to explain the CO emission, taking into account the differences in the adopted distance and velocity. However, the discrepancy still exists between these mass loss rates and those derived from the H₂O emission, the latter being consistently higher. The difference in derived mass loss rates from CO and H₂O may simply be due to the abundances assumed for each species.

We obtained the full scan SWS spectrum of W Hya in which we modelled the absorption bands due to various molecules. The derived excitation temperatures of different bands suggest that they originate from different layers of gas, from $300 \mathrm{K}$ to $3000 \mathrm{K}$. However, some molecules ex-

ist over a wide range of temperatures and cannot really be stratified within a particular zone, particularly those which show multiple absorption bands. The exception may be OH which is a product of water molecules which are dissociated by shocks in the inner part of the envelope.

We also observed CO_2 bands in emission using SWS FP and were able to resolve the three strongest Q branches in the mid-IR. The resulting rotation diagram yields an excitation temperature of $\sim 450 \mathrm{K}$ while the CO_2 absorption at $14.97 \mu \mathrm{m}$ requires a gas of $1\,000 \mathrm{K}$ to explain the shift of the peak absorption bluewards. The stretching mode at $4.25 \mu \mathrm{m}$ also requires a layer of hot gas to explain its width. We find from simple calculation based on the rotation diagram that the vibrationally excited levels may not be in LTE. Whether the CO_2 absorption and emission lines can indeed be explained by temperature stratification in an extended dynamical atmosphere needs to be investigated in a more detailed radiation transfer model.

Acknowledgements. We are grateful to our referee, Dr. J Alcolea for constructive comments which improves the discussion in our paper. This research has made use of the SIMBAD database, operated at CDS Strasbourg, France.

References

Anders E., & Grevesse N., 1989, Geochimica et Cosmochimica Acta 53, 197

Banks A.J., & Clary D.C., 1987, J. Chem. Phys. 86, 802

Barlow M.J., Nguyen-Q-Rieu, Truong-Bach et al., 1996, A&A 315, L241

Bedijn P.J., 1987, A&A 186, 136

Begemann B., Dorschner J., Henning T., et al., 1997, A&A 476, 199

Bujarrabal V., Gómez-González J., & Planesas P., 1989, A&A 219, 256

Cami J., PhD thesis, University of Leiden

 Cami J., Yamamura I., de Jong T., et al., 2000, A&A 360, 562
 Cernicharo J., Alcolea J., Baudry A., Gonz'alez-Alfonso E., 1997, A&A 319, 607

Chapman J.M., Sivagnanam P., Cohen R.J., & Le Squeren A.M., 1994, MNRAS 268, 475

Clark F.O., Troland T.H., & Miller J.S., 1985, ApJ 289, 756
Clegg P., Ade P.A.R., Armand C., et al., 1996, A&A 315, L38
de Graauw T., Haser L.N., Beintema D.A., et al., 1996 A&A 315, L49

Draine B.T., & Lee H.M., 1984, ApJ 285, 89

ESA, 1997, the Hipparcos and Tycho Catalogues, ESA SP-1200 Gilman R.C., 1972 ApJ 178, 423

Goldreich P., & Scoville N., 1976, ApJ 205, 144

Groenewegen M.A.T., 1994, A&A 290, 531

Haisch B.M., 1979, A&A 72, 161

Harwit M., & Bergin E.A., 2002, ApJ 565, L105

Hawkins G.W., 2000, A&A 229, L5

Helmich F.P., 1996, PhD thesis (Leiden)

Hinkle K.H., Hall D.N.B., & Ridgway S.T., 1982 ApJ 252, 697
 Höfner S., Jørgensen U.G., Loidl R., Aringer B., 1998, A&A 340, 497

Justtanont K., Feuchtgruber H., de Jong T., et al., 1998, A&A 330, L17 Justtanont K., Skinner C.J., & Tielens A.G.G.M., 1994, ApJ 435, 852

Justtanont K., & Tielens A.G.G.M., 1992, ApJ 389, 400

Kemper F., et al., 2003, A&A, submitted

Kessler M.F., Steinz J.A., Anderegg M.E., et al., 1996, A&A 315, L27

Knapp G.R., & Morris M., 1985, ApJ 292, 640

Kozasa T., & Sogawa H., 1998, Ap&SS 255, 437

Lane A.P., Johnston K.J., Bowers P.F., Spencer J.H., & Diamond P.J., 1987, ApJ 323, 756

Laor A., & Draine B.T., 1993, ApJ 402, 441

Le Floch A., 1991, Mol Phys 72, 133

Lebzelter T., Kiss L.L., Hinkle K.H., 2000 A&A 361, 167

Markwick A.J., & Millar T.J., 2000, A&A 359, 1162

Mathis J.S., Rumple W., & Norsieck K.H., 1977, ApJ 217, 425

Nercessian E., Guilloteau S., Omont A., & Benayoun J.J., 1989 A&A 210, 225

Neufeld D.A., Chen W., Melnick G.J. et al., 1996, A&A 315, L237

Olofsson H., Gonzales-Delgado D., Kerschbaum F., Schöier F.L., 2002, A&A 391, 1053

Onaka T., de Jong T., & Willem F.J., 1989, A&A 218, 169

Posch T., Kerschbaum F., Mutschke H., et al., 1999, A&A 352, 609

Reid M.J., Menten K.M., 1990, ApJ 360, L51

Rothman L.S., Rinsland C.P., Goldman S.T., et al., 1998, J Quantitative Spectroscopy and Radiative Transfer 68, 665

Ryde N., Eriksson K., & Gustafsson B., 1999, A&A 341, 579

Schwartz P.R., Zuckerman B., & Bologna J.M., 1982, ApJ 256, L55

Skinner C.J., Justtanont K., Tielens A.G.G.M., Betz A.L., & Boreiko R.T., 1999, MNRAS 302, 296

Sloan G.C., LeVan P.D., & Little-Merenin I.R., 1996, ApJ 463, 310

Speck A.K., Barlow M.J., Sylvester R.J., & Hofmeister A.M., 2000, A&AS 146, 437

Spencer J.H., Johnston K.J., Moran J.M., et al., 1979, ApJ 230, 449

Stencel R.E., Nuth J.A., Little-Marenin I.R., & Little S.J., 1990, ApJ 350, L45

Szymczak M., Cohen R.J., & Richards A.M.S., 1998, MNRAS 297, 1151

Tielens A.G.G.M., 1990, in From Miras to Planetary Nebulae: which path for stellar evolution?, eds, Mennessier M.O., Omomt A., France: Editions Frontières, p. 186

Toon O.B., Pollack J.B., & Khare B.N., 1976, J. Geophys. Res. 81, 5733

Tsuji T., Ohnaka K., Yamamura I., 1997, A&A 320, L1

Wannier P.G., & Sahai R., 1986, ApJ 311, 335

Willacy K., & Millar T.J., 1997, A&A 324, 237

Wilson W.J., Schwartz P.R., Neugebauer G., Harvey P.M., & Becklin E.E., 1972, ApJ 177, 523

Yamamura I., de Jong T., Onaka T., Cami J., & Waters L.B.F.M., 1999, A&A 341, L9

Zubko V., & Elitzur M., 2000, ApJ 554, L137