

Evolution of Neutral Gas at High Redshift – Implications for the Epoch of Galaxy Formation

L. J. Storrie-Lombardi^{1,*}, R. G. McMahon¹, & M. J. Irwin²

¹*Institute of Astronomy, Madingley Road, Cambridge CB3 0HA*

²*Royal Greenwich Observatory, Madingley Road, Cambridge CB3 0EZ*

* *current address: Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101 USA*
email: lisa@ociw.edu (LJSL); rgm@ast.cam.ac.uk (RGM); mike@ast.cam.ac.uk (MJI)

28 November 2017

ABSTRACT

Though observationally rare, damped Ly α absorption systems dominate the mass density of neutral gas in the Universe. Eleven high redshift damped Ly α systems covering $2.8 < z < 4.4$ were discovered in 26 QSOs from the APM $z > 4$ QSO Survey, extending these absorption system surveys to the highest redshifts currently possible. Combining our new data set with previous surveys we find that the cosmological mass density in neutral gas, Ω_g , does not rise as steeply prior to $z \sim 2$ as indicated by previous studies. There is evidence in the observed Ω_g for a flattening at $z \sim 2$ and a possible turnover at $z \sim 3$. When combined with the decline at $z > 3.5$ in number density per unit redshift of damped systems with column densities $\log N_{\text{HI}} \geq 21$ atoms cm^{-2} , these results point to an epoch at $z \gtrsim 3$ prior to which the highest column density damped systems are still forming. We find that over the redshift range $2 < z < 4$ the total mass in neutral gas is marginally comparable with the total visible mass in stars in present day galaxies. However, if one considers the total mass visible in stellar disks alone, i.e. excluding galactic bulges, the two values are comparable. We are observing a mass of neutral gas comparable to the mass of visible disk stars. Lanzetta, Wolfe & Turnshek found that $\Omega(z \approx 3.5)$ was twice $\Omega(z \approx 2)$, implying a much larger amount of star formation must have taken place between $z=3.5$ and $z=2$ than is indicated by metallicity studies. This created a ‘cosmic G-dwarf problem’. The more gradual evolution of Ω_g we find alleviates this. These results have profound implications for theories of galaxy formation.

Key words: cosmology—galaxies: evolution—galaxies: formation—quasars: absorption lines

1 INTRODUCTION

While the baryonic content of spiral galaxies that are observed in the present epoch is concentrated in stars, in the past this must have been in the form of gas. The principal gaseous component in spiral galaxies is HI which has led to surveys for absorption systems detected by the damped lines they produce (Wolfe et al. 1986 [WTSC]; Lanzetta et al. 1991 [LWTLMH]; Lanzetta, Wolfe, & Turnshek 1995 [LWT], Wolfe et al. 1995). Damped Ly α absorption systems have neutral hydrogen column densities of $N_{\text{HI}} > 2 \times 10^{20}$ atoms cm^{-2} and they dominate the baryonic mass contributed by HI. We extend the earlier work on damped Ly α systems to higher redshifts using twenty-six QSOs from the APM Damped Ly α Survey (Storrie-Lombardi et al. 1996 [SMIH], Storrie-Lombardi, Irwin & McMahon 1996 [SIM]), with eleven candidate or confirmed damped Ly α absorption

systems covering the redshift range $2.8 \leq z \leq 4.4$ (8 with $z > 3.5$). These data more than triple the redshift path surveyed at $z > 3$ and allow the first systematic study up to $z = 4.7$.

2 EVOLUTION OF Ω_G – BARYONS IN NEUTRAL GAS

The mean cosmological mass density contributed by Ly α absorbers can be estimated as

$$\langle \Omega_g \rangle = \frac{H_0 \mu m_H}{c \rho_{\text{crit}}} \int_{N_{\text{min}}}^{\infty} N f(N, z) dN \quad (1)$$

giving the current mass density in units of the current critical density (LWTLMH). μ is the mean molecular weight of the gas which is taken to be 1.3 (75% H and 25% He by

arXiv:astro-ph/9608147v1 22 Aug 1996

mass), m_{H} is the mass of the hydrogen atom, ρ_{crit} is the current critical mass density, N_{min} is the low end of the HI column density range being investigated, and $f(N, z)$ is the column density distribution function. Unfortunately $f(N, z)$ is not a simple function and its evolution with redshift is difficult to accurately quantify (LWT, SIM). The integral in equation 1 can be estimated using

$$\int_{N_{\text{min}}}^{\infty} N f(N, z) dN = \frac{\sum_i N_i(\text{HI})}{\Delta X}, \quad (2)$$

where ΔX is the absorption distance interval. The absorption distance X is used to remove the redshift dependence in the sample and put everything on a comoving coordinate scale. If the population of absorbers is nonevolving (i.e. the product of their space density multiplied by their cross-section does not change with redshift) they have a constant number density per unit absorption distance. In a standard Friedmann universe X is defined as

$$X(z) = \begin{cases} \frac{2}{3}[(1+z)^{3/2} - 1] & \text{if } q_0 = 0.5; \\ \frac{1}{2}[(1+z)^2 - 1] & \text{if } q_0 = 0. \end{cases} \quad (3)$$

(Bahcall & Peebles 1969; cf. Tytler 1987). The errors in Ω_g are also difficult to estimate without knowing $f(N, z)$. LWTLMH used the standard error in the distribution of N_{HI} which yields zero error if all the column densities in a bin are the same. We have estimated the fractional variance in Ω_g by comparing the observed distribution of $f(N, z)$ with the equivalent Poisson sampling process. This gives

$$\left(\frac{\Delta\Omega_g}{\Omega_g}\right)^2 = \sum_{i=1}^p N_i^2 / \left(\sum_{i=1}^p N_i\right)^2 \quad (4)$$

and $1/\sqrt{p}$ fractional errors if all the column densities included in a bin are equal. To address uncertainties in $f(N, z)$ we also calculated a maximum likelihood estimate of the errors in the HI column density. We used the power law with an exponential turnover form of the column density distribution function, i.e. the gamma-distribution from SIM

$$f(N, z) = (f_*/N_*) (N/N_*)^{-\beta} e^{-N/N_*}, \quad (5)$$

with $\log N_* = 21.63 \pm 0.35$, $\beta = 1.48 \pm 0.30$, and $f_* = 1.77 \times 10^{-2}$. Unlike a pure power law this form has a finite integral mass. The maximum-likelihood estimates of the errors agree well with the fractional variance.

LWT found that Ω_g inferred from studies of damped systems rises with increasing redshift for $0.008 < z < 3.5$. For the range $3.0 < z < 3.5$ which included 4 damped systems and was the the highest redshift bin in the study, $\Omega(z \approx 3.5)$ was twice $\Omega(z \approx 2)$. This implied a much larger amount of star formation must have taken place between $z=3.5$ and $z=2$ than is indicated by metallicity studies. Specifically, Pettini et al. (1994) measured a low mean metallicity in damped Ly α systems at $z \approx 2.2$ inferring that they are observed prior to the bulk of star formation in the disk. The LWT result also implied that the bulk of stars in nearby galaxies should be metal poor whereas only a small fraction of disk stars in the solar neighbourhood are metal poor. This result presented a ‘cosmic G-dwarf problem’ similar to the ‘G-dwarf problem’ described in Schmidt (1963) that comes about if bright and faint stars formed at the same rate and there was no accretion on to the disk of the Galaxy. They concluded that the characteristic epoch of metal produc-

tion in galaxies occurred after the characteristic epoch of star formation and that damped Ly α absorbers might trace disk as well as spheroid evolution. Fall & Pei (1993; and Pei & Fall 1995) have argued that obscuration caused by dust in damped Ly α systems could lead to significant underestimates of the neutral gas fraction at all redshifts, particularly in the range $1 \leq z \leq 2$, also possibly explaining these results.

To further investigate these issues, the high redshift confirmed and candidate damped Ly α systems discovered in the APM Damped Ly α Survey (SMIH, SIM) have been combined with lower redshift samples (WTSC, LWTLMH, LWT) to study the evolution of Ω_g over the range $0.008 < z < 4.7$. The combined data set includes 44 absorbers in 366 quasars. The results are tabulated in table 1 and shown in figure 1(a) for $q_0=0$ and 1(b) $q_0=0.5$ ($H_0=50$) following the format presented in Wolfe et al. (1995). The solid bins include the entire data set and the dashed bins exclude the new high redshift APM data*. The region Ω_{star} is the $\pm 1\sigma$ range for the mass density in stars in nearby galaxies (Gnedin & Ostriker 1992). The point at $z=0$ is the value inferred from 21 cm emission from local galaxies (Fall & Pei 1993; Rao & Briggs 1993). The most striking result is Ω_g does not rise as steeply prior to $z \sim 2$ as indicated by previous studies. There is now evidence for a flattening in Ω_g at $z \sim 2$ and a possible turnover at $z \sim 3$. This result, combined with the decline at $z > 3.5$ in number density per unit redshift of damped systems with column densities $\log N_{\text{HI}} \geq 21$ atoms cm^{-2} (SIM) points to an epoch at $z \gtrsim 3$ prior to which the largest damped systems are still forming. The decrease in number density at high redshift of the highest column density absorbers is in marked contrast to the more numerous lower column density systems, e.g. Lyman-limit systems ($N_{\text{HI}} \sim 10^{18}$ atoms cm^{-2}) (Storrie-Lombardi et al. 1994; Stengler-Larrea et al. 1995) and Ly α forest absorbers ($N_{\text{HI}} \sim 10^{13} - 10^{15}$ cm^{-2}) (Williger et al. 1994).

The inclusion of the APM survey data for $z > 3$ reduces significantly the value previously found for Ω_g in the bin $3 < z < 3.5$. Only one damped system is added to the existing four in this redshift range, but the absorption distance is doubled, which comes in to the calculation of Ω_g in the denominator in equation 2 (see table 2). The additional redshift path added by the APM survey is shown graphically by the sensitivity function in figure 6 of SMIH. We find that over the redshift range $2 < z < 4$ the total mass in neutral gas (Ω_g) is marginally comparable with the total visible mass in stars in present day galaxies (Ω_{star}) for $q_0=0.5$. However, if one considers the total mass visible in stellar disks alone, i.e. excluding galactic bulges, the two values are comparable. Using the result from Schechter & Dressler (1987) that galactic disks and bulges contribute equally to the mass density of the Universe, we are observing a mass of neutral gas comparable to the mass of visible disk stars, i.e. $\Omega_g \sim \Omega_{\text{disk-stars}}$. We note that the uncertainty in the total mass in visible stars in the local Universe is comparable with our estimates of the mass in neutral gas at $z > 2$. Given this, and the fact that we do not know if damped systems are the precursors to galactic disks, bulges, or both, these results are difficult to interpret. If we make a plausible correction

* Excluding the new APM data effectively yields the data set analyzed in LWT.

for obscuration by dust as advocated by Pei & Fall (1995) the Ω_g points shown in figure 1(b) would migrate to the positions of the open circles shown in figure 2. More work is needed to determine the severity of dust obscuration in optically selected QSO surveys. An estimated 20% correction for the neutral gas not in damped Ly α systems is shown by the open squares.

3 IMPLICATIONS FOR GALAXY FORMATION THEORIES

The shape of the Ω_g curve has been used by numerous authors to constrain theories of galaxy formation and cosmological models (Klypin et al. 1995; Kauffmann & Charlot 1994; Ma & Bertschinger 1994; Mo & Miralde-Escudé 1994). They have found that cold+hot dark matter (CHDM) models are incompatible with the previous results from the damped Ly α systems at $z \sim 3$ as they predict too few galactic halos. These models need to be reevaluated now that the value of $\Omega_g(z=3)$ indicated from our new observations has changed significantly. For example, (Klypin et al. 1995) show a peak in Ω_g at $z=3$ for various CHDM models. Though the error bars on Ω_g are still far too large to accurately discriminate between details of cosmological models, the overall shape has implications for structure formation.

4 SUMMARY

Combining the new data from the APM high redshift survey extends studies of the cosmological mass density in neutral gas, Ω_g , to $z=4.7$. We find evidence for a flattening in Ω_g at $z \sim 2$ and a possible turnover at $z \sim 3$. Though the turnover is not formally significant, when combined with the decline at $z > 3.5$ in number density of damped systems with column densities $\log N_{\text{HI}} \geq 21$ atoms cm^{-2} , these results point to an epoch prior to which the largest damped systems are still forming.

Previous studies indicated that $\Omega(z \approx 3.5)$ was twice $\Omega(z \approx 2)$, implying a much larger amount of star formation must have taken place between $z=3.5$ and $z=2$ than is indicated by metallicity studies. The more gradual evolution we find in Ω_g alleviates this problem. The results are consistent with observations of damped Ly α systems at $z \approx 2.2$ that show a low mean metallicity, suggesting that they are observed prior to the bulk of star formation in the disk. We have also made an estimated correction to Ω_g to account for bias in optically selected quasar samples due to possible obscuration by dust in foreground absorbers. Theories of galaxy formation and constraints on cosmological models utilising Ω_g should be reevaluated in light of the new observational results presented here. The error bars are still very large at high redshift and to differentiate between a peak versus a flattening in the Ω_g curve, larger samples of bright $z > 3.5$ quasars are needed to discover damped Ly α systems with $z > 3$.

Acknowledgments

We acknowledge fruitful discussions with Art Wolfe, Max Pettini and Mike Fall. RGM thanks the Royal Society for support. LSL acknowledges support from an Isaac

Newton Studentship, the Cambridge Overseas Trust, and a University of California President's Postdoctoral Fellowship. We thank the PATT for time awarded to do the observations with the William Herschel Telescope that made this work possible. We thank the referee, Ken Lanzetta, for his comments.

REFERENCES

- Bahcall J.N., Peebles P.J.E. 1969, ApJ, 156, L7
 Fall, S.M. & Pei, Y.C. 1993, ApJ, 402, 479
 Gnedin, N.Y. & Ostriker, J.P. 1992, ApJ, 400, 1
 Kauffmann, G., Charlot, S. 1994, ApJ, 430, L97
 Klypin, A., Borgani, S., Holtzman, J., Primack, J. 1995 ApJ, 444, 1
 Lanzetta, K.M., Wolfe, A.M., Turnshek, D.A., Lu, L., McMahon, R.G. & Hazard, C. 1991, ApJS, 77, 1 [LWTLMH]
 Lanzetta, K.M., Wolfe, A.M. & Turnshek, D.A., 1995, ApJ, 440, 435 [LWT]
 Ma, C.P., Bertschinger, E. 1994, ApJ, 434, L25
 Mo, H.J., Miralde-Escudé, J. 1994, ApJ, 430, L25
 Pei, Y.C. & Fall, S.M. 1995, ApJ, 454, 69
 Pettini, M., Smith, L.J., Hunstead, R.W. & King, D.L. 1994, ApJ, 426, 79
 Rao, S. & Briggs, F. 1993, ApJ, 419, 515
 Schechter, P.L., & Dressler, A. 1987, AJ, 94, 563
 Schmidt, M. 1963, ApJ, 137, 758
 Stengler-Larrea E.A., et al., Boksenberg A., Steidel C.C., Sargent W.L.W., Bahcall J.N., Bergeron J., Hartig G.F., Januzzi B.T., Kirhakos S., Savage B.D., Schneider D.P., Turnshek D.A., Weymann R.J., 1995, ApJ, 444, 64
 Storrie-Lombardi, L.J., McMahon, R.G., Irwin, M.J., & Hazard, C. 1994, ApJ, 427, L13
 Storrie-Lombardi, L.J., McMahon, R.G., Irwin, M.J. & Hazard, C. 1996, ApJ, 468, 128 [SMIH]
 Storrie-Lombardi, L.J., Irwin, M.J. & McMahon, R.G. 1996, MNRAS, in press [SIM]
 Tytler D. 1987, ApJ, 321, 49
 Wolfe, A.M., Lanzetta, K.M., Foltz, C.B. & Chaffee, F.H. 1995, ApJ, 454, 698
 Wolfe, A.M., Turnshek, D.A., Smith, H.E. & Cohen R.D. 1986, ApJS, 61, 249 [WTSC]
 Williger, G.M., Baldwin, J.A., Carswell, R.F., Cooke, A.J., Hazard, C., Irwin, M.J., McMahon, R.G., & Storrie-Lombardi, L.J. 1994, ApJ, 428, 574

Table 1. Data for Figures

Bin	DLA	# of		$q_0 = 0$		$q_0 = 0.5$		
Redshift	$\langle z \rangle$	Δz	DLA	QSO	ΔX	Ω_g	ΔX	Ω_g
Range			in bin			$[\times h_{50}10^3]$		$[\times h_{50}10^3]$
.008-1.5	0.64	47.8	4	186	73.1	0.56 ± 0.32	58.5	0.70 ± 0.40
1.5-2.0	1.89	27.9	4	126	79.5	1.21 ± 0.71	47.1	2.05 ± 1.19
2.0-3.0	2.40	120.2	22	176	415.9	1.50 ± 0.49	223.4	2.79 ± 0.91
3.0-3.5	3.17	24.3	5	82	102.0	1.48 ± 0.72	49.8	3.04 ± 1.48
3.5-4.7	4.01	19.2	9	32	93.8	0.85 ± 0.34	42.5	1.87 ± 0.75
Dashed bins, excluding high redshift data.								
2.0-3.0	2.38	114.6	21	154	394.8	1.56 ± 0.52	212.6	2.90 ± 0.96
3.0-3.5	3.19	11.8	4	56	48.9	2.79 ± 1.47	24.0	5.68 ± 3.00


Table 2. Redshift Path and Absorption Distance

Data Set	Δz	ΔX	ΔX
		($q_0 = 0$)	($q_0 = 0.5$)
$3.0 < z < 3.5$			
APM Damped Ly α Survey	12.5	53.1	25.8
WTSC + LWTLMH + LWT	11.8	48.9	24.0
Combined	24.3	102.0	49.8
$z > 3$			
APM Damped Ly α Survey	30.5	141.2	65.5
WTSC + LWTLMH + LWT	13.0	54.6	26.7
Combined	43.5	195.8	92.2
$0 < z < 4.7$			
APM Damped Ly α Survey	36.1	162.3	76.5
WTSC + LWTLMH + LWT	203.4	602.1	344.8
Combined	239.5	764.4	421.3


Figure 1. The mean cosmological mass density in neutral gas, Ω_g , contributed by damped Ly α absorbers for $0.008 \leq z \leq 4.7$ for (a) $q_0=0$ and (b) $q_0=0.5$. The solid bins include the combined data set and the dashed bins exclude the new APM high redshift data. The region Ω_{star} is the $\pm 1\sigma$ range for the mass density in stars in nearby galaxies (Gnedin & Ostriker 1992). The point at $z=0$ is the value inferred from 21 cm emission from local galaxies (Fall & Pei 1993; Rao & Briggs 1993). These results are tabulated in table 1.

Figure 2. The mean cosmological mass density in neutral gas, Ω_g , contributed by damped Ly α absorbers for $0.008 \leq z \leq 4.7$. The solid bins are the combined data set shown in figure 1(b) for $q_0 = 0.5$. The circles show the observed data points corrected for possible dust obscuration using values determined from the closed-box/outflow models shown in figure 4(b) of Pei & Fall (1995). The squares add an estimated 20% correction for neutral gas not in damped Ly α absorbers. The region Ω_{star} is the $\pm 1\sigma$ range for the mass density in stars in nearby galaxies (Gnedin & Ostriker 1992). The point at $z=0$ is the value inferred from 21 cm emission from local galaxies (Fall & Pei 1993; Rao & Briggs 1993).

(a) $q_0=0$



(b) $q_0=0.5$



$q_0=0.5$

