Unconventional Integer Quantum Hall effect in graphene

V.P. Gusynin^{1*} and S.G. Sharapov^{2†}

¹ Bogolyubov Institute for Theoretical Physics, Metrologicheskaya Str. 14-b, Kiev, 03143, Ukraine

² Department of Physics and Astronomy, McMaster University, Hamilton, Ontario, Canada, L8S 4M1

(Dated: November 26, 2024)

Monolayer graphite films, or graphene, have quasiparticle excitations that can be described by 2+1 dimensional Dirac theory. We demonstrate that this produces an unconventional form of the quantized Hall conductivity $\sigma_{xy} = -(2e^2/h)(2n+1)$ with $n = 0, 1, \ldots$, that notably distinguishes graphene from other materials where the integer quantum Hall effect was observed. This unconventional quantization is caused by the quantum anomaly of the n = 0 Landau level and was discovered in recent experiments on ultrathin graphite films.

PACS numbers: 73.43.Cd,71.70.Di,81.05.Uw

The quantum Hall effect (QHE) is one of the most remarkable phenomena in condensed matter discovered in the second half of the 20th century. The basic experimental fact characterizing QHE is that the diagonal electric conductivity of a two-dimensional electron system in a strong magnetic field is vanishingly small $\sigma_{xx} \to 0$, while the non-diagonal conductivity is quantized in multiples of e^2/h : $\sigma_{xy} = -\nu e^2/h$, where ν is an integer (the integer quantum Hall effect (IQHE)) or a fractional number (the fractional QHE). In a recent paper¹ the fabrication of free-standing monocrystalline graphite films with thickness down to a single atomic layer was reported. This new material, called graphene, possesses truly remarkable properties such as excellent mechanical characteristics, scalability to the nanometer sizes, and the ability to sustain huge $(> 10^8 A/cm^2)$ electric currents. By using the electric field $effect^1$, it is possible to change the carrier concentration in samples by tens times and even to change the carrier type from electron to hole when the sign of applied gate voltage is altered. All this make graphene a promising candidate for applications in future micro- and nanoelectronics.

On the theoretical side, the linear, Dirac-like, spectrum of quasiparticle excitations (up to energies of the order of 1000 K) and the pseudospin degeneracy make graphene a unique truly two-dimensional "relativistic" electronic system. The thinnest graphite films can be described by a low-energy (2+1) dimensional effective massless Dirac theory². Of special interest are the properties of graphene in a magnetic field. The important differences between the Dirac and Schrödinger theories may be observed in thermodynamic and magnetotransport measurements 1,3,4,5,6 . For instance, the phase of de Haas van Alphen and Shubnikov de Haas oscillations for Dirac quasiparticles is shifted 7,8,9,10 compared to the phase of non-relativistic quasiparticles. Moreover, the Dingle and temperature factors in the amplitude of oscillations explicitly depend on the carrier density in the case of a Dirac-like spectrum 7,8 .

Because of the large value of the cyclotron gap, it is expected that the QHE in this material can be observed for much higher temperatures and lower magnetic fields than in conventional semiconductors. Therefore it is naturally to ask whether the fundamental difference between the properties of Landau levels (LL) (see Eqs. (5) and (10) below) in the Dirac and Schrödinger theories can be observed experimentally in the Hall conductivity? The purpose of this letter is to show that the Dirac-like dynamics of graphene results in an unconventional form of the Hall quantization

$$\sigma_{xy} = -\frac{2e^2}{h}(2n+1), \quad n = 0, 1, \dots$$
 (1)

We argue that the quantization rule (1) is caused by the quantum anomaly of the $n = 0 \ LL$, i.e. by the fact that it has a twice smaller degeneracy than the levels with n > 0 and its energy does not depend on the magnetic field¹¹. Remarkably this quantization is observed experimentally¹⁰ for ultrathin graphite films which exhibit the behavior expected for ideal 2D graphene.

We begin with the Lagrangian density of noninteracting quasiparticles in a single graphene sheet that in the continuum limit reads 2

$$\mathcal{L} = \sum_{\sigma=\pm 1} \bar{\Psi}_{\sigma} \left[i\gamma^0 (\hbar\partial_t - i\mu_{\sigma}) + iv_F \gamma^i (\hbar\partial_i - i\frac{e}{c}A_i) \right] \Psi_{\sigma},$$
(2)

where $\Psi_{\sigma} = (\psi_{1\sigma}(t, \mathbf{r}), \psi_{2\sigma}(t, \mathbf{r}))$ is the four-component Dirac spinor combined from two spinors $\psi_{1\sigma}, \psi_{2\sigma}$ [corresponding to **K** and **K'** points of the Fermi surface, respectively] that describe the Bloch states residing on the two different sublattices of the biparticle hexagonal lattice of the graphene sheet, and $\sigma = \pm 1$ is the spin. In Eq. (2) γ^{μ} with $\mu = 0, 1, 2$ are $4 \times 4 \gamma$ matrices belonging to a reducible representation in 2 + 1, $\bar{\Psi}_{\sigma} = \Psi_{\sigma}^{\dagger} \gamma_0$ is the Dirac conjugated spinor, -e < 0 is the electron charge, v_F is the Fermi velocity. We set $k_B = 1$, but kept Planck constant $\hbar = h/2\pi$.

The external magnetic field **B** is applied perpendicularly to the plane along the positive z axis and the vector potential is taken in the symmetric gauge $\mathbf{A} = (-B/2y, B/2x)$. In contrast to the truly relativistic $(3+1) \operatorname{case}^{12}$, the Zeeman interaction term still has to be explicitly added to the Lagrangian (2), because it originates from nonrelativistic many-body theory. This can

be done by considering spin splitting $\mu_{\sigma} = \mu - \sigma \mu_B B$ of the chemical potential μ , where $\mu_B = e\hbar/(2mc)$ is the Bohr magneton. However, for realistic values of $v_F \propto 10^5 \mathrm{m/s}$ in graphene the distance between LL is very large compared to the Zeeman splitting⁸, so that in what follows we will not consider this term and simply multiply all relevant expressions by 2 to count the spin degeneracy. While simple tight-binding calculations made for the hexagonal lattice of a single graphene sheet predict that $\mu = 0$, the real picture is more complicated and the actual value of μ can be nonzero due to finite doping and/or disorder. Moreover, nonzero and even tunable value of μ [including the change of the character of carriers, either electrons or holes] is possible in electric-field doping experiments^{1,4,10}. In our notations $\mu > 0$ corresponds to electrons and accordingly to the positive gate voltage.

Using the Kubo formalism and modeling the LL by Lorentzians with a constant width Γ the following expression for the diagonal conductivity was obtained in Refs.^{8,13}

$$\sigma_{xx}(B,\mu,\Gamma) = \frac{2e^2}{\hbar} \int_{-\infty}^{\infty} d\omega [-n'_F(\omega-\mu)] \mathcal{A}_{xx}(\omega,B,\Gamma),$$
(3)

where $n_F(\omega) = 1/[\exp(\omega/T) + 1]$ is the Fermi distribution and the function \mathcal{A}_{xx} that incorporates the effect of all LL is given by Eq. (11) of Ref.⁸. Now this result is extended for the Hall conductivity and we derive a general analytical expression for $\sigma_{xy}(B, \mu, \Gamma)$.¹⁴ The resulting dependence $\sigma_{xy}(\mu)$ is shown in Fig. 1, where one sees that the plateaux of σ_{xy} follow Eq. (1). This agrees with the latest experimental results¹⁰ and resemble earlier theoretical predictions¹⁵. However, to demonstrate result (1)



FIG. 1: (Color online) The Hall conductivity σ_{xy} measured in e^2/h units as a function of chemical potential μ for two different values of Γ for T = 3K and B = 1T. We use $\hbar v_F^2 eB/c \rightarrow (4.5 \times 10^4 \text{K}^2)B(\text{T}).$

in the most transparent way it is useful to write down a simpler conventional representation^{13,16} for σ_{xy} obtained

in the clean limit $\Gamma \to 0$:

$$\sigma_{xy} = -\frac{ec\rho}{B} \equiv -\frac{e^2 \mathrm{sgn}(eB) \mathrm{sgn}\mu}{\pi\hbar} \nu_B. \tag{4}$$

Here we introduced the filling factor of LL, $\nu_B = \pi \hbar c |\rho|/|eB|$ with ρ being the carrier imbalance ($\rho \equiv n_e - n_h$, where n_e and n_h are the densities of "electrons" and "holes", respectively). This filling factor can be represented as a sum over LL

$$M_n = \sqrt{\Delta^2 + 2n\hbar v_F^2 |eB|/c}, \quad n = 0, 1, \dots$$
 (5)

of the Dirac theory:

$$\operatorname{sgn}\mu\nu_{B} = \frac{1}{2} \left[\tanh\frac{\mu+\Delta}{2T} + \tanh\frac{\mu-\Delta}{2T} + \operatorname{tanh}\frac{\mu-\Delta}{2T} + 2\sum_{n=1}^{\infty} \left(\tanh\frac{\mu+M_{n}}{2T} + \tanh\frac{\mu-M_{n}}{2T} \right) \right], \quad (6)$$

where we separated out the level with n = 0 because its degeneracy is only half of the degeneracy of the levels with n > 0. To illustrate this rather peculiar property of the Dirac theory in a perspicuous way, we included in M_n and ν_B the mass (excitonic gap) Δ which was discussed recently to explain some experiments^{13,17}. Our consideration of σ_{xy} is in fact independent of the presence of Δ , so in what follows we set $\Delta = 0$. A zero value of Δ is expected for noninteracting quasiparticles on the hexagonal lattice of graphene.

The first equality in Eq. (4) corresponds to a classical straight line $\sigma_{12} \propto \nu_B$. As discussed, for example, in Ref.¹⁸, this line emerges from two step function dependences, viz. $\mu(n)$ and $\sigma_{12}(\mu)$. Indeed, using $\tanh(\omega/2T) = \operatorname{sgn}(\omega)$ for $T \to 0$, we obtain from Eqs. (4) and (6) (compare with Ref.¹⁶)

$$\sigma_{xy} = -\frac{2e^2 \operatorname{sgn}(eB) \operatorname{sgn}\mu}{h} \left(1 + 2\left[\frac{\mu^2 c}{2\hbar |eB| v_F^2}\right]\right), \quad (7)$$

where [x] denotes the integer part of x. The usual argumentation (see e.g. Ref.¹⁸) for the occurrence of the IQHE states that in the presence of disorder the dependence of $\sigma_{12}(\mu)$ remains the same, while $\mu(n)$ becomes a smooth function. The classical (4) and quantum (7) Hall conductivities coincide only for the fillings, $\nu_B = 2n + 1$ (see Fig. 2). The odd integer rather than integer fillings that produces the quantization rule (1) appears due to the above-mentioned halved degeneracy of the n = 0 LL. Another interesting feature of Eq. (7) (see also Figs. 1 and 2) is that $\sigma_{xy} = \pm 2e^2/h$ for the fillings $\nu_B < 1$ and it crosses 0 only when μ changes sign. On the contrary, in a conventional IQHE $\sigma_{xy} = 0$ for $\nu_B < 1$.

Although Eqs. (4) - (7) are obtained in the clean limit and using a simple bare bubble expression for conductivity, our main result (1) is model independent and is only based on the n = 0 level anomaly.



FIG. 2: (Color online) The Hall conductivity σ_{xy} measured in e^2/h units as a function of the filling ν_B . The straight line corresponds to a classical dependence (4). The parameters are the same as in Fig. 1.

Now we rewrite Eqs. (4)-(6) in terms of the Fermi distribution

$$\sigma_{xy} = -\frac{2e^2 \operatorname{sgn}(eB)}{h} \sum_{n=0}^{\infty} (2n+1) \times [n_F(M_n - \mu) + n_F(-M_n - \mu) - n_F(M_{n+1} - \mu) - n_F(-M_{n+1} - \mu)]$$
(8)

to compare it with Eq. (18) of Ref.¹⁹ that was obtained for an ideal two-dimensional electron gas

$$\sigma_{xy} = -\frac{e^2}{h} \sum_{n=0}^{\infty} (n+1) [n_F(\omega_n^{\text{nonrel}}) - n_F(\omega_{n+1}^{\text{nonrel}})] \quad (9)$$

with nonrelativistic spectrum

$$\omega_n^{\text{nonrel}} = \frac{e\hbar B}{mc} \left(n + \frac{1}{2} \right). \tag{10}$$

There is a commonsense reasoning^{1,20} that graphene is a two-band [the first band would corresponds to the electrons with $\omega_n = M_n - \mu$ and the second band, to the holes with $\omega_n = -M_n - \mu$], two-valley [corresponding to **K** and **K'** points of graphen's Fermi surface] semiconductor with zero gap Δ between the bands. Accordingly its Hall conductivity can be directly obtained from (9) by summing over all these bands and valleys

$$\sigma_{xy}^{\text{semicond}} = -\frac{2e^2 \text{sgn}(eB)}{h} \sum_{n=0}^{\infty} 2(n+1) \times [n_F(M_n - \mu) + n_F(-M_n - \mu) - n_F(M_{n+1} - \mu) - n_F(-M_{n+1} - \mu)], \qquad (11)$$

where we also counted spin degeneracy. It is easy to see that Eqs. (8) and (11) correspond to two completely different Hall conductivity quantization rules, viz. Eq. (8) which correctly counts the degeneracy of the n = 0 level produces Eq. (1), while the semiconducting analogy (11) leads to

$$\sigma_{xy}^{\text{semicond}} = -\frac{4e^2}{h}n, \qquad n = 0, 1, \dots$$
 (12)

Here we assumed that $e, B, \mu > 0$. Although previous experimental observations supported the picture based on Eq. (12), the latest experiments made on thin films¹⁰ are in accord with the unconventional Hall quantization (1). This shows that in an applied magnetic field the semiconducting interpretation of graphene's band structure that led us to Eq. (11) becomes invalid (see also Ref^{21}). The drastic difference between Eqs. (1) and (12) is caused by the above-mentioned fact that the lowest LL in Dirac theory is special and has twice smaller degeneracy than the levels with n > 0, because depending on the sign eB it is occupied either by electrons or holes, while higher levels contain both electrons and holes 11,22 . In the nonrelativistic theory when the Lande factor $q \neq 2$ all Landau levels have the same degeneracy 12 . It turns out that graphene for which the valence and conduction bands intersect in discrete points 23 , is reasonably well described by the Dirac formalism which naturally embodies the n = 0 level anomaly.

We now consider the phenomenon of quantum magnetic oscillations in graphene which is closely related to the quantization of σ_{xy} and discuss the specific of the n = 0 level. The de Haas van Alphen and Shubnikov de Haas effects in graphene were studied in Refs.^{7,8,9}. In particular, in Ref.⁸ it was shown that the oscillatory part of the diagonal conductivity (3) is given by

$$\sigma_{xx} \propto \sum_{k=1}^{\infty} \cos\left[\frac{\pi k\mu^2}{\hbar v_F^2 |eB|/c}\right] R_T(k) R_D(k) R_s(k), \quad (13)$$

where R_T , R_D and R_s are respectively the temperature, Dingle and spin factors. Using the relationship $\mu^2 = \pi \hbar^2 v_F^2 |\rho|$ valid for $T = \Gamma = B = 0^{13}$ one can check that the minima of the diagonal conductivity (3) occur at the fillings $\nu_B = 2n + 1$ giving an indication of the possible positions of the plateaux in the IQHE¹⁰. [Note that in thick films the minima of σ_{xx} occur at integer fillings⁴.] Obviously for $\mu = 0$ there is no oscillations of σ_{xx} , the conductivity $\sigma_{xx}(\mu = 0) = 2e^2/(\pi^2\hbar)$ becomes a field independent universal²⁴ quantity that is another distinctive feature of the n = 0 level anomaly.

Although the quantization (1) can be understood by considering noninteracting Dirac quasiparticles placed in an external magnetic field, even this simple model reveals other unusual properties¹¹ intimately related to nontrivial dynamics of quasiparticles from the n = 0 level. For example, the U(4) symmetry of the Lagrangian (2) is spontaneously broken down to $U(2) \times U(2)$ at $\mu = 0$ in non-zero magnetic field even in the absence of additional interaction between fermions¹¹ thus leading to the emergence of the chiral condensate $\langle \bar{\Psi}\Psi \rangle$. Including many body effects such as an attractive interaction between quasiparticles, could further generate a gap for quasiparticles like the above mentioned gap Δ (see e.g. Refs.^{13,17}). Fortunately in the case of the IQHE the presence of the condensate does not affect our consideration. On the other hand, a possible gap generation for the fermions from the lowest LL might become important for the fractional quantum Hall effect and this issue certainly deserves further experimental and theoretical study.

To conclude, we have shown that the integer numbers associated with quantized Hall conductivity in graphene have an unusual pattern $\sigma_{xy}h/e^2 = 2, 6, 10, 14...$ We argued that it is related to the fact that a theoretical

- * Electronic address: vgusynin@bitp.kiev.ua
- [†] Electronic address: sharapov@bitp.kiev.ua
- ¹ K.S. Novoselov, A.K. Geim, S.V. Morozov, D. Jiang, Y. Zhang, S.V. Dubonos, I.V. Grigorieva, and A.A. Firsov, Science **306**, 666 (2004); K.S. Novoselov, A.K. Geim, S.V. Morozov, S.V. Dubonos, Y. Zhang, and D. Jiang, cond-mat/0410631.
- ² G.W. Semenoff, Phys. Rev. Lett. **53**, 2449 (1984).
- ³ Y. Kopelevich, J.H.S. Torres, R.R. da Silva, F. Mrowka, H. Kempa, and P. Esquinazi, Phys. Rev. Lett. **90**, 156402 (2003).
- ⁴ S.V. Morozov, K.S. Novoselov, D. Jiang, A.A. Firsov, S.V. Dubonos, A.K. Geim, cond-mat/0505319.
- ⁵ C. Berger, Z. Song, T. Li, X. Li, A.Y. Ogbazghi, R. Feng, Z. Dai, A.N. Marchenkov, E.H. Conrad, P.N. First, and W.A. de Heer, J. Phys. Chem. B **108**, 19912 (2004).
- ⁶ Y. Zhang, J.P. Small, M.E.S. Amori, and P. Kim, Phys. Rev. Lett. **94**, 176803 (2005).
- ⁷ S.G. Sharapov, V.P. Gusynin, and H. Beck, Phys. Rev. B 69, 075104 (2004).
- ⁸ V.P. Gusynin and S.G. Sharapov, Phys. Rev. B **71**, 125124 (2005).
- ⁹ I.A. Luk'yanchuk and Y. Kopelevich, Phys. Rev. Lett. **93**, 166402 (2004).
- ¹⁰ A.K. Geim, private communication.
- ¹¹ V.P. Gusynin, V.A. Miransky, and I.A. Shovkovy, Phys. Rev. Lett. **73**, 3499 (1994); Phys. Rev. D **52**, 4718 (1995).
- 12Note that in the relativistic (3 + 1) Dirac theory the Zeeman term is built in the formalism. When the relativistic eigenenergy, $E(n, \sigma = \pm, k_z = 0) = [m^2 c^4 +$ $mc^{2}\hbar|eB|/(mc)(2n+1+\sigma)|^{1/2}$ is separated into the Pauli term with $\sigma \mu_B B$ and the diamagnetic term, the nonrelativistic LL with the same degeneracy and the spectrum given by Eq. (10) emerge. Each nonrelativisitic LL suffers a Zeeman splitting $\pm 1/2g\mu_B B$, so that it doubles the number of LL. Here we included the Lande factor g which plays an important role in solids. For g = 2, there is an accidental (from a nonrelativistic point of view) degeneracy of LL, so that each level with n > 0 is filled by both spin down and spin up electrons from the n-1 level. The n=0 LL is special and remains nondegenerate, so that we come to the picture considered in this paper. In our case, however, an "accidental" degeneracy of n > 0 LL is not related to the spin degree of freedom and occurs due to the presence

description of graphene is based on 2 + 1 dimensional Dirac theory, where the lowest Landau level has half of the higher Landau levels degeneracy.

We are indebted to A. Geim for showing us his experimental results prior to publication and for stimulating discussions. We also thank J.P. Carbotte, V.M. Loktev and V.A. Miransky for useful discussions and W.A. de Heer, P. Kim for informing us about their latest results. S.G.Sh. was supported by the Natural Science and Engineering Council of Canada (NSERC) and by the Canadian Institute for Advanced Research (CIAR).

of two sublattices in graphene.

- ¹³ E.V. Gorbar, V.P. Gusynin, V.A. Miransky, and I.A. Shovkovy, Phys. Rev. B 66, 045108 (2002).
- ¹⁴ The corresponding analytical expression for $\mathcal{A}_{xy}(\omega, B, \Gamma)$ is simple, but rather lengthy, V.P. Gusynin and S.G. Sharapov, in preparation.
- ¹⁵ Similar figures also appear in Y. Zheng and T. Ando, Phys. Rev. B **65**, 245420 (2002), but with a twice smaller size of the steps of σ_{xy} , and without making a link between the unusual behavior of σ_{xy} and the quantum anomaly of the n = 0 level.
- ¹⁶ A.M.J. Schakel, Phys. Rev. D **43**, 1428 (1991).
- ¹⁷ D.V. Khveshchenko, Phys. Rev. Lett. 87, 206401; *ibid.* 87, 246802 (2001).
- ¹⁸ M. Jan
 ³en, O. Veihweger, U. Fastenrath, and J. Hajdu, Introduction to the Theory of the Integer Quantum Hall Effect, edited by J. Hajdu, VCH, Weinheim, 1994.
- ¹⁹ M. Jonson and S. M. Girvin, Phys. Rev. B **29**, 1939 (1984).
- 20 M.S. Dresselhaus and G. Dresselhaus, Adv. Phys. $\mathbf{51},~1$
- (2002).
- ²¹ T. Ando, J. Phys. Soc. Jpn. **74**, 777 (2005).
- ²² M.H. Johnson and B.A. Lippmann, Phys. Rev. **76**, 828 (1949).
- ²³ The points near which the electrons have a linear spectrum and are described by a two-level Hamiltonian are called in the literature¹⁶ diabolic points . There is a theorem of H.B. Nielsen and M. Ninomiya, Nucl. Phys. **B185**, 20 (1981); **B 193**, 173 (1981) that asserts that in condensed matter systems diabolic points come in parity-inavariant pairs. Accordingly the Dirac Lagrangian (2) which embedes a pair of such points, written using a parity preserving 4×4 reducible representation of γ matrices is rather natural.
- ²⁴ This universality has the same origin as the universality considered in E. Fradkin, Phys. Rev. B **33**, 3263 (1986) for degenerate semiconductors and in P.A. Lee, Phys. Rev. Lett. **71**, 1887 (1993) for *d*-wave superconductors. Moreover in the case of graphene where the quasiparticles have not only linear disperion relation similar to the Bogolyubov quasiparticles in *d*-wave superconductor, but also couple to the vector potential in the conventional QED₂₊₁ way, the conductivity $\sigma_{xx}(\mu = 0)$ is expected to be field independent¹³.