Photon-meson transition form factors of light pseudoscala r mesons

Bo-Wen Xiao

Department of Physics, Peking University, Beijing 100871, China Department of Physics, Columbia University, New York, 10027, USA

Bo-Qiang Ma [∗](#page-0-0)

CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China Department of Physics, Peking University, Beijing 100871, China

Abstract

The photon-meson transition form factors of light pseudoscalar mesons π^0 , η , and η' are systematically calculated in a light-cone framework, which is applicable as a light-cone quark model at low Q^2 and is also physically in accordance with the light-cone pQCD approach at large Q^2 . The calculated results agree with the available experimental data at high energy scale. We also predict the low Q^2 behaviors of the photon-meson transition form factors of π^0 , η and η' , which are measurable in $e + A$ (Nucleus) $\rightarrow e + A + M$ process via Primakoff effect at JLab and DESY.

PACS numbers: 14.40.Aq; 12.39.Ki; 13.40.Gp; 13.60.Le

[∗]Corresponding author; Electronic address: mabq@phy.pku.edu.cn

I. INTRODUCTION

The meson-photon and photon-meson transition form factors contain interesting physics concerning the QCD structure of both photons and mesons. The pion-photon transition form factor provides a very simple example for the perturbative QCD (pQCD) analysis to exclusive processes, and was first analyzed by Brodsky and Lepage $[1]$ at large Q^2 . It has been shown [\[2\]](#page-19-1) that the applicability of pQCD can be extended to lower Q^2 around a few GeV^2 by taking into the transverse momentum contributions in both hard scattering amplitude and pion wave function. In our recent study [\[3\]](#page-19-2) within light-cone quark model, it is shown that the pion-photon transition form factor is identical to the photon-pion transition form factor when taking into account only QCD and QED contributions. Therefore the formalism that applies to the pion-photon transition form factor is also applicable to the photon-pion transition form factor. Taking the minimal quark-antiquark Fock states of both the photon and pion as their wave functions, we could calculate the photon-pion transition form factor by using the Drell-Yan-West assignment. This framework is applicable at low $Q²$ as a light-cone quark model approach, and it is also physically in accordance with the light-cone pQCD approach at large Q^2 . Thus we can describe the photon-pion form factors at both low Q^2 and high Q^2 within a same framework. The purpose if this work is to apply this framework [\[3\]](#page-19-2) for a systematic description of the photon-meson transition form factors of pseudoscalar mesons π^0 , η , and η' , at both $Q^2 \to 0$ and $Q^2 \to \infty$ limits, and to make predictions in a wide Q^2 range.

The photon-meson transition form factor $\gamma^* \gamma \to M$ can be realized in $e + e \to e + e + M$ or $e + A(\text{Nucleus}) \rightarrow e + A + M$ processes. The $\gamma^* \gamma \rightarrow M$ transition form factors of π^0 , η , and η' at medium to high Q^2 have been measured at Cornell [\[4\]](#page-19-3) and at DESY [\[5\]](#page-19-4) through the $e^+ + e^- \rightarrow e^+ + e^- + M$ process, while the latter process $e + A$ (Nucleus) \rightarrow $e + A + M$ is convenient to provide measurement of the photon-meson transition form factors at low Q^2 . Moreover, high precision measurements of the electromagnetic properties of these pseudoscalar mesons via Primakoff effect are proposed by PrimEx Collaboration at the Thomas Jefferson National Accelerator Facility (JLab) [\[6\]](#page-19-5), which would give the experimental value of transition form factors $F_{\gamma^*\gamma \to M}(Q^2)$ of π^0 , η , and η' at low Q^2 (0.001 – 0.5 GeV^2), and lead to a clarification on the obvious disagreement between the former Primakoff experiment and collider cases in the measurements of $\Gamma(\eta \to \gamma\gamma)$ and a more

precise determination of the η - η' mixing angle. Similar measurements can be also performed by HERMES Collaboration at Deutsche Elektronen-Synchrotron (DESY) [\[7](#page-19-6)]. Therefore, theoretical predictions at low Q^2 are necessary and essential for comparison with future experimental measurements.

It is well known that the physical η and η' states dominantly consist of a flavor $SU(3)$ octet η_8 and singlet η_0 in the $SU(3)$ quark model, respectively. The usual mixing scheme reads:

$$
\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \eta_8 \\ \eta_0 \end{pmatrix} . \tag{1}
$$

Using different sets of experimental data, we recalculate the value of the mixing angle θ by employing the limiting method developed by Cao-Signal [\[8\]](#page-19-7). Our results are also compatible with other approaches [\[9\]](#page-19-8) for the mixing angle and scheme.

In general, people use the chiral perturbation theory [\[10](#page-19-9)] or some other methods [\[11\]](#page-19-10) which deal with current quark masses in order to take the chiral symmetry and chiral anomaly into account, since the chiral symmetry predominates the $\pi^0(\eta, \eta')\gamma\gamma$ vertex at large Q^2 [\[12\]](#page-19-11), and chiral anomaly determines the $\pi^0(\eta, \eta')$ transition form factors at $Q^2 = 0$ (Eqs. [\(30](#page-10-0)[-33\)](#page-10-1)). In addition, the chiral perturbation theory is also very useful and effective in discussing the η and η' mixing properties [\[13\]](#page-19-12). Since we are consistently using the valence quark masses in the light-cone treatment to the form factor calculation, it is not very applicable to start with current quark mass within the chiral symmetry and investigate the chiral limits in the transition form factor computation. However, our main purpose of this paper is to employ the new light-cone $\gamma \to q\overline{q}, s\overline{s}$ wave functions [\[3,](#page-19-2) [14,](#page-19-13) [15](#page-19-14)] to compute the transition form factors of the light mesons. Moreover, we considered the chiral symmetry when we choose η and η' mixing scheme, and took the chiral limit approximation when we try to determine and fix the parameters. Therefore our results respect the chiral symmetry and its breaking at some extent. Phenomenologically, we could give the predictions of the η and η' mixing angle within the light-cone formalism, as well as the photon-meson transition form factor which is applicable at both low and high energy scales.

This paper is organized as follows. In section 2 we present the formalism for the photonmeson transition form factor using the minimal quark-antiquark Fock states of the photon and pion as wave functions. In section 3, we will introduce the η - η' mixing scheme used in our calculation. In section 4, we calculate systematically the photon-meson transition

form factors of π^0 , η , and η' , and show that the calculated results agree with the available experimental data at medium to large Q^2 scale. We also predict the low Q^2 behaviors of the photon-meson transition form factors of π^0 , η , and η' , which are measurable in $e +$ $A(\text{Nucleus}) \rightarrow e + A + M$ process via Primakoff effect at JLab and DESY. In section 5, we present a brief summary.

II. FORMALISM OF PHOTON-MESON TRANSITION FORM FACTOR

We work in the light-cone formalism [\[16\]](#page-19-15), which provides a convenient framework for the relativistic description of hadrons in terms of quark and gluon degrees of freedom, and for the application of perturbative QCD to exclusive processes. The transition form factor $F_{\gamma^*\gamma\to M}$ $(M = \pi^0, \eta, \text{ and } \eta')$, in which an on-shell photon is struck by one off-shell photon and decays into a meson, as schematically shown in Fig. [1,](#page-4-0) is defined by the $\gamma^* \gamma M$ vertex,

$$
\Gamma_{\mu} = -ie^2 F_{\gamma^* \gamma \to M}(Q^2) \varepsilon_{\mu \nu \rho \sigma} p_M^{\nu} \epsilon^{\rho} q^{\sigma}, \qquad (2)
$$

in which q is the momentum of the off-shell photon, $Q^2 = -q^2 = \mathbf{q}_{\perp}^2 - q^+q^- = \mathbf{q}_{\perp}^2$ is the squared four momentum transfer of the virtual photon, and ϵ is the polarization vector of the on-shell photon. We choose the light-cone frame

$$
\begin{cases}\nP = (P^+, \frac{q^2 + \mathbf{q}_\perp^2}{P^+}, \mathbf{0}_\perp), \\
P' = (P'^+, \frac{M^2}{P'^+}, \mathbf{q}_\perp), \\
q = (0, \frac{Q^2}{P^+}, \mathbf{q}_\perp), \\
p_1 = (xP^+, \frac{\mathbf{k}_\perp^2 + m^2}{xP^+}, \mathbf{k}_\perp) \\
p_2 = ((1-x)P^+, \frac{\mathbf{k}_\perp^2 + m^2}{(1-x)P^+}, -\mathbf{k}_\perp), \\
p'_1 = (xP'^+, \frac{\mathbf{k}_\perp^2 + m^2}{xP^+}, \mathbf{k}'_\perp).\n\end{cases} (3)
$$

Instead of calculating the diagram directly, we introduce the quark-antiquark wave function of the photon [\[3](#page-19-2)] by calculating the matrix elements of

$$
\frac{\overline{u}(p_1^+, p_1^-, \mathbf{p}_{1\perp})}{\sqrt{p_1^+}} \gamma \cdot \epsilon \frac{v(p_2^+, p_2^-, \mathbf{p}_{2\perp})}{\sqrt{p_2^+}},\tag{4}
$$

which are the numerators of the wave functions corresponding to each constituent spin S^z configuration. The two boson polarization vectors in light-cone gauge are $\epsilon^{\mu} = (\epsilon^+ =$ $(0, \epsilon^{-}, \epsilon_{\perp}),$ where $\epsilon_{\perp \uparrow, \downarrow} = \pm \frac{1}{\sqrt{2}}$ $\frac{1}{2}(\hat{\mathbf{x}} \pm \hat{\mathbf{y}})$. To satisfy the Lorentz condition $k_{photon} \cdot \epsilon = 0$, the

FIG. 1: The diagram for the contribution to the transition form factor $F_{\gamma^*\gamma \to M}$. The arrows indicate the particle moving directions.

polarizations have the relation $\epsilon^- = \frac{2 \epsilon_+ \cdot k_+}{k^+}$ with k_{hoton} , thus we have

$$
\begin{cases}\n\Psi_R^{\uparrow}(x, \mathbf{k}_{\perp}, \uparrow, \downarrow) = -\frac{\sqrt{2}(k_1 + ik_2)}{1 - x} \varphi_{\gamma}, \,\, [l^z = +1] \\
\Psi_R^{\uparrow}(x, \mathbf{k}_{\perp}, \downarrow, \uparrow) = +\frac{\sqrt{2}(k_1 + ik_2)}{x} \varphi_{\gamma}, \,\, [l^z = +1] \\
\Psi_R^{\uparrow}(x, \mathbf{k}_{\perp}, \uparrow, \uparrow) = -\frac{\sqrt{2}m}{x(1 - x)} \varphi_{\gamma}, \qquad [l^z = 0] \\
\Psi_R^{\uparrow}(x, \mathbf{k}_{\perp}, \downarrow, \downarrow) = 0,\n\end{cases} \tag{5}
$$

in which:

$$
\varphi_{\gamma} = \frac{e_q}{D} = \frac{e_q}{\lambda^2 - \frac{m^2 + \mathbf{k}_{\perp}^2}{x} - \frac{m^2 + \mathbf{k}_{\perp}^2}{1 - x}},\tag{6}
$$

where λ is the photon mass and equals to 0. Each configuration satisfies the spin sum rule: $J^z = S_q^z + S_{\overline{q}}^z + l^z = +1$. Therefore, the quark-antiquark Fock-state for the photon $(J^z = +1)$ has the four possible spin combinations:

$$
\left| \Psi_{\gamma}^{\uparrow} \left(P^{+}, \mathbf{P}_{\perp} \right) \right\rangle = \int \frac{\mathrm{d}^{2} \mathbf{k}_{\perp} \mathrm{d}x}{16\pi^{3}} \times \left[\Psi_{R}^{\uparrow}(x, \mathbf{k}_{\perp}, \uparrow, \downarrow) \left| x P^{+}, \mathbf{k}_{\perp}, \uparrow, \downarrow \right\rangle + \Psi_{R}^{\uparrow}(x, \mathbf{k}_{\perp}, \downarrow, \uparrow) \left| x P^{+}, \mathbf{k}_{\perp}, \downarrow, \uparrow \right\rangle \right] \times \left[\Psi_{R}^{\uparrow}(x, \mathbf{k}_{\perp}, \uparrow, \uparrow) \left| x P^{+}, \mathbf{k}_{\perp}, \uparrow, \uparrow \right\rangle + \Psi_{R}^{\uparrow}(x, \mathbf{k}_{\perp}, \downarrow, \downarrow) \left| x P^{+}, \mathbf{k}_{\perp}, \downarrow, \downarrow \right\rangle \right]. \tag{7}
$$

The quark-antiquark Fock-state wave function of the pion is also derived [\[3](#page-19-2)] by using the relativistic field theory treatment of the interaction vertex along with the idea in [\[17](#page-20-0), [18\]](#page-20-1). In the light-cone frame of pion,

$$
\begin{cases}\nP = (P^+, \frac{M^2}{P^+}, \mathbf{0}_{\perp}), \\
p_1 = (xP^+, \frac{\mathbf{p}_{1\perp}^2 + m^2}{xP^+}, \mathbf{p}_{1\perp}) \\
p_2 = ((1-x)P^+, \frac{\mathbf{p}_{2\perp}^2 + m^2}{(1-x)P^+}, \mathbf{p}_{2\perp}),\n\end{cases}
$$
\n(8)

we can obtain the four components of the spin wave function by calculating the matrix elements of

$$
\frac{\overline{v}(p_2^+, p_2^-, -\mathbf{k}_\perp)}{\sqrt{p_2^+}} \gamma_5 \frac{u(p_1^+, p_1^-, \mathbf{k}_\perp)}{\sqrt{p_1^+}},\tag{9}
$$

from which we have

$$
\begin{cases}\n\frac{\overline{v}_{\downarrow}}{\sqrt{p_2^+}} \gamma_5 \frac{u_{\uparrow}}{\sqrt{p_1^+}} = -\frac{2mP^+}{4mx(1-x)P^{+2}},\\ \n\frac{\overline{v}_{\downarrow}}{\sqrt{p_2^+}} \gamma_5 \frac{u_{\uparrow}}{\sqrt{p_1^+}} = +\frac{2mP^+}{4mx(1-x)P^{+2}},\\ \n\frac{\overline{v}_{\uparrow}}{\sqrt{p_2^+}} \gamma_5 \frac{u_{\uparrow}}{\sqrt{p_1^+}} = +\frac{2(k_1+ik_2)P^+}{4mx(1-x)P^{+2}},\\ \n\frac{\overline{v}_{\downarrow}}{\sqrt{p_2^+}} \gamma_5 \frac{u_{\downarrow}}{\sqrt{p_1^+}} = +\frac{2(k_1-ik_2)P^+}{4mx(1-x)P^{+2}},\n\end{cases} (10)
$$

where m is the mass of the quark. After the normalization, we can obtain light-cone representation for the spin structure of the pion, which is the minimal Fock-state of the pion light-cone wave function:

$$
\begin{cases}\n\Psi_{\pi L}(x, \mathbf{k}_{\perp}, \uparrow, \downarrow) = -\frac{m}{\sqrt{2(m^2 + \mathbf{k}_{\perp}^2)}} \varphi_{\pi}, \,\, [l^z = 0] \\
\Psi_{\pi L}(x, \mathbf{k}_{\perp}, \downarrow, \uparrow) = +\frac{m}{\sqrt{2(m^2 + \mathbf{k}_{\perp}^2)}} \varphi_{\pi}, \,\, [l^z = 0] \\
\Psi_{\pi L}(x, \mathbf{k}_{\perp}, \uparrow, \uparrow) = +\frac{k_1 + ik_2}{\sqrt{2(m^2 + \mathbf{k}_{\perp}^2)}} \varphi_{\pi}, \,\, [l^z = -1] \\
\Psi_{\pi L}(x, \mathbf{k}_{\perp}, \downarrow, \downarrow) = +\frac{k_1 - ik_2}{\sqrt{2(m^2 + \mathbf{k}_{\perp}^2)}} \varphi_{\pi}, \,\, [l^z = +1]\n\end{cases} (11)
$$

in which we may employ the Brodsky-Huang-Lepage (BHL) prescription [\[19\]](#page-20-2),

$$
\varphi_{\pi}(x,\mathbf{k}) = A \exp\left[-\frac{1}{8\beta^2} \frac{\mathbf{k}_{\perp}^2 + m^2}{x(1-x)}\right],\tag{12}
$$

for the momentum space wave function, which is a non-relativistic solution of the Bethe-Salpeter equation in an instantaneous approximation in the rest frame for meson. Each configuration satisfies the spin sum rule: $J^z = S_q^z + S_{\overline{q}}^z + l^z = 0$. Hence, the Fock expansion of the two particle Fock-state for the pion has these four possible spin combinations:

$$
\langle \Psi_{\pi} (P^{+}, \mathbf{P}_{\perp} = \mathbf{0}_{\perp}) | = \int \frac{d^{2} \mathbf{k}_{\perp} dx}{16 \pi^{3}} \times \left[\Psi_{\pi L} (x, \mathbf{k}_{\perp}, \uparrow, \downarrow) \langle x P^{+}, \mathbf{k}_{\perp}, \uparrow, \downarrow | + \Psi_{\pi L} (x, \mathbf{k}_{\perp}, \downarrow, \uparrow) \langle x P^{+}, \mathbf{k}_{\perp}, \downarrow, \uparrow \right] + \Psi_{\pi L} (x, \mathbf{k}_{\perp}, \uparrow, \uparrow) \langle x P^{+}, \mathbf{k}_{\perp}, \uparrow, \uparrow | + \Psi_{\pi L} (x, \mathbf{k}_{\perp}, \downarrow, \downarrow) \langle x P^{+}, \mathbf{k}_{\perp}, \downarrow, \downarrow | \right].
$$
\n(13)

There are two higher helicity $(\lambda_1 + \lambda_2 = \pm 1)$ components in the expression of the light-cone spin wave function of the pion besides the ordinary helicity $(\lambda_1 + \lambda_2 = 0)$ components. Such higher helicity components come from the Melosh-Wigner rotation in the light-cone quark model [\[20,](#page-20-3) [21](#page-20-4)], and the same effect plays an important role to understand the proton "spin puzzle" in the nucleon case [\[22](#page-20-5), [23](#page-20-6)]. One may also state that these higher helicity components contain contribution from orbital angular moment from a relativistic viewpoint [\[24\]](#page-20-7).

In addition, we would like to add some more remarks on the Gaussian-type wavefunction of the BHL prescription that we employ above. As a matter of fact, the Gaussian wave function is a non-relativistic solution of the Bethe-Salpeter equation in an instantaneous approximation in the rest frame of the meson as the space wave function. The BHL wavefunction Eq. [\(12\)](#page-5-0) is an extension from a non-relativistic wavefunction into a relativistic form by using the Brodsky-Huang-Lepage Ansatz [\[19](#page-20-2)], and we can consider it as an approximate wavefunction that respects the Lorentz invariance in the light-cone formalism. However, it works phenomenologically well in a lot of calculations $(e.g., [3, 21, 25, 26])$ $(e.g., [3, 21, 25, 26])$. Moreover, Donnachie, Gravelis, and Shaw [\[25](#page-20-8)] indicated that the other four possible different space wave functions have similar analytical properties with the BHL wavefunction when the parameter β is small (The β is equal to P_F in their paper, the small β is corresponding to the ρ and φ mesons cases). However, they also illustrated that the BHL wavefunction is better than the other four wave functions in the high β situation (for the J/ψ meson). Hence, it gives us the idea that the the BHL wavefunction may be an appropriate choice that we could have right now. (Noticing that the space wave functions for the vector mesons are the same with those for the pseudoscalar mesons, the argument that made by Donnachie *et al* is also valid for π^0 and other pseudoscalar mesons.)

For the physical state of π^0 , one should also take into account the color and flavor degrees of freedom into account [\[1,](#page-19-0) [2](#page-19-1)]

$$
|\Psi_{\pi^0}\rangle = \sum_a \frac{\delta_b^a}{\sqrt{n_c}} \frac{1}{\sqrt{2}} \left[\left| u^a \overline{u}^b \right\rangle - \left| d^a \overline{d}^b \right\rangle \right],\tag{14}
$$

where a and b are color indices, $n_c = 3$ is the number of colors, and now $\left| q^a \overline{q}^b \right\rangle$ contains the full spin structure shown above. So we can get the photon-meson transition form factor of the pion:

$$
F_{\gamma\gamma^* \to \pi}(Q^2) = \frac{\Gamma^+}{-ie^2(\epsilon_{\perp} \times \mathbf{q}_{\perp})p_{\pi}'} = 2\sqrt{3}(e_u^2 - e_d^2) \int_0^1 dx \int \frac{d^2 \mathbf{k}_{\perp}}{16\pi^3} \varphi_{\pi}(x, \mathbf{k}'_{\perp})
$$

$$
\left\{ \frac{m}{x\sqrt{m^2 + \mathbf{k}'_{\perp}^2}} \times \left[\frac{1}{-\lambda^2 + \frac{m^2 + \mathbf{k}_{\perp}^2}{x} + \frac{m^2 + \mathbf{k}_{\perp}^2}{1 - x^2}} \right] + (1 \leftrightarrow 2) \right\}
$$
(15)

$$
= 4\sqrt{3}(e_u^2 - e_d^2) \int_0^1 dx \int \frac{d^2 \mathbf{k}_{\perp}}{16\pi^3} \left[\varphi_{\pi}(x, \mathbf{k}'_{\perp}) \frac{m}{x\sqrt{m^2 + \mathbf{k}'_{\perp}^2}} \right]
$$

$$
\times \frac{1}{\frac{m^2 + \mathbf{k}_{\perp}^2}{x} - \frac{m^2 + \mathbf{k}_{\perp}^2}{1 - x}} ,
$$
(16)

in which $\mathbf{k}'_{\perp} = \mathbf{k}_{\perp} + (1-x)\mathbf{q}_{\perp}$ after considering the Drell-Yan-West assignment [\[27](#page-20-10)], and λ $(= 0)$ is the mass of photon.

III. THE η - η' MIXING SCHEMES

In fact, there are two popular mixing schemes for η and η' . Feldmann *et al* [\[9](#page-19-8)] suggested the mixing scheme based on the quark flavor basis $q\overline{q} = (u\overline{u} + d\overline{d})/\sqrt{2}$ and $s\overline{s}$,

$$
\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \eta_q \\ \eta_s \end{pmatrix}, \qquad (17)
$$

and

$$
\begin{pmatrix} f_{\eta}^q & f_{\eta}^s \\ f_{\eta'}^q & f_{\eta'}^s \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} f_q & 0 \\ 0 & f_s \end{pmatrix},
$$
(18)

where ϕ is the mixing angle. The $q\overline{q}$ -ss mixing only introduces one mixing angle in the mixing of the decay constants.

On the other hand, people also use the mixing scheme based on the basis of η_8 and η_0 mixing for η and η' . In the $SU(3)$ quark model, the physical η and η' states dominantly consist of a flavor $SU(3)$ octet $\eta_8 = \frac{1}{\sqrt{2}}$ $\frac{1}{6}(u\overline{u} + d\overline{d} - 2s\overline{s})$ and a singlet $\eta_0 = \frac{1}{\sqrt{s}}$ $\overline{3}(u\overline{u}+dd+s\overline{s}),$ respectively. The usual mixing scheme reads:

$$
\begin{cases}\n|\eta\rangle = \cos\theta |\eta_8\rangle - \sin\theta |\eta_0\rangle, \\
|\eta'\rangle = \sin\theta |\eta_8\rangle + \cos\theta |\eta_0\rangle,\n\end{cases}
$$
\n(19)

in which θ is the mixing angle. For the calculation of the decay constants of the η_8 and η_0 mixing, Feldmann-Kroll indicate that two mixing angle scheme could be better from their former investigations[\[28\]](#page-20-11):

$$
\begin{pmatrix} f_{\eta}^8 & f_{\eta}^0 \\ f_{\eta'}^8 & f_{\eta'}^0 \end{pmatrix} = \begin{pmatrix} \cos \theta_8 & -\sin \theta_0 \\ \sin \theta_8 & \cos \theta_0 \end{pmatrix} \begin{pmatrix} f_8 & 0 \\ 0 & f_0 \end{pmatrix}.
$$
 (20)

In addition, one could find that these two schemes could be related by the following equation through the mixing angles finally:

$$
\theta = \phi - \arctan \frac{1}{\sqrt{3}}.\tag{21}
$$

From the point view of the chiral symmetry and the $SU(3)$ symmetry as well as their breaking mechanisms, we find that the η_8 and η_0 mixing scheme may be more reasonable and physical.

First of all, let us have a brief review on the chiral symmetry and its breaking which have underlying relationship with the π^0 , η and η' mesons [\[6](#page-19-5)]. In the chiral symmetry limit, it is well-known that the Lagrangian has the $SU(3)_L \times SU(3)_R \times U(1)_B \times U(1)_A$ symmetry, but the absence of this symmetry in the ground state (the QCD vacuum) leads to the chiral symmetry spontaneously breaking into $SU(3) \times U(1)_B$ symmetry. Because there are 8 spontaneously broken continuous symmetries (there are 9 when taking into account the chiral anomaly which is associated with the the $U(1)_A$ symmetry breaking), there are 8 massless Goldstone Bosons (which finally are identified as meson octet) and 1 massive particle (which is known as η_0) according to the Goldstone's theorem and chiral anomaly, respectively. The massless octet includes the meson π_8^0 and η_8 . Together with η_0 , they mix into massive mesons π^0 , η and η' during the explicit $SU(3)$ symmetry breaking after introducing the quark mass term into the Lagrangian.

From the above discussion, we may reach a physical intuitive idea that it is natural and straightforward to use the η_0 and η_8 mixing scheme as a direct result of the $SU(3)$ symmetry breaking if we assume that the π_8^0 does not mix with η_0 and η_8 at all. From this point of view, the introduction of η_0 and η_8 is more reasonable than η_q and η_s .

Moreover, since it is well-known that pion, kaon, and η_8 belong to the same group of octet mesons in the $SU(3)$ symmetry limit, their parameters should be the same except the quark masses. In this sense, one may relate the decay constants of η and η' , to pion and kaon in the $\eta_8 - \eta_0$ mixing scheme. The CLEO Collaboration[\[4\]](#page-19-3) reported their pole fit results as $\Lambda_{\pi} = 776 \pm 10 \pm 12 \pm 16$ MeV , $\Lambda_{\eta} = 774 \pm 11 \pm 16 \pm 22$ MeV , and $\Lambda_{\eta\prime} = 859 \pm 9 \pm 18 \pm 20$ MeV . These results imply that the nonperturbative properties of π and η are very similar. In addition, the absolute value of θ is small and $\cos \theta | \eta_8 \rangle$ is the leading order in the $\eta_8 - \eta_0$ mixing scheme of the η . They are consistent with the basic physical intuition that both π and η_8 are in the $SU_f(3)$ octet and are pseudo-massless Goldstone particles. Therefore, that is why the authors of [\[29\]](#page-20-12) take the parameters of η_8 as equal to pion, such as $b_8 = b_\pi$ in their paper. From a strict sense, if pion, kaon, and η_8 are in the same group of octet mesons, the mass of m_q , m_s , and $\beta_8 = \beta_\pi$ in the BHL wave function should be the same in the calculations of the π , η and η' transition form factors.

Therefore, we employ the intuitive $\eta_8-\eta_0$ mixing scheme in the calculations of the π , η and η' transition form factors by using the uniform parameters, which shows that the $SU(3)$ symmetry limit works well in this work.

In practice, we utilize the $SU_f(3)$ broken form of wave functions for flavor octet η_8 and singlet η_0 :

$$
|\eta_8\rangle = \frac{1}{\sqrt{6}}(u\overline{u} + d\overline{d})\phi_8^q(x, \mathbf{k}_\perp) - \frac{2}{\sqrt{6}}s\overline{s}\phi_8^s(x, \mathbf{k}_\perp),\tag{22}
$$

$$
|\eta_0\rangle = \frac{1}{\sqrt{3}} (u\overline{u} + d\overline{d}) \phi_0^q(x, \mathbf{k}_\perp) + \frac{1}{\sqrt{3}} s\overline{s} \phi_0^s(x, \mathbf{k}_\perp), \tag{23}
$$

in which we use Gaussian wave function of the BHL prescription:

$$
\phi_8^q(x, \mathbf{k}_{\perp}) = A_8 \exp\left[-\frac{m_q^2 + \mathbf{k}_{\perp}^2}{8\beta_8^2 x(1-x)}\right],\tag{24}
$$

$$
\phi_8^s(x, \mathbf{k}_{\perp}) = A_8 \exp\left[-\frac{m_s^2 + \mathbf{k}_{\perp}^2}{8\beta_8^2 x (1 - x)} \right],\tag{25}
$$

$$
\phi_0^q(x, \mathbf{k}_{\perp}) = A_0 \exp\left[-\frac{m_q^2 + \mathbf{k}_{\perp}^2}{8\beta_0^2 x (1 - x)}\right],\tag{26}
$$

$$
\phi_0^s(x, \mathbf{k}_{\perp}) = A_0 \exp\left[-\frac{m_s^2 + \mathbf{k}_{\perp}^2}{8\beta_0^2 x (1 - x)} \right],\tag{27}
$$

and $q\bar{q}$ and $s\bar{s}$ are the spin parts of the wave functions which are similar to the pion with all possible spin states.

Moreover, it is convenient to use the method for the $\eta_8-\eta_0$ mixing scheme which was developed by Cao-Signal [\[8](#page-19-7)] in obtaining the mixing angle θ and the decay constants. In this treatment, we can get θ , f_8 and f_0 directly without involving θ_8 and θ_0 . In the $\eta_8-\eta_0$ mixing scheme, we have:

$$
F_{\gamma\gamma^* \to \eta}(Q^2) = F_{\gamma\gamma^* \to \eta_8}(Q^2) \cos\theta - F_{\gamma\gamma^* \to \eta_0}(Q^2) \sin\theta,
$$
\n(28)

$$
F_{\gamma\gamma^* \to \eta\prime}(Q^2) = F_{\gamma\gamma^* \to \eta_8}(Q^2)\sin\theta + F_{\gamma\gamma^* \to \eta_0}(Q^2)\cos\theta.
$$
 (29)

While for the π^0 case, we have:

$$
\Gamma(\pi^0 \to \gamma \gamma) = \frac{\pi \alpha^2 m_{\pi^0}^3}{4} |F_{\gamma \gamma^* \to \pi}(0)|^2,\tag{30}
$$

$$
\Gamma(\pi^0 \to \gamma \gamma) = \frac{\alpha^2 m_{\pi^0}^3}{64\pi^3} \frac{1}{f_\pi^2}.
$$
\n(31)

Generalizing these equations to η_8 and $\eta_0,$ we have

$$
\Gamma(\eta \to \gamma\gamma) = \frac{\pi\alpha^2 m_\eta^3}{4} |F_{\gamma\gamma^* \to \eta}(0)|^2 = \frac{\alpha^2 m_\eta^3}{64\pi^3} \left(\frac{\cos\theta}{\sqrt{3}f_8} - \frac{2\sqrt{2}\sin\theta}{\sqrt{3}f_0}\right)^2,\tag{32}
$$

$$
\Gamma(\eta' \to \gamma\gamma) = \frac{\pi\alpha^2 m_{\eta'}^3}{4} |F_{\gamma\gamma^* \to \eta'}(0)|^2 = \frac{\alpha^2 m_{\eta'}^3}{64\pi^3} \left(\frac{\sin\theta}{\sqrt{3}f_8} + \frac{2\sqrt{2}\cos\theta}{\sqrt{3}f_0}\right)^2.
$$
 (33)

Thus we could get:

$$
\rho_1 = \frac{F_{\gamma\gamma^* \to \eta}(0)}{F_{\gamma\gamma^* \to \eta\prime}(0)} = \frac{\tan \theta_{08} - \tan \theta}{1 + \tan \theta_{08} \times \tan \theta},\tag{34}
$$

$$
= \tan (\theta_{08} - \theta) , \qquad (35)
$$

in which we let $\tan \theta_{08} = \frac{f_0}{\sqrt{8}}$ $\frac{f_0}{8f_8}$. Along with the same idea by taking the $Q^2 \to \infty$ limit, we could have:

$$
\rho_2 = \frac{F_{\gamma\gamma^* \to \eta}(Q^2 \to \infty)}{F_{\gamma\gamma^* \to \eta\prime}(Q^2 \to \infty)} = \frac{F_{\gamma\gamma^* \to \eta_8}(Q^2 \to \infty)\cos\theta - F_{\gamma\gamma^* \to \eta_0}(Q^2 \to \infty)\sin\theta}{F_{\gamma\gamma^* \to \eta_8}(Q^2 \to \infty)\sin\theta + F_{\gamma\gamma^* \to \eta_0}(Q^2 \to \infty)\cos\theta}
$$

=
$$
\frac{1 - 8\tan\theta_{08} \times \tan\theta}{\tan\theta + 8\tan\theta_{08}},
$$
(36)

in which we have $\lim_{Q^2 \to \infty} Q^2 F_8(Q^2) = \frac{2f_8}{\sqrt{3}}$ $\frac{\xi_8}{3}$ and $\lim_{Q^2 \to \infty} Q^2 F_0(Q^2) = \frac{4\sqrt{3}}{\sqrt{3}}$ $\frac{\sqrt{2f_0}}{\sqrt{2}}$ $\frac{^{2}10}{3}$. CLEO [\[4\]](#page-19-3) proposed that the $\gamma\gamma^*\to M$ transition form factors could be approximated by:

$$
F_{\gamma\gamma^* \to M}(Q^2) = F_{\gamma\gamma^* \to M}(0) \times \frac{1}{1 + Q^2/\Lambda_M^2},
$$
\n(37)

thus we obtain:

$$
\rho_2 = \rho_1 \frac{\Lambda_\eta^2}{\Lambda_{\eta\prime}^2}.\tag{38}
$$

Finally one obtains:

$$
\tan \theta = \frac{-9(\rho_1 + \rho_2) + \sqrt{81(\rho_1 - \rho_2)^2 + 32(\rho_1 \rho_2 + 1)^2}}{2(8 - \rho_1 \rho_2)},
$$
\n(39)

$$
\frac{f_0}{f_8} = \sqrt{8} \tan(\theta + \arctan \rho_1),\tag{40}
$$

and gets

$$
f_8 = \frac{1}{4\sqrt{3}\pi^2 \left[F_{\gamma\gamma^* \to \eta}(0)\cos\theta + F_{\gamma\gamma^* \to \eta\prime}(0)\sin\theta\right]},\tag{41}
$$

$$
f_0 = \frac{\sqrt{8}}{4\sqrt{3}\pi^2 \left[F_{\gamma\gamma^* \to \eta'}(0)\sin\theta - F_{\gamma\gamma^* \to \eta}(0)\cos\theta\right]},\tag{42}
$$

by using the above results.

IV. $\gamma^*\gamma \to \eta$ AND $\gamma^*\gamma \to \eta$ TRANSITION FORM FACTORS

There have been many different approaches to discuss the photon-meson transition form factors of light pseudoscalar mesons π^0 , η , and η' , such as the light-cone perturbation theory by Cao-Huang-Ma [\[2,](#page-19-1) [29](#page-20-12)], the light-front quark model by Hwang and Choi-Ji [\[26\]](#page-20-9), QCD sum rule calculation by Radyushkin-Ruskov [\[30\]](#page-20-13), and also other approaches et al. [\[31\]](#page-20-14). We now perform a systematic calculation of these transition form factors in the light-cone framework just presented in section 2. The advantage of this new framework is that the predictions should be applicable at both low and high energy scales.

Similar to the pion transition form factor and from Eq. [\(22\)](#page-9-0) and Eq. [\(23\)](#page-9-0), we can get η_8 and η_0 transition form factors:

$$
F_{\gamma\gamma^* \to \eta_8}(Q^2) = 4(e_u^2 + e_d^2) \int \frac{\mathrm{d}x \mathrm{d}^2 \mathbf{k}_\perp}{16\pi^3} \frac{m_q}{x\sqrt{m_q^2 + \mathbf{k}_\perp^{\prime 2}}} \phi_8^q(x, \mathbf{k}_\perp') \frac{x(1-x)}{m_q^2 + \mathbf{k}_\perp^2}
$$

$$
-8e_s^2 \int \frac{\mathrm{d}x \mathrm{d}^2 \mathbf{k}_\perp}{16\pi^3} \frac{m_s}{x\sqrt{m_s^2 + \mathbf{k}_\perp^{\prime 2}}} \phi_8^s(x, \mathbf{k}_\perp') \frac{x(1-x)}{m_s^2 + \mathbf{k}_\perp^2}, \qquad (43)
$$

$$
F_{\gamma\gamma^* \to \eta_0}(Q^2) = 4\sqrt{2}(e_u^2 + e_d^2) \int \frac{\mathrm{d}x \mathrm{d}^2 \mathbf{k}_\perp}{16\pi^3} \frac{m_q}{x\sqrt{m_q^2 + \mathbf{k}_\perp^{\prime 2}}} \phi_0^q(x, \mathbf{k}_\perp') \frac{x(1-x)}{m_q^2 + \mathbf{k}_\perp^2}
$$

$$
+4\sqrt{2}e_s^2 \int \frac{\mathrm{d}x \mathrm{d}^2 \mathbf{k}_\perp}{16\pi^3} \frac{m_s}{x\sqrt{m_s^2 + \mathbf{k}_\perp^{\prime 2}}} \phi_0^s(x, \mathbf{k}_\perp') \frac{x(1-x)}{m_s^2 + \mathbf{k}_\perp^{\prime 2}}, \qquad (44)
$$

in which $\mathbf{k}'_{\perp} = \mathbf{k}_{\perp} + (1-x)\mathbf{q}_{\perp}$ after considering the Drell-Yan-West assignment, and then we get $F_{\gamma\gamma^*\to\eta}(Q^2)$ and $F_{\gamma\gamma^*\to\eta\prime}(Q^2)$ in the $\eta_8\text{-}\eta_0$ mixing scheme

$$
F_{\gamma\gamma^* \to \eta}(Q^2) = F_{\gamma\gamma^* \to \eta_8}(Q^2)\cos\theta - F_{\gamma\gamma^* \to \eta_0}(Q^2)\sin\theta,\tag{45}
$$

$$
F_{\gamma\gamma^* \to \eta\prime}(Q^2) = F_{\gamma\gamma^* \to \eta_8}(Q^2)\sin\theta + F_{\gamma\gamma^* \to \eta_0}(Q^2)\cos\theta. \tag{46}
$$

A. Numerical calculations

First of all, we would like to determine the mixing angle θ and decay constants of f_8 and f_0 by employing Eq. [\(39\)](#page-11-0) to Eq. [\(42\)](#page-11-1) with two different sets of experimental data which may cast some light on the clarification of the obvious current disagreement between the former Primakoff experiments and collider results in the measurements of $\Gamma(\eta \to \gamma\gamma)$, and then give more reasonable predictions on the mixing angle θ . From the Particle Data Group book [\[32\]](#page-21-0), we get:

$$
\Gamma(\pi^0 \to \gamma\gamma) = 7.74 \pm 0.54 \text{ eV},\tag{47}
$$

$$
\Gamma(\eta \to \gamma\gamma) = 0.46 \pm 0.04 \text{ keV},\tag{48}
$$

$$
\Gamma(\eta' \to \gamma\gamma) = 4.29 \pm 0.15 \text{ keV},\tag{49}
$$

and

$$
m_{\pi^0} = 134.9766 \pm 0.0006 \text{ MeV},\tag{50}
$$

$$
m_{\eta} = 547.30 \pm 0.12 \text{ MeV},\tag{51}
$$

$$
m_{\eta\prime} = 957.78 \pm 0.14 \text{ MeV}.\tag{52}
$$

We can get $\theta = -14.7^{\circ} \pm 2.0^{\circ}$, $f_0 = 1.13 \pm 0.08 f_{\pi}$ and $f_8 = 0.97 \pm 0.07 f_{\pi}$. However, $\Gamma(\eta \to \gamma \gamma) = 0.511 \pm 0.026$ keV if we do not include the Primakoff production measurement of $\eta \to \gamma \gamma$ (Γ($\eta \to \gamma \gamma$) = 0.324 ± 0.046 keV) which obviously disagrees with other collider measurement. Therefore, we obtain $\theta = -16.1^{\circ} \pm 1.5^{\circ}$, $f_0 = 1.11 \pm 0.08 f_{\pi}$ and $f_8 = 0.95 \pm 0.05 f_{\pi}$ 0.07 f_{π} . Moreover, we find that the mixing angle $\phi = \theta + \arctan \frac{1}{\sqrt{2}}$ $\frac{1}{3}$ = 38.6° is compatible with [\[9\]](#page-19-8) which gives the phenomenological value of the mixing angle $\phi = 39.3^{\circ} \pm 1.0^{\circ}$ from eight decay and scattering processes. The mixing independent ratio R can be defined as follow:

$$
R \equiv \frac{M_{\pi}^3}{\Gamma(\pi \to \gamma \gamma)} \left[\frac{\Gamma(\eta \to \gamma \gamma)}{M_{\eta}^3} + \frac{\Gamma(\eta' \to \gamma \gamma)}{M_{\eta'}^3} \right]
$$
(53)

$$
= \frac{1}{3} \left(\frac{f_{\pi}^2}{f_8^2} + 8 \frac{f_{\pi}^2}{f_0^2} \right). \tag{54}
$$

The current experimental value of R which was given in the proposal of the PrimEx Collab-oration [\[6](#page-19-5)] at JLab is $R_{\text{exp}} = 2.5 \pm 0.5 \text{(stat)} \pm 0.5 \text{(syst)}$. We can get $R = 2.45$ and $R = 2.54$ respectively by using the above two sets of the parameters. With the latter set of the fitted value of the mixing angle θ and decay constants of f_8 and f_0 as the input parameters, we can fix the left seven parameters by the following nine constraints.

In the formulas of the transition form factor $F_{\gamma\gamma^*\to P}(Q^2)$ $(P = \pi, \eta_8, \eta_0)$, the parameters are the normalization constants A_{π} , A_8 and A_0 , the harmonic scale $\beta_{\pi} = \beta_8$ and β_0 , and the quark masses $m_q = m_u = m_d$ and m_s . In order to take a numerical calculation of the transition form factor $F_{\gamma\gamma^*\to M}(Q^2)$ and compare it with the available experimental data, we need to employ nine constraints to fix those seven parameters above. Thus, we can determine all these seven parameters in the transition form factor uniquely.

1. The decay widths of π , η and η' [\[4,](#page-19-3) [32\]](#page-21-0):

$$
F_{\pi\gamma}(0) = \sqrt{\frac{4}{\alpha^2 \pi M_\pi^3} \Gamma(\pi \to \gamma \gamma)} = 0.274 \pm 0.010 \text{ GeV}^{-1}, \quad 0.274 \text{ GeV}^{-1}, \tag{55}
$$

$$
F_{\eta\gamma}(0) = \sqrt{\frac{4}{\alpha^2 \pi M_\eta^3} \Gamma(\eta \to \gamma\gamma)} = 0.273 \pm 0.009 \text{ GeV}^{-1}, \quad 0.277 \text{ GeV}^{-1}, \tag{56}
$$

$$
F_{\eta\prime\gamma}(0) = \sqrt{\frac{4}{\alpha^2 \pi M_{\eta\prime}^3} \Gamma(\eta\prime \to \gamma\gamma)} = 0.342 \pm 0.006 \text{ GeV}^{-1}, \quad 0.343 \text{ GeV}^{-1}. \tag{57}
$$

2. The $Q^2 \to \infty$ limiting behavior of $Q^2 F_{\gamma\gamma^* \to P}(0) F_{\gamma\gamma^* \to P}(Q^2)$ [\[1,](#page-19-0) [8](#page-19-7), [33](#page-21-1)]:

$$
\lim_{Q^2 \to \infty} \pi^2 Q^2 F_{\gamma \gamma^* \to \pi}(0) F_{\gamma \gamma^* \to \pi}(Q^2) = \frac{1}{2}, \quad 0.49,\tag{58}
$$

$$
\lim_{Q^2 \to \infty} 3\pi^2 Q^2 F_{\gamma\gamma^* \to \eta_8}(0) F_{\gamma\gamma^* \to \eta_8}(Q^2) = \frac{1}{2}, \quad 0.48,\tag{59}
$$

$$
\lim_{Q^2 \to \infty} 3\pi^2 Q^2 F_{\gamma\gamma^* \to \eta_0}(0) F_{\gamma\gamma^* \to \eta_0}(Q^2) = 4, 3.99. \tag{60}
$$

3. The $Q^2 \to \infty$ limiting behavior of $Q^2 F_{\gamma\gamma^* \to P}(Q^2)$ [\[1](#page-19-0), [8](#page-19-7), [33](#page-21-1)]:

$$
\lim_{Q^2 \to \infty} Q^2 F_{\gamma \gamma^* \to \pi} (Q^2) = 2f_{\pi} = 184.8 \pm 0.2 \text{ MeV}, \quad 184.8 \text{ MeV}, \tag{61}
$$

$$
\lim_{Q^2 \to \infty} Q^2 F_{\gamma \gamma^* \to \eta_8}(Q^2) = \frac{2}{\sqrt{3}} f_8 = 101 \pm 7 \text{ MeV}, \quad 95 \text{ MeV}, \tag{62}
$$

$$
\lim_{Q^2 \to \infty} Q^2 F_{\gamma \gamma^* \to \eta_0}(Q^2) = \frac{4\sqrt{2}}{\sqrt{3}} f_0 = 334 \pm 15 \text{ MeV}, \quad 332 \text{ MeV}, \tag{63}
$$

in which the weak decay constant $f_{\pi} = 92.4 \text{ MeV}$ is defined [\[34\]](#page-21-2) from $\pi \to \mu \nu$ decay.

These constraints are not completely independent, but are necessary since some of them are free from uncertainties, for example, Eqs. $(59-60)$ are free from the decay constants f_0 and f_8 . Combined with consideration of other properties of the pion [\[3](#page-19-2)], we can obtain $m_q = 200 \text{ MeV}, m_s = 550 \text{ MeV}, \beta_\pi = \beta_8 = 410 \text{ MeV}, \beta_0 = 475 \text{ MeV}, A_\pi = 0.0475 \text{ MeV}^{-1},$ $A_8 = 0.0331 \text{ MeV}^{-1}$, and $A_0 = 0.0440 \text{ MeV}^{-1}$. Among these 7 parameters, 3 of them $(m_q,$ A_{π} and β_{π}) are the same in our previous work [\[3](#page-19-2)] and have already been fixed, only the other 4 are new parameters. These three parameters satisfy Eqs. [\(55\)](#page-13-1), [\(58\)](#page-13-0) and [\(61\)](#page-14-0) very well. Then we fix the 4 new parameters by using the four equations Eqs. [\(56\)](#page-13-1), [\(57\)](#page-13-1), [\(59\)](#page-13-0) and [\(60\)](#page-13-0). Since the parameter fixing scheme is somehow unique, numerical results of these parameters do not have much room to vary, and not surprisingly we find these fixed 7 parameters give very good prediction for Eqs. [\(62\)](#page-14-0)-[\(63\)](#page-14-0). Reversely, we can compute the values of the above nine constraints by using the above seven fixed parameters, and we also provide the fitted values at the end of each equation. Therefore, after this simple parameter fixing scheme, we could start to calculate the transition form factor for these mesons.

The results are in good agreement with the experimental data which we have listed above. Moreover, it is interesting to notice that the masses of the light-flavor quarks (the up quarks and down quarks) from the above constrains are just in the correct range (e.g., 200 \sim 300 MeV) of the constituent quark masses from more general considerations. Naturally, the transition form factor results emerging from this assumption are in quite good agreement with the experimental data.

Fig. 2 indicates that the theoretical values of the photon-pion $(\gamma\gamma^*\to\pi)$ transition form factors in the case of low Q^2 fit the experimental data well. One may consider this work as a light-cone version of relativistic quark model [\[20](#page-20-3), [21\]](#page-20-4), which should be valid in the low-energy scale about $Q^2 \leq 2 \text{ GeV}^2$. However, it is also physically in accordance with the light-cone perturbative QCD approach [\[1,](#page-19-0) [2](#page-19-1)], which is applicable at the high-energy scale of

FIG. 2: Theoretical prediction of $(4\pi\alpha)^2 \frac{m_\pi^3}{64\pi} |F_{\gamma\gamma^*\to\pi}(Q^2)|^2$ calculated with the pion wave function in the BHL prescription compared with the experimental data. The data for the transition form factor are taken from Ref. [\[5](#page-19-4)].

FIG. 3: Theoretical prediction of $Q^2|F_{\gamma\gamma^*\to\pi}(Q^2)|$ calculated with the pion wave function in the BHL prescription compared with the experimental data. The data for the transition form factor are taken from Ref. [\[4](#page-19-3)] and Ref. [\[5](#page-19-4)].

FIG. 4: Theoretical prediction of $(4\pi\alpha)^2 \frac{m_\eta^3}{64\pi} |F_{\gamma\gamma^* \to \eta}(Q^2)|^2$ compared with the experimental data in the low energy scale. The data for the transition form factor are taken from Ref. [\[5\]](#page-19-4) and Ref. [\[35](#page-21-3)].

FIG. 5: Theoretical prediction of $Q^2|F_{\gamma\gamma^*\to\eta}(Q^2)|$ compared with the experimental data. The data for the transition form factor are taken from Ref. [\[4](#page-19-3)], Ref. [\[5](#page-19-4)] and Ref. [\[35](#page-21-3)].

FIG. 6: Theoretical prediction of $(4\pi\alpha)^2 \frac{m_{\eta\prime}^3}{64\pi} |F_{\gamma\gamma^* \to \eta\prime}(Q^2)|^2$ compared with the experimental data in the low energy scale. The data for the transition form factor are taken from Ref. [\[4\]](#page-19-3), Ref. [\[5](#page-19-4)], Ref. [\[35\]](#page-21-3) and Ref. [\[36](#page-21-4)].

FIG. 7: Theoretical prediction of $Q^2|F_{\gamma\gamma^*\to\eta'}(Q^2)|$ compared with the experimental data. The data for the transition form factor are taken from Ref. [\[4](#page-19-3)], Ref. [\[5](#page-19-4)], Ref. [\[35\]](#page-21-3) and Ref. [\[36](#page-21-4)].

 $Q^2 > 2 \text{ GeV}^2$. The reason is that the hard-gluon exchange between quark and antiquark of the meson, which should be generally considered at high $Q²$ for exclusive processes, is not necessary to be incorporated in the light-cone perturbative QCD approach for pion-photon transition form factor [\[1](#page-19-0), [2](#page-19-1)]. As a result, there is no wonder that our predictions for the transition form factor at high Q^2 also agree with the experimental data at high energy scale in Fig. 3.

Fig. 4 and Fig. 5 show that our predictions for the $\gamma^*\gamma \to \eta$ transition form factors agree with the experimental data in the low and high energy scale, respectively. In addition, the numerical results of $\gamma^* \gamma \to \eta'$ transition form factor also give good fit of the experiments both in the low and moderately high energy scale in Fig. 6 and Fig. 7. The prediction that we have made for the low Q^2 (0.001 – 0.5 GeV²) behaviors of the photon-meson transition form factors of π^0 , η and η' are measurable in $e + A(Nucleus) \rightarrow e + A + M$ process via Primakoff effect at JLab and DESY.

Generally speaking, the medium to high Q^2 behavior of the transition form factors should include leading-twist order (so-called pQCD picture) and NLO corrections [\[37,](#page-21-5) [38\]](#page-21-6), but we only take the leading order into account in this literature. However, we find that our results for the leading order of the transition form factors fit the experimental data at small Q^2 well and are also physically consistent with the light-cone pQCD approach at large Q^2 .

V. CONCLUSION

The light-cone formalism provides a convenient framework for the relativistic description of hadrons in terms of quark and gluon degrees of freedom, and for the application of perturbative QCD to exclusive processes. With the minimal Fock-state expansions of the pion and photon wave functions from the light-cone representation of the spin structure of the pseudoscalar meson and photon vertexes, we investigate the photon-meson transition form factors by adopting the Drell-Yan-West assignment to get the light-cone framework that works at both low Q^2 and high Q^2 . We employ the experimental values of the decay widths of π , η and η' , the limiting behavior of $\lim_{Q^2 \to \infty} Q^2 F_{\gamma^* \gamma \to M}(Q^2) F_{\gamma^* \gamma \to M}(0)$ $(M = \pi, \eta_8, \eta_0)$, and the limiting behavior of $Q^2 F_{\gamma^* \gamma \to M}(Q^2)$ as the nine constrains to fix those seven parameters in the π , η_8 , and η_0 wave functions. With the fixed π , η_8 , and η_0 wave functions, we find that our numerical predictions for the photon-meson transition form factors of light pseudoscalar

mesons π , η , and η' agree with the experimental data at both low and high energy scale, in a wide region comparing to previous studies.

Acknowledgements We acknowledge the helpful and encouraging communications with Liping Gan. This work is partially supported by National Natural Science Foundation of China under Grant Numbers 10025523 and 90103007. It is also supported by Hui-Chun Chin and Tsung-Dao Lee Chinese Undergraduate Research Endowment (Chun-Tsung Endowment) at Peking University.

- [1] G.P. Lepage and S.J. Brodsky, Phys. Rev. D 22, 2157 (1980).
- [2] F.-G. Cao, T. Huang, and B.-Q. Ma, Phys. Rev. D 53, 6582 (1996).
- [3] B.-W. Xiao and B.-Q. Ma, Phys. Rev. D 68, 034020 (2003).
- [4] CLEO Collaboration, J. Gronberg et al., Phys. Rev. D 57, 33 (1998).
- [5] CELLO Collaboration, H.-J. Behrend et al., Z. Phys. C 49, 401 (1991).
- [6] L. Gan, private communications, and proposal by PrimEx Collaboration at JLab.
- [7] HERMES Collaboration, K. Ackerstaff et al., Nucl. Instr. and Meth. A 417, 230 (1998).
- [8] F.-G. Cao and A.I. Signal, Phys. Rev. D 60, 114012 (1999).
- [9] T. Feldmann, P. Kroll, and B. Stech, Phys. Lett. B 449, 339 (1999).
- [10] J.L. Goity, A.M. Bernstein and B.R. Holstein, Phys. Rev. D 66, 076014 (2002).
- [11] P. Maris and P.C. Tandy, Phys. Rev. C65, 045211 (2002); F.J. Llanes-Estrada and S.R. Cotanch, Nucl. Phys. A 697, 303 (2002).
- [12] Ta-Pei Cheng and Ling-Fong Li, "Gauge theory of elementary particle physics" (Claredon: Oxford, 1984).
- [13] N. Beisert and B. Borasoy Eur. Phys. J. A 11, 329 (2001); P. Herrera-Siklody, J.L. Latorre, P. Pascual and J. Taron, Phys. Lett. B 419, 326 (1998). Ll. Ametller, J. Bijnens, A. Bramon, and F. Cornet, Phys. Rev. D 45, 986 (1992).
- [14] V.V. Anisovich, L.G. Dakhno, V.N. Markov, V.A. Nikonov, A.V. Sarantsev, [hep-ph/0410361;](http://arxiv.org/abs/hep-ph/0410361) A.V. Anisovich, V.V. Anisovich, L.G. Dakhno, V.A. Nikonov, A.V. Sarantsev. [hep-ph/0406320.](http://arxiv.org/abs/hep-ph/0406320)
- [15] S. Gieseke and C.F. Qiao, Phys. Rev. D 61, 074028 (2000).
- [16] S.J. Brodsky, in Lectures on Lepton Nucleon Scattering and Quantum Chromodynamics, edited

by A. Jaffe, D. Ruelle (Birkh \overline{a} user, Boston, 1982), p.255;

S.J. Brodsky and G.P. Lepage, in Perturbative Quantum Chromodynamics, edited by A.H. Mueller, (World Scientific, Singapore, 1989), p.93;

S.J. Brodsky, H-C. Pauli, and S.S. Pinsky, Phys. Rep. 301, 299 (1998).

- [17] S.J. Brodsky and S.D. Drell, Phys. Rev. D 22, 2236 (1980).
- [18] S.J. Brodsky, D.S. Hwang, B.-Q. Ma, and I. Schmidt, Nucl. Phys. B 593, 311 (2001).
- [19] S.J. Brodsky, T. Huang, and G.P. Lepage, in Particles and Fields-2, Proceedings of the Banff Summer Institute, Banff, Alberta, 1981, edited by A.Z. Capri and A.N. Kamal (Plenum, New York, 1983), p. 143.
- [20] B.-Q. Ma, Z. Phys. A 345, 321 (1993); T. Huang, B.-Q. Ma, and Q.-X. Shen, Phys. Rev. D 49, 1490 (1994); B.-Q. Ma and T. Huang, J. Phys. G: Nucl. Part. Phys. 21, 765 (1995); F.-G. Cao, J. Cao, T. Huang, and B.-Q. Ma, Phys. Rev. D 55, 7107 (1997).
- [21] B.-W. Xiao, X. Qian, and B.-Q. Ma, Eur. Phys. J. A 15, 523 (2002).
- [22] B.-Q. Ma, J.Phys.G: Nucl. Part. Phys. 17, L53 (1991); B.-Q. Ma and Q.-R. Zhang, Z. Phys. C 58, 479 (1993).
- [23] B.-Q. Ma, Phys. Lett. B 375, 320 (1996); B.-Q. Ma and A. Schäfer, Phys. Lett. B 378, 307 (1996); B.-Q. Ma, I. Schmidt, and J. Soffer, Phys. Lett. B 441, 461 (1998); B.-Q. Ma, I. Schmidt, and J.-J. Yang, Eur. Phys. J. A 12, 353 (2001).
- [24] B.-Q. Ma and I. Schmidt, Phys. Rev. D 58, 096008 (1998).
- [25] A. Donnachie, J. Gravelis, and G. Shaw, Phys. Rev. D 63, 114013 (2001).
- [26] C.-W. Hwang, Eur. Phys. J. C 19, 105 (2001); H.M. Choi and C.R. Ji, Nucl. Phys. A 618, 291 (1997).
- [27] S.D. Drell and T.-M. Yan, Phys. Rev. Lett. 24, 181 (1970); G. West, Phys. Rev. Lett. 24, 1206 (1970).
- [28] T. Feldmann and P. Kroll, Eur. Phys. J. C 5, 327 (1998).
- [29] J. Cao, F.-G. Cao, T. Huang, and B.-Q. Ma, Phys. Rev. D 58, 113006 (1998).
- [30] A.V. Radyushkin and R. Ruskov, Phys. Lett. B 374, 173 (1996).
- [31] See, e.g., R. Jakob, P. Kroll, and M. Raulfs, J. Phys.G: Nucl. Part. Phys. 22, 45 (1996); V.V. Anisovich, D.I. Melikhov, and V.A. Nikonov, Phys. Rev. D 55, 2918 (1997);

I.V. Musatov and A.V. Radyushkin, Phys. Rev. D 56, 2713 (1997).

- [32] Particle Data Group, K. Hagiwara et al., Phys. Rev. D 66, 010001 (2002).
- [33] S.J. Brodsky and G.P. Lepage, Phys. Rev. D 24, 1808 (1981).
- [34] C. Caso et al., Eur. Phys. J. C 3, 1 (1998).
- [35] TPC/Two-Gamma Collaboration, H. Aihara et al., Phys. Rev. Lett. 64, 172 (1990).
- [36] L3 Collaboration, M. Acciarri et al., Phys. Lett. B 418, 399 (1998).
- [37] E. Braaten, Phys. Rev. D 28, 524 (1983).
- [38] P. Kroll and K. Passek-Kumericki, Phys. Rev. D 67, 054017 (2003).