# New gauge bosons from the littlest Higgs model and the process $e^+e^- \rightarrow t\bar{t}$

Chong-Xing Yue, Lei Wang, Jian-Xing Chen

Department of Physics, Liaoning Normal University, Dalian 116029. P.R.China \*

July 1, 2018

### Abstract

In the context of the littlest  $\operatorname{Higgs}(LH)$  model, we study the process  $e^+e^- \to t\bar{t}$ . We find that the new gauge bosons  $Z_H$  and  $B_H$  can produce significant correction effects on this process, which can be further enhanced by the suitably polarized beams. In most of the parameter space preferred by the electroweak precision data, the absolute value of the relative correction parameter  $R_{B_H}$  is larger than 5%. As long as  $1TeV \leq M_{Z_H} \leq 1.5TeV$  and  $0.3 \leq c \leq 0.5$ , the absolute value of the relative correction parameter  $R_{Z_H}$  is larger than 5%. With reasonable values of the parameters of the LH model, the possible signals of the new gauge bosons  $B_H$ and  $Z_H$  can be detected via the process  $e^+e^- \to t\bar{t}$  in the future LC experiments with the c.m. energy  $\sqrt{S} = 800GeV$ .  $B_H$  exchange and  $Z_H$  exchange can generate significantly corrections to the forward-backward asymmetry  $A_{FB}(t\bar{t})$  only in small part of the parameter space.

<sup>\*</sup>E-mail:cxyue@lnnu.edu.cn

### I. Introduction

Although the standard model(SM) that bases on the gauge group  $SU(2)_L \times U(1)_Y$  has been successful in describing the physics of electroweak interactions, the mechanism of the electroweak symmetry breaking(EWSB) and the origins of the masses of the elementary fermions are still unknown. Furthermore, its scalar sector suffers from the problems of triviality and unnaturalness, etc. Thus, it is quite possible that the SM is only an effective theory valid below some high energy scale. New physics(NP) should exist at energy scales around TeV.

Recently, a kind of theory for EWSB was proposed to solve the hierarchy between the TeV scale of possible NP and the electroweak scale v = 246GeV, which is known as "little Higgs models" [1,2,3]. The key feature of these models is that the Higgs boson is a pseudo-Goldstone boson of a global symmetry which is spontaneously broken at some higher scale f and thus is naturally light. EWSB is induced by a Coleman-Weinberg potential, which is generated by integrating out the heavy degrees of freedom. This type of models can be regarded as one of the important candidates of the NP beyond the SM.

A high energy  $e^+e^-$  linear collider(*LC*) will offer an opportunity to make precision measurement of the properties of the electroweak gauge bosons, top quarks, Higgs bosons and also to constrain *NP* [4]. In the *LC* experiments, top quark pairs are mainly produced from the S-channel exchange of the *SM* gauge bosons  $\gamma$  and *Z* via the process  $e^+e^- \rightarrow t\bar{t}$ [5]. The total cross section is of the order of 1pb, so that top quark pairs will be produced at large rates in a clean environment at *LC*. If we assume that the integrated luminosity  $\mathcal{L}_{int}$  is about  $100fb^{-1}$ , there will be several times  $10^4$  top quark pairs to be generated in the future *LC* experiments. Furthermore, the *QCD* and *EW* corrections to the process  $e^+e^- \rightarrow t\bar{t}$  are small and decrease as the centre-of-mass(c.m.) energy  $\sqrt{S}$  increasing. The option of longitudinally polarized beams can help to improve the measurement precision and reduce background in search for *NP*. Thus, theoretical calculations of new particles contributions to the process  $e^+e^- \rightarrow t\bar{t}$  are of much interest for testing of *NP*.

In general, the new gauge bosons are heavier than the current experimental limits on direct searches. However, these new particles may produce virtual effects on some physical

observable, which may be detected in the present or future high energy experiments. In Ref.[6], we discussed the possible of detecting the new gauge bosons  $Z_H$  and  $B_H$  predicted by the littlest Higgs(LH) model [1] in the future LC experiments with the c.m. energy  $\sqrt{S} = 500 GeV$  and the integrating luminosity  $\pounds_{int} = 340 f b^{-1}$  and both beams polarized via considering their contributions to the processes  $e^+e^- \rightarrow f\bar{f}$  with  $f = \tau, \mu, b$  and c. Since the masses of these fermions are largely smaller than the c.m. energy  $\sqrt{S}$ , we have neglected the masses of these fermions in our numerical estimations. Our results show that the new gauge bosons  $Z_H$  and  $B_H$  can indeed produce significant contributions to these process in most of the parameter space preferred by the electroweak precision data, which might be observable in the future LC experiments. The aim of this paper is to consider the contributions of the  $Z_H$  and  $B_H$  to the process  $e^+e^- \to t\bar{t}$  and discuss whether these new particles can be detected via this process in the future LC experiments with the c.m. energy  $\sqrt{S} = 800 GeV$  and the integrating luminosity  $\pounds_{int} = 580 f b^{-1}$ . We find that the absolute value of the relative correction parameter  $R_{B_H}$  generated by  $B_H$ exchange is larger than 8% in most of the parameter space of the LH model preferred by the electroweak precision data. As long as  $1TeV \leq M_{Z_H} \leq 1.5TeV$  and  $0.3 \leq c \leq 0.5$ , the absolute value of  $R_{Z_H}$  is larger than 5%. If we assume that the initial electron and positron beams are suitably polarized, the absolute values of the relative correction parameters  $R_{B_H}$  and  $R_{Z_H}$  can be enhanced. Thus, with reasonable values of the parameters of the LH model, the possible signals of the new gauge bosons  $B_H$  and  $Z_H$  can be detected in the future LC experiments with the c.m. energy  $\sqrt{S} = 800 GeV$ , which is similar to the conclusions given in Ref.[6]. We further calculate the contributions of these new gauge bosons to the forward-backward asymmetry  $A_{FB}(t\bar{t})$ . We find that they can generate significantly corrections to the forward-backward asymmetry  $A_{FB}(t\bar{t})$  only in small part of the parameter space.

In section II, we give the formula of the contributions of new gauge bosons  $B_H$  and  $Z_H$ to the process  $e^+e^- \rightarrow t\bar{t}$  and estimate the values of the relative corrections parameters  $R_{B_H} = \sigma^{B_H}(t\bar{t})/\sigma^{SM}(t\bar{t})$  and  $R_{Z_H} = \sigma^{Z_H}(t\bar{t})/\sigma^{SM}(t\bar{t})$ . The dependence of the relative correction parameters  $R_{B_H}$  and  $R_{Z_H}$  on the initial beam polarization is discussed in section III. In section IV, we calculate the contributions of these new gauge bosons to the forwardbackward asymmetry  $A_{FB}(t\bar{t})$ . Our conclusions and discussions are given in section V.

### II. Corrections of the new gauge bosons $B_H$ and $Z_H$ to the process $e^+e^- \rightarrow t\bar{t}$

The *LH* model [1] is one of the simplest and phenomenologically viable models, which realizes the little Higgs idea. It consists of a non-linear  $\sigma$  model with a global SU(5)symmetry, which is broken down to its subgroup SO(5) by a vacuum condensate  $f \sim \Lambda s/4\pi \sim TeV$ . At the same time, the locally gauged group  $SU(2)_1 \times U(1)_1 \times SU(2)_2 \times U(1)_2$  is broken to its diagonal subgroup  $SU(2) \times U(1)$ , identified as the *SM* electroweak gauge group. This breaking scenario gives rise to four massive gauge bosons  $B_H$ ,  $Z_H$ , and  $W_H^{\pm}$ , which might produce characteristic signatures at the present and future high energy collider experiments [7,8,9].

Taking account of the gauge invariance of the Yukawa coupling and the U(1) anomaly cancellation, the coupling expressions of the gauge bosons  $B_H$  and  $Z_H$  to ordinary particles, which are related to our calculation, can be written as [7]:

$$g_V^{B_H ee} = \frac{3e}{4C_w s' c'} (c'^2 - \frac{2}{5}), \qquad \qquad g_A^{B_H ee} = \frac{e}{4C_w s' c'} (c'^2 - \frac{2}{5}); \qquad (1)$$

$$g_V^{B_H tt} = \frac{e}{2C_w s' c'} \left[\frac{5}{6} \left(\frac{2}{5} - c'^2\right) - \frac{1}{5} x_L\right], \ g_A^{B_H tt} = \frac{e}{2C_w s' c'} \left[\frac{1}{2} \left(\frac{2}{5} - c'^2\right) - \frac{1}{5} x_L\right];$$
(2)

$$g_V^{Z_H ee} = -\frac{ec}{4S_w s}, \qquad \qquad g_A^{Z_H ee} = \frac{ec}{4S_w s}; \qquad (3)$$

$$g_V^{Z_H tt} = \frac{ec}{4S_w s}, \qquad \qquad g_A^{Z_H tt} = -\frac{ec}{4S_w s}. \tag{4}$$

Where  $S_w = sin\theta_w$ ,  $\theta_w$  is the Weinberg angle. Using the mixing parameters  $c(s = \sqrt{1 - c^2})$ and  $c'(s' = \sqrt{1 - c'^2})$ , we can represent the SM gauge coupling constants as  $g = g_1 s = g_2 c$ and  $g' = g'_1 s' = g'_2 c'$ . The mixing angle parameter between the SM top quark t and the vector-like quark T is defined as  $x_L = \lambda_1^2/(\lambda_1^2 + \lambda_2^2)$ , in which  $\lambda_1$  and  $\lambda_2$  are the Yukawa coupling parameters.

Global fits to the eletroweak precision data produce rather severe constraints on the parameter space of the *LH* model [10]. However, if the *SM* fermions are charged under  $U(1)_1 \times U(1)_2$ , the constraints become relaxed. The scale parameter  $f = 1 \sim 2TeV$  is allowed for the mixing parameters c, c', and  $x_L$  in the ranges of  $0 \sim 0.5, 0.62 \sim 0.73$ , and

 $0.3 \sim 0.6$ , respectively [11]. In this case, the masses of  $B_H$  and  $Z_H$  are allowed in the ranges of  $300 GeV \sim 900 GeV$  and  $1TeV \sim 3TeV$ , respectively. Thus, we will take the  $Z_H$  mass  $M_{Z_H}$ ,  $B_H$  mass  $M_{B_H}$  and the mixing parameters c, c' and  $x_L$  as free parameters in our calculation.



Figure 1: The relative correction parameter  $R_{B_H}$  as a function of the  $B_H$  mass  $M_{B_H}$  for different values of the mixing parameters c' and  $x_L$ .

For the SM, top quark pair  $t\bar{t}$  can be produced in sufficient abundance in the LC experiments. The main production mechanism proceed at the Born level by the S-channel annihilation of an initial electron-position pair into virtual photon or neutral gauge boson

Z, and their subsequent splitting into top quark pairs,  $e^+e^- \rightarrow \gamma, Z \rightarrow t\bar{t}$ . For the LH model, the  $B_H$  exchange and  $Z_H$  exchange can also produce the top quark pairs. The production cross sections can be written as:

$$\begin{split} \sigma^{B_{H}}(t\bar{t}) &= \frac{N_{c}^{f}\beta}{8\pi S} \{ (1 - \frac{\beta^{2}}{3}) \frac{4}{3} e^{2} g_{V}^{B_{H}ee} g_{V}^{B_{H}tt} \frac{S(M_{B_{H}}^{2} - S)}{(S - M_{B_{H}}^{2})^{2} + M_{B_{H}}^{2} \Gamma_{B_{H}}^{2}} \\ &+ [(g_{V}^{B_{H}ee})^{2} + (g_{A}^{B_{H}ee})^{2}] [(1 - \frac{\beta^{2}}{3})](g_{V}^{B_{H}tt})^{2} + (g_{A}^{B_{H}t})^{2}] - (1 - \beta^{2})(g_{A}^{B_{H}tt})^{2}] \\ &\frac{S^{2}}{(S - M_{B_{H}}^{2})^{2} + M_{B_{H}}^{2} \Gamma_{B_{H}}^{2}} + (g_{V}^{Zee} g_{V}^{B_{H}ee} + g_{A}^{Zee} g_{A}^{B_{H}ee}) \\ &[(1 - \frac{\beta^{2}}{3})(g_{V}^{Ztt} g_{V}^{B_{H}tt} + g_{A}^{Ztt} g_{A}^{B_{H}t}) - (1 - \beta^{2})(g_{A}^{B_{H}t})(g_{A}^{Ztt})] \\ &\frac{2S^{2}[(S - M_{Z}^{2})(S - M_{B_{H}}^{2}) + M_{Z}\Gamma_{Z}M_{B_{H}}\Gamma_{B_{H}}]}{[(S - M_{Z}^{2})^{2} + M_{Z}^{2}\Gamma_{Z}^{2}][(S - M_{B_{H}}^{2})^{2} + M_{B_{H}}^{2}\Gamma_{B_{H}}^{2}]} \},$$
(5)  
$$\sigma^{Z_{H}}(t\bar{t}) = \frac{N_{c}^{f}\beta}{8\pi S} \{(1 - \frac{\beta^{2}}{3})\frac{4}{3}e^{2}g_{V}^{ZHee} g_{V}^{ZHet} \frac{S(M_{Z_{H}}^{2} - S)}{(S - M_{Z_{H}}^{2})^{2} + M_{Z}^{2}}\Gamma_{Z_{H}}^{2}} \\ &+ [(g_{V}^{ZHee})^{2} + (g_{A}^{ZHee})^{2}][(1 - \frac{\beta^{2}}{3})[(g_{V}^{ZHt})^{2} + (g_{A}^{ZHt})^{2}] - (1 - \beta^{2})(g_{A}^{ZHt})^{2}] \\ &\frac{S^{2}}{(S - M_{Z_{H}}^{2})^{2} + M_{Z_{H}}^{2}}\Gamma_{Z_{H}}^{2}} + (g_{V}^{Zee} g_{V}^{ZHee} + g_{A}^{Zee} g_{A}^{ZHee}) \\ [(1 - \frac{\beta^{2}}{3})(g_{V}^{ZH} g_{V}^{ZHt} + g_{A}^{ZH} g_{A}^{ZHt}) - (1 - \beta^{2})(g_{A}^{ZHt})(g_{A}^{ZHt})] \\ &\frac{2S^{2}[(S - M_{Z_{H}}^{2})(S - M_{Z_{H}}^{2}) + M_{Z}}{[(S - M_{Z_{H}}^{2})^{2} + M_{Z}^{2}}\Gamma_{Z_{H}}^{2}]} \}$$
(6)

with

$$g_V^{Zee} = \frac{e}{4S_w C_w} (-1 + 4S_w^2), \qquad \qquad g_A^{Zee} = \frac{e}{4S_w C_w}$$
(7)

$$g_V^{Ztt} = \frac{e}{4S_w C_w} (1 - \frac{8}{3}S_w^2), \qquad \qquad g_A^{Ztt} = \frac{e}{4S_w C_w}, \qquad (8)$$

where  $\beta = \sqrt{1 - \frac{4m_t^2}{S}}$ ,  $m_t$  is the top quark mass.  $\Gamma_i$  represent the total decay widths of the gauge bosons  $Z, Z_H$ , and  $B_H$ .  $\Gamma_{Z_H}$  and  $\Gamma_{B_H}$  have been given in Ref.[6]. From above equations, we can see that  $\sigma^{B_H}(t\bar{t})$  mainly dependents the free parameters  $M_{B_H}$ , c' and  $x_L$ , while  $\sigma^{Z_H}(t\bar{t})$  only dependents the free parameters c and  $M_{Z_H}$ , which is differently from those for the process  $e^+e^- \to f\bar{f}$  with  $f = \tau, \mu, b$  and c. In that case, the contributions of the gauge bosons  $B_H$  is independent of the mixing parameter  $x_L$ . Thus, in this paper, we will take the mixing parameters c, c' and  $x_L$  as free parameters. Certainly, due to the mixing between the gauge bosons Z and  $Z_H$ , the SM tree-level couplings  $Ze\bar{e}$  and  $Zt\bar{t}$  receive corrections at the order of  $v^2/f^2$ , which can also produce contributions to the production cross section of the process  $e^+e^- \rightarrow t\bar{t}$ . However, the contributions are suppressed by the factor  $v^4/f^4$ , which are smaller than those of  $B_H$  or  $Z_H$ . Thus, we have neglected this kind of corrections in above equations.



Figure 2: The relative correction parameter  $R_{Z_H}$  as a function of the  $Z_H$  mass  $M_{Z_H}$  for three values of the mixing parameter c.

To see the correction effects of  $B_H$  exchange and  $Z_H$  exchange on the  $t\bar{t}$  production cross section, we plot the relative correction parameters  $R_{B_H} = \sigma^{B_H}(t\bar{t})/\sigma^{SM}(t\bar{t})$  and  $R_{Z_H} = \sigma^{Z_H}(t\bar{t})/\sigma^{SM}(t\bar{t})$  as functions of  $M_{B_H}$  and  $M_{Z_H}$  in Fig.1 and Fig.2, respectively. From these figures, we can see that the gauge boson  $Z_H$  decreases the  $SM t\bar{t}$  production cross section  $\sigma^{SM}(t\bar{t})$  in all of the parameter space, which satisfies the electroweak precision constraints. In most part of the parameter space, the absolute value of the relative correction parameter  $R_{Z_H}$  is smaller than 5%, which is very difficult to be detected in the future LC experiments. This is consistent with the contributions of  $Z_H$  to the process  $e^+e^- \to f\bar{f}$ , which has been studied in Ref.[6]. However, for the gauge boson  $B_H$ , it is not this case. For  $M_{B_H} \leq 800 GeV$ ,  $B_H$  exchange produce positive corrections to the  $t\bar{t}$  production cross section  $\sigma^{SM}(t\bar{t})$  and the value of  $R_{B_H}$  increase as  $M_{B_H}$ ,  $x_L$  and c' increasing. For  $800 GeV < M_{B_H} \leq 900 GeV$ ,  $B_H$  exchange decrease the cross section  $\sigma^{SM}(t\bar{t})$  and the absolute of  $R_{B_H}$  increase as  $M_{B_H}$  decreasing and  $x_L$ , c' increasing. The peak of the  $R_{B_H}$  resonance emerges when the  $B_H$  mass  $M_{B_H}$  is approximately equal to the c.m. energy  $\sqrt{S} = 800 GeV$ . In most part of the parameter space, the absolute value of  $R_{B_H}$  is larger than 8%. Thus, the virtual effects of  $B_H$  on the process  $e^+e^- \to t\bar{t}$  should be easy detected in the future LC experiment with  $\sqrt{S} = 800 GeV$  and  $\pounds_{int} = 580 fb^{-1}$ .

## III. The dependence of the relative correction parameters $R_{B_H}$ and $R_{Z_H}$ on the electron and positron beam polarization

An *LC* has a large potential of the discovery of new particles and is well suited for the precise analysis of *NP* beyond the *SM*. At present, the existing proposals are designed with high luminosity of about  $\mathcal{L}_{int} = 340 f b^{-1}$  at  $\sqrt{S} = 500 GeV$  and  $\mathcal{L}_{int} = 580 f b^{-1}$  at  $\sqrt{S} = 800 GeV$  [4]. An important tool of an *LC* is the use of polarized beams. Beam polarization is not only useful for a possible reduction of the background, but might also serve as a possible tool to disentangle different contributions to the signal and lead to substantial enhancement of the produce cross sections of some processes [12]. To see whether the contributions of the new gauge bosons  $B_H$  and  $Z_H$  to the process  $e^+e^- \to t\bar{t}$  can indeed be detected, we discuss the dependence of the relative correction parameters  $R_{B_H}$  and  $R_{Z_H}$  on the initial electron and positron beam polarization in this section.

Considering the polarization of the initial electron and positron beams, the cross section of the process  $e^+e^- \rightarrow t\bar{t}$  can be generally written as:

$$\sigma(t\bar{t}) = (1 + P_e)(1 - P_{\bar{e}})(\sigma_{RR}(t\bar{t}) + \sigma_{RL}(t\bar{t})) + (1 - P_e)(1 + P_{\bar{e}})(\sigma_{LL}(t\bar{t}) + \sigma_{LR}(t\bar{t})), \quad (9)$$

where  $P_e$  and  $P_{\bar{e}}$  are the degrees of longitudinal electron and position polarization, respectively.  $\sigma_{ij}$  are the chiral cross sections of this process. The relative correction parameters  $R_{B_H}$  and  $R_{Z_H}$  are plotted as functions of  $M_{B_H}$  and  $M_{Z_H}$  for c' = 0.65,  $x_L = 0.5$ , c = 0.3 and different beam polarizations in Fig.3 and Fig.4, respectively. In these two figures, we have used the solid line, dashed line, and dotted line to represent  $(P_e, P_{\bar{e}})=(0,0)$ , (0.8, -0.6), and (-0.8, 0.6), respectively. Our calculation results show that the absolute values of  $R_{B_H}[R_{Z_H}]$  for  $(P_e, P_{\bar{e}}) = (0.8, 0.6)[(-0.8, -0.6)]$  are smaller than those for  $(P_e, P_{\bar{e}}) = (0, 0)$ . Thus, in Fig.3 and Fig.4 we do not plot these lines.



Figure 3: The relative correction parameter  $R_{B_H}$  as a function of the  $B_H$  mass  $M_{B_H}$  for  $c' = 0.65, x_L = 0.5, \text{ and } (P_e, P_{\bar{e}}) = (0, 0), (0.8, -0.6), (-0.8, 0.6).$ 

From Fig.3 and Fig.4 we can see that the suitably polarized beams can indeed enhance the virtual effects of the new gauge bosons  $B_H$  and  $Z_H$  on the process  $e^+e^- \rightarrow t\bar{t}$ . In the whole parameter space preferred by the electroweak precision data, the value of  $R_{B_H}$  for  $(P_e, P_{\bar{e}}) = (0.8, -0.6)$  is larger than that for  $(P_e, P_{\bar{e}}) = (0, 0)$ , while the absolute values of  $R_{Z_H}$  for  $(P_e, P_{\bar{e}}) = (-0.8, 0.6)$  is larger than that for  $(P_e, P_{\bar{e}}) = (0, 0)$ . Varying the values of the free parameters c',  $x_L$ , and c does not change this conclusion. So, in Fig.3 and Fig.4 we have taken these parameters for fixed values  $x_L = 0.5$ , c' = 0.65, and c = 0.3. Certainly, the values of  $R_{B_H}$  and  $R_{Z_H}$  change as the values of these parameters varying. For example, for  $0.3 \le c \le 0.5$  and  $1TeV \le M_{Z_H} \le 2TeV$ , the absolute value of  $R_{Z_H}$  for  $(P_e, P_{\bar{e}}) = (-0.8, 0.6)$  is larger than 6%. The absolute of  $R_{B_H}$  for  $(P_e, P_{\bar{e}}) = (0.8, -0.6)$  is larger than 5% for  $x_L = 0.5$ ,  $0.68 \le c' \le 0.73$  and  $500GeV < M_{B_H} \le 900GeV$ , but for  $x_L = 0.6$  its value is larger than 5% for  $0.65 \le c' \le 0.73$  and  $450GeV \le M_{B_H} \le 900GeV$ . Thus, using the suitably polarization of the initial electron and positron beams, it is more easy to detect the possible signals of the new gauge bosons  $B_H$  and  $Z_H$  in the future LC experiments.



Figure 4: The relative correction parameter  $R_{Z_H}$  as a function of the  $Z_H$  mass  $M_{Z_H}$  for c = 0.3 and  $(P_e, P_{\bar{e}}) = (0, 0), (0.8, -0.6), (-0.8, 0.6).$ 

### IV. Gauge bosons $B_H$ , $Z_H$ and the forward-backward asymmetry $A_{FB}(t\bar{t})$

The events generated by the process  $e^+e^- \rightarrow f\bar{f}$  can be characterized by the momentum direction of the emitted fermion. If we assume that the final state fermion travels forward(F) or backward(B) with respect to the electron beam, than the forward-backward asymmetry can be defined as:

$$A_{FB} = \frac{\sigma_F - \sigma_B}{\sigma_F + \sigma_B},\tag{10}$$

which is easier to be measured because only the identification of the charge of the fermion and the measurement of its direction are needed [13]. It can be measured for all tagged flavors and inclusively for hadrons. Thus, it is needed to calculate the contributions of  $B_H$  exchange and  $Z_H$  exchange to the forward-backward asymmetry  $A_{FB}(t\bar{t})$ .



Figure 5: The relative correction parameter  $R'_{B_H}$  as a function of  $M_{B_H}$  for different values of the mixing parameters c' and  $x_L$ .

The total formula of  $A_{FB}(t\bar{t})$  for the new gauge bosons  $B_H$  and  $Z_H$  including the contributions of the SM gauge bosons  $\gamma$  and Z can be written as:

$$A_{FB}^{B_{H}}(t\bar{t}) = \frac{M_{2}^{B_{H}}(t\bar{t})}{M_{1}^{B_{H}}(t\bar{t})}, \qquad A_{FB}^{Z_{H}}(t\bar{t}) = \frac{M_{2}^{Z_{H}}(t\bar{t})}{M_{1}^{Z_{H}}(t\bar{t})},$$
(11)

where

$$\begin{split} M_{2}^{Bn}(t\bar{t}) &= \beta \{ \frac{2e^{2}}{3} g_{A}^{2ee} g_{A}^{2tt} \frac{4S(M_{Z}^{2}-S)}{(S-M_{Z}^{2})^{2} + M_{Z}^{2}\Gamma_{Z}^{2}} \\ &+ \frac{2e^{2}}{3} g_{A}^{Bnee} g_{A}^{Bnut} \frac{4S(M_{Du}^{2}-S)}{(S-M_{Du}^{2})^{2} + M_{Du}^{2}\Gamma_{Du}^{2}} \\ &+ g_{V}^{Vec} g_{A}^{Zec} g_{V}^{Ze} g_{A}^{Zu} \frac{8S^{2}}{(S-M_{Z}^{2})^{2} + M_{Du}^{2}\Gamma_{Z}^{2}} \\ &+ g_{V}^{Bnee} g_{A}^{Bnee} g_{V}^{Bntt} g_{A}^{Bntt} \frac{8S^{2}}{(S-M_{Z}^{2})^{2} + M_{Du}^{2}\Gamma_{Z}^{2}} \\ &+ g_{V}^{Bnee} g_{A}^{Bnee} g_{V}^{Bntt} g_{A}^{Bntt} \frac{8S^{2}}{(S-M_{Du}^{2})^{2} + M_{Du}^{2}\Gamma_{Z}^{2}} \\ &+ (g_{V}^{Vee} g_{A}^{Dec} g_{V}^{Bntt} g_{A}^{Bntt} \frac{8S^{2}}{(S-M_{Du}^{2})^{2} + M_{Du}^{2}\Gamma_{Z}^{2}} \\ &+ (g_{V}^{Vee} g_{A}^{Bnee} g_{V}^{Bntt} g_{A}^{Bntt} \frac{8S^{2}}{(S-M_{Du}^{2})^{2} + M_{Du}^{2}\Gamma_{Z}^{2}} \\ &+ (g_{V}^{Vee} g_{A}^{Dec} g_{V}^{Bntt} g_{A}^{Bntt} \frac{8S^{2}}{(S-M_{Du}^{2})^{2} + M_{Du}^{2}\Gamma_{Z}^{2}} \\ &+ (g_{V}^{Vee} g_{A}^{Bnee} g_{V}^{Bntt} g_{A}^{Bntt} \frac{8S^{2}}{(S-M_{Du}^{2})^{2} + M_{Du}^{2}\Gamma_{Z}^{2}} ] \\ &+ (g_{V}^{Vee} g_{A}^{Dec} g_{V}^{Bnt} g_{A}^{Du} + g_{A}^{T} g_{A}^{Du} g_{A}^{Du} + g_{A}^{T} g_{A}^{Du} g_{A}^{Du} g_{A}^{Du} \\ &+ (g_{V}^{Vee} g_{A}^{Ane} g_{V}^{2} + M_{Z}^{2}\Gamma_{Z}^{2} + \frac{8e^{2}}{3} (1 - \frac{\beta^{2}}{3}) g_{V}^{Suee} g_{V}^{Sut} \frac{2S(M_{Z}^{2} - S)}{(S - M_{Du}^{2})^{2} + M_{Du}^{2}\Gamma_{Du}^{2}} \\ &+ (g_{V}^{Bnee} g_{V}^{Du} g_{V}^{Du} g_{V}^{Du} g_{V}^{Du} g_{V}^{Du} g_{V}^{Du} g_{A}^{Du} g_{A}^{D$$

$$\begin{split} M_{1}^{Z_{H}}(t\bar{t}) &= \left\{ \frac{16e^{4}}{9} \left(1 - \frac{\beta^{2}}{3}\right) + \frac{8e^{2}}{3} \left(1 - \frac{\beta^{2}}{3}\right) g_{V}^{Zee} g_{V}^{Ztt} \frac{2S(M_{Z}^{2} - S)}{(S - M_{Z}^{2})^{2} + M_{Z}^{2} \Gamma_{Z}^{2}} \right. \\ &\left. \left[ (g_{V}^{Zee})^{2} + (g_{A}^{Zee})^{2} \right] \left[ 4(1 - \frac{\beta^{2}}{3}) \left[ (g_{V}^{Ztt})^{2} + (g_{A}^{Zt})^{2} \right] - 4(1 - \beta^{2}) (g_{A}^{Ztt})^{2} \right] \times \right. \\ &\left. \frac{S^{2}}{(S - M_{Z}^{2})^{2} + M_{Z}^{2} \Gamma_{Z}^{2}} + \frac{8e^{2}}{3} \left(1 - \frac{\beta^{2}}{3}\right) g_{V}^{Z_{H}ee} g_{V}^{Z_{H}tt} \frac{2S(M_{Z_{H}}^{2} - S)}{(S - M_{Z_{H}}^{2})^{2} + M_{Z}^{2} \Gamma_{Z_{H}}^{2}} \right. \\ &\left. + \left[ (g_{V}^{Z_{H}ee})^{2} + (g_{A}^{Z_{H}ee})^{2} \right] \left[ 4(1 - \frac{\beta^{2}}{3}) \left[ (g_{V}^{Z_{H}tt})^{2} + (g_{A}^{Z_{H}tt})^{2} \right] - 4(1 - \beta^{2}) (g_{A}^{Z_{H}tt})^{2} \right] \right. \\ &\left. \frac{S^{2}}{(S - M_{Z_{H}}^{2})^{2} + M_{Z_{H}}^{2} \Gamma_{Z_{H}}^{2}} + \left( g_{V}^{Zee} g_{V}^{Z_{H}ee} + g_{A}^{Zee} g_{A}^{Z_{H}ee} \right) \right. \\ &\left. \left[ 4(1 - \frac{\beta^{2}}{3}) (g_{V}^{Ztt} g_{V}^{Z_{H}tt} + g_{A}^{Ztt} g_{A}^{Z_{H}tt}) - 4(1 - \beta^{2}) (g_{A}^{Ztt}) (g_{A}^{Ztt}) \right] \right. \\ &\left. \frac{2S^{2} [(S - M_{Z}^{2})(S - M_{Z_{H}}^{2}) + M_{Z} \Gamma_{Z} M_{Z_{H}} \Gamma_{Z_{H}}}]}{[(S - M_{Z}^{2})^{2} + M_{Z}^{2} \Gamma_{Z}^{2}] [(S - M_{Z_{H}}^{2})^{2} + M_{Z}^{2} \Gamma_{Z}^{2}]} \right\}. \end{split}$$

In above equations, we have assumed that the initial electron and positron beams are not polarized.

To see whether the new gauge bosons  $B_H$  and  $Z_H$  can produce significant deviations from the SM prediction value for  $A_{FB}(t\bar{t})$ , we plot the relative correction parameters  $R'_{B_H} = \delta A_{FB}^{B_H}(t\bar{t})/A_{FB}^{SM}(t\bar{t})$  and  $R'_{Z_H} = \delta A_{FB}^{Z_H}(t\bar{t})/A_{FB}^{SM}(t\bar{t})$  as functions of  $M_{B_H}$  and  $M_{Z_H}$ in Fig.5 and Fig.6, respectively. From Fig.5 and Fig.6 we can see that, in most of the parameter space preferred by the electroweak precision data, the absolute values of the relative correction parameters  $R'_{B_H}$  and  $R'_{Z_H}$  are smaller than 5%. The absolute values of  $R'_{Z_H}$  is larger than 5% only for the mixing parameter c = 0.5 and  $1TeV \leq M_{Z_H} \leq$ 1.4TeV.  $B_H$  exchange makes the deviation of the forward-backward asymmetry  $A_{F_B}(t\bar{t})$ from its SM value may be positive or negative, which depends on the  $B_H$  mass  $M_{B_H}$ . The resonance peak can emerge for  $M_{B_H} \approx 800GeV$ . Furthermore, the absolute value of  $R'_{B_H}$  increases as the mixing parameters c' and  $x_L$  increasing. For  $c' \geq 0.71$ ,  $x_L \geq 0.5$ , and  $600GeV \leq M_{B_H} \leq 1000GeV$ , the absolute value of  $R'_{B_H}$  is larger than 5%, which might be detected in the future LC experiments. However, for  $c' \leq 0.68$  and  $x_L \leq 0.4$ , except for a small region near  $M_{B_H} = 800GeV$ , the absolute value of  $R'_{B_H}$  is smaller than 5%.

Similar to above calculation, we can obtain the corrections of  $B_H$  exchange and  $Z_H$ 

exchange to the forward-backward symmetry  $A_{FB}(f\bar{f})$  with  $f=\mu$ ,  $\tau$ , b, or, c. From the coupling formula of the new gauge bosons  $B_H$  and  $Z_H$  to differently fermions given in Ref.[7], we can surmise that the conclusions are similar to those for  $A_{FB}(t\bar{t})$ . We have confirmed this expectation through explicit calculation. Certainly, the contributions of  $B_H$  exchange to  $A_{FB}(f\bar{f})$  mainly dependent on the free parameters  $M_{B_H}$  and c', while the contributions of  $B_H$  exchange to  $A_{FB}(t\bar{t})$  mainly dependent on the free parameters  $M_{B_H}$  and c', while  $M_{B_H}$ , c', and  $x_L$ .



Figure 6: The relative correction parameter  $R'_{Z_H}$  as a function of  $M_{Z_H}$  for three values of the mixing parameter c.

### V. Conclusions and discussins

An *LC* will be an ideal machine for precisely testing the *SM* and probing *NP* beyond the *SM*. Some kinds of *NP* predict the existence of new particles that will be manifested as a rather spectacular resonance in the *LC* experiments if the achievable c.m. energy  $\sqrt{S}$  is sufficient. Even if their masses exceed the c.m. energy  $\sqrt{S}$ , the *LC* experiments also retain an indirect sensitivity through a precision study their virtual corrections to observables.

It is widely believed that the top quark, with a mass of the order of the electroweak scale, will be a sensitive probe into NP beyond the SM. The quantum correction effects of the new particles to some SM processes involving top quark are more important than those for lighter fermions. Thus, the top quark plays a key role in the quest for deviations of observables from their SM predictions. On the other hand, top quark pairs can be copiously produced mainly through the process  $e^+e^- \rightarrow t\bar{t}$  in the future LC experiments. So, in this paper, we discuss and calculate the corrections of the new gauge bosons  $B_H$ and  $Z_H$  predicted by the LH model to the production cross section  $\sigma(t\bar{t})$  and the forwardbackward asymmetry  $A_{FB}(t\bar{t})$  of the process  $e^+e^- \rightarrow t\bar{t}$ .

The LH model has all essential features of the little Higgs models. So, in this paper, we give our numerical results in the context of the LH model, although many alternatives have been proposed [2,3]. We find that the new gauge bosons  $Z_H$  and  $B_H$  can produce significant correction effects on the process  $e^+e^- \rightarrow t\bar{t}$ , which can be further enhanced by the suitably polarized beams. In most of the parameter space f =  $1 TeV \sim 2 TeV, c'$  =  $0.62 \sim 0.73, c = 0.1 \sim 0.5$ , and  $x_L = 0.3 \sim 0.6$ , which consistent with the electroweak precision data, the absolute value of the relative correction parameter  $R_{B_H}$  generated by  $B_H$  exchange is larger than 5%. As long as  $1TeV \leq M_{Z_H} \leq 1.5TeV$ , and  $0.3 \leq c \leq 0.5$ , the absolute value of  $R_{Z_H}$  is larger than 5%. Thus, we can say that, with reasonable values of the parameters in the LH model, the possible signals of the new gauge bosons  $B_H$  and  $Z_H$  can be detected via the process  $e^+e^- \to t\bar{t}$  in the future LC experiments with the c.m. energy  $\sqrt{S} = 800 GeV$ . However,  $B_H$  exchange and  $Z_H$  exchange can only generate very small corrections to the forward-backward asymmetry  $A_{FB}(t\bar{t})$  in most of the parameter space. It is possible that, in very small range of the parameter space, the possible signals of  $B_H$  and  $Z_H$  might be detected via measuring the deviations of  $A_{FB}(t\bar{t})$ from its SM prediction.

The couplings of the new gauge boson  $B_H$  to fermions are quite model dependent, which depend on the choice of the fermion U(1) charges under the two U(1) groups. The U(1) charges of the SM fermions are constrained by requiring that the Yukawa couplings are gauge invariant and maintaining the usual SM hypercharge assignment. Combining the gauge invariance of the Yukawa couplings with the U(1) anomaly-free can fix all of the U(1) charge values. In this paper, we have used the couplings of the  $B_H$  to fermions, which come from this kind of choice. Certainly, this is only one example of all possible U(1) charge assignments. In other little Higgs models, several alternatives for the U(1) charge choice exist[2, 3, 10], the numerical results for the new gauge boson  $B_H$  obtained in this paper might be changed.

### Acknowledgments

This work was supported in part by the National Natural Science Foundation of China under the grant No.90203005 and No.10475037 and the Natural Science Foundation of the Liaoning Scientific Committee(20032101).

### References

- [1] N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, **JHEP 0207**(2002)034.
- [2] N. Arkani-Hamed, A. G. Cohen and H. Georgi, *Phys. Lett.* B513(2001)232; N. Arkani-Hamed, A. G. Cohen, T. Gregoire and J. G. Wacker, JHEP 0208(2002)020;
  N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, T. Gregoire and J. G. Wacker, JHEP 0208(2002)021; I. Low, W. Skiba and D. Smith, *Phys. Rev.* D66(2002)072001; M. Schmaltz, *Nucl. Phys. Proc. Suppl.* 117(2003)40; D. E. Kaplan and M. Schmaltz, JHEP 0310(2003)039.
- [3] J. G. Wacker, hep-ph/0208235; S. Chang and J. G. Wacker, Phys. Rev. D69(2004)035002; W. Skiba and J. Terning, Phys. Rev. D68(2003)075001; S. Chang, JHEP 12(2003)051; M. Schmaltz, JHEP 0408(2002)056.
- [4] T. Abe et al. [American Linear Collider Group], hep-ex/0106057; J. A. Aguilar-Saavedra et al. [ECFA/DESY Physics Working Group ], hep-ph/0106315; K. Abe et al. [ECFA linear Collider Working Group Collaboration], hep-ph/0109166.
- [5] J. Jersak, E. Laermann, P. M. Zerwas, *Phys. Rev.* D25(1982)1218; Erratum *Phys. Rev.* D36(1987)310; W. Beenakker, S. C. vander Marck, W. Hollik, *Nucl. Phys.* B365(1991)24; W. Beenkker, A. Denner, A. Kraft, *Nucl. Phys.* B410(1993)219; K. G. Chetyrkin, J. H. Kühn, M. Steinhauser, *Nucl. Phys.* B482(1996)213; R. Harlander, M. Steinhauser, *Eur. Phys. J.* C2(1998)151.
- [6] Chong-Xing Yue, Wei Wang, Feng Zhang, hep-ph/0409066.
- [7] T. Han, H. E. Logan, B. McElrath and L. T. Wang, *Phys. Rev.* D67(2003)095004.
- [8] M. Perelstein, M. E. Peskin and A. Pierce, *Phys. Rev.* D69(2004)075002.
- [9] G. Burdman, M. Perelstein and A. Pierce, *Phys. Rev. Lett.* **90** (2003) 241802; T. Han.
  H. E. Logen, B. McElrath and L. T. Wang, *Phys. lett.* B563(2003)191; Chong-Xing
  Yue, Shunzhi Wang, Dongqi Yu, *Phys. Rev.* D68(2003)115004; S. Chang, H.-J. He,

Phys. lett. B 586(2004)95; S. C. Park and J. Song, Phys. Rev. D69(2004)115010;
G. A. Gonzalez-Sprinberg, R. Martinez, and J. Alexis Rodriguez, hep-ph/0406178;
Gi-Chol Cho and Aya Omote, Phys. Rev. D70(2004)057701; Jaeyong Lee, JHEP
0412(2004)065; Chong-Xing Yue, Wei Wang, Phys. Rev. D71(2005)015002.

- [10] J. L. Hewett, F. J. Petriell and T. G. Rizzo, JHEP 0310(2003)062; C. Csaki et al., *Phys. Rev.* D67(2003)115002.
- [11] C. Csaki et al., *Phys. Rev.* D68(2003)035009; T. Gregoire, D. R. Smith and J. G. Wacker, *Phys. Rev.* D69(2004)115008; Mu-chun chen and S. Dawson, *Phys. Rev.* D70 (2004)015003; Chong-Xing Yue and Wei Wang, *Nucl. Phys.* B683(2004)48.
- [12] G. Moortgat-Pick and H. M. Steiner, Eur. Phys. J. C6(2001)1; J. Erler, et al., hep-ph/0112070; G. Moortgat-Pick, et al., hep-ex/0303234; G. Moortgat-Pick, hep-ph/0410118
- [13] M. Martinez, R. Miquel, L. Rolandi, and R. Tenchini, *Rev. Mod. Phys.* **71**(1999)575;
   S. F. Novaes, *hep-ph*/**0001283**.