Growth of sumsets in abelian semigroups^{*}

Melvyn B. Nathanson Department of Mathematics Lehman College (CUNY) Bronx, New York 10468 Email: nathansn@alpha.lehman.cuny.edu

Let S be an abelian semigroup, written additively, that contains the identity element 0. Let A be a nonempty subset of S. The cardinality of A is denoted |A|. For any positive integer h, the sumset hA is the set of all sums of h not necessarily distinct elements of A. We define $hA = \{0\}$ if h = 0. Let A_1, \ldots, A_r , and B be nonempty subsets of S, and let h_1, \ldots, h_r be nonnegative integers. We denote by

$$B + h_1 A_1 + \dots + h_r A_r \tag{1}$$

the set of all elements of S that can be represented in the form $b + u_1 + \cdots + u_r$, where $b \in B$ and $u_i \in h_i A_i$ for all $i = 1, \ldots, r$. If the sets A_1, \ldots, A_r , and B are finite, then the sumset (1) is finite for all h_1, \ldots, h_r . The growth function of this sumset is

$$\gamma(h_1,\ldots,h_r) = |B + h_1 A_1 + \cdots + h_r A_r|.$$

For example, let S be the additive semigroup of nonnegative integers \mathbf{N}_0 , and let A_1, \ldots, A_r , and B be nonempty, finite subsets of \mathbf{N}_0 , normalized so that $0 \in B \cap A_1 \cap \cdots \cap A_r$ and $gcd(A_1 \cup \cdots \cup A_r) = 1$. Let $b^* = max(B)$ and $a_i^* = max A_i$ for $i = 1, \ldots, r$. Han, Kirfel, and Nathanson [1, 5] determined the asymptotic structure of the sumset $B + h_1A_1 + \cdots + h_rA_r$. They proved that there exist integers c and d and finite sets $C \subseteq [0, c-2]$ and $D \subseteq [0, d-2]$ such that

$$B + h_1 A_1 + \dots + h_r A_r = C \cup [c, b^* + \sum_{i=1}^r a_i^* h_i - d] \cup \left(b^* + \sum_{i=1}^r a_i^* h_i - D \right).$$

for $\min(h_1, \ldots, h_r)$ sufficiently large. This implies that the growth function is eventually a multilinear function of h_1, \ldots, h_r , that is, there exists an integer Δ such that

$$|B + h_1 A_1 + \dots + h_r A_r| = a_1^* h_1 + \dots + a_r^* h_r + b^* + 1 - \Delta$$

 $^{^{*}}$ Supported in part by grants from the PSC–CUNY Research Award Program and the NSA Mathematical Sciences Program.

for $\min(h_1, \ldots, h_r)$ sufficiently large. The explicit determination of the sets C and D is a difficult unsolved problem in additive number theory. In the case r = 1, it is called the *linear diophantine problem of Frobenius*. For a survey of finite sumsets in additive number theory, see Nathanson [6].

The theorem about sums of finite sets of integers generalizes to sums in an arbitrary abelian semigroup S. We shall prove that if A_1, \ldots, A_r , and B are finite, nonempty subsets of S, then the growth function $\gamma(h_1, \ldots, h_r)$ is eventually polynomial, that is, there exists a polynomial $p(z_1, \ldots, z_r)$ such that

$$\gamma(h_1, \dots, h_r) = |B + h_1 A_1 + \dots + h_r A_r| = p(h_1, \dots, h_r)$$

for min (h_1, \ldots, h_r) sufficiently large. The case r = 1 is due to Khovanskii [3, 4]. We use his method to extend the result to the case $r \ge 2$. The idea of the proof is to show that the growth function is the Hilbert function of a suitably constructed module graded by the additive semigroup \mathbf{N}_0^r of r-tuples of nonnegative integers.

We need the following result about Hilbert functions. Let R be a finitely generated \mathbf{N}_0^r -graded connected commutative algebra over a field E. Then $R = \bigoplus_{h \in \mathbf{N}_0^r} R_h$. Suppose that R is generated by s homogeneous elements y_1, \ldots, y_s with $y_i \in R_{\delta_i}$, that is, the degree of y_i is deg $y_i = \delta_i \in \mathbf{N}_0^r$. Let Mbe a finitely generated \mathbf{N}_0^r -graded R-module. For $h = (h_1, \ldots, h_r) \in \mathbf{N}_0^r$, we define the Hilbert function

$$H(M,h) = \dim_E \left(M_{(h_1,\dots,h_r)} \right).$$

For $z = (z_1, \ldots, z_r)$, we define

$$z^h = z_1^{h_1} \cdots z_r^{h_r}.$$

Consider the formal power series

$$F(M,z) = \sum_{h \in \mathbf{N}_0^r} H(M,h) z^h.$$

Then there exists a vector β with integer coordinates and a polynomial $P(M, z) = P(M, z_1, \dots, z_h)$ with integer coefficients such that

$$F(M,z) = \frac{z^{\beta}P(M,z)}{\prod_{i=1}^{s}(1-z^{\delta_i})}.$$

(This is Theorem 2.3 in Stanley [7, p. 33]).

Theorem 1 Let A_1, \ldots, A_r , and B be finite, nonempty subsets of an abelian semigroup S. There exists a polynomial $p(z_1, \ldots, z_r)$ such that

$$|B + h_1 A_1 + \dots + h_r A_r| = p(h_1, \dots, h_r)$$

for all sufficiently large integers h_1, \ldots, h_r .

Proof. For $i = 1, \ldots, r$, let

$$A_i = \{a_{i,1}, \ldots, a_{i,k_i}\},\$$

where

$$|A_i| = k_i \ge 1.$$

We introduce a variable $x_{i,j}$ for each i = 1, ..., r and $j = 1, ..., k_i$. Fix a field E. We begin with the polynomial ring

$$R = E[x_{1,1}, \dots, x_{r,k_r}]$$

in the $s = k_1 + \cdots + k_r$ variables $x_{i,j}$. The algebra R is connected since it is an integral domain (cf. Hartshorne [2, Exercise 2.19, p. 82]). For each r-tuple $(h_1, \ldots, h_r) \in \mathbf{N}_0^r$ we let

$$R_{(h_1,\ldots,h_r)}$$

be the vector subspace of R consisting of all polynomials that are homogeneous of degree h_i in the variables $x_{i,1}, \ldots, x_{i,k_i}$. In particular, $E = R_{(0,\ldots,0)}$. Then

$$R = \bigoplus_{(h_1,\dots,h_r) \in \mathbf{N}_0^r} R_{(h_1,\dots,h_r)}.$$

The multiplication in the algebra R is consistent with this direct sum decomposition in the sense that

$$R_{(h_1,\dots,h_r)}R_{(h'_1,\dots,h'_r)} \subseteq R_{(h_1+h'_1,\dots,h_r+h'_r)},$$

and so R is graded by the semigroup \mathbf{N}_0^r .

Next we construct an \mathbf{N}_0^r -graded R-module M. To each r-tuple $(h_1, \ldots, h_r) \in \mathbf{N}_0^r$ we associate a finite-dimensional vector space $M_{(h_1,\ldots,h_r)}$ over the field E in the following way. To each element

$$u \in B + h_1 A_1 + \dots + h_r A_r$$

we assign the symbol

$$[u, h_1, \ldots, h_r].$$

Let $M_{(h_1,\ldots,h_r)}$ be the vector space consisting of all *E*-linear combinations of these symbols. Then

$$\dim_E M_{(h_1,\dots,h_r)} = |B + h_1 A_1 + \dots + h_r A_r|.$$
 (2)

Let

$$M = \bigoplus_{(h_1,\dots,h_r)\in\mathbf{N}_0^r} M_{(h_1,\dots,h_r)}.$$

This is an \mathbf{N}_0^r -graded vector space over E.

To make M a module over the algebra R, we must construct a bilinear multiplication $R \times M \to M$. We define the product of the variable $x_{i,j} \in R$ and the basis element $[u, h_1, \ldots, h_r] \in M$ as follows:

$$x_{i,j}[u, h_1, \dots, h_r] = [u + a_{i,j}, h_1, \dots, h_{i-1}, h_i + 1, h_{i+1}, \dots, h_r].$$

This makes sense since

$$u \in B + h_1 A_1 + \dots + h_i A_i + \dots + h_r A_r$$

and so

$$u + a_{i,j} \in B + h_1 A_1 + \dots + (h_i + 1) A_i + \dots + h_r A_r$$

This induces a well-defined multiplication of elements of M by polynomials in R since, if i < i',

$$\begin{aligned} x_{i',j'} \left(x_{i,j}[u,h_1,\ldots,h_r] \right) \\ &= x_{i',j'}[u+a_{i,j},h_1,\ldots,h_i+1,\ldots,h_r] \\ &= [u+a_{i,j}+a_{i',j'},h_1,\ldots,h_i+1,\ldots,h_{i'}+1,\ldots,h_r] \\ &= [u+a_{i',j'}+a_{i,j},h_1,\ldots,h_i+1,\ldots,h_{i'}+1,\ldots,h_r] \\ &= x_{i,j}[u+a_{i',j'},h_1,\ldots,h_{i'}+1,\ldots,h_r] \\ &= x_{i,j} \left(x_{i',j'}[u,h_1,\ldots,h_r] \right). \end{aligned}$$

The case $i \ge i'$ is similar. Note that this is the only place where we use the commutativity of the semigroup S. It follows that M is an R-module. Moreover,

$$R_{(h_1,\dots,h_r)}M_{(h'_1,\dots,h'_r)} \subseteq M_{(h_1+h'_1,\dots,h_r+h'_r)},$$

and so M is a graded R-module. Furthermore, the finite set

$$\{[b,0,\ldots,0]:b\in B\}\subseteq M$$

generates M as an R-module.

Since $x_{i,j} \in R_{\delta_{i,j}}$, where $\deg(x_{i,j}) = \delta_{i,j}$ is the *r*-tuple whose *i*-th coordinate is 1 and whose other coordinates are 0, and since

$$\frac{1}{(1-z_i)^{k_i}} = \sum_{h_i=0}^{\infty} \binom{h_i + k_i - 1}{k_i - 1} z_i^{h_i},$$

we have

$$F(M,z) = \sum_{h \in \mathbf{N}_0^r} H(M,h) z^h$$
$$= \frac{z^\beta P(M,z)}{\prod_{i=1}^r \prod_{j=1}^{k_i} (1-z^{\delta_{i,j}})}$$
$$= \frac{z^\beta P(M,z)}{\prod_{i=1}^r (1-z_i)^{k_i}}$$

$$= z^{\beta} P(M, z) \prod_{i=1}^{r} \sum_{h_i=0}^{\infty} {\binom{h_i + k_i - 1}{k_i - 1}} z_i^{h_1}$$
$$= z^{\beta} P(M, z) \sum_{h=(h_1, \dots, h_r) \in \mathbf{N}_0^r} \prod_{i=1}^{r} {\binom{h_i + k_i - 1}{k_i - 1}} z^h$$

This implies that the Hilbert function H(M, h) is a polynomial in h_1, \ldots, h_r for $\min(h_1, \ldots, h_r)$ sufficiently large. By (2), the growth function is the Hilbert function of M. This completes the proof.

References

- S. Han, C. Kirfel, and M. B. Nathanson. Linear forms in finite sets of integers. *Ramanujan J.*, 2:271–281, 1998.
- [2] R. Hartshorne. Algebraic Geometry, volume 52 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1977.
- [3] A. G. Khovanskii. Newton polyhedron, Hilbert polynomial, and sums of finite sets. *Funktsional. Anal. i Prilozhen.*, 26:276–281, 1992.
- [4] A. G. Khovanskii. Sums of finite sets, orbits of commutative semigroups, and Hilbert functions. *Funktsional. Anal. i Prilozhen.*, 29:102–112, 1995.
- [5] M. B. Nathanson. Sums of finite sets of integers. Amer. Math. Monthly, 79:1010–1012, 1972.
- [6] M. B. Nathanson. Additive Number Theory: Inverse Problems and the Geometry of Sumsets, volume 165 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1996.
- [7] R. P. Stanley. Combinatorics and Commutative Algebra. Birkhäuser, Boston, 1983.