

On the Algebraic Bethe Ansatz for XXX spin chain: creation operators "beyond the equator".

A.E. Kovalsky

*MIPT, Dolgoprudny, Moscow reg.,
IHEP, Protvino, Moscow reg., Russia,*

G.P. Pronko

*IHEP, Protvino, Moscow reg., Russia,
International Solvay Institute, Brussels, Belgium*

Abstract

Considering the XXX spin-1/2 chain in the framework of the Algebraic Bethe Ansatz (ABA) we make the following short comment: the product of the creation operators corresponding to the recently found solution of the Bethe equations "on the wrong side of the equator" [1] is just zero (not only its action on the pseudovacuum).

Consider the periodic XXX spin-1/2 chain with N sites in the framework of ABA (see for example [2]). Let us introduce the Lax operator acting in the two-dimensional local quantum space $h_n = \mathbb{C}^2$ and in the two-dimensional auxiliary space $V = \mathbb{C}^2$:

$$L_n(x) = \begin{pmatrix} x + is_n^3 & is_n^- \\ is_n^+ & x - is_n^3 \end{pmatrix}, \quad (1)$$

where s^i are operators of spin 1/2, x is an arbitrary complex number (the spectral parameter). The monodromy matrix is the ordered product over all sites:

$$T(x) = L_N(x)L_{N-1}(x)\dots L_1(x) = \begin{pmatrix} A(x) & B(x) \\ C(x) & D(x) \end{pmatrix}, \quad (2)$$

where $A(x), B(x), C(x), D(x)$ are operators acting in the full quantum space $H = \otimes \prod_{n=1}^N h_n$. In the framework of ABA one looks for the eigenvectors of the transfer matrix

$$\hat{t}(x) = \text{tr}T(x) = A(x) + D(x)$$

in the form

$$\Phi(\{x\}) = B(x_1)B(x_2)\dots B(x_l)\Omega, \quad (3)$$

where $\Omega = \prod_{n=1}^N \omega_n$, $s_n^+ \omega_n = 0$. It follows from the intertwining relations for the monodromy matrices that vector (3) will be an eigenvector of the transfer matrix when the parameters x_1, \dots, x_l satisfy the Bethe equations:

$$\left(\frac{x_j + i/2}{x_j - i/2} \right)^N = \prod_{k \neq j}^l \frac{x_j - x_k + i}{x_j - x_k - i} \quad (j = 1, 2, \dots, l). \quad (4)$$

Let us denote vectors of the form (3) with parameters x_j ($j = 1, 2, \dots, l$) satisfying the system (4) by $\Phi(\{x\}_B)$. It is well known that vectors $\Phi(\{x\}_B)$ are the highest weights vectors with respect to $SU(2)$ generated by J^i i.e.

$$J^+ \Phi(\{x\}_B) = 0 \quad (5)$$

and

$$J^3 \Phi(\{x\}_B) = (N/2 - l) \Phi(\{x\}_B), \quad (6)$$

where J^3 , J^\pm are operators of the total spin. It is clear that if $l > N/2$ then $\Phi(\{x\}_B) = 0$. The solution $\{x\}$ of (4) with $l \leq N/2$ defines the polynomial $q(x)$ of the degree l , whose roots are $\{x\}$:

$$q(x) = \prod_{j=1}^l (x - x_j). \quad (7)$$

Let $t(x)$ be the eigenvalue of transfer matrix $\hat{t}(x)$ corresponding to the eigenvector $\Phi(\{x\}_B)$ i.e. $\hat{t}(x)\Phi(\{x\}_B) = t(x)\Phi(\{x\}_B)$. It is a polynomial of degree N . Then the polynomials $t(x)$ and $q(x)$ satisfy the Baxter equation [4] (we consider the case of the simple roots):

$$t(x)q(x) = (x - i/2)^N q(x + i) + (x + i/2)^N q(x - i). \quad (8)$$

In the paper [1] was shown that there exist the polynomial $p(x)$ of degree $N - l + 1$ with roots satisfying the Bethe equation (4)¹ and

$$t(x)p(x) = (x - i/2)^N p(x + i) + (x + i/2)^N p(x - i), \quad (9)$$

with the same $t(x)$ as in (8). Actually there exists the one-parametric family of such polynomials

$$p(x, \alpha) = p(x) + \alpha q(x), \quad (10)$$

so there is the one-parametric family of sets of parameters ($\{x\}$)—the zeroes of $p(x, \alpha)$ which belongs to the "beyond the equator" case. Let us denote these zeroes as $x_i(\alpha)$ (it is clear that the zeroes of the polynomial (10) depends on α). Now consider the creation operator

$$\mathbf{B}(\alpha) = B(x_1(\alpha))B(x_2(\alpha))\dots B(x_{N-l+1}(\alpha))$$

corresponding to the beyond the equator case. The following statement is valid:

Theorem.

$$\mathbf{B}(\alpha) = 0 \quad (11)$$

Proof.

Consider its action on the basis constructed using the Bethe vectors (in the case of finite $\{x\}$ Bethe vectors are the highest weights; to obtain the rest eigenvectors we use the operator J^- (it commutes with the transfer matrix), which also can be considered as a creation operator, since $B(x) = x^{N-1}(J^- + o(1/x))$ when $x \rightarrow \infty$; on the hypothesis of the completeness of the Bethe ansatz see [2, 3]). We immediately see that its action is zero due to $[B(x), B(y)] = 0$, $[B(x), J^-] = 0$ and $\mathbf{B}(\alpha)\Omega = 0$. So, the action of $\mathbf{B}(\alpha)$ is zero onto each vector of the basis, then

$$\mathbf{B}(\alpha) = 0. \quad (12)$$

¹See also the interesting discussion of the "beyond equator" solution in [5].

To see that this fact is nontrivial let us consider the concrete examples. Let us first analyse the structures of the product of B -operators. We have the following commutation relation:

$$[B(x), J^3] = B(x), \quad (13)$$

so for the product of l B -operators

$$e^{\beta J^3} B(x_1) \dots B(x_l) e^{-\beta J^3} = e^{-l\beta} B(x_1) \dots B(x_l) \quad (14)$$

and we see that each term of expansion of this product necessarily contains the products of l -operators s_j^- with different j , since $(s^-)^2 = 0$ (so if $l > N$ this product is zero). At $N = 6$, product of four B -operators ($l = 4 > 3$ i.e. this is the "beyond the equator" case): $B(x_1)B(x_2)B(x_3)B(x_4)$, as was shown by the explicit construction of this product, contains terms, proportional to $s_2^+ s_1^- s_3^- s_4^- s_5^- s_6^-$, $s_3^+ s_1^- s_2^- s_4^- s_5^- s_6^-$, $s_4^+ s_1^- s_2^- s_3^- s_5^- s_6^-$ and $s_5^+ s_1^- s_2^- s_3^- s_4^- s_6^-$ with nonzero coefficients—polynomials in x_1, x_2, x_3, x_4 . The appearance of such terms is not excluded by (14). Its action on the vacuum Ω is zero identically, so if x_1, x_2, x_3, x_4 satisfy the system (4), then these terms is zero, not only terms, whose action on the vacuum Ω is nonzero, for example the term proportional to $s_1^- s_2^- s_3^- s_4^-$. Such solutions do exist, for example roots of the polynomial

$$p(x) = x^4 - \frac{6}{\sqrt{13}} x^3 + x^2 - \frac{9}{16} \quad (15)$$

satisfy system (4). This solution corresponds to the total spin $J = 0$, the eigenvalue of the transfer matrix

$$t(x) = 2x^6 + \frac{9}{2} x^4 + \frac{23}{8} x^2 - \frac{3}{\sqrt{13}} x - \frac{1}{32}$$

the corresponding eigenvector can be constructed, using the roots of the polynomial

$$q(x) = x^3 + \frac{1}{12} x + \frac{1}{4\sqrt{13}}. \quad (16)$$

and the one-parametric family $\mathbf{B}(\alpha) = 0$ corresponds to the roots of the polynomial

$$p(x, \alpha) = x^4 - \frac{6}{\sqrt{13}} x^3 + x^2 - \frac{9}{16} + \alpha \left(x^3 + \frac{1}{12} x + \frac{1}{4\sqrt{13}} \right). \quad (17)$$

The products of l operators $B(x)$ with $l > N/2$ considered above are not used for the construction of the eigenvectors of the transfer matrix. However, we would like to emphasize that there is another important example, when the product of B -operators corresponding to the case $l \leq N/2$ is zero:

$$B(-i/2)B(i/2) = 0.$$

This product corresponds to the polynomial

$$q(x) = x^2 + 1/4$$

and

$$t(x) = (x + i/2)(x - 3/2i)^{N-1} + (x - i/2)(x + 3/2i)^{N-1}$$

and at even $N \geq 4$ corresponds to some eigenvector of the transfer matrix. In the paper [6] was shown how one can use the ABA to construct this eigenvector corresponding to this exceptional solution. If we consider the following vector $B(-i/2 + \epsilon + 2i\epsilon^N)B(i/2 + \epsilon)\Omega$ at $\epsilon \rightarrow 0$, then

$$B(-i/2 + \epsilon + 2i\epsilon^N)B(i/2 + \epsilon)\Omega = \epsilon^N \Phi(\{-i/2, i/2\}) + O(\epsilon^{N+1})$$

where $\Phi(\{-i/2, i/2\})$ is the desired eigenvector. The proof of our statement holds true in this case too, since we use again only B -operators to construct this eigenvector.

This work was supported in part by grants of RFBR 00-15-96645, 01-02-16585, 01-01-00201, CRDF MO-011-0, of the Russian Ministry on the education E00-3.3-62 and INTAS 00-00561.

References

- [1] Pronko G.P., Stroganov Yu.G.,
Bethe Equations "on the Wrong Side of Equator",
J.Phys.A: Math.Gen. **32** 2333-40, 1999.
e-Print Archive: hep-th/9808153.
- [2] Faddeev L. D.,
How Algebraic Bethe Ansatz works for integrable model.
1995 *UMANA* **40** 214,
e-Print Archive: hep-th/9605187.
- [3] A.N. Kirillov,
Combinatorial identities and completeness of states for
the generalized Heisenberg magnet,
Zap. Nauch. Sem. LOMI, 1984, v.131, p.88-105.
- [4] Baxter R.J. Stud.Appl.Math, L51-69, 1971; Ann.Phys. N.Y., v.70, 193-228, 1972;
Ann.Phys. N.Y., v.76, 1-71, 1973.
- [5] R.J. Baxter,
Completeness of the Bethe ansatz for the six and eight-vertex models
e-Print Archive: cond-mat/0111188
- [6] L. V. Avdeev, A. A. Vladimirov,
On the exceptional solutions of the Bethe-Ansatz equations.
Theor. Math. Phys. **69**: 1071, 1987.